

UNCLASSIFIED

AD NUMBER
AD868355
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; APR 1970. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Release Branch/TID, Frederick, MD 21701.
AUTHORITY
BDRL ltr, 29 Sep 1971


THIS PAGE IS UNCLASSIFIED

STATEMENT OF
AND RECORD OF
IN COAST GUARD

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of Dept. of Army, Fort Detrick, ATTN: Technical Release Branch/TID, Frederick, Maryland 21701

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151



DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 595

SEPARATION OF DONOR AND RECIPIENT BACTERIA
BY COLUMN CHROMATOGRAPHY

Robert M. Zsigray

George E. Fulk

William D. Lawton

Medical Bacteriology Division
BIOLOGICAL SCIENCES LABORATORIES

Project 1B061102B71A

April 1970

ABSTRACT

When donor and recipient strains of Escherichia coli were added to columns containing Cellux-P (a cation-exchange cellulose), more than 80% of the female cells passed through the column but only 11% or less of the male cells were eluted. However, when donor strains were blended prior to their addition to the column, the majority of these cells were eluted. These results indicated that the filamentous appendages termed F pili (which are removed by blending) were responsible for the adherence of donor cells to the cellulose.

I. INTRODUCTION*

Phenotypic variations are known to exist between the two mating types of bacteria. In 1956, Mucedano and Conolly¹ reported that donor and recipient strains of *Escherichia coli* K-12 differed in their acid agglutination point and in their affinity for basic dyes. In all instances, the recipient (F^-) organisms were found to be more electromotively charged than the donor (F^+) strains. Subsequently, the existence of the F^+ antigen was demonstrated in Hfr and F^+ strains;² this antigen was found to correspond to the small filamentous appendage termed F pili.³⁻⁵ In addition, Smith and Lederberg⁶ have demonstrated the existence of a periodate-sensitive site(s) on the surface of Hfr and F^+ cells.

Because these phenotypic differences are expressed at or near the cell periphery and appear to alter the surface charge, we devised a one-step column chromatographic method for separating the two mating types.

II. MATERIALS AND METHODS

A. BACTERIAL STRAINS

The organisms used in this study and their sources are listed in Table 1. Strain W1485 F^- is an acridine orange - cured derivative of W1485 (Falkow). We obtained F^- strains from X^{646} and HB 11 after spontaneous segregation of the F -lac episome.

TABLE 1. *ESCHERICHIA COLI* STRAINS

Strain	Sex Type	Source
<i>E. coli</i> K12 W1485	F^+	Falkow
<i>E. coli</i> W1485	F^-	Falkow
<i>E. coli</i> W1485	Hfr	Falkow
<i>E. coli</i> Hfr Hayes	Hfr	Jacob and Weisman ⁷
<i>E. coli</i> X^{646}	F^+ (pro A^+ pro B^+ lac ⁺)	Curtiss III
<i>E. coli</i> X^{646}	F^-	Authors
<i>E. coli</i> B HB 11	F' lac	Boyer
<i>E. coli</i> B HB 11	F^-	Authors

* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

B. MEDIA

Liquid medium was Difco brain heart infusion broth (BHI). Solid media consisted of BHI with the addition of 1.5% agar (Difco) or lactose medium,* to facilitate the assay of F'_{lac}^+ and F'_{lac}^- clones.

C. COLUMN PREPARATION

One gram of Cellux-P (Bio-Rad*), a cation-exchange cellulose, was permitted to swell in 100 ml of 0.1 M ethylenediaminetetraacetic acid (EDTA), pH 6.8. A small piece of glass wool was inserted in the base of a glass column (15 by 300 mm) for support, and 20 ml of the swollen cellulose were transferred to the column and allowed to settle by gravity. The flow rate after packing was approximately 3 ml/min.

D. GROWTH CONDITIONS AND COLUMN ELUTION

Each of the strains listed in Table 1 was grown aerobically for 18 hours in 5 ml of BHI at 37 C. Each culture was diluted with an equal volume of 0.1 M EDTA, pH 6.8, and 1.0 ml of this mixture was added to the column, allowed to drain into the resin, and then followed with 10 ml of 0.1 M EDTA, pH 6.8. The eluate was collected and diluted in 0.15 M NaCl and the appropriate dilutions were plated and incubated overnight at 37 C. Viable counts also were made from a portion of each culture that did not go through the column.

E. RECONSTRUCTION EXPERIMENTS

Equal portions (0.5 ml) of strains x^{646} and $x^{646} p^-$ were mixed in 1.0 ml of 0.1 M EDTA, pH 6.8. One milliliter of this mixture was immediately added to a column and eluted. The eluate was collected and appropriate dilutions were plated on lactose medium.

* Bio-Rad Laboratories, 32nd and Griffin Ave., Richmond, Calif.

III. RESULTS

When individual overnight cultures of female or male bacteria were diluted in 0.1 M EDTA and added to the cation-exchange cellulose, more than 80% of the female cells passed through the column but only 11% or less of the male cells were eluted (Table 2). The only exception was strain W1485 carrying Hfr . This result was not unusual because most organisms that harbor R factors are of the repressed type.¹⁰

TABLE 2. ELUTION OF DONOR AND RECIPIENT BACTERIA FROM CELLEX-P COLUMNS

Strain	Mating Type	Total Colony-Forming Units Recovered		Recovery, %
		Before Column	After Column	
<i>E. coli</i> Hfr Hayes	Male	5.3×10^8	1.5×10^7	2.8
<i>E. coli</i> W1485	Male	1.8×10^9	3.0×10^7	1.7
<i>E. coli</i> W1485 F^-	Female	1.0×10^9	9.2×10^8	92.0
<i>E. coli</i> x^{646}	Male	5.6×10^8	5.5×10^7	9.8
<i>E. coli</i> $\text{x}^{646} \text{F}^-$	Female	5.6×10^8	4.9×10^8	84.5
<i>E. coli</i> B HB 11	Male	1.7×10^9	1.9×10^8	11.2
<i>E. coli</i> B HB 11 F^-	Female	2.0×10^9	1.8×10^9	90.0
<i>E. coli</i> W1485 RTF	Male	2.6×10^8	2.1×10^5	80.8

A. SEPARATION OF MATING TYPES

When mixtures of strain x^{646} and $\text{x}^{646} \text{F}^-$ were added to Cellex-P and eluted, a significant separation of the two mating types was obtained. The results of a typical experiment (Table 3) showed that 90% of the donor cells within the mixture were retained by the column and approximately 90% of the total F^- cells were eluted.

TABLE 3. SEPARATION OF DONOR FROM RECIPIENT CELLS

Strain	Total Colony-Forming Units Recovered from Mixture		Recovery, %
	Before Column	After Column	
$\text{x}^{646} (\text{lac}^+)$	1.4×10^8	1.4×10^8	10.0
$\text{x}^{646} \text{F}^- (\text{lac}^-)$	1.7×10^9	1.5×10^9	88.2

B. SPECIFICITY OF DONOR BINDING

Because periodate treatment is known to interfere with Hfr and F⁺ virility by interacting with carbohydrate at the cell surface,⁷ periodate might alter the overall charge of the donor cell and change its affinity for Cellax-P. To test this possibility, an overnight culture of Hfr Hayes was treated with periodate according to the procedure described by Smatch and Lederberg⁷ before the culture was added to the column. Such treatment did not significantly alter the elution pattern of the donor strain.

C. ROLE OF F PILI

An obvious possibility responsible for the selective retention of donor cells was the F pili produced by donor cells but not produced by recipient cells. Because F pili regeneration takes place readily at 37 C⁸ but not at 5 C,⁹ an overnight culture of strain W1485 was cooled to and blended at 5 C for 5 minutes in a Sorvall Omni-Mixer, and a sample was added to the resin and eluted. An unblended sample served as a control.

The results of the experiment (Table 4) showed that without blending, 89% of the donor cells were retained by the column, but after blending, only 28% of the donor cells were retained, indicating that the F pili were responsible for the male population adhering to the column.

TABLE 4. EFFECT OF BLENDING ON ELUTION OF *E. COLI* W1485 FROM CELLEX-P COLUMNS

Treatment		Total Number of Colony-Forming Units Recovered	Recovery, %
Blended	Column		
-	-	3.6×10^8	
-	+	4.0×10^7	11.1
+	+	2.6×10^8	72.2

To determine the viability of donor cells retained by the resin, the Cellax-P-donor mixture was removed from the column with 8 ml of broth and cooled to and blended at 5 C; appropriate dilutions were plated. Recoveries ranging from 70 to 90% were obtained, indicating that Cellax-P had little or no effect on donor viability.

IV. DISCUSSION

The main result of this investigation is discovery of the enrichment of one mating type (F^-) over the other (Hfr , F^+ , or F'). When donor and recipient cells were added (individually or in mixtures) to columns containing Cellux-F, 90% of the donor cells were retained by the cation-exchange cellulose. On the other hand, approximately 90% of the recipient cells were eluted.

The elution pattern of donor strains can be altered by high-speed blending, indicating that the male-specific appendages termed F pili are probably responsible for the retention of the males by the column. The F pili may be physically entangled within rather than ionically adsorbed to the cellulose, because increases in salt concentration or changes in pH did not alter the elution pattern of donor bacteria.

Strain W1485 Hfr did not exhibit the same retention characteristics as the other donors (Table 2), a result that can be explained on the basis of F piliation. Those organisms that contain a functional episome (one that is derepressed) are capable of producing F pili that, in turn, cause the cells to be retained by the cellulose. On the other hand, organisms that harbor a repressed episome fail to produce these appendages¹⁰ and are eluted. Because of a lack of agglutination of W1485 Hfr cells with antisera to F pili, this strain appears to contain a repressed F factor.

Because Cellux-F appears selectively to separate male cells from female cells (or from repressed male cells), this procedure may prove to be advantageous for (i) isolating rare derepressed cells from a repressed culture, (ii) determining the expression period for F pili formation, and (iii) enriching for recipient cells immediately after they have acquired genetic material from donor strains.

LITERATURE CITED

1. MacLeod, G.A.; Cosolli, E. 1956. Surface properties correlated with sex compatibility in Escherichia coli. J. Gen. Microbiol. 15:121-132.
2. Orskov, I.; Orskov, F. 1960. An antigen termed f^+ occurring in f^+ E. coli strains. Acta Pathol. Microbiol. Scand. 48:37-44.
3. Einton, C.C. 1965. The structure, function, synthesis and genetic control of bacterial pili and a molecular model for DNA and RNA transport in gram-negative bacteria. Trans. N.Y. Acad. Sci. 27: 1003-1054.
4. Ishihashi, H. 1967. F pilus as f^+ antigen. J. Bacteriol. 93:379-389.
5. Koelle, P.; Orskov, I. 1967. The identity of the f^+ antigen and the cellular receptor for the RFA phage fr. Mol. Gen. Genet. 99:109-114.
6. Lenn, A.M.; Maynard, E.; Maynard, G.C.; Batta, E. 1967. Sex pili and the classification of sex factors in Enterobacteriaceae. Nature 216:343-344.
7. Smith, P.H.A.; Lederberg, J. 1961. Inhibition by periodate of mating in Escherichia coli K-12. Proc. Nat. Acad. Sci. U.S. 47:86-90.
8. Lawton, W.D.; Morris, B.C.; Burrows, T.W. 1968. Gene transfer in strains of Pasteurella pseudotuberculosis. J. Gen. Microbiol. 52:25-34.
9. Jacob, F.; Wollman, E. 1961. Sexuality and the genetics of bacteria. Academic Press, New York.
10. Maynard, E.; Maynard, G.C.; Batta, E. 1968. Phylogenetic relationships of drug-resistance factors and other transmissible bacterial plasmids. Bacteriol. Rev. 32:55-83.