

AWARD NUMBER: W81XWH-13-0115

TITLE: The Parkinson Transcriptome Project: Roadmap to Personalized Care

PRINCIPAL INVESTIGATOR: Clemens R. Scherzer, MD

RECIPIENT: The Brigham and Women's Hospital, Inc
Boston, MA 02115-6110

REPORT DATE: August 2015

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution
Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE August 2015		2. REPORT TYPE Final		3. DATES COVERED 15 May 2013 - 14 May 2015	
4. TITLE AND SUBTITLE The Parkinson Transcriptome Project: Roadmap to Personalized Care				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-13-1-0115	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Dr. Clemens Scherzer E-Mail: cscherzer@rics.bwh.harvard.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Brigham and Women's Hospital, Inc. 75 Francis St. Boston, MA 02115-6110				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT This study provides a comprehensive catalog of gene expression - both protein-coding and non-coding - specifically in dopamine neurons <i>in situ</i> in human brain. Going beyond traditional messenger RNA sequencing, <i>all</i> polyadenylated <i>and</i> non-polyadenylated transcripts over 50 bases were deeply sequenced using ribo-depleted RNA from dopamine neurons laser-captured from human post-mortem brains of patients with Parkinson's disease using laser-capture RNA sequencing. We show unequivocally that non-coding RNAs are abundantly transcribed in human dopamine neurons and highlight significant changes in the expression of both protein-coding and non-coding transcripts in patients with clinical and neuropathological Parkinson's disease. Expression Quantitative Trait Locus analysis indicates that the expression of many non-coding and protein-coding transcripts in nigral dopamine neurons is associated with genetic variants. Finally, these pilot phase data support the feasibility of measuring select RNAs linked to Parkinson's in vulnerable dopamine neurons as possible biomarkers in blood.					
15. SUBJECT TERMS Parkinson's disease, non-coding RNA, RNA sequencing, expression Quantitative Trait Locus, biomarkers					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
Unclassified	Unclassified	Unclassified	Unclassified	26	19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
1. Introduction	4
2. Keywords	4
3. Overall Project Summary	5
4. Key Research Accomplishments	9
5. Conclusion	9
6. Publications, Abstracts, and Presentations	11
7. Inventions, Patents and Licenses	12
8. Reportable Outcomes	12
9. Other Achievements	13
10. References	13
11. Appendix	13

1. Introduction

The recent discovery of a hidden universe of tens of thousands of non-protein-coding RNAs is revolutionizing our thinking about the flow of genetic information that shapes the complexity of the human brain and underlies its diseases. A revised view suggests parallel processing from DNA to protein-coding mRNAs as well as non-coding RNAs that are themselves active (without translation into proteins) --- with important biologic functions as regulators and biosensors.

We hypothesize that in complex neurodegenerative diseases such as Parkinson's genetic and epigenetic risks (from aging and environmental exposures) exert their disease-causing effects in part through a concerted modulation of transcriptional programs that include non-coding and protein-coding RNA functions. The aims of this pilot study were: 1, Establishing a catalog of tens of thousands previously unexplored, non-coding RNAs and assess their association with Parkinson's precisely in the preferentially vulnerable cell-type --- dopaminergic neurons laser-captured from human substantia nigra --- using massively parallel sequencing and unlimited *ab initio* reconstruction of the whole transcriptome 2, Beginning to delineate the genetic variability that underlies differences in non-coding RNA abundance and processing in human nigral dopamine neurons using transcriptome genetics. 3, Evaluating PD-relevant transcripts as potential biomarkers in a pilot study of blood samples from patients with PD and controls enrolled in the Harvard Biomarker Study.

In this study we were able to provide a first look at the "dark universe" of thousands of short and long non-coding RNAs expressed in the dopamine neuron of normal brains and in Parkinson's disease. Moreover, the study begins to clarify the role of genetic variants in dopamine neurons and to translate transcripts associated with PD-neuropathology. Finally, we provide proof of the concept that select RNAs associated with Parkinson's in vulnerable dopamine neurons can be translated into personalized, dual-use, peripheral biomarkers. Validated biomarkers will be clinically relevant for identifying PD-vulnerable warfighters and civilians, and for monitoring neurotoxin exposure and therapeutic response.

2. Keywords

Parkinson's disease, non-coding RNA, RNA sequencing, expression Quantitative Trait Locus, biomarkers

3. Overall project summary

Aim 1. Delineating non-coding RNAs associated with Parkinson's in nigral dopamine neurons (Fig. 1)




Fig. 1. Non-coding RNAs in dopamine neurons of human brain are associated with Parkinson's disease.

33% of all unique RNAs robustly detected in nigral dopamine neurons are long or short non-protein-coding RNAs (ncRNAs) (**Fig. 1a**). In control brains, 8,493 ncRNAs many of which had never before been described in human brain were identified by *ab initio* transcriptome reconstruction (**Fig. 1b**, stacked colored bar). These include short ncRNAs (e.g. 17 snoRNAs and 11 snRNAs) as well as

thousands of long, regulatory, ncRNAs (e.g. 2,846 pseudogenes, 2,806 lincRNAs, and 2,454 natural antisense RNAs).

121 non-coding RNAs (**Fig. 1c, d**) were statistically significantly differentially expressed in dopamine neurons laser-captured from substantia nigra pars compacta of 19 patients with PD compared to 62 controls without PD with genome-wide significance (adjusted for multiple testing using the false discovery rate method) and adjusting for the pertinent covariates of age, gender, RIN number, postmortem interval, and batch). These PD-associated non-coding RNAs included 77 expressed pseudogenes, 22 lincRNAs, 17 antisense transcripts, as well as sense-intronic, sense-overlapping, and other non-coding RNAs. Moreover, we identified 1,178 protein-coding mRNAs significantly over- or under-expressed in dopamine neurons captured from brains of patients with PD (**Fig. 1c, d**). The scatter plot in Fig. 1d visualizes non-coding and coding RNAs more than three-fold differentially expressed in dopamine neurons of patients with PD compared to controls. Non-coding RNAs are shown in magenta. The lincRNA, *LINC00674* and the expressed pseudogene *UQRBP1* are highlighted in Fig. 1d and in the box plots in Fig. 1e. Moreover, expression changes in two genes, which when mutated cause familial PD, *LRRK2* and *DNAJC6*^{1,2} (**Fig. 1 d, e**) were significant in dopamine neurons of patients with PD compared to controls.

Expression of the dopamine neuron-specific *ALDH1A1* gene is lost in neurons from patients with Parkinson's (Fig. 2).




Fig. 2. Expression of the dopamine neuron-specific *ALDH1A1* gene (a) is lost in dopamine neurons from patients with PD (b).

We compared the RNA content of dopamine neurons and pyramidal neurons laser-captured from control brains. Variance- and sex- adjusted differential expression analysis was performed. 115 genes were exclusively transcribed in dopamine neurons and not in pyramidal neurons with $P < 10^{-6}$. **Fig. 2a** shows the top-ranked 22 dopamine neuron-specific genes (black) and the top 5 pyramidal neuron-specific genes (magenta). Confirming the cell-type precision of our method, these dopamine-neuron specific transcripts included the established parts of the dopamine metabolism machinery, the vesicular monoamine transporter 2 (*VMAT2*), dopamine receptor D2 (*DRD2*), tyrosine hydroxylase (*TH*), and dopamine decarboxylase (*DDC*). Surprisingly, the by far most abundant dopamine neuron-specific transcript however was Aldehyde Dehydrogenase 1A1 (*ALDH1A1*). *ALDH1A1* was significantly underexpressed in dopamine neurons of cases with PD compared to dopamine neurons of controls ($P = 10^{-5}$; **Fig. 2b**). *ALDH1A1* mediates the oxidation of toxic dopamine-3,4-dihydroxyphenylacetaldehyde (DOPAL) into non-toxic 3,4-dihydroxyphenylacetic acid (DOPAC). In mice *ALDH1A1* overexpression suppresses A53T- α -synuclein-induced dopaminergic neurodegeneration and *ALDH1A1* knockout enhances dopaminergic neurodegeneration in model mice³. *ALDH1A1* expression might also be reduced in blood of patients with PD⁴.

Aim 2. Delineating the genetic variants regulating non-coding RNAs in nigral dopamine neurons. In order to elucidate the relationship between genetic variation and gene expression, we have developed a strategy by combining PEER⁵ and Matrix-eQTL⁶ to conduct expression quantitative trait loci (eQTL) analysis. In brief, we normalized the gene expression by adjusting both known and

hidden factors learnt from the Bayesian framework PEER, then used Matrix-eQTL to call significant *cis*-eQTL on the residuals of all genes for both genotyped and imputed SNPs.




Fig. 3. Genetic variants regulate non-coding RNAs specifically in dopamine neurons.

30 expressed pseudogenes, 18 antisense genes, and 15 lincRNAs. 2,602 eSNPs were associated with *RP11-707O23.5* expression, a pseudogene of *MAPK8IP1* (mitogen-activated protein kinase 8 interacting protein 1). 51 eSNPs overlapped with PD GWAS risk SNPs in the *MAPT* susceptibility locus. Importantly, eQTL analysis suggests that these variants (although physically located in the proximity of the *MAPT* locus) do not regulate expression of the *MAPT* gene, but instead of three other genes physically located nearby (i.e. *RP11-707O23.5* with FDRs as low as 1.08^{-4} , *KANSL1-AS1* with FDR of 0.013, and *LRRC37A2* with FDR of 0.0014).

When looking at the regression of gene expression on genotype, we observed distinct patterns relating minor allele dosage to expression. For example, expression levels of the lincRNA *AP000350.5* negatively correlated with minor allele dosage of eSNP rs5760099 (**Fig. 3a**), suggesting a repressive effect of the minor allele. By contrast, minor allele dosage of the genetic variant rs7955007 was associated with increased expression of *RP11-726G1.1* (a processed transcript that overlaps with a piRNA; **Fig. 3b**), suggesting an activating, additive effect. We also observed eSNP-gene expression pairs with an apparently dominant, dosage-independent relation (**Fig. 3c**). For example, expression levels of the antisense transcript *TMLHE-AS1* showed a binary increase associated with the presence (vs. absence) of the minor allele for rs73569697, but not with its dosage.

Aim 3. Evaluating PD neuropathology-linked RNAs as personalized blood biomarkers

Gene	Mean mRNA counts in blood of patients with PD (SD)	Mean mRNA counts in blood of healthy controls (SD)	Difference of means in blood (%)	<i>P</i> -value* (in blood)	<i>P</i> -value# (in dopamine neurons)
<i>SNCA</i>	2,117 (1,732)	3,042 (2,030)	- 30.4%	3.8×10^{-15}	0.02
<i>LRRK2</i>	10,531 (3,451)	9,352 (3,159)	12.6%	4.7×10^{-08}	0.001
<i>CHMP3</i>	2,259 (495)	2,135 (442)	5.8%	4.5×10^{-05}	0.01
<i>COMT</i>	390 (81)	405 (97)	- 3.7%	0.02	0.002

*, *P* value derived from generalized linear models analysis adjusting for age at baseline, gender, RNA Integrity Number, white blood cell counts, red blood cell counts, and platelet counts. #, *P* value comparing transcript counts in dopamine neurons of cases vs. controls adjusting for age, gender, RNA Integrity Number, and batch number.

We evaluated the feasibility of translating transcripts associated with PD neuropathology in dopamine neurons of human brain (from Aim 1) into potentially clinically useful biological markers detectable in peripheral blood.

We performed a cross-sectional study using blood samples from cases representing untreated de novo PD as well as treated early-stage PD and age- and sex-matched healthy controls enrolled in

the Harvard Biomarker Study (HBS)⁷⁻¹² --- a study of >2,500 participants with detailed clinical phenotypes and longitudinal biospecimens collection co-founded and co-directed by Dr. Scherzer. PD patients met stringent UK PD Brain Bank Society's (UKPDBBS) diagnostic criteria and were at an early-stage of the disease as defined by a Modified Hoehn & Yahr stage ≤ 3 assessed by a movement disorders specialist. This single-center study followed stringent rules of evidence for molecular marker




Fig. 4. Four transcripts that were abnormally expressed in dopamine neurons from brains of patients with PD were initially translated into potential blood biomarkers. mRNA counts in peripheral blood of controls (HC), untreated, de novo PD (d-PD), and PD patients on treatment (t-PD) are shown. Error bars, standard error.

discovery and validation and was made possible by our existing collection of high-quality RNA already extracted from blood samples of >2,500 individuals enrolled in HBS.

Excitingly, in this proof-of-concept pilot we were already able to translate four transcripts that are pathologically expressed in laser-captured dopamine neurons of patients with PD into peripheral, easily

accessible possible blood biomarkers. These transcripts are α -synuclein (*SNCA*) and *LRRK2*, *COMT*, and *CHMP3* (Table 1 and Fig. 4). These four transcripts were statistically significantly differentially expressed in dopamine neurons of patients with PD compared to controls based on our laser-capture, total RNAseq method (from Aim 1; Table 1) as well as in peripheral blood samples from the Harvard Biomarker Study (HBS) on the precise Nanostring platform (Fig. 4). The *P* values for associations in blood as well as in dopamine neurons are shown in Table 1. Importantly, two of these four transcripts are also genetically linked to familial and sporadic PD: α -synuclein (*SNCA*) and *LRRK2*.

The *SNCA* transcript was further evaluated in neuroimaging-supported, early stage PD on the qPCR, microarray, and NanoString platform in the PROBE and PPMI study, in addition to HBS. We found that *SNCA* transcripts in circulating blood cells were reduced in early stage, untreated, and dopamine transporter neuroimaging-supported Parkinson's disease (PD) in three independent regional, national, and international populations representing 500 cases and 363 controls and on three analog and digital platforms with *P* = 0.0001 in meta-analysis. Individuals with *SNCA* transcripts in the lowest quartile of counts had an odds ratio for PD of 2.45 compared to individuals in the highest quartile. Disease-relevant transcript isoforms were low even near disease onset. Importantly, low *SNCA* transcript abundance predicted cognitive decline in patients with Parkinson's during up to five years of longitudinal follow-up. *These important findings on SNCA were published in the prestigious journal Brain in August of 2015 (Locascio et al., Brain, 2015; the reprint is attached in the Appendix).*

The other three transcripts - *LRRK2*, *COMT*, and *CHMP3* - significantly associated with early stage PD in blood (Fig. 4) are also directly relevant to the neuropathology with significant expression changes in dopamine neurons (Table 1). *LRRK2* is genetically strongly implicated in both familial and sporadic PD. *COMT* transcripts encode catechol-O-methyl-transferase, an essential gene in the metabolism of L-dopa. Finally, the vacuolar Protein Sorting 24 Homolog (*CHMP3*) was pathologically expressed in our RNAseq analysis of dopamine neurons from patients with PD. It is a novel PD-associated transcript, to our knowledge previously not linked to Parkinson's. Interestingly, the gene *CHMP3* encodes a protein that sorts transmembrane proteins into lysosomes/vacuoles via the multivesicular body pathway. It is part of endosomal sorting complexes required for transport (ESCRT). This pathway is implicated in the pathogenesis of sporadic PD¹³.

This study provides proof of the concept that select RNAs associated with Parkinson's in vulnerable dopamine neurons can be translated into personalized, dual-use, peripheral biomarkers.

4. Key research accomplishments

- This study provides the first comprehensive catalog of gene expression - both protein-coding and non-coding - in a specific cell type, not in cultured cells or mice, but *in situ* in human brain.
- Going beyond traditional mRNA sequencing, *all* polyadenylated *and* non-polyadenylated transcripts over 50 bases were ultra deeply sequenced using ribo-depleted RNA from dopamine neurons laser-captured from human post-mortem brains of patients with PD and controls using a new method we developed, termed laserRNAseq.
- A vast universe of annotated as well as novel non-coding RNAs expressed in dopamine neurons was discovered suggesting a more diverse and much more complex transcriptional architecture than previously imagined.
- The expression of 121 non-coding transcripts and more than a thousand protein-coding transcripts in nigral dopamine neurons was significantly associated with Parkinson's Lewy body neuropathology.
- Expression of the gene *ALDH1A1* that plays a critical role in scavenging toxic dopamine metabolites is lost in dopaminergic neurons laser-captured from patients with PD
- This study provides proof of the concept that select RNAs associated with Parkinson's in vulnerable dopamine neurons can be translated into personalized, dual-use, peripheral biomarkers. Specifically, we were able to translate four transcripts (*SNCA*, *LRRK2*, *COMT* and *CHMP3*) that were abnormally transcribed in laser-captured dopamine neurons of PD brains into potential blood biomarkers for patients with Parkinson's.

5. Conclusion

This study provides a first look at the “dark universe” of tens of thousands of short and long non-coding RNAs expressed in the human dopamine neuron of controls and in Parkinson's disease. We show unequivocally that non-coding RNAs are abundantly transcribed in dopamine neurons of human brain. We identify highly significant changes in the expression of both protein-coding and non-coding transcripts in Parkinson's disease. These changes indicate key roles for non-coding RNAs and for the protein-coding familial PD gene *DNAJC6* and *ALDH1A1* in the survival of dopamine neurons in Parkinson's. *ALDH1A1* mediates the oxidation of toxic dopamine-3,4-dihydroxyphenylacetaldehyde (DOPAL) into non-toxic 3,4-dihydroxyphenylacetic acid (DOPAC). In mice *ALDH1A1* overexpression suppresses A53T- α -synuclein-induced dopaminergic neurodegeneration and *ALDH1A1* knockout enhances dopaminergic neurodegeneration³.

By translating four transcripts (that were found to be pathologically expressed in dopamine neurons of patients with PD) into potential biomarkers accessible in blood of patients and controls, we provide proof of concept for the idea that the activity of select PD-relevant genes can be precisely monitored in blood tests. In support of this view we show robust evidence indicating significant associations between *SNCA*, *LRRK2*, *COMT*, and *CHMP3* mRNA counts in blood and Parkinson's using precise digital gene expression analysis with molecular barcodes on the NanoString platform. Additionally, previous reports indicate that *ALDH1A1* expression might also be reduced in blood of patients with PD⁴.

In future steps, we will continue to delineate all regions of the human genome actively transcribed in dopamine neurons. We will fully decode all non-coding RNAs active in dopamine neurons. We will extend our investigation from known non-coding RNAs annotated in the GENECODE database to the characterization of thousand of novel, previously not described non-coding RNAs emerging from these analyses (such as novel transcribed enhancers or eRNAs and new classes of

RNA biotypes). We will increase statistical power and scientific knowledge by extending the current observations to larger numbers of high-quality brains, to pre-clinical Lewy body neuropathology, and to additional brain cell-types (e.g. pyramidal cells). Genetically, as our sample size becomes larger we will be increasingly powered to uncover how genetic risk variants causally contribute to the degeneration of dopamine neurons via dysregulation of cell-type-specific transcription. Clinically, we will continue to translate PD-associated transcripts into peripheral tests in increasingly large⁸ and definitive studies. Moreover, we aim to assemble validated transcriptional and genetic markers into polygenic test panels with improved sensitivity and specificity. Validated biomarkers will be clinically relevant for identifying PD-vulnerable warfigthers and civilians, and for monitoring neurotoxin exposure and therapeutic response in clinical trials and eventually, clinical practice.

6. Publications since submission of the original application (chronologic order; *, denotes corresponding or co-corresponding author)

- *1. Unrecognized vitamin D₃ deficiency is common in Parkinson's: Harvard Biomarker Study. Ding H, Dhima K, Locascio J, Sohur US, Wills A-M, Flaherty AW, Unni VK, Hung AY, Selkoe DJ, Schwarzschild MA, Schlossmacher MG, Sudarsky LR, Growdon JH, Iverson AJ, Hyman BT, **Scherzer CR**. *Neurology*, 2013; 81: 1531-7. Perspective in *Nature Reviews Neurology*, 2014; 10: 8-9. doi: 10.1212/01.wnl.0000449750.81263.7d. PMID: 24799519. (Impact Factor = 8).
2. Specific splice variants are associated with Parkinson's disease. JA Santiago, **Scherzer CR**, Harvard Biomarker Study, and JA Potashkin. *Mov Disord*, 2013; 28:1724-7. doi: 10.1002/mds.25635. PMID: 24108702. (Impact Factor = 5.7)
3. A novel method for detecting 7-methyl guanine reveals aberrant methylation levels in Huntington disease. Thomas B, Matson S, Chopra V, Sun L, Sharma S, Hersch S, Rosas HD, **Scherzer CR**, Ferrante R, Matson W. *Anal Biochem*. 2013; 436: 112-20. doi: 10.1016/j.ab.2013.01.035. PMID:23416183.
- *4. Metallothioneins as dynamic markers for brain disease in lysosomal disorders. Cesani M, Cavalca E, Macco R, Leoncini G, Terreni MR, Lorioli L, Furlan R, Comi G, Doglioni C, Zacchetti D, Sessa M, **Scherzer CR**, Biffi A*, *Ann of Neurol*, 2014; 75: 127-37. doi: 10.1002/ana.24053. PMID:24242821. (Impact Factor = 10)
5. Motor impulsivity in Parkinson disease: Associations with COMT and DRD2 polymorphisms. Ziegler DA, Ashourian P, Wonderlick JS, Sarokhan AK, Prelec D, **Scherzer CR**, Corkin S. Motor impulsivity in Parkinson disease: Associations with COMT and DRD2 polymorphisms, *Scand J Psychol.*, 2014; 55:278-86. doi: 10.1111/sjop.12113.PMID:24749760.
6. MHC-I expression renders catecholaminergic neurons susceptible to T-cell-mediated degeneration. Cebrian C, Zucca FA, Mauri P, Steinbeck JA, Studer L, **Scherzer CR**, Kanter E, Budhu S, Mandelbaum J, Vonsattel JP, Luigi Zecca, Loike JD, Sulzer D. *Nat Commun*, 2014; 5: 3633. doi: 10.1038/ncomms4633. PMID:24736453. (Impact Factor = 11.5)
7. High prevalence of NMDA receptor IgA/IgM antibodies in different dementia types. Doss S, Wandinger KP, Hyman BT, Panzer JA, Synofzik M, Dickerson B, Mollenhauer B, **Scherzer CR**, Iverson AJ, Finke C, Schöls L, Müller Vom Hagen J, Trenkwalder C, Jahn H, Höltje M, Biswal BB, Harms L, Ruprecht K, Buchert R, Höglinger GU, Oertel WH, Unger MM, Körtvélyessy P, Bittner D, Priller J, Spruth EJ, Paul F, Meisel A, Lynch DR, Dirnagl U, Endres M, Teegen B, Probst C, Komorowski L, Stöcker W, Dalmau J, Prüss H. *Ann Clin Transl Neurol*. 2014; 1: 822-32. doi: 10.1002/acn3.120. PMID:25493273.
8. Network analysis identifies SOD2 mRNA as a potential biomarker for Parkinson's disease. Santiago JA, **Scherzer CR**, Potashkin JA. *PLoS One*. 2014; 9: e109042. doi: 10.1371/journal.pone.0109042. eCollection 2014. PMID:2527975.
- *9. Association between α -synuclein blood transcripts and early, neuroimaging-supported Parkinson's disease. Locascio JJ, Eberly S, Liao Z, Liu G, Hoessing AN, Duong K, Trisini-Lipsanopoulos A, Dhima K, Hung AY, Flaherty AW, Schwarzschild MA, Hayes MT, Wills AM, Shivraj Sohur U, Mejia NI, Selkoe DJ, Oakes D, Shoulson I, Dong X, Marek K, Zheng B, Iverson A, Hyman BT, Growdon JH, Sudarsky LR, Schlossmacher MG, Ravina B, **Scherzer CR**. *Brain*. 2015 Jul 28. pii: awv202. [Epub ahead of print] PMID: 26220939. (Impact Factor = 9)

Presentations

- 2013 Can BIG DATA solve big questions?
Movement Disorders Conference, Department of Neurology
Massachusetts General Hospital
Boston, MA
- 2014 Can BIG data help neurological disease?
Massachusetts General Hospital, J. Philip Kistler Stroke Center Meeting
Massachusetts General Hospital, Boston, MA
- 2014 From BIG data to personalized medicine for Parkinson's
Icahn School of Medicine at Mount Sinai
New York, NY
- 2014 Mapping transcriptome space in Parkinson's: from the known world to an expanding universe
Molecular Pathology Seminar
Johns Hopkins University, Baltimore, MD
- 2014 The PD Brain Map: How does the human genome function in dopamine neurons?
NIH/NINDS Parkinson's Disease Genetics and Systems Biology Workshop
Bethesda, MD
- 2014 Human genome function in Parkinson's
Neurology Department, Invited Lecture
Goethe University, Frankfurt, Germany
- 2015 Parkinson's disease: How Big Data will enable a future personal medicine
Genzyme
Framingham, MA

Abstracts

- 2013 The PD Brain Map Project: Mapping the Transcriptional Architecture of Dopamine Neurons in Human Brain
Annual Meeting of the American Society for Human Genetics
Boston, MA
- 2015 Functional characterization of the human brain transcriptome in Parkinson's disease
European Molecular Biology Laboratory Meeting
Heidelberg, Germany
- 2015 BRAINCORE: How Does the Human Genome Function in Specific Brain Neurons?
Annual Meeting of the American Society for Human Genetics
Baltimore, MD

7. Inventions, Patents and Licenses - Nothing to report

8. Reportable Outcomes - Nothing to report

9. Other Achievements

2015 - Scientific Advisory Board, American Parkinson Disease Association

2015 Neurotoxin Exposure Treatment PD Vision Setting Panel, U.S. Department of Defense

10. References

1. Bras JM, Singleton A. Genetic susceptibility in Parkinson's disease. *Biochim Biophys Acta* 2009;1792:597-603.
2. Singleton AB, Farrer MJ, Bonifati V. The genetics of Parkinson's disease: progress and therapeutic implications. *Mov Disord* 2013;28:14-23.
3. Liu G, Yu J, Ding J, et al. Aldehyde dehydrogenase 1 defines and protects a nigrostriatal dopaminergic neuron subpopulation. *J Clin Invest* 2014.
4. Molochnikov L, Rabey JM, Dobronevsky E, et al. A molecular signature in blood identifies early Parkinson's disease. *Mol Neurodegener* 2012;7:26.
5. Stegle O, Parts L, Piipari M, Winn J, Durbin R. Using probabilistic estimation of expression residuals (PEER) to obtain increased power and interpretability of gene expression analyses. *Nat Protoc* 2012;7:500-7.
6. Shabalín AA. Matrix eQTL: ultra fast eQTL analysis via large matrix operations. *Bioinformatics* 2012;28:1353-8.
7. Mohammadi D. The Harvard Biomarker Study's big plan. *Lancet Neurol* 2013;12:739-40.
8. Locascio JJ, Eberly S, Liao Z, et al. Association between alpha-synuclein blood transcripts and early, neuroimaging-supported Parkinson's disease. *Brain* 2015.
9. Ziegler DA, Ashourian P, Wonderlick JS, et al. Motor impulsivity in Parkinson disease: associations with COMT and DRD2 polymorphisms. *Scand J Psychol* 2014;55:278-86.
10. Ding H, Dhima K, Lockhart KC, et al. Unrecognized vitamin D3 deficiency is common in Parkinson disease: Harvard Biomarker Study. *Neurology* 2013.
11. Pankratz N, Beecham GW, DeStefano AL, et al. Meta-analysis of Parkinson's disease: identification of a novel locus, RIT2. *Ann Neurol* 2012;71:370-84.
12. Ding H, Sarokhan AK, Roderick SS, et al. Association of SNCA with Parkinson: replication in the Harvard NeuroDiscovery Center Biomarker Study. *Mov Disord* 2011;26:2283-6.
13. Kurashige T, Takahashi T, Yamazaki Y, et al. Localization of CHMP2B-immunoreactivity in the brainstem of Lewy body disease. *Neuropathology* 2013;33:237-45.

11. Appendix

Association between α -synuclein blood transcripts and early, neuroimaging-supported Parkinson's disease

Locascio JJ, Eberly S, Liao Z, Liu G, Hoelsing AN, Duong K, Trisini-Lipsanopoulos A, Dhima K, Hung AY, Flaherty AW, Schwarzschild MA, Hayes MT, Wills AM, Shivraj Sohur U, Mejia NI, Selkoe DJ, Oakes D, Shoulson I, Dong X, Marek K, Zheng B, Iverson A, Hyman BT, Growdon JH, Sudarsky LR, Schlossmacher MG, Ravina B, and **Scherzer CR**.

Brain. 2015 Jul 28. pii: awv202. [Epub ahead of print]

Association between α -synuclein blood transcripts and early, neuroimaging-supported Parkinson's disease

Joseph J. Locascio,^{1,2,*} Shirley Eberly,^{3,*} Zhixiang Liao,^{1,4,*} Ganqiang Liu,^{1,4} Ashley N. Hoising,^{1,4,5} Karen Duong,^{1,4,5} Ana Trisini-Lipsanopoulos,^{1,4,5} Kaltra Dhima,^{1,4,5} Albert Y. Hung,² Alice W. Flaherty,^{2,6} Michael A. Schwarzschild,² Michael T. Hayes,⁷ Anne-Marie Wills,^{2,5} U. Shivraj Sohur,² Nicté I. Mejia,² Dennis J. Selkoe,^{4,7} David Oakes,³ Ira Shoulson,⁸ Xianjun Dong,^{1,4} Ken Marek,⁹ Bin Zheng,^{1,4} Adrian Ivinson,^{4,5} Bradley T. Hyman,^{2,5} John H. Growdon,² Lewis R. Sudarsky,⁷ Michael G. Schlossmacher,¹⁰ Bernard Ravina¹¹ and Clemens R. Scherzer^{1,2,4,5,7}

*These authors contributed equally to this work.

There are no cures for neurodegenerative diseases and this is partially due to the difficulty of monitoring pathogenic molecules in patients during life. The Parkinson's disease gene α -synuclein (SNCA) is selectively expressed in blood cells and neurons. Here we show that SNCA transcripts in circulating blood cells are paradoxically reduced in early stage, untreated and dopamine transporter neuroimaging-supported Parkinson's disease in three independent regional, national, and international populations representing 500 cases and 363 controls and on three analogue and digital platforms with $P < 0.0001$ in meta-analysis. Individuals with SNCA transcripts in the lowest quartile of counts had an odds ratio for Parkinson's disease of 2.45 compared to individuals in the highest quartile. Disease-relevant transcript isoforms were low even near disease onset. Importantly, low SNCA transcript abundance predicted cognitive decline in patients with Parkinson's disease during up to 5 years of longitudinal follow-up. This study reveals a consistent association of reduced SNCA transcripts in accessible peripheral blood and early-stage Parkinson's disease in 863 participants and suggests a clinical role as potential predictor of cognitive decline. Moreover, the three independent biobank cohorts provide a generally useful platform for rapidly validating any biological marker of this common disease.

- 1 Neurogenomics Lab and Parkinson Personalized Medicine Program, Harvard Medical School and Brigham and Women's Hospital, Cambridge, MA 02139, USA
- 2 Department of Neurology, Massachusetts General Hospital, Boston, MA 02114, USA
- 3 Department of Biostatistics and Computational Biology, University of Rochester Medical Center, Rochester, NY 14642, USA
- 4 Ann Romney Centre for Neurologic Diseases, Brigham and Women's Hospital, Boston, MA 02115, USA
- 5 Biomarkers Program, Harvard NeuroDiscovery Center, Boston, MA 02115, USA
- 6 Department of Psychiatry, Massachusetts General Hospital, Boston, MA 02114, USA
- 7 Department of Neurology, Brigham and Women's Hospital, Boston, MA 02115, USA
- 8 Program for Regulatory Science and Medicine, Department of Neurology, Georgetown University, Washington, DC 20007, USA
- 9 Institute for Neurodegenerative Disorders, New Haven, CT 06510, USA
- 10 Program in Neuroscience, Ottawa Hospital Research Institute, University of Ottawa, Ottawa, Ontario K1H8M5, Canada
- 11 Voyager Therapeutics, Clinical Development, Cambridge, MA 02142, USA

Correspondence to: Clemens R. Scherzer, MD,
Neurogenomics Lab and Parkinson Personalized Medicine Program,

Received November 14, 2014. Revised May 12, 2015. Accepted May 25, 2015.

© The Author (2015). Published by Oxford University Press on behalf of the Guarantors of Brain. All rights reserved.

For Permissions, please email: journals.permissions@oup.com

Harvard Medical School and Brigham and Women's Hospital,
65 Landsdowne Street, Suite 307A,
Cambridge, MA 02139, USA
E-mail: cscherzer@rics.bwh.harvard.edu

Keywords: α -synuclein; biomarker; gene expression; biobank; mitochondria; cognitive decline

Abbreviations: DAT = dopamine transporter; HBS = Harvard Biomarker Study; MMSE = Mini-Mental State Examination; PPMI = Parkinson's Progression Markers Initiative; PROBE = Prognostic Biomarkers in Parkinson's Disease Study

Introduction

Nearly 5 million people have Parkinson's disease, a sporadic, progressive neurodegenerative disorder linked to a complex genetic architecture and environmental exposures. Over the next 15 years, this number will almost double to 9 million patients worldwide (Dorsey *et al.*, 2007). Already today, the costs to public health are enormous—estimated at \$10.8 billion annually in the USA alone (O'Brien *et al.*, 2009). There are no cures or disease-modifying therapies for Parkinson's disease, and this may be due in part to our inability to monitor biological markers of early disease. Currently, patients with Parkinson's disease are diagnosed, cared for, and assigned to clinical trials based on medical history and physical exam.

To translate research progress into the clinic, simple, dynamic biomarkers are needed that are rooted in genetic discoveries. Traditionally, Parkinson's disease has been approached from a clinical perspective, as a homogeneous movement disorder typified by the triad of resting tremor, bradykinesia and rigidity. Based on this clinical view of the disease, it was postulated that a single laboratory or imaging biomarker for Parkinson's disease would be developed to serve as a universal diagnostic. This goal has been elusive. More recently, a human genome-inspired view of Parkinson's disease has provided unequivocal evidence for a diversity of genetic variants, each exacting a small, but definitive effect, underlying the clinical phenotype (Nalls *et al.*, 2014). It is becoming clearer that Parkinson's disease is aetiologically, genetically, pathologically and even clinically more heterogeneous than previously implied (Scherzer and Feany, 2004; Scherzer *et al.*, 2004; Forman *et al.*, 2005). In light of this diversity the traditional expectation of a 'one size fits all' laboratory diagnostic that can detect all patients and answer all clinical questions may need to be revised. Personalized biomarkers, tailored to patient endophenotypes and specific clinical applications, are likely to emerge. They may more appropriately describe the molecular architecture of the disease.

What are the practical applications that such biomarkers may serve, if they cannot be used as a universal diagnostic? Prognostic and mechanism-directed biomarkers will help to stratify patients for clinical trials. Risk markers are required for identifying high-risk individuals prior to the onset of overt clinical symptoms, when disease-modifying interventions will be most effective. Pharmacodynamic markers are required to determine whether an investigational drug

engages the intended target. In cancer drug development, biomarker use has increased average phase III trial success by 34% (Hayashi *et al.*, 2013) and shortened total drug development timelines (Chin *et al.*, 2011). Emerging CSF tests for Parkinson's disease (Mollenhauer *et al.*, 2011) and dopamine transporter imaging are relatively invasive or time-consuming and expensive. Blood-based tests are attractive, but no disease-causing molecule has been rigorously validated.

Accumulation of α -synuclein protein (encoded by the *SNCA* gene) in the brain is a hallmark of Parkinson's disease (Braak *et al.*, 2002). For nearly two decades since its characterization as an electric ray protein, the Parkinson's disease gene α -synuclein was thought of as 'neuron-specific' (Maroteaux *et al.*, 1988), but this view was disproven with the discovery of high levels of α -synuclein expression in blood cells, particularly those of erythroid lineage (Nakai *et al.*, 2007; Scherzer *et al.*, 2008; Maitta *et al.*, 2011). Here we tested whether levels of *SNCA* transcripts (mRNAs) in circulating blood cells might serve to identify individuals with increased risk of Parkinson's disease. We evaluated whether levels of α -synuclein gene (*SNCA*) transcripts are associated with the presence of early-stage, clinically and neuroradiologically confirmed Parkinson's disease in a platform of three independent biomarker study cohorts.

Materials and methods

Brief methods are summarized below. See the online [Supplementary material](#) for detailed clinical, laboratory and statistical experimental methods for the Harvard Biomarker Study (HBS), Prognostic Biomarkers in Parkinson's Disease Study (PROBE) and Parkinson's Progression Markers Initiative (PPMI), respectively, and for the meta-analysis across three studies.

Biospecimen collections

Briefly, the studies were designed to minimize bias from sample processing by collecting, handling and analysing specimens of cases and controls in a standardized manner following defined operating procedures and according to rules of evidence (Ransohoff, 2004, 2005; Hennecke and Scherzer, 2008). Cases and controls were processed in parallel by technicians blinded to diagnosis to avoid bias due to 'run order' of samples. Venous whole blood was collected in PAXgeneTM tubes.

Blood and PAXgene™ reagents were mixed by gently inverting 8–10 times.

RNA isolation and quality control

RNA was extracted according to the PAXgene™ Blood RNA kit manual extraction protocol including DNase treatment. RNA quality was determined with the RNA 6000 Nano Chip on an Agilent 2100 Bioanalyzer (Agilent Technologies) and the RNA Integrity Number package (Imbeaud *et al.*, 2005).

Quantitative PCR and microarray expression analyses

Quantitative PCR was performed similar to Scherzer *et al.* (2008). Microarray procedures were performed as described (Zheng *et al.*, 2010) using Illumina HumanHT-12v3 Expression BeadChips. The PROBE microarray data set has been submitted to the GEO database (accession number GSE57475).

NanoString assay performance

Probes were designed according to the manufacturer's design principles (Geiss *et al.*, 2008), including screening for inter- and intra-reporter and capture probe interactions, and selection for probes with optimal melting temperatures (Geiss *et al.*, 2008). Direct counts of the target RNAs were measured in 125 ng of RNA by digital expression analysis based on NanoString technology (without reverse transcription into cDNA). Probes for the target and control RNAs were multiplexed and assayed on the nCounter Digital Analyzer. The laboratory running the assay was blinded to the diagnosis. No-template (negative) controls containing water substituted for template were run and no signal was detected. To avoid run-order bias, samples of cases or controls were randomly assigned to plates. To control for plate-to-plate variation and drift, equal amounts of Human Universal Reference RNA were included at the beginning, within, and at the end of the entire experiment. Transcript counts (excluding SNCA-007) measured in reference RNA were highly correlated with $R^2 > 0.999$ both within one plate and in-between different plates, thus excluding drift as a potential source of bias in the experiment. Furthermore, 5% of participants' samples were randomly resampled to verify the retest reliability (technical precision). The average R^2 was 0.98 for these correlations. The NanoString data are accessible through the PPMI website (<http://www.ppmi-info.org>).

Results

Harvard Biomarker Study: SNCA mRNA abundance in early-stage Parkinson's disease

We first evaluated relative SNCA mRNA abundance in a case-control study nested in the HBS using precise, kinetic, quantitative PCR based on fluorogenic 5' nuclease chemistry (quantitative PCR). We specifically designed the HBS as

Table 1 The HBS: patient characteristics

	Patients with Parkinson's disease (n = 222)	Controls (n = 183)	P-value
Age at phlebotomy (years, SD)	67.05 (9.53)	68.44 (10.23)	0.16
Gender			
Male (n, %)	130 (58.6)	74 (40.4)	0.0003
Female (n, %)	92 (41.4)	109 (59.6)	
Clinical findings (n, %)			
UKPDBBS ^a	207 (93.2)		
Resting tremor	164 (73.87)		
Bradykinesia	221 (99.59)		
Rigidity	219 (98.65)		
Postural imbalance	109 (49.10)		
Asymmetric onset	175 (78.83)		
Medications (n, %)			
Levodopa	170 (76.58)		
Dopamine replacement medications	201 (90.5)		
De novo	21 (9.5%)		
Disease severity			
Disease duration (years) (mean, SD)	4.8 (4.5)		
Hoehn and Yahr (mean, SD)	2.1 (0.6)		
UPDRS Part I: mentation, behaviour, mood	1.5 (1.5)		
UPDRS Part II: ADLs	8.7 (5.2)		
UPDRS Part III: motor examination	18.5 (9.0)		
UPDRS total score	30.9 (13.6)		

^aUnited Kingdom Parkinson's Disease Brain Bank Society diagnostic criteria. ADL = activities of daily living.

a clinical biomarker study with rigorous, predefined collection and processing protocols. Cases and controls had similar ages, but patients with Parkinson's disease were more likely to be males (Table 1). Cases were at an early stage of the disease, with a mean modified Hoehn and Yahr stage of 2.1. A large majority of cases (201 of 222, 90.5%) were on medications that ameliorate the dopamine deficiency caused by the degeneration of neurons in the substantia nigra, while 9.5% (21 of 222) were untreated, *de novo* patients. The controls were recruited from the same source population. Case and control samples were collected, processed, and analysed in parallel. Samples were required to meet stringent quality control criteria in order to enter the study including a RNA Integrity Number (RIN) (Auer *et al.*, 2003) threshold of ≥ 7.3 indicating high RNA quality. The two groups had excellent RNA quality with mean RINs of 8 ± 0.4 versus 8.05 ± 0.4 [mean $\pm$ standard deviation (SD)], respectively.

Relative SNCA transcript abundance in venous blood was significantly lower for patients with Parkinson's disease than for controls (difference of 20%; P -value = 0.0004;

Table 2 Blood SNCA expression is associated with early-stage clinical and DAT-neuroimaging supported Parkinson's disease in three independent populations and on three independent assay platforms

	Cases	Controls	Difference of means	P-value*
HBS				
Quantitative PCR platform ^c				
n (total n = 405)	222	183		
Unadjusted relative SNCA expression (mean, SD)	51.5 (1.97)	64.2 (1.8)	−20 ^e	0.0004
Adjusted relative SNCA expression ^d (mean, 95% CI)	49 (45–54)	61 (55–68)	−17 ^e	0.003
PROBE				
Quantitative PCR platform ^c				
n (total n = 118)	76	42		
Unadjusted relative SNCA expression (mean, SD)	75.6 (2.27)	96.3 (1.7)	−22 ^e	0.025
Adjusted relative SNCA expression ^d (mean, 95% CI)	72 (61–85)	96 (76–122)	−25 ^e	0.046
Microarray platform				
n (total n = 142)	93	49		
ILMN-PROBE1 ^a Unadjusted SNCA expression (mean, SD)	2846 (1962)	3469 (1622)	−18	0.01
ILMN-PROBE1 ^a Adjusted SNCA expression ^d (mean, 95% CI)	2452 (2167–2775)	3098 (2635–3644)	−21	0.02
ILMN-PROBE2 ^b Unadjusted SNCA expression (mean, SD)	681 (473)	823 (432)	−17	0.02
ILMN-PROBE2 ^b Adjusted SNCA expression ^d (mean, 95% CI)	578 (508–657)	730 (616–864)	−21	0.03
PPMI				
Digital gene expression platform				
n (total n = 340)	202	138		
Unadjusted analyses				
E3E4-SNCA transcript counts (mean, SD)	13 458 (7812)	15 959 (12 576)	−16	0.08
E4E6-SNCA transcript counts (mean, SD)	2826 (1860)	3496 (3,119)	−19	0.04
Long 3'UTR-SNCA transcript counts (mean, SD)	573 (387)	781 (1013)	−27	0.007
SNCA-007 transcript counts (mean, SD)	23 (16)	25 (21)	−8	0.6
Adjusted models ^d				
Adjusted E4E6-SNCA transcript counts (mean, 95% CI)	2409 (2196–2643)	2784 (2495–3107)	−13	0.046
Adjusted long 3'UTR-SNCA transcript counts (mean, 95% CI)	485 (440–534)	578 (515–647)	−16	0.02

^aILMN-PROBE1 indicates Illumina probe ILMN1701933 that targets the 3'UTR of SNCA.

^bILMN-PROBE2 indicates Illumina probe ILMN1766165 that targets exon 5 of SNCA.

^cAll statistical analyses for quantitative PCR assays were conducted using the natural logarithm of Δ CT values. For presentation purposes in Table 2 and Figs 1 and 2, a linearizing transformation of mean Δ CTs was performed. Values shown are in arbitrary units.

^dAdjusted for covariates of gender, and white and red blood cell counts.

^eIndicates per cent reduction in expression in cases compared to controls determined using the comparative CT method.

*Indicates significance analysis for all measurement values was performed on log-transformed data.

Table 2 and Fig. 1A). *Post hoc* analysis of the subgroup of untreated, *de novo* cases showed consistent results with a >20% lower mean relative SNCA transcript abundance in the *de novo* cases compared to the controls (P -value = 0.01). Counts of the four endogenous reference genes (Fig. 1B) and their geometric mean (that was used to normalize for input RNA) were virtually identical in cases and controls (Fig. 1C). Importantly, complete blood counts in cases and controls could not account for the reduction in relative SNCA mRNA abundance in patients consistent with previous reports by others (Kasten et al., 2010). Specifically, white blood cell counts (Fig. 1D) and platelet counts (Fig. 1F) were unchanged between cases and controls, while red blood cell counts were marginally higher in cases than in controls (P = 0.03; Fig. 1E).

General linear model analysis was performed adjusting for the covariates of counts of white and red blood cells, and gender. In this covariate-adjusted analysis the mean relative abundance of SNCA expression was 17% lower in cases than in controls with P = 0.003.

Odds ratios (OR) for the lowest (first) quartile relative to the highest (fourth) quartile for blood SNCA transcript abundance were associated with Parkinson's disease status in the unadjusted analysis and remained associated following adjustments for the covariates of age, hours at 4°C, red and white blood cells, and platelets (Model 1) or for gender (Model 2) with odds ratios ranging from 2.14 (95% CI, 1.10–4.14) to 2.15 (95% CI, 1.2–3.82) (Table 3).

PROBE study

Replication of low SNCA mRNA abundance in dopamine transporter imaging-supported Parkinson's disease

To be of use in clinical trials, it must be feasible to measure a biomarker in a multicentre study design. We thus further evaluated SNCA mRNA levels in the USA-wide, multicentre PROBE study (Supplementary Table 1). There were no significant baseline inequalities for age, gender, and white and red blood cell counts. Cases had a mean




Figure 1 HBS: reduced *SNCA* mRNA abundance in early-stage Parkinson's disease. (A) Mean *SNCA* expression was significantly lower in 222 patients with early-stage clinical Parkinson's disease (PD) compared to 183 controls (HC) without neurologic disease enrolled in the HBS using precise, quantitative PCR (unadjusted difference of 20%; P -value of 0.0004). * P -value = 0.01 for untreated, *de novo* cases versus controls. Box plots visualize first, third quartiles and means; the ends of the whiskers represent the lowest (or highest) value still within 1.5-times the interquartile range. (B) Expression values of the four endogenous reference genes used in the HBS were virtually identical in cases and controls. Cycle threshold (CT) values (standard error) are shown for cases (crimson bars) and controls (grey bars). (C) The geometric mean of these four endogenous reference genes was used to normalize for input RNA. (D–F) Importantly, complete blood counts in cases and controls could not account for the reduction in relative *SNCA* mRNA abundance in patients. Specifically, white blood cell counts (D, WBC) and platelet counts (F) did not significantly differ between cases and controls, while red blood cell counts (RBC) were marginally higher in cases than in controls (E; mean red blood cell, 4.6 versus 4.5, $P = 0.03$). Bar graphs show means and standard errors.

Hoehn and Yahr stage of 2.0, a mean total UPDRS score of 30.6, and mean disease duration of 5.5 years. The clinical diagnosis was supported by dopamine transporter (DAT) imaging with ^{123}I - β -CIT. A subset of 76 cases with Parkinson's disease and 42 controls with available high-quality RNA were analysed by quantitative PCR. The mean relative abundance of *SNCA* expression was 22% lower in cases than in controls with $P = 0.025$ in the unadjusted analysis (Table 2 and Fig. 2A). This remained significant after adjusting for the covariates of white and red blood cells, and gender with $P = 0.046$. Quantitative PCR allows for precise relative, but not absolute quantification. Thus, the difference in *SNCA* mRNA abundance in cases relative to controls found in PROBE was comparable to the 20% difference seen in HBS (i.e. in Figs 1A and 2A), while the absolute expression values (in arbitrary units) were not directly comparable due to different calibrators.

We confirmed these expression changes using two *SNCA* probes represented on a secondary, microarray expression platform. Results for both probes were available for 93

cases and 49 controls enrolled in PROBE and confirmed that relative blood *SNCA* expression was lower in cases compared to controls (Table 2 and Fig. 2B and C). Results for ILMN-PROBE1 confirmed an association between *SNCA* expression and Parkinson's disease adjusting for the covariates of white and red blood cell counts, and gender with $P = 0.02$. The association was further confirmed by ILMN-PROBE2 with $P = 0.03$, again adjusting for the same covariates.

Individuals with *SNCA* transcript abundance in the lowest quartile of values had an OR for Parkinson's disease of 4.5 (95% CI, 1.3–15.0) compared to individuals in the highest quartile of values (Table 3). The corresponding odds ratios for the two microarray probes were 4.4 (95% CI, 1.4–14.5) and 4.6 (95% CI, 1.5–13.7), respectively. Thus, measuring blood *SNCA* expression was feasible in a multicentre study and reduced blood *SNCA* expression in Parkinson's disease was confirmed in this independent population on two distinct analysis platforms.

Table 3 The odds ratio for Parkinson's disease prevalence is increased in individuals with blood SNCA expression in the lowest quartile of values compared to individuals with SNCA expression in the highest quartile of values in three study populations

	Quartile			
	1	2	3	4
HBS				
Relative SNCA mRNA abundance	5–37 ^a	37–60 ^a	60–91 ^a	91–294 ^a
Parkinson's disease (<i>n</i> = 222), <i>n</i> /total (%)	67/222 (30)	58/222 (26)	50/222 (23)	47/222 (21)
Healthy control (<i>n</i> = 183), <i>n</i> /total (%)	34/183 (18.6)	43/183 (23.5)	51/183 (27.9)	55/183 (30)
Unadjusted OR, (95% CI) ^b	2.31 (1.31–4.1)	1.58 (0.91–2.75)	1.15 (0.66–1.99)	1
P-value	0.0036	0.11	0.63	
Model 1, OR (95% CI) ^c	2.14 (1.10–4.14)	1.86 (1.01–3.42)	1.07 (0.6–1.92)	1
P-value	0.024	0.046	0.82	
Model 2, OR (95% CI) ^d	2.15 (1.2–3.82)	1.61 (0.92–2.83)	1.12 (0.64–1.96)	1
P-value	0.0096	0.098	0.69	
PROBE^e				
Relative SNCA mRNA abundance	16–52 ^a	56–79 ^a	84–119 ^a	119–2702 ^a
Parkinson's disease (<i>n</i> = 76), <i>n</i> /total (%)	24/76 (32)	20/76 (26)	17/76 (22)	15/76 (20)
Healthy control (<i>n</i> = 42), <i>n</i> /total (%)	5/42 (12)	10/42 (24)	13/42 (31)	14/42 (33)
OR (95% CI) ^{b,f}	4.5 (1.3–15.0)	1.9 (0.7–5.3)	1.2 (0.4–3.4)	1
P-value	0.02	0.90	0.25	
PPMI				
Long-3'UTR-SNCA transcript counts	59–328	335–497	505–786	796–10861
Parkinson's disease (<i>n</i> = 202), <i>n</i> /total (%)	58/202 (29)	48/202 (24)	55/202 (27)	41/202 (20)
Healthy control (<i>n</i> = 138), <i>n</i> /total (%)	27/138 (19.6)	37/138 (26.8)	30/138 (21.7)	44/138 (31.9)
Unadjusted OR, 95% CI) ^b	2.3 (1.2–4.3)	1.4 (0.8–2.5)	1.97 (1.1–3.6)	1
P-value	0.009	0.28	0.03	
Adjusted OR (95% CI) ^g	2.1 (1.1–4.0)	1.4 (0.8–2.6)	2.1 (1.1–3.9)	1
P-value	0.02	0.25	0.02	

^aAll statistical analyses for quantitative PCR assays were conducted using Δ CT values. For presentation purposes the ranges of each quartile of values shown were linearly transformed, e.g. 2^{lowest Δ CT} and 2^{highest Δ CT}.

^bUnadjusted odds ratio (OR) and confidence intervals (CI) were calculated using logistic regression models comparing the risk of each of quartiles 1 to 3 to the reference quartile 4 (highest relative SNCA transcript abundance).

^cModel 1 consists of covariates age, hours in 4°C, red blood cells, platelets, and white blood cells in addition to the SNCA quartiles.

^dModel 2 consists of covariate sex in addition to the SNCA quartiles.

^ePROBE results on quantitative PCR platform are shown, similar results were obtained on the microarray platform.

^fNo adjustments were indicated.

^gModel consists of the covariate white blood cells in addition to the quartiles of long 3'UTR-SNCA counts.

PPMI

Reduced abundance of disease-relevant SNCA isoforms already near disease onset

These associations prompted us to further extend and verify our findings in patients near disease onset, when individual motor symptoms are just beginning to manifest. Two hundred and two cases with *de novo* motor Parkinson's disease, with less than 2 years of disease and with confirmed loss of dopaminergic substantia nigra terminals by DAT imaging and 138 age- and sex-matched controls without neurologic disease and with normal DAT imaging, who met stringent RNA quality-control criteria (RIN $\geq$ 7.3) were included. Key eligibility criteria for PPMI require the presence of asymmetric resting tremor or asymmetric bradykinesia, a supportive DAT scan, and a diagnosis of Parkinson's disease for 2 years or less, without current dopamine replacement medication (Parkinson Progression Marker Initiative, 2011). Patients therefore

were clinically at a very early stage and did not yet necessarily fulfil the standard, full clinical diagnostic criteria of the UK Parkinson's Disease Society Brain Bank that require bradykinesia and a second mandatory clinical cardinal sign and three supportive clinical criteria to be present for diagnosis (Hughes *et al.*, 1992). Cases and controls were frequency matched for age and gender (Supplementary Table 2). The PPMI population was remarkable for the virtual absence of Parkinson's disease medications, and the equal distribution of potentially confounding variables of age, gender, and blood cell counts among cases and controls (Supplementary Table 2 and Fig. 3G–I).

The advent of digital expression analysis with molecular barcodes (Geiss *et al.*, 2008; Malkov *et al.*, 2009) allowed us to count the abundance of SNCA transcripts directly in RNA from blood cells, without the need for reverse transcription or PCR amplification. No signal was detected in controls lacking template. To avoid run-order bias, samples of cases or controls were randomly assigned to plates.




Figure 2 USA-wide, multicentre PROBE study: replication of low SNCA mRNA abundance in DAT imaging-confirmed Parkinson's disease. (A) We further evaluated this association in 76 cases with DAT imaging-confirmed Parkinson's disease and 42 controls enrolled in the PROBE study. A 22% reduction in levels of blood-SNCA expression was seen in cases compared to controls on the quantitative PCR platform ($P = 0.025$). (B and C) These results were technically replicated on a second, independent microarray expression platform that includes two distinct probes for SNCA. Results for each of the two probes confirmed that relative blood-SNCA expression was lower in cases compared to controls. Box plots visualize first, third quartiles and means; the ends of the whiskers represent the lowest (or highest) value still within 1.5-times the interquartile range.

Transcript counts (excluding SNCA-007) measured in reference RNA were highly correlated with $R^2 > 0.999$ both within one plate and in-between different plates, thus excluding drift as a potential source of bias. Furthermore, 5% of participants' samples were randomly resampled to verify the retest reliability (technical precision). The average R^2 was 0.98 for these correlations.

Alternative SNCA transcript isoform usage has been suggested as a potential convergent mechanism in Parkinson's disease pathology (Rhinn *et al.*, 2012). We therefore designed multiple probes (Supplementary Fig. 1) to target the boundaries of exon 3 and exon 4 (henceforth termed E3E4-SNCA transcript isoform), transcripts specifically with a long 3'UTR region (which may redirect α -synuclein towards mitochondria; Rhinn *et al.*, 2012), transcripts that skip exon 5 (termed E4E6-SNCA transcript isoform) and are translated into a truncated, aggregation-prone 112 amino acid protein (Murray *et al.*, 2003), or the rare short SNCA-007 transcript isoform that comprises exons 1–4. Relative counts of long 3'UTR-SNCA transcripts were reduced by 27% in cases with Parkinson's disease compared to controls in the unadjusted analysis with $P = 0.007$ (Table 2 and Fig. 3A). This was confirmed with a second digital molecular probe directed at a different target sequence in long-3'UTR-SNCA transcripts. The relation between long-3'UTR-SNCA transcripts and Parkinson's disease remained significant with a P -value of 0.02 after adjusting for the covariates of white blood cell, red blood cell, and gender (Table 2). Counts of exon 5-skipping transcripts were also associated with Parkinson's disease in unadjusted and adjusted analyses with P -values of 0.04 and 0.046, respectively (Table 2 and Fig. 3B). Counts for E3E4-SNCA transcripts were 16% lower in cases compared to controls but did not reach significance. Counts of the rare SNCA-007 transcript did not significantly differ between cases and controls. To estimate

whether SNCA expression is specially associated with Parkinson's disease or whether Parkinson's disease-linked transcripts are generally perturbed in circulating blood cells of patients with Parkinson's disease, we examined the expression of loci linked to familial Parkinson's disease [*PARK7* (also known as *DJ-1*) and *PARK15* (also known as *FBXO7*)] and *ZNF746* (implicated in mediating the effects of mutant *PARK2*; Shin *et al.*, 2011). Furthermore, the *QDPR* gene produces an essential cofactor for dopamine biosynthesis and was also evaluated. These Parkinson's disease-linked transcripts were similarly expressed in cases and controls with no or minimal fold-changes of $<5\%$ and non-significant P -values [although one of five *PARK15/FBXO7* isoforms tested (*FBXO7-001*) showed a marginal trend with $P = 0.06$]. For example, mean counts for *PARK7* mRNAs were 1144 in *de novo* cases and 1160 in controls (1.5% change, $P = 0.4$; Fig. 3C). Counts (Fig. 3D and E) and geometric mean (Fig. 3F) of the six endogenous reference genes (used to normalize for input RNA) were nearly identical in cases and controls. Cases and controls did not significantly differ in white cells, red cells, and platelets counts (Fig. 3G–I).

Individuals with long-3'UTR-SNCA transcript counts in the lowest quartile of values had an OR > 2 for Parkinson's disease compared to individuals in the highest quartile of values in both the unadjusted analysis and analyses adjusted for white blood cell count, the only covariate retained in a step-wise backward elimination procedure (Table 3).

Risk of Parkinson's disease associated with relative SNCA transcript abundance

Clinically useful biological markers of complex diseases can have modest areas under the curve (AUC) and a wide range




Figure 3 PPMI: disease-relevant *SNCA* isoforms are already reduced near disease onset. (A and B) *SNCA* isoforms were probed on a highly robust digital expression platform in 202 cases with *de novo* motor Parkinson's disease (PD), with less than 2 years of disease and with confirmed loss of dopaminergic substantia nigra terminals by DAT imaging and 138 age- and sex-matched controls (HC) without neurological disease and normal DAT imaging participating in PPMI. Unadjusted counts of disease-relevant *SNCA* transcript isoforms with long 3' UTR (A) or skipping exon 5 (B), were reduced by 27% and 19%, respectively, in *de novo* cases compared to controls ($P = 0.007$ and 0.04 , respectively). (C) By contrast mean counts for the gene *PARK7* (*DJ-1*) mutated in rare autosomal recessive Parkinson's disease were not significantly different in *de novo* cases with sporadic Parkinson's disease compared to controls (difference of 1.5%). Box plots visualize first, third quartiles and means; the ends of the whiskers represent the lowest (or highest) value still within 1.5-times the interquartile range. (D–F) Reference genes were stably expressed in blood cells of cases compared to controls. Counts of the six endogenous reference genes used in PPMI (D and E) were virtually identical in the 202 cases and 138 controls. Mean counts (standard error) are shown for cases (orange bars) and controls (grey bars). (F) The geometric mean of these six endogenous reference genes was used to normalize for input RNA. It was nearly identical in cases and controls. (G–I) White (WBC) and red (RBC) blood cell, and platelet counts did not differ significantly between cases and controls. Bar graphs show means and standard errors.

of values that overlap in subjects with a clinical disease and in those without it (Manolio, 2003). Consistent with this notion, *SNCA* transcripts showed AUCs for Parkinson's disease in the range of a possible risk marker. The AUC for *SNCA* transcripts as a predictor of Parkinson's disease was 0.6 in HBS, ranged from 0.63 to 0.66 in PROBE (for quantitative PCR and microarray probes, respectively), and was 0.58 in PPMI.

A meta-analysis was made possible by the similar collection protocols between the cohorts and allowed for powerfully combining the three studies to estimate the association between *SNCA* transcript abundance and risk of early-stage Parkinson's disease. It increased the sample size to 500 cases and 363 controls (total of 863). *SNCA* variables estimated on quantitative PCR and digital expression platforms were standardized and ranked. The covariates of




Figure 4 Risk of Parkinson's is associated with relative SNCA transcript abundance. Individuals with *SNCA* transcript levels in the lowest quartile of values had a summary odds ratio for Parkinson's disease of 2.45 (1.65, 3.64) compared to individuals with levels in the highest quartile in a meta-analysis across the three studies. In the Forrest plot squares represent the estimate of effect, i.e. the OR for each individual study, the size of the square corresponds to the size of the study, and bars around the square represent the 95% CI of the odds ratio. The diamond indicates the summary estimate; the width (the horizontal tips) of the diamond represents the 95% CI of the summary OR.

white and red blood cell count, gender, and age were adjusted for in the meta-analysis. For PPMI we first conservatively included expression measures for the generic probe (E3E4-*SNCA*) that targets most *SNCA* transcripts. In a second iteration, we included the specific probe targeting long 3'UTR-*SNCA*. In both scenarios the association between *SNCA* blood transcripts and early Parkinson's disease was highly significant with $P = 0.0001$ and $P < 0.0001$, respectively. Overall, individuals with *SNCA* transcript levels in the lowest quartile had a highly significant OR for Parkinson's disease of 2.18 (96% CI, 1.47–3.22) and 2.45 (96% CI, 1.65–3.64; Fig. 4) for each of the two scenarios, respectively. Cochran's Q statistic equalled 2.67 ($P = 0.26$), giving no evidence of heterogeneity among the three studies.

Taken together these results indicate that reduced *SNCA* mRNA levels in circulating cells are associated with Parkinson's disease, even near disease onset and in patients with DAT-imaging supported loss of dopaminergic nigral projections. Exploration of specific transcript isoforms using digital transcript counting indicated that *SNCA* transcripts containing a long 3'UTR may be particularly relevant for this association.

SNCA transcript abundance predicts cognitive decline

Predictors of cognitive decline would be valuable for stratifying clinical trials and for advancing clinical care (Chen-Plotkin, 2014). We have tracked Mini-Mental State Examination (MMSE) scores in the HBS and the PROBE populations during multiple longitudinal in-person follow-up visits for ~5 years. To evaluate whether *SNCA* transcript levels measured at enrolment can longitudinally

predict cognitive prognosis in Parkinson's disease, we analysed the 222 patients with Parkinson's disease that were longitudinally followed in the HBS. In HBS, MMSE scores were obtained during in-person study visits at enrolment, after 12, 24, and 60 months in the study. There was overall a significant relation between *SNCA* transcript abundance at enrolment and MMSE scores during the longitudinal follow-up period with $P = 0.0007$. We then tested whether low versus high *SNCA* expression status measured at enrolment could predict longitudinal cognitive decline in these patients. The 25% of patients with Parkinson's disease with the highest *SNCA* transcript levels at enrolment ($n = 55$; termed henceforth 'SNCA high expressors') and the 25% of patients with the lowest *SNCA* transcript levels at enrolment ($n = 55$; termed 'SNCA low expressors') were compared (Fig. 5). Notably, at enrolment high and low *SNCA* expressors had nearly identical mean (SD) MMSE scores of 28.37 (1.95) versus 28.53 (2.28) ($P = 0.3$). Moreover, the two groups showed no statistically significant differences in other clinical characteristics at enrolment, including gender, age, disease duration, total Unified Parkinson's Disease Rating Scale (UPDRS) score, Hoehn and Yahr stage, medications, and years of education (Supplementary Table 3). Notably, low *SNCA* expressors showed accelerated longitudinal cognitive decline compared to high expressors with $P = 0.01$ in the longitudinal mixed random and fixed effects model analysis (Locascio and Atri, 2011) adjusting for age, gender, disease duration upon enrolment, and years of education (Fig. 5A).

We then asked whether low and high expressor status can also predict cognitive decline in PROBE. Prospective longitudinal clinical data were available for 65 of the 76 patients in PROBE with relative *SNCA* abundance measured by quantitative PCR at enrolment. Again, the 25% of patients with Parkinson's disease with the highest *SNCA* transcript levels at enrolment into PROBE ($n = 16$ in PROBE) and the 25% of patients with the lowest *SNCA* transcript levels at enrolment ($n = 16$ in PROBE) were tested. High and low *SNCA* expressors had similar MMSE scores of 29.19 (0.98) versus 28.44 (2.78) at enrolment ($P = 0.95$) (Supplementary Table 2). Mixed effects analysis adjusting for the covariates of age, gender, disease duration upon enrolment, and years of education indicated a similar trend as observed in the larger HBS population, but this did not reach statistical significance, likely because of insufficient power (Fig. 5B). PPMI was not included in the longitudinal analysis because the multi-year longitudinal follow-up data needed are not yet available. Finally, to increase power, we performed a mixed random and fixed effects meta-analysis across the low and high expressors from each of the HBS and PROBE populations. The combined analysis robustly confirmed *SNCA* expression status as predictor of longitudinal cognitive decline in Parkinson's disease with $P = 0.005$ after adjusting for pertinent covariates and for study (Fig. 5C). Similar analyses were performed to evaluate whether *SNCA* expression can also predict decline in longitudinal total UPDRS scores.




Figure 5 *SNCA* transcript abundance predicts cognitive decline in patients with Parkinson's disease during longitudinal follow-up. **(A)** The 25% patients with Parkinson's disease with the highest *SNCA* transcript levels at enrolment into HBS ($n = 55$; 'SNCA high expressors') and the 25% of patients with the lowest *SNCA* transcript levels at enrolment in HBS ($n = 55$; 'low SNCA expressors') were compared. Illustrative mean scores on the MMSE across time predicted from the estimated fixed effect parameters in the mixed random and fixed effects model analysis (Locascio and Atri, 2011) are shown for SNCA high expressors and SNCA low expressors. Low SNCA expressors showed accelerated longitudinal cognitive decline compared to high expressors with $P = 0.01$ adjusting for age, gender, disease duration upon enrolment, and years of education. **(B)** Mixed effects model analysis for the 25% SNCA high expressors ($n = 16$) versus the 25% low expressors in PROBE ($n = 16$) adjusting for covariates indicated a similar, but non-significant trend. **(C)** In the combined analysis, SNCA expression status was robustly confirmed as a predictor of longitudinal cognitive decline in Parkinson's disease with $P = 0.005$ after adjusting for pertinent covariates and study. Illustrative mean scores on the MMSE across time predicted by the fitted model in the longitudinal analysis for Parkinson's disease patients in the lowest quartile of SNCA expression levels are shown as red triangles; values for Parkinson's disease patients in the highest quartile of SNCA expression levels are represented as blue circles (solid lines indicate predicted values for subjects with a mean value of disease duration at enrolment; large-dashed lines indicate those for individuals with 1 SD longer disease duration at enrolment; and short-dashed lines indicate those for subjects with 1 SD shorter disease duration at enrolment).

These analyses were inconclusive and will require further evaluation in larger, longitudinally followed patient populations.

Discussion

We show here that *SNCA* transcript counts in circulating blood cells are reduced in early Parkinson's disease—even in the landmark PPMI study that enrolled patients before they met standard clinical diagnostic criteria. The association between *SNCA* expression and Parkinson's disease was generalizable across platforms and across populations (Table 2 and Figs 1–3). It was observed in studies from a single health care system (HBS), from 22 USA sites (PROBE), and from 24 USA and international sites (PPMI). This association remained significant when adjusted for confounding variables such as age, gender, white and red blood cell counts, Parkinson's disease medications, or sample handling.

Biological markers of processes that lead to complex diseases such as Parkinson's disease are not simply present or absent, but have a wide range of values that overlap in subjects with a clinical disease and in those without it (Manolio, 2003). For example, total cholesterol has modest specificity and sensitivity for predicting cardiovascular disease (with an area under the receiver operating

curve of 0.59 in the Women's Health Study; Ridker et al., 2000)—well short of the >90% specificity required for a *bona fide* diagnostic test. Nonetheless, biological markers of complex diseases can have powerful clinical applications. Cholesterol metabolism-related biomarkers have transformed cardiology. They enable early intervention with cholesterol-lowering medications because of the favourable risk/benefit ratio (low side-effect profile combined with clear benefits for reducing debilitating cardiovascular events). Levels of these commonly used biomarkers are also valuable in tracking therapeutic response and patient stratification. Thus, if confirmed in prospective studies, *SNCA* blood transcripts could be developed into future markers of risk, of patient stratification for clinical trials, and of response to novel therapeutics (particularly those targeting α -synuclein production or clearance).

We have begun to explore a clinical application for this biological marker as a predictor of cognitive decline. Cognitive decline is one of the most debilitating manifestations of disease progression in Parkinson's and is closely correlated with the presence of α -synuclein protein-positive Lewy body burden throughout the neocortex (Braak et al., 2002). Predictors of cognitive decline would be valuable for stratifying clinical trials and for advancing clinical care (Chen-Plotkin, 2014). Several previous studies have examined the association between cognitive decline and biological markers in cross-sectional studies (Mollenhauer et al.,

2014) or with a single longitudinal follow-up after 1–2 years (Siderowf *et al.*, 2010; Chen-Plotkin *et al.*, 2011). Longitudinal studies with repeated measurements provide more independent information than a single measurement (Hedeker and Gibbons, 2006; Locascio and Atri, 2011). In our study low SNCA transcript abundance at enrolment predicted cognitive decline in patients with Parkinson's disease that were cognitively reassessed during multiple longitudinal visits over up to 5 years of follow-up with a *P*-value of 0.005 in the combined analysis of HBS and PROBE studies.

Beyond these translational implications, the association here observed offers an immediate clue towards unravelling the biological function of α -synuclein in health and disease. α -Synuclein had been thought of as a neuron-specific gene for decades, but this view was turned on its head with the discovery of high levels of α -synuclein present in blood cells, particularly in reticulocytes (circulating immature red blood cells) and during terminal erythroid differentiation and in megakaryocytes (Scherzer *et al.*, 2008; Maitta *et al.*, 2011), with low or no expression in other tissues (Maroteaux *et al.*, 1988). In erythroid cells SNCA expression is regulated by the GATA-1 transcription factor in synchrony with machinery essential for incorporating iron into heme within mitochondria (Scherzer *et al.*, 2008) thereby pointing at a biological link between three nodes of Parkinson's disease: α -synuclein, mitochondria (Zheng *et al.*, 2010), and iron (Rhodes and Ritz, 2008). Our study suggests that this link might be perturbed in early disease stages. Alternative α -synuclein transcript isoform usage has been suggested as a potential convergent mechanism in Parkinson's disease pathology (Rhinn *et al.*, 2012). 3'UTR ends of mRNAs are critical for targeting transcripts to specific subcellular compartments and for translational control (Andreassi and Riccio, 2009). Specific transcript isoforms of SNCA with an extended 3'UTR impact accumulation of α -synuclein protein, which appears redirected away from synaptic terminals and towards mitochondria (Rhinn *et al.*, 2012). This might be relevant to disease processes as mitochondrial dysfunction is implicated in the onset of Parkinson's disease neuropathology (Zheng *et al.*, 2010). By contrast, deletion of exon 5 of the SNCA gene results in the translation of a truncated 112 amino acid protein that is prone to aggregation and increases seeding efficiency (Murray *et al.*, 2003).

In sporadic Parkinson's disease, α -synuclein protein (and iron) accumulates in the brain and peripherally α -synuclein aggregates are found in skin, glands, intestines and epicardium (reviewed in Jellinger, 2011). In a few dozen patients a triplication of the SNCA locus causes monogenic familial Parkinson's disease. The direction of the change observed here was therefore surprising, even paradoxical, at first glance. How can it be that SNCA blood transcript levels are consistently reduced in each of the three studies? Ample precedent exists for paradoxical changes in peripheral markers of neurological diseases. In Parkinson's disease, extracellular α -synuclein protein levels are reduced in CSF for

reasons unknown (Mollenhauer *et al.*, 2011). In Wilson's disease, copper accumulates in the basal ganglia, but serum levels are paradoxically low. In Alzheimer's disease, amyloid- β_{42} levels are reduced in CSF (Hansson *et al.*, 2006). We hypothesize that intracellular SNCA transcript levels in clinically manifest, sporadic Parkinson's disease may be reduced as a consequence of intracellular α -synuclein protein accumulation (due to impaired clearance or other mechanisms) that leads to a feedback repression of transcription—in a homeostatic effort to ameliorate the 'log jam' of excessive α -synuclein protein products. In an alternate view, because SNCA is so highly expressed in reticulocytes, reduced SNCA expression could be a marker of generally abnormal reticulocyte biogenesis or function in early-stage Parkinson's disease. We performed association studies that do not imply causality and that were not designed to distinguish between these alternative hypotheses. However, neither ours nor previous studies (Kasten *et al.*, 2010) found evidence for an association between anaemia and Parkinson's disease. By contrast, anaemia during early life, preceding disease onset by up to 20–29 years (Savica *et al.*, 2009), and multiple recent blood donations have been correlated to increased risk of Parkinson's disease (Logroscino *et al.*, 2006).

The ease and non-invasiveness of an α -synuclein blood test is appealing. Previous attempts to search for α -synuclein blood tests centred on α -synuclein plasma protein, involved small cohorts, and were inconclusive (for review see Chahine *et al.*, 2014). That approach poses considerable technical challenges as plasma can be easily contaminated with α -synuclein released from erythrocytes ruptured during sample transport and centrifugation. Our studies leapfrogged this technical challenge. Patient blood was directly collected into an FDA-approved system for preserving 'intracellular RNA from whole blood for RT-PCR used in molecular diagnostic testing' (without a need for plasma separation) where chemicals immediately stabilize the *in vivo* gene transcription profile by reducing *in vitro* RNA degradation and minimizing gene induction (Pahl and Brune, 2002a, b; Rainen *et al.*, 2002; Kim *et al.*, 2014). The HBS, PROBE and PPMI studies were purposefully designed to minimize the threats of bias from sample collection, processing and storage (Ransohoff, 2004). Case and control samples for HBS and PPMI were collected, processed and analysed in parallel using standardized procedures for each step. In contrast to other studies, RNA integrity, complete blood counts, and several processing and quality indicators were carefully measured for all three studies.

Although each of the three biomarker studies here presented has limitations when taken on its own, these individual constraints are compensated for by the collective strengths. For example, while the PROBE and HBS study could not completely rule out an undue influence of Parkinson's disease medications on SNCA expression, this question was fully addressed by exclusively recruiting *de novo* cases in PPMI. We clearly show that this is not simply a marker of sick patients; instead, the association

is already detectable in individuals who are very early in the disease course and highly functional. Clinical diagnostic certainty was enhanced through DAT neuroimaging.

Beyond revealing an association between lowered *SNCA* expression and Parkinson's disease, this study presents a generally useful resource of three purpose-designed biobanks and linked clinical data. This platform of biobanks can now be used for accelerating the exploration of nearly any biofluid marker for Parkinson's disease by decreasing the time from discovery to replication from multiple years down to the assay run-time.

Acknowledgements

We thank all study participants, their families, and friends for their support and participation, and our study coordinators.

Harvard Biomarker Study. Co-Directors: Harvard NeuroDiscovery Center: Clemens R. Scherzer, Bradley T. Hyman, Adrian J. Iverson; Investigators and Study Coordinators: Harvard NeuroDiscovery Center: Ana Trisini-Lipsanopoulos, Kaltra Dhima, Stephen Bayer, Kaitlin C. Lockhart; Brigham and Women's Hospital: Lewis R. Sudarsky, Michael T. Hayes, Reisa Sperling; Massachusetts General Hospital: John H. Growdon, Michael A. Schwarzschild, Albert Y. Hung, Alice W. Flaherty, Deborah Blacker, Anne-Marie Wills, U. Shivraj Sohur, Vivek K. Unni, Nite I. Mejia, Anand Viswanathan, Stephen N. Gomperts, Vikram Khurana, Mark W. Albers, Kyleen E. Swords, Rebecca K. Rudel; University of Ottawa: Michael G. Schlossmacher; Scientific Advisory Board: Massachusetts General Hospital: John H. Growdon, Brigham and Women's Hospital: Dennis J. Selkoe, Reisa Sperling; Harvard School of Public Health: Alberto Ascherio; Data Coordination: Harvard NeuroDiscovery Center: Thomas Yi, Massachusetts General Hospital: Joseph J. Locascio; Biobank Management Staff: Harvard NeuroDiscovery Center: Karen Duong, Zhixiang Liao, Ashley N. Hoising.

PROBE Study. PROBE Steering Committee: Voyager Therapeutics: Bernard Ravina; Brigham and Women's Hospital: Clemens Scherzer, University of Ottawa: Michael Schlossmacher, Avid Radiopharmaceuticals: Andrew Siderowf, University of Rochester: David Oakes; Institute for Neurodegenerative Disorders: Kenneth Marek; Georgetown University: Ira Shoulson. The investigators who contributed to the PROBE cohort and collection of clinical data and blood samples are listed in the Appendix of Ravina et al., *A Longitudinal Program for Biomarker Development in Parkinson's Disease: A Feasibility Study, Movement Disorders*, Vol. 24, No. 14, 2009, pp. 2081–2090.

PPMI Study. The PPMI data used in this study were obtained from the Parkinson's Progression Markers Initiative (PPMI) database (www.ppmi-info.org/data). For up-to-date information on the study, visit www.ppmi-info.org.

Funding

This study was made possible by NIH grants U01 NS082157 (C.R.S.), U01 NS082080 (C.R.S.) the Harvard NeuroDiscovery Center (to A.I., C.R.S., and B.T.H.), U.S. Department of Defense grants W81XWH-1-0007 (B.R.) and W81XWH-13-1-0115 (C.R.S.), and the Michael J. Fox Foundation (C.R.S. and M.G.S.). Additional funding was provided by the M.E.M.O. Hoffman Foundation (C.R.S.) and U.S. Department of Defense grant W81XWH-11-1-0150 (M.A.S.).

The Harvard Biomarker Study is supported by the Harvard NeuroDiscovery Center (HNDC), the Parkinson's Disease Biomarkers Program (PDBP) grant U01 NS082157 of the NINDS, and the Massachusetts Alzheimer's Disease Research Center (ADRC) P50 AG005134 grant of the National Institute on Aging.

The PROBE cohort was funded by Department of Defense Neurotoxin Exposure Treatment Parkinson's Research Program (W23RRYX7022N606), NINDS PD-DOC (NS050095), the Parkinson's Disease Foundation (New York, NY), with initial funding from Cephalon Inc and Lundbeck for the parent PRECEPT clinical trial and follow-up PostCEPT cohort, from which the nested PROBE cohort was derived.

PPMI, a public-private partnership, is funded by the Michael J. Fox Foundation for Parkinson's Research and funding partners, including Abbott, Avid, Biogen Idec, Bristol-Myers Squibb, Covance, Elan, GE Healthcare, Genentech, GlaxoSmithKline, Lilly, Merck, Meso Scale Discovery, Pfizer, Roche, UCB.

The funders had no role in design or conduct of the study, collection, management, analysis and interpretation of the data, and preparation, review, or approval of the manuscript. The PPMI Data and Publications Review Committee reviewed the manuscript.

Supplementary material

Supplementary material is available at *Brain* online.

References

- Andreassi C, Riccio A. To localize or not to localize: mRNA fate is in 3'UTR ends. *Trends Cell Biol* 2009; 19: 465–74.
- Auer H, Lyianarachchi S, Newsom D, Klisovic MI, Marcucci G, Kornacker K. Chipping away at the chip bias: RNA degradation in microarray analysis. *Nat Genet* 2003; 35: 292–3.
- Braak H, Del Tredici K, Bratzke H, Hamm-Clement J, Sandmann-Keil D, Rub U. Staging of the intracerebral inclusion body pathology associated with idiopathic Parkinson's disease (preclinical and clinical stages). *J Neurol* 2002; 249 (Suppl 3): III/1–5.
- Chahine LM, Stern MB, Chen-Plotkin A. Blood-based biomarkers for Parkinson's disease. *Parkinsonism Relat Disord* 2014; 20 (Suppl 1): S99–103.
- Chen-Plotkin AS. Unbiased approaches to biomarker discovery in neurodegenerative diseases. *Neuron* 2014; 84: 594–607.

- Chen-Plotkin AS, Hu WT, Siderowf A, Weintraub D, Goldmann Gross R, Hurtig HI, et al. Plasma epidermal growth factor levels predict cognitive decline in Parkinson disease. *Ann Neurol* 2011; 69: 655–63.
- Chin L, Andersen JN, Futreal PA. Cancer genomics: from discovery science to personalized medicine. *Nat Med* 2011; 17: 297–303.
- Dorsey ER, Constantinescu R, Thompson JP, Biglan KM, Holloway RG, Kieburtz K, et al. Projected number of people with Parkinson disease in the most populous nations, 2005 through 2030. *Neurology* 2007; 68: 384–6.
- Forman MS, Lee VM, Trojanowski JQ. Nosology of Parkinson's disease: looking for the way out of a quagmire. *Neuron* 2005; 47: 479–82.
- Geiss GK, Bumgarner RE, Birditt B, Dahl T, Dowidar N, Dunaway DL, et al. Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol* 2008; 26: 317–25.
- Hansson O, Zetterberg H, Buchhave P, Londos E, Blennow K, Minthon L. Association between CSF biomarkers and incipient Alzheimer's disease in patients with mild cognitive impairment: a follow-up study. *Lancet Neurol* 2006; 5: 228–34.
- Hayashi K, Masuda S, Kimura H. Impact of biomarker usage on oncology drug development. *J Clin Pharm Ther* 2013; 38: 62–7.
- Hedeker D, Gibbons RD. Longitudinal data analysis. Hoboken, NJ: Wiley; 2006.
- Hennecke G, Scherzer CR. RNA biomarkers of Parkinson's disease: developing tools for novel therapies. *Biomark Med* 2008; 2: 41–53.
- Hughes AJ, Daniel SE, Kilford L, Lees AJ. Accuracy of clinical diagnosis of idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* 1992; 55: 181–4.
- Imbeaud S, Graudens E, Boulanger V, Barlet X, Zaborski P, Eveno E, et al. Towards standardization of RNA quality assessment using user-independent classifiers of microcapillary electrophoresis traces. *Nucleic Acids Res* 2005; 33: e56.
- Jellinger KA. Synuclein deposition and non-motor symptoms in Parkinson disease. *J Neurol Sci* 2011; 310: 107–11.
- Kasten M, Tadic V, Klein C, Rocca WA, Savica R, Eric Ahlsgog J, et al. Anemia or low hemoglobin levels preceding Parkinson disease: a case-control study. *Neurology* 2010; 74: 1655; author reply 1656.
- Kim JH, Jin HO, Park JA, Chang YH, Hong YJ, Lee JK. Comparison of three different kits for extraction of high-quality RNA from frozen blood. *Springerplus* 2014; 3: 76.
- Locascio JJ, Atri A. An overview of longitudinal data analysis methods for neurological research. *Dement Geriatr Cogn Dis Extra* 2011; 1: 330–57.
- Logroscino G, Chen H, Wing A, Ascherio A. Blood donations, iron stores, and risk of Parkinson's disease. *Mov Disord* 2006; 21: 835–8.
- Maitta RW, Wolgast LR, Wang Q, Zhang H, Bhattacharyya P, Gong JZ, et al. Alpha- and beta-synucleins are new diagnostic tools for acute erythroid leukemia and acute megakaryoblastic leukemia. *Am J Hematol* 2011; 86: 230–4.
- Malkov VA, Serikawa KA, Balantac N, Watters J, Geiss G, Mashadi-Hosseini A, et al. Multiplexed measurements of gene signatures in different analytes using the Nanostring nCounter Assay System. *BMC Res Notes* 2009; 2: 80.
- Manolio T. Novel risk markers and clinical practice. *N Engl J Med* 2003; 349: 1587–9.
- Maroteaux L, Campanelli JT, Scheller RH. Synuclein: a neuron-specific protein localized to the nucleus and presynaptic nerve terminal. *J Neurosci* 1988; 8: 2804–15.
- Mollenhauer B, Locascio JJ, Schulz-Schaeffer W, Sixel-Doring F, Trenkwalder C, Schlossmacher MG. alpha-Synuclein and tau concentrations in cerebrospinal fluid of patients presenting with parkinsonism: a cohort study. *Lancet Neurol* 2011; 10: 230–40.
- Mollenhauer B, Rochester L, Chen-Plotkin A, Brooks D. What can biomarkers tell us about cognition in Parkinson's disease? *Mov Disord* 2014; 29: 622–33.
- Murray IV, Giasson BI, Quinn SM, Koppaka V, Axelsen PH, Ischiropoulos H, et al. Role of alpha-synuclein carboxy-terminus on fibril formation *in vitro*. *Biochemistry* 2003; 42: 8530–40.
- Nakai M, Fujita M, Waragai M, Sugama S, Wei J, Akatsu H, et al. Expression of alpha-synuclein, a presynaptic protein implicated in Parkinson's disease, in erythropoietic lineage. *Biochem Biophys Res Commun* 2007; 358: 104–10.
- Nalls MA, Pankratz N, Lill CM, Do CB, Hernandez DG, Saad M, et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet* 2014; 46: 989–93.
- O'Brien JA, Ward A, Michels SL, Tziveleakis S, Brandt NJ. Economic burden associated with Parkinson disease. *Drug Benefit Trends* 2009; 21: 179–90.
- Pahl A, Brune K. Gene expression changes in blood after phlebotomy: implications for gene expression profiling. *Blood* 2002a; 100: 1094–5.
- Pahl A, Brune K. Stabilization of gene expression profiles in blood after phlebotomy. *Clin Chem* 2002b; 48: 2251–3.
- Parkinson Progression Marker Initiative. The Parkinson Progression Marker Initiative (PPMI). *Prog Neurobiol* 2011; 95: 629–35.
- Rainen L, Oelmueller U, Jurgensen S, Wyrich R, Ballas C, Schram J, et al. Stabilization of mRNA expression in whole blood samples. *Clin Chem* 2002; 48: 1883–90.
- Ransohoff DF. Rules of evidence for cancer molecular-marker discovery and validation. *Nat Rev Cancer* 2004; 4: 309–14.
- Ransohoff DF. Bias as a threat to the validity of cancer molecular-marker research. *Nat Rev Cancer* 2005; 5: 142–9.
- Rhinn H, Qiang L, Yamashita T, Rhee D, Zolin A, Vanti W, et al. Alternative alpha-synuclein transcript usage as a convergent mechanism in Parkinson's disease pathology. *Nat Commun* 2012; 3: 1084.
- Rhodes SL, Ritz B. Genetics of iron regulation and the possible role of iron in Parkinson's disease. *Neurobiol Dis* 2008; 32: 183–95.
- Ridker PM, Hennekens CH, Buring JE, Rifai N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000; 342: 836–43.
- Savica R, Grossardt BR, Carlin JM, Icen M, Bower JH, Ahlsgog JE, et al. Anemia or low hemoglobin levels preceding Parkinson disease: a case-control study. *Neurology* 2009; 73: 1381–7.
- Scherzer CR, Feany MB. Yeast genetics targets lipids in Parkinson's disease. *Trends Genet* 2004; 20: 273–7.
- Scherzer CR, Grass JA, Liao Z, Pepivani I, Zheng B, Eklund AC, et al. GATA transcription factors directly regulate the Parkinson's disease-linked gene alpha-synuclein. *Proc Natl Acad Sci USA* 2008; 105: 10907–12.
- Scherzer CR, Jensen RV, Gullans SR. Simplifying complex neurodegenerative diseases by gene chip analysis. In: Freese A, Simeone FA, Leone P, Janson C, editors. *Principles of molecular neurosurgery*. Basel: Karger; 2004. p. 246–57.
- Shin JH, Ko HS, Kang H, Lee Y, Lee YI, Pletinkova O, et al. PARIS (ZNF746) repression of PGC-1alpha contributes to neurodegeneration in Parkinson's disease. *Cell* 2011; 144: 689–702.
- Siderowf A, Xie SX, Hurtig H, Weintraub D, Duda J, Chen-Plotkin A, et al. CSF amyloid [beta] 1-42 predicts cognitive decline in Parkinson disease. *Neurology* 2010; 75: 1055–61.
- Zheng B, Liao Z, Locascio JJ, Lesniak KA, Roderick SS, Watt ML, et al. PGC-1alpha, a potential therapeutic target for early intervention in Parkinson's disease. *Sci Transl Med* 2010; 2: 52ra73.