



Effects of Corrugated Temperature Sheets on Optical Propagation

Andreas Muschinski
NORTHWEST RESEARCH ASSOCIATES INC.

12/11/2015
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/ RTB1
Arlington, Virginia 22203
Air Force Materiel Command

REPORT DOCUMENTATION PAGE		Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Executive Services, Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.			
1. REPORT DATE (DD-MM-YYYY) 13-01-2016		2. REPORT TYPE Final Performance	
		3. DATES COVERED (From - To) 01-07-2012 to 30-06-2015	
4. TITLE AND SUBTITLE Effects of Corrugated Temperature Sheets on Optical Propagation Along Quasi-horizontal Paths in the Stably Stratified Atmosphere		5a. CONTRACT NUMBER FA9550-12-C-0070	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER 61102F	
6. AUTHOR(S) Andreas Muschinski		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NORTHWEST RESEARCH ASSOCIATES INC. 4118 148 AVE NE REDMOND, WA 98052 US		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AF Office of Scientific Research 875 N. Randolph St. Room 3112 Arlington, VA 22203		10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/AFOSR RTB1	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT A DISTRIBUTION UNLIMITED: PB Public Release			
13. SUPPLEMENTARY NOTES			
14. ABSTRACT Optical propagation through the clear atmosphere is affected by small-scale refractive-index fluctuations which are caused mainly by temperature fluctuations. In the stably stratified atmosphere, these temperature fluctuations are the result of a combination of (1) more or less homogeneous and isotropic turbulence and (2) non-turbulent, quasi-horizontal interfaces, or "sheets". Collocated in-situ and optical field measurements conducted in the atmospheric surface layer confirmed that angle-of-arrival fluctuations and irradiance fluctuations observed with large-aperture telescopes (36 cm aperture diameter) are consistent with theoretical predictions based on Taylor's frozen-turbulence hypothesis and the geometrical-optics approximation. Short-term (less than a few seconds) fluctuations are dominated by turbulence while longer-term fluctuations are dominated by horizontally extended sheets. Direct numerical simulations of isotropic turbulence showed very good agreement with the turbulence spectrum predicted by Hill's 1978 model. A theoretical model of corrugated sheets was developed and analyzed.			
15. SUBJECT TERMS Propagation through turbulence			

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Andreas Muschinski
Unclassified	Unclassified	Unclassified	UU		19b. TELEPHONE NUMBER (Include area code) 413-253-0129

1.

1. Report Type

Final Report

Primary Contact E-mail

Contact email if there is a problem with the report.

andreas@nwra.com

Primary Contact Phone Number

Contact phone number if there is a problem with the report

303-415-9701

Organization / Institution name

NorthWest Research Associates

Grant/Contract Title

The full title of the funded effort.

Effects of corrugated temperature sheets on optical propagation along quasi-horizontal paths in the stably stratified atmosphere

Grant/Contract Number

AFOSR assigned control number. It must begin with "FA9550" or "F49620" or "FA2386".

FA9550-12-C-0070

Principal Investigator Name

The full name of the principal investigator on the grant or contract.

Andreas Muschinski

Program Manager

The AFOSR Program Manager currently assigned to the award

Julie Moses

Reporting Period Start Date

07/01/2012

Reporting Period End Date

06/30/2015

Abstract

Optical propagation through the clear atmosphere is affected by small-scale refractive-index fluctuations which are caused mainly by temperature fluctuations. In the stably stratified atmosphere, these temperature fluctuations are the result of a combination of (1) more or less homogeneous and isotropic turbulence and (2) non-turbulent, quasi-horizontal interfaces, or "sheets". Collocated in-situ and optical field measurements conducted in the atmospheric surface layer confirmed that angle-of-arrival fluctuations and irradiance fluctuations observed with large-aperture telescopes (36 cm aperture diameter) are consistent with theoretical predictions based on Taylor's frozen-turbulence hypothesis and the geometrical-optics approximation. Short-term (less than a few seconds) fluctuations are dominated by turbulence while longer-term fluctuations are dominated by horizontally extended sheets. Direct numerical simulations of isotropic turbulence showed very good agreement with the turbulence spectrum predicted by Hill's 1978 model. A theoretical model of corrugated sheets was developed and analyzed.

Distribution Statement

This is block 12 on the SF298 form.

Distribution A - Approved for Public Release

Explanation for Distribution Statement

If this is not approved for public release, please provide a short explanation. E.g., contains proprietary information.

SF298 Form

Please attach your [SF298](#) form. A blank SF298 can be found [here](#). Please do not password protect or secure the PDF. The maximum file size for an SF298 is 50MB.

[sf 298 Muschinski AFOSR 20151207.pdf](#)

Upload the Report Document. File must be a PDF. Please do not password protect or secure the PDF . The maximum file size for the Report Document is 50MB.

[AFOSR_finalreport_20151207_AM.pdf](#)

Upload a Report Document, if any. The maximum file size for the Report Document is 50MB.

Archival Publications (published) during reporting period:

Tichkule, S. and A. Muschinski, 2012: Optical anemometry based on the temporal cross-correlation of angle-of-arrival fluctuations obtained from spatially separated light sources. Appl. Opt., 51, 5272-5282.

Tichkule, S. and A. Muschinski, 2014: Effects of wind-driven telescope vibrations on measurements of turbulent angle-of-arrival fluctuations. Appl. Opt., 53, 4651-4660.

Muschinski, A., 2015: Temperature variance dissipation equation and its relevance for optical turbulence modeling. J. Opt. Soc. Am. A, 32, 2195-2200.

Muschinski, A. and S. M. de Bruyn Kops, 2015: Investigation of Hill's optical turbulence model by means of direct numerical simulation. J. Opt. Soc. Am. A, 32, 2423-2430.

Changes in research objectives (if any):

Change in AFOSR Program Manager, if any:

Extensions granted or milestones slipped, if any:

AFOSR LRIR Number

LRIR Title

Reporting Period

Laboratory Task Manager

Program Officer

Research Objectives

Technical Summary

Funding Summary by Cost Category (by FY, \$K)

	Starting FY	FY+1	FY+2
Salary			
Equipment/Facilities			
Supplies			
Total			

Report Document

Report Document - Text Analysis

Report Document - Text Analysis

Appendix Documents

2. Thank You

E-mail user

Dec 07, 2015 22:05:06 Success: Email Sent to: andreas@nwra.com

Effects of corrugated temperature sheets on optical propagation along quasi-horizontal paths in the stably stratified atmosphere

Final Report, Contract #FA9550-12-C-0070

AFOSR, *Remote Sensing and Imaging Physics* program

by **Andreas Muschinski**

(December, 2015)

Project Summary

Optical propagation through the clear atmosphere is affected by small-scale refractive-index fluctuations which are caused mainly by temperature fluctuations. In the stably stratified atmosphere, these temperature fluctuations are the result of a combination of (1) more or less homogeneous and isotropic turbulence and (2) non-turbulent, quasi-horizontal interfaces, or “sheets”. Collocated in-situ and optical field measurements conducted in the atmospheric surface layer confirmed that angle-of-arrival fluctuations and irradiance fluctuations observed with large-aperture telescopes (36 cm aperture diameter) are consistent with theoretical predictions based on Taylor’s frozen-turbulence hypothesis and the geometrical-optics approximation. Short-term (less than a few seconds) fluctuations are dominated by turbulence while longer-term fluctuations are dominated by horizontally extended sheets. Direct numerical simulations of isotropic turbulence showed very good agreement with the turbulence spectrum predicted by Hill’s 1978 model. A theoretical model of corrugated sheets was developed and analyzed.

Contact:

Dr. Andreas Muschinski

NorthWest Research Associates, Inc. (NWRA), Boulder Office

3380 Mitchell Lane

Boulder, CO 80301

Email andreas@nwra.com

Tel. (303) 415-9701 ext. 228

1 Background

The modern physics of optical propagation through the turbulent atmosphere, pioneered by Tatarskii and coworkers in the 1950s and 1960s (Tatarskii, 1961, 1971), continues to be the conceptual basis for progress in various science and engineering disciplines, such as remote sensing of the optically clear atmosphere, astronomy, free-space optical communication, directed-energy technology, and terrestrial and extraterrestrial imaging and surveillance (e.g., Ishimaru, 1978; Strohbehn, 1978; Rytov et al., 1989; Roggemann and Welsh, 1996; Wheelon, 2001, 2003; Andrews and Phillips, 2005; Schmidt, 2010; Sasiela, 2007; Korotkova, 2014).

During the last decade or so, effects of anisotropic turbulence on optical propagation have become an important research focus within the optical-propagation community. So-called “non-Kolmogorov turbulence” models (e.g., Toselli et al., 2008; Toselli and Korotkova, 2015) have been introduced that deviate from the classical $-11/3$ power law predicted by the Obukhov-Corrsin theory of scalar turbulence. One goal of this project has been to develop, as an alternative to the “non-Kolmogorov turbulence” models, a statistical model of non-turbulent, non-overturning, corrugated interfaces, or “sheets” that have long been known to be ubiquitous in the stably stratified atmosphere (e.g., Doviak and Zrnić, 1984; Muschinski, 2004; Dalaudier et al., 1994; Muschinski and Wode, 1998).

2 Project Outcomes

In the following, we summarize the main project outcomes.

2.1 Observations

2.1.1 Optical measurements of cross-wind velocity

The temporal cross-correlation function of the angle-of-arrival (AOA) fluctuations of two optical waves propagating through atmospheric turbulence carries information regarding the average wind velocity transverse to the propagation path. In Tichkule and Muschinski (2012), we presented and discussed two estimators for the retrieval of the path-averaged, beam-transverse, horizontal wind velocity, v_t . Both methods retrieve v_t from the temporal cross-correlation function of AOA fluctuations obtained from two closely spaced, light-emitting diodes (LEDs). The first method relies on the time delay of the peak (TDP) of the cross-correlation function, and the second method exploits its slope at zero lag (SZL). Over a 9 h period during which v_t varied between -1.3 m s^{-1} and 2.0 m s^{-1} , the maximum rms difference between optically retrieved and in-situ measured, 10-s estimates of v_t was found to be 0.18 m s^{-1} for the TDP estimator and 0.23 m s^{-1} for the SZL estimator.

2.1.2 Wind-induced telescope vibrations

Turbulence in the atmospheric refractive-index field causes optical angle-of-arrival (AOA) fluctuations that can be used for atmospheric remote sensing of various parameters, including wind velocities and the optical refractive-index turbulence structure parameter, C_n^2 . If AOA measurements are contaminated by wind-induced telescope vibrations, the underlying retrieval algorithms

may fail. In order to study the effects of wind-driven telescope vibrations on optical-turbulence measurements, we conducted a field experiment in which we exposed two small telescopes deliberately to the wind (Tichkule and Muschinski, 2014). We measured AOA fluctuations of visible light propagating along a horizontal, 174 m long path 1.7 m above flat terrain, and we used fast-response ultrasonic anemometers to measure the wind velocity at multiple locations along the path. We found (1) that the AOA turbulence spectra were contaminated by multiple resonance peaks, (2) that the resonance frequencies were independent of the wind speed, and (3) that the AOA variance associated with the dominating vibration mode was proportional to the fourth power of the wind speed.

2.2 Theory

2.2.1 Homogeneous and isotropic turbulence

The three-dimensional (3D) spectrum $\Phi(\boldsymbol{\kappa})$ of the turbulent air temperature fluctuations is a key quantity for the physics of optical propagation through the turbulent atmosphere. The standard model, which was derived in the 1950s by Tatarskii from the Obukhov-Corrsin theory of homogeneous and isotropic turbulence, is $\Phi(\boldsymbol{\kappa}) = 0.033 C_T^2 \kappa^{-11/3} h(\kappa l_0)$, where $\kappa = |\boldsymbol{\kappa}|$ is the wave number, C_T^2 is the temperature structure parameter, l_0 is the inner temperature scale, and $h(\kappa l_0)$ is a universal function that approaches 1 for wave numbers in the inertial range and drops to zero for $\kappa l_0 \gg 1$. Certain performance characteristics of optical systems, such as the scintillation index for small receiving apertures, depend sensitively on the functional form of $h(y)$ at $y \approx 1$. During the last 70 years, the optical-turbulence community has developed and applied various heuristic $h(y)$ models. There is a constraint that any valid $h(y)$ model has to fulfill: $\int_0^\infty h(y) y^{1/3} dy = (27/10)\Gamma(1/3) = 7.233$. This constraint is a dimensionless form of the spectral temperature variance dissipation equation, which follows directly from first-principle fluid mechanics. In Muschinski (2015), we showed that Tatarskii’s cut-off (Tatarskii, 1961) and Gaussian (Tatarskii, 1971) models fulfill this constraint while three more recent models, including the widely used Andrews model (Andrews, 1992), do not. The dissipation constraint can be used to “re-calibrate” the coefficients in these models.

2.2.2 Statistical model of corrugated temperature sheets

It is a standard assumption in the theory of optical propagation through the turbulent atmosphere that the refractive-index fluctuations $n_1(\mathbf{x})$ are statistically isotropic. It is well known, however, that $n_1(\mathbf{x})$ in the free atmosphere and in the nocturnal boundary layer is often strongly anisotropic, even at the smallest scales. We introduced and analyzed (Muschinski, 2016) a model atmosphere characterized by corrugated but non-overturning refractive-index interfaces, or “sheets,” such that $n_1(\mathbf{x}) = v[z - h(x, y)]$, where $v(z)$ is a random function with a 1D spectrum $V(\kappa_z)$ that describes the vertical microstructure, and $h(x, y)$ is a random function of the horizontal coordinates x and y , which characterizes the local vertical displacement of the vertical microstructure. On the basis of some natural simplifying assumptions, we derived a closed-form expression for the 3D spectrum $\Phi(\boldsymbol{\kappa})$. This 3D spectrum differs from commonly used spectra in that the 3D spectral density decreases horizontally in the form of a Gaussian function, not as a power law. We show that if the vertical 1D spectrum follows a power law, then the horizontal 1D spectra follow the same power law but at a lower spectral level.

2.3 Computer simulations

For almost four decades, Hill’s “Model 4” (Hill, 1978) has played a central role in research and technology of optical turbulence. Based on Batchelor’s generalized Obukhov-Corrsin theory of scalar turbulence, Hill’s model predicts the dimensionless function $h(\kappa l_0, \text{Pr})$ that appears in Tatarskii’s well-known equation for the three-dimensional refractive-index spectrum in the case of homogeneous and isotropic turbulence, $\Phi_n(\kappa) = 0.033 C_n^2 \kappa^{-11/3} h(\kappa l_0, \text{Pr})$. In Muschinski and de Bruyn Kops (2015), we investigated Hill’s model by comparing numerical solutions of Hill’s differential equation with scalar spectra estimated from direct numerical simulation (DNS) output data. Our DNS solved the Navier-Stokes equation for the three-dimensional velocity field and the scalar transport equation for the scalar field on a numerical grid containing 4096^3 grid points. Two independent DNS runs were analyzed, one with the Prandtl number $\text{Pr} = 0.7$ and a second run with $\text{Pr} = 1.0$. We found very good agreement between $h(\kappa l_0, \text{Pr})$ estimated from the DNS output data and $h(\kappa l_0, \text{Pr})$ predicted by the Hill model. We found that the height of the Hill bump is $1.79 \text{Pr}^{1/3}$, implying that there is no bump if $\text{Pr} < 0.17$. Both the DNS and the Hill model predict that the viscous-diffusive “tail” of $h(\kappa l_0, \text{Pr})$ is exponential, not Gaussian.

References Cited

- Andrews, L. C., 1992: An analytical model for the refractive index power spectrum and its application to optical scintillations in the atmosphere. *J. Mod. Opt.*, **39**, 1849–1853.
- Andrews, L. C. and R. L. Phillips, 2005: *Laser beam propagation through random media*. 2d ed., SPIE Press, Bellingham, Washington, 783 pp.
- Dalaudier, F., C. Sidi, M. Crochet, and J. Vernin, 1994: Direct evidence of “sheets” in the atmospheric temperature field. *J. Atmos. Sci.*, **51**, 237–248.
- Doviak, R. J. and D. S. Zrnić, 1984: Reflection and scatter formula for anisotropically turbulent air. *Radio Sci.*, **19**, 325–336.
- Hill, R. J., 1978: Models of the scalar spectrum for turbulent advection. *J. Fluid Mech.*, **88**, 541–562.
- Ishimaru, A., 1978: *Wave propagation and scattering in random media, Vol. 2, Scattering in random media*. Academic Press, Boston, Massachusetts, 251–572 pp.
- Korotkova, O., 2014: *Random light beams: theory and applications*. CRC Press, Boca Raton, 361 pp.
- Muschinski, A., 2004: Local and global sampling of clear-air Doppler radar signals. *Radio Sci.*, **39**, RS1008.
- Muschinski, A., 2015: Temperature variance dissipation equation and its relevance for optical turbulence modeling. *J. Opt. Soc. Am. A*, **32**, 2195–2200.
- Muschinski, A., 2016: Optical propagation through non-overturning, corrugated temperature sheets in the atmosphere. *J. Opt. Soc. Am. A*, **33**, in preparation for journal submission.
- Muschinski, A. and S. M. de Bruyn Kops, 2015: Investigation of Hill’s optical turbulence model by means of direct numerical simulation. *J. Opt. Soc. Am. A*, **32**, 2423–2430.

- Muschinski, A. and C. Wode, 1998: First in situ evidence for coexisting submeter temperature and humidity sheets in the lower free troposphere. *J. Atmos. Sci.*, **55**, 2893–2906.
- Roggemann, M. C. and B. M. Welsh, 1996: *Imaging through turbulence*. CRC Press, Boca Raton, Florida, 320 pp.
- Rytov, S. M., Y. A. Kravtsov, and V. I. Tatarskii, 1989: *Principles of statistical radio physics – 4. Wave propagation through random media*. Springer, Berlin, Germany, 187 pp.
- Sasiela, R. J., 2007: *Electromagnetic wave propagation in turbulence*. 2d ed., SPIE Press, Bellingham, Washington, USA, 367 pp.
- Schmidt, J. D., 2010: *Numerical simulation of optical wave propagation*. SPIE Press, Bellingham, Washington, 196 pp.
- Strohbehn, J. W., (Ed.) , 1978: *Laser beam propagation in the atmosphere*, Topics in Applied Physics, Vol. 25. Springer, Berlin, 325 pp.
- Tatarskii, V. I., 1961: *Wave propagation in a turbulent medium*. McGraw-Hill, New York, 285 pp.
- Tatarskii, V. I., 1971: *The effects of the turbulent atmosphere on wave propagation*. Israel Program for Scientific Translation, Jerusalem, Israel, 472 pp.
- Tichkule, S. and A. Muschinski, 2012: Optical anemometry based on the temporal cross-correlation of angle-of-arrival fluctuations obtained from spatially separated light sources. *Appl. Opt.*, **51**, 5272–5282.
- Tichkule, S. and A. Muschinski, 2014: Effects of wind-driven telescope vibrations on measurements of turbulent angle-of-arrival fluctuations. *Appl. Opt.*, **53**, 4651–4660.
- Toselli, I., L. C. Andrews, R. L. Phillips, and V. Ferrero, 2008: Free-space optical system performance for laser beam propagation through non-Kolmogorov turbulence. *Opt. Eng.*, **47**, 026 003.
- Toselli, I. and O. Korotkova, 2015: General scale-dependent anisotropic turbulence and its impact on free space optical communication system performance. *J. Opt. Soc. Am. A*, **32**, 1017–1025.
- Wheelon, A. D., 2001: *Electromagnetic scintillation — I. Geometrical optics*. Cambridge University Press, Cambridge, United Kingdom, 455 pp.
- Wheelon, A. D., 2003: *Electromagnetic scintillation — II. Weak scattering*. Cambridge University Press, Cambridge, United Kingdom, 440 pp.