

Award Number: W81XWH-11-1-0715

TITLE: Early Diagnosis of Clear Cell Kidney Cancer via VHL/HIF Pathway-Regulated Circulating microRNA

PRINCIPAL INVESTIGATOR: Allan Pantuck, MD

CONTRACTING ORGANIZATION: University of California, Los Angeles
Los Angeles, CA 90095

REPORT DATE: May 2016

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to the Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE May 2016			2. REPORT TYPE Annual		3. DATES COVERED 1 Sep 2014 - 14 Apr 2016	
4. TITLE AND SUBTITLE Early Diagnosis of Clear Cell Kidney Cancer via VHL/HIF Pathway- Regulated Circulating microRNA					Sa. CONTRACT NUMBER	
					Sb. GRANT NUMBER W81XWH-11-1-0715	
					Sc. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Allan J. Pantuck, MD Muneesh Tewari, MD, PhD E-Mail: anantuck@mednet.ucla.edu					Sd. PROJECT NUMBER	
					Sf. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, Los Angeles Los Angeles, CA 90095					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The over-arching objective of this research project is to identify a panel of diagnostic miRNAs that are measurable in serum and will be able to identify kidney cancer in its earliest stages. We hypothesized that serum levels of miR-210 would be significantly elevated in patients with clear cell renal carcinoma and that a panel of serum miRNAs could be defined and optimized for use as a diagnostic tool in patients presenting with a renal mass. To achieve this objective, we proposed three Specific Aims to test our hypotheses. Institutional IRB approvals have been obtained (Specific Aim 1, task 1). Retrospectively collected samples have been transferred and analyzed (Specific Aim 1, task 3). Prospective sample collection has been completed (Specific Aim 3, task 1). Task 2 will be completed by Initiating PI (Tewari) during the remainder of this no-cost grant extension necessitated by his relocation to the University of Michigan.						
15. SUBJECT TERMS None Listed						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 13	19a. NAME OF RESPONSIBLE PERSON USAMRMC	
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)	

Table of Contents

Page

Introduction.....	1
Body.....	2 - 3
Key Research Accomplishments.....	4
Reportable Outcomes.....	4
Conclusion.....	4
Year Three Progress Report Addendum:.....	5
References.....	None
Appendices.....	None
Supporting Data.....	6 - 10

DOD Idea Award: Collaborative Option – Final Progress Report

Early Diagnosis of Clear Cell Kidney Cancer via VHL/HIF Pathway-Regulated Circulating microRNA

Award Number W81XWH-11-1-0715

Initiating PI: Muneesh Tewari, MD, PhD (University of Michigan)

Partnering PI: Allan Pantuck, MD (University of California, Los Angeles)

INTRODUCTION:

The over-arching objective of this research project is to identify a panel of diagnostic miRNAs that are measurable in serum and will be able to identify kidney cancer in its earliest stages. We hypothesized that serum levels of miR-210 would be significantly elevated in patients with clear cell renal carcinoma and that a panel of serum miRNAs could be defined and optimized for use as a diagnostic tool in patients presenting with a renal mass. To achieve this objective we proposed three Specific Aims to test our hypotheses. As previously reported, in Year 1 we focused on Aim 1 in which we evaluated the utility of miR-210 as a marker for clear cell renal carcinoma. In an initial sample set of retrospective samples we found significant differences in the level of miR-210 in clear cell renal cases as compared to control serum samples. We also made significant progress in establishing digital PCR for miRNA detection in this sample set. During the Year 2 reporting period, we utilized our sample set in global miRNA profiling studies as well as in confirmation studies of our previous miR-210 data using digital PCR. We finished analysis of our profiling data to identify the most likely candidates for a marker panel for use with prospectively collected samples. Since the last progress report, sample collection was completed for the validation studies proposed in Aim 3, which represents the final reportable outcome from the UCLA portion of the grant. The final sub-aim of the grant, *unselected prospectively collected blood specimens for miRNA panel cross-validation*, will be completed by the grant Initiating PI (Tewari) during his final period of no-cost extension and will be reported in his final report.

BODY:

Specific Aim 1: Evaluate the HIF-induced miRNA, miR-210, as a circulating marker for clear cell renal carcinoma detection in a retrospective cohort of patients presenting with a renal mass.

Tasks 1-6: Completed in Year 1

Task 7: Digital PCR analysis of miR-210 and control miRNAs in retrospectively collected specimens. (Tewari/FHCRC; months 15-18)

AND

Task 8: Data analysis to determine performance of miR-210 in retrospective samples and to compare results of qRT-PCR vs. Digital PCR for measurement of circulating miRNA. (Tewari and McIntosh/FHCRC; months 18-21)

As described in our previous report we extracted RNA from a retrospectively collected specimen set consisting of serum samples from 20 non-RCC controls and 42 RCC cases. Using

qRT-PCR we determined the level of miR-210 in these samples. As previously reported, 14 of our case samples proved to be unusable. Consequently, we proceeded to analyze miR-210 levels in the 20 controls and 28 cases. We reported a significant difference in the levels of miR-210 between cases and controls, with case samples having higher levels of miR-210 (**Figure 1**, left panel). During the current reporting period we utilized these same RNAs to examine the level of miR-210 using digital PCR. **Figure 1** presents the digital PCR data (right panel) as compared to the previously reported qRT-PCR data (left panel). As shown, we confirmed by digital PCR a significant difference in the level of miR-210 in case samples relative to controls, with cases having on average about 2.5-fold greater serum miR-210 than controls. Receiver Operating Curve (ROC) analysis demonstrated the ability of serum miR-210 to distinguish cases from controls and the AUC was in fact slightly better with digital PCR than with standard qRT-PCR. We have further subdivided the data so that clear cell histology cases are analyzed separately from non-clear cell cases, and compared separately to controls. qRT-PCR data is presented on the left while digital PCR data is presented on the right in **Figure 2**. We find the greatest difference in miR-210 levels to be between clear cell and control samples, although miR-210 levels are also higher overall even in non-clear cell kidney cancer patients compared with controls, suggesting that this marker could be useful for more comprehensive detection of kidney cancer than had been originally hypothesized. This conclusion at this point is tempered by the limited number of non-clear cell samples studied (i.e., n=5, as compared to 23 clear cell cases).

Specific Aim 2: Discover circulating miRNA markers that are complementary to miR-210 for clear cell renal carcinoma detection, for assembly into a biomarker panel.

Task 1: Global profiling of miRNAs in a subset of retrospective blood specimens, described in Aim 1, using microfluidic Taqman miRNA qRT-PCR array platform. (Tewari/FHCRC; months 12-18)

AND

Task 2: Data analysis to identify the best performing miRNA biomarkers from Task 1, assessed both individually and in combination with miR-210. (Tewari & McIntosh/FHCRC; months 19-21)

For this Task we used miRNA profiling qRT-PCR arrays (Exiqon, Inc.) to globally profile miRNA expression in our sample sets with the aim of identifying additional circulating miRNAs that may serve as diagnostic markers for clear cell renal carcinoma. Using the same RNA as in Aim 1, a universal reverse transcription reaction for use with the Exiqon miRNA array panels was performed for each RNA sample. Expression of 378 human miRNAs was profiled on these arrays by qRT-PCR using locked nucleic acid-based primers to achieve high technical sensitivity and specificity. Of these, 19 miRNAs were undetected in all samples. Quality control of the array data was achieved using a number of spike-in and inter-plate calibrators provided with the arrays. As demonstrated in **Figure 3**, with the exception of a few UniSP6 outliers, the controls performed well and we did not observe any significant batch or edge effects which could negatively affect the quality of the data. We then performed differential miRNA expression analysis and the results of top differentially abundant serum miRNAs between cases and controls are shown in **Figure 4**. This data has been normalized using the median UniSP6 value and undetermined samples have been set to a Ct of 40. Depicted in **Figure 4A** are the top 5 miRNAs overexpressed in kidney cancer cases relative to control samples, chosen on the basis of greatest fold difference in expression (i.e., greatest difference in Ct values of cases vs. control). These miRNAs, all of which show a greater than 4-fold elevation in cases include miR-133a, -193b, -452, -1 and -365. In **Figure 4B** the top 5 overexpressed miRNAs as determined by t-test p-value are displayed. These include three of the miRNAs with the greatest

fold difference (miR-133a, -452 and -1) from **Figure 4A**, as well as two additional miRNAs (miR-143 and -218). We also analyzed the profiling data with respect to miR-210 (**Figure 5**). While our results do suggest a difference between cases and controls, the differences are not as striking as seen using the same samples in individual Taqman qPCR and digital PCR assays (Aim 1). This finding is not completely unexpected, as we frequently find individual Taqman assays utilizing assay specific reverse transcription primers to show superior performance than assays performed in a bulk reaction using a universal primer, as is required for miRNA profiling. The analysis of prospectively collected samples in Aim 3 will ultimately provide the best validation of serum miR-210 and the seven additional markers identified to date.

Task 3: Confirmation of up to 10 top-ranking miRNAs identified in Task 2. (Tewari/FHCRC; months 22-26)

At this point we are planning to use the validation set in Aim 3 for both confirmation and validation, as it will provide the most rigorous confirmation of the candidate biomarker panel that is emerging from Aim 2.

Specific Aim 3: Prospectively validate a circulating miRNA panel for the pre-operative diagnosis of clear cell renal carcinoma in patients with renal mass.

Task 1: Unselected, prospective blood collection from 150 patients presenting with renal mass (IRB approval and Material Transfer Agreement to be obtained under Aim 1, Task 1 above). (Pantuck/UCLA; months 7-29)

Prospective collection of serum samples has been completed at UCLA by Dr. Pantuck at UCLA.

Task 2: Transfer of unselected prospectively collected blood specimens for miRNA panel cross-validation. (Pantuck/UCLA; months 7-29)

We are ready to ship the sample collection and perform cross-validation of our miRNA marker panel as planned. We plan to transfer prospectively collected samples from UCLA to University of Michigan once Dr. Tewari is ready to complete task 2. In order to avoid confounding batch effects that we have found in our experience can occur if specimens are not analyzed together.

KEY RESEARCH ACCOMPLISHMENTS:

- We have confirmed and validated our previous qRT-PCR data via digital PCR, establishing that clear cell renal carcinoma serum samples contain elevated levels of miR-210.
- We have profiled the global expression level of 378 human miRNAs in our sample set and identified 7 miRNAs, in addition to miR-210, that are substantially elevated in our case samples relative to control samples.
- We have completed our prospective blood collection (by Dr. Pantuck at UCLA) for biomarker panel validation.

REPORTABLE OUTCOMES:

- Manuscripts/Abstracts/Presentations: We recently published the following manuscript: Hindson CM, Chevillet JR, Briggs H, Gallichotte E, Ruf IK, Hindson BJ, Vessella RL, Tewari M. Absolute quantification of low abundance targets by droplet digital PCR versus analog real-time PCR. *Nature Methods*, doi: 10.1038/nmeth.2633, e-published ahead of print, 2013.
- Patents and licenses applied for and/or issued: None in the current reporting period
- Development of cell lines, tissue or serum repositories: None in the current reporting period
- Informatics such as databases and animal models, etc.: None in the current reporting period
- Funding applied for based on work supported by this award: None in the current reporting period
- Employment or research opportunities applied for and/or received based on experience/training supported by this award: None in the current reporting period

CONCLUSION:

In summary, our efforts to date have been focused on evaluating the utility of circulating miR-210 as a diagnostic marker for clear cell renal carcinoma and on identifying additional differentially expressed miRNAs which may be assembled into a diagnostic screening panel. These efforts were initiated on serum from a retrospective cohort of patients that included 42 RCC patients, of which 23 were of the clear cell histology, and 20 control individuals. Analysis of miR-210 levels in this sample set revealed a significantly higher level of miR-210 in serum from the RCC sample set as compared to the control set. These results, generated using standard qRT-PCR, were further confirmed using digital PCR. The most significant difference in miR-210 levels we obtained occurred in the comparison of clear cell cases to controls, although non-clear cell case also had higher levels of miR-210 than controls, suggesting that this marker may have value in distinguishing kidney cancer cases more broadly from controls. Global miRNA expression profiling identified several additional candidate serum miRNA markers which are present at higher abundance in RCC cases, which together with miR-210 represent a panel of markers that we plan to evaluate in Specific Aim Task 2 of this project using a prospective blood specimen set collected by Dr. Pantuck at UCLA. With completion of this final Task, we will

complete our overall goal of discovering and validating a circulating miRNA panel for blood-based RCC diagnosis.

Year Three Progress Report Addendum:

We have requested a no-cost extension for our project, "Early Diagnosis of Clear Cell Kidney Cancer via VHL/HIF Pathway-Regulated Circulating microRNA" (W81WH-11-1-0715). This project is a collaborative award with proposal log number CA100606 under PI Dr. Muneesh Tewari (W81XWH-11-1-0714). The reason for the extension request is that during year three of the grant period, Dr. Tewari relocated his employment and laboratory from University of Washington to University of Michigan, necessitating a delay in his completion of the sample analysis required to complete the grant. He has also requested his own no-cost extension and is in the process of negotiating the transfer of the primary award of W81XWH-11-1-0714 with an extension of time for one year. The new proposed period of performance end date will be 30 April 2017.

No additional funds are required or requested, however we will continue to collect additional samples until Dr. Tewari is ready to analyze the samples, and our unexpended balance of \$387.45 will be used for supplies during our no cost extension period. We are confident that with the no-cost extension, we will be able to meet the project objectives and to obtain meaningful results.

Final Progress Report Addendum:

We have completed our prospective blood collection (by Dr. Pantuck at UCLA) for biomarker panel validation.

SUPPORTING DATA:




Figure 1: miR-210 levels in 28 RCC cases vs. 20 control serum samples with normalization using spiked-in cel-miR-39 RNA oligonucleotide. **Left, top panel:** miR-210 levels as measured by standard qRT-PCR. Y axis has been inverted to indicate that high CT values=low copy number and low CT values=high copy number. **Right, top panel:** miR-210 copy number as measured using digital PCR. **Bottom panels:** show the corresponding Receiver Operating Characteristic (ROC) curves. Area Under the Curve (AUC) for qRT-PCR is 0.85 and for digital PCR is 0.88.




Figure 2: Further analysis of data presented in Fig.1 subdivided into clear cell and non-clear cell renal carcinoma and control samples. As in Fig. 1, left panel depicts qRT-PCR data while right panel depicts droplet digital PCR data, and bottom panels show ROC curves. "Area" refers to AUC of the ROC curve.





Figure 3: Global miRNA expression profiling was performed on our case-control sample set. Depicted here are the values obtained for control RNAs used to normalize miRNA expression data from sample-to-sample and plate-to-plate. Sample order is indicative of the order in which samples were processed (randomly mixed cases and controls).

A.



B.




Figure 4: Global miRNA expression profiling reveals a number of miRNAs overexpressed in RCC cases relative to control serum samples. (A) Data presented for the top 5 miRNAs which show the greatest absolute differences in abundance between average case and average control CT values. (B) Data presented for the top 5 miRNAs overexpressed in cases as defined by lowest t-test p-value (unadjusted).




Figure 5: miR-210 levels in RCC case and control samples as determined by global miRNA profiling. **Left panel:** Raw CT values obtained for miR-210. **Right panel:** CTs for miR-210 following correction of data via normalization using numbers obtained for UniSp6.