

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.						
1. REPORT DATE (DD-MM-YYYY) 10/20/2017		2. REPORT TYPE Poster		3. DATES COVERED (From - To) 10/20/2017-10/23/2017		
4. TITLE AND SUBTITLE The Pediatric Airway: Cleft Palate Complications				5a. CONTRACT NUMBER		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Maj Steven J Durning				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBSA-Lackland, TX 78236-9908 210-292-7141				8. PERFORMING ORGANIZATION REPORT NUMBER 17394		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBSA-Lackland, TX 78236-9908 210-292-7141				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.						
13. SUPPLEMENTARY NOTES American Society of Anesthesiologists Annual Meeting 2017 (ASA), Boston MA 20-23 Oct 2017						
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			Clarice Longoria	
					19b. TELEPHONE NUMBER (Include area code) 210-292-7141	



Life vs Limb: The Compartment Syndrome Conflict

Joseph Wray MD, Steve Durning MD, Vincent Diaz MD, and Adrienne Cummings MD
San Antonio Uniformed Services Health Educations Consortium (SAUSHEC), San Antonio, TX



Case Presentation

- A 24-year-old, male patient, status post MVC rollover in Mexico, was transferred to San Antonio Military Medical Center by air. His major injuries comprised of right tibia and fibula fractures, multiple rib fractures, and liver laceration. Diarrhea and vomitus positive for blood, and was transfused 2 units of PRBCs en-route. He was also noted to have elevated potassium of 7.5 mEq/L.
- His only known past medical history consisted of alcoholic cirrhosis, otherwise unknown further medical, surgical, allergy history, NPO status.
- Upon arrival he was noted to be GCS 15, obese, agitated, diaphoretic and pale, with vitals showing hypotension and tachycardia
- Ultrasound revealed a positive FAST exam and a cirrhotic liver.
- Abdomen was tender to palpation throughout and right lower leg was noted to be extremely swollen with indeterminate pulses.
- The trauma surgery team wanted to proceed with both emergent ex-lap and subsequent fasciotomies given the clinical diagnosis of compartment syndrome.
- Given hemodynamic instability, he was emergently intubated in the trauma bay and transported to the OR for exploratory laparotomy and fasciotomy.

Preoperative and Intraoperative Considerations

- Presumed full stomach - RSI with rocuronium to avoid succinylcholine induced hyperkalemic response
- Class 3 Shock by ATLS Designation - large bore IVs, arterial line, SASAM
- Elevated potassium, likely from prolonged extrication and transport from MVC in Mexico, possibly secondary to lactic acidosis from exsanguination and tissue ischemia or K⁺ sparing diuretics for treatment of alcoholic cirrhosis. Avoidance of K⁺ sparing diuretics and maintenance of MAP to avoid further ischemia
- Multiple lower long bone fractures with tense lower extremity compartments
- Unknown allergies, medications, past medical and surgical history

Signs and symptoms of Hyperkalemia

- Clinical manifestations usually when ≥ 7.0 mEq/L
- Muscle weakness or paralysis - Ascending muscle weakness that begins with legs and progresses to trunk and arms¹ mimicking Guillain-Barre syndrome² and rare respiratory muscle weakness³
- Cardiac manifestations - EKG changes - (Tall peaked T waves with shortened QT, progressing to PR and QRS lengthening, concluding with ventricular standstill)
- Reduced urinary acid excretion - Interferes with renal ammonium (NH₄⁺) excretion and further contributes to metabolic acidosis seen in trauma patients

Intraoperative Management of Hyperkalemia

- Stabilize the membrane - Calcium chloride (500-1000mg over 3 mins) 3x elemental calcium than calcium gluconate (however can cause local irritation) Repeated every 30-60 minutes because results are shortlived⁴
- Remove excess potassium from body - loop/thiazide diuretics vs. dialysis
 - Up to 40mg of IV furosemide in diuretic naive patients
 - Avoid in patients with severe renal failure (consider dialysis or CRRT in these patients)
- Drive potassium into the cell
 - IV insulin - Enhances activity of Na-K-ATPase pump function in skeletal muscle
 - Give insulin alone if serum glucose is >250 mg/dL
 - 10 units insulin with 500 mL of 10% dextrose IV
 - Beta-2 agonist - enhances Na-K-ATPase pump and has additive effect with insulin
 - Expected K drop 0.5 to 1.5 mEq/L (1.2 to 1.5mEq/L with insulin)
 - Given in 4-8x dose used for bronchodilation (8-16 puffs)⁵
 - Hyperventilation - shifts H⁺ out of cells, driving K⁺ into cells
- Treat reversible causes of hyperkalemia
 - ACEi, ARBs, chronic heparin therapy, NSAIDs inhibit aldosterone release and reduce urinary potassium excretion
 - Metabolic acidosis, beta blockers, and insulin deficiency cause increased potassium release from cells

Compartment Syndrome Implications

- Risk Factors - long bone fractures, most commonly tibia (75% of cases)
- Delay in fracture resolution or closed reduction can increase pressures
- Blood flow cannot meet demands of surrounding tissue, venous outflow is reduced, tissue edema ensues with further rise in pressure
- Signs: pain, pallor, paresthesia, paralysis, with late sign-pulselessness
- Gold standard for diagnosis is compartment pressure
- The definitive treatment is emergent fasciotomy (compartment release)
- Release causes washout of electrolytes (namely K⁺), myoglobin, inflammatory cytokines, and can result in rhabdomyolysis and DIC

Case Management

- Despite optimal medical management (hyperventilation, albuterol, furosemide, insulin with glucose, calcium) patient's potassium ranged between 6.8 and 7.0.
- His abdomen was packed and his right lower leg noted to be extremely swollen with indeterminate compartment pressures and marginally Doppler-able pulses
- A discussion between the lead trauma surgeon and anesthesia team regarding the risks of compartment release while patient remained hyperkalemic despite maximal management, resulted in halt of further surgery and transport to ICU for emergent dialysis for return to OR once potassium normalized.
- This conflict, exemplifies the need for good communication between anesthesiologist and surgeons and ultimately led to not only sparing of the limb, but potentially saving the patient's life.

References:

1. Finch CA, Sawyer CG, Flynn JM. Clinical syndrome of potassium intoxication. *Am J Med* 1946; 1:337.
2. Evers S, Engelen A, Karsch V, Hund M. Secondary hyperkalemic paralysis. *J Neurol Neurosurg Psychiatry* 1998; 64:249.
3. Bell H, Hayes WL, Vosbrugh J. Hyperkalemic paralysis due to adrenal insufficiency. *Arch Intern Med* 1965; 115:418.
4. Braun HA, Van Horne R, Bettinger JC, Bellet S. The influence of hypocalcemia induced by sodium ethylenediamine tetraacetate on the toxicity of potassium; an experimental study. *J Lab Clin Med* 1955; 46:544.
5. Allon M, Copkney C. Albuterol and insulin for treatment of hyperkalemia in hemodialysis patients. *Kidney Int* 1990; 38:869.
6. Mabwire NT, Malahias M, Hindocha S, Khan W, Juma A. Acute compartment syndrome of the limbs: current concepts and management. *Open Orthop J* 2012. 6:535-43.
7. Kanj WW, Gunderson MA, Carrigan RB, Sankar WN. Acute compartment syndrome of the upper extremity in children: diagnosis, management, and outcomes. *J Child Orthop*. 2013 Jun. 7 (3):225-33.