



AFRL-RX-WP-JA-2017-0501

**pH-DEPENDENT SPIN STATE POPULATION AND ¹⁹F
NMR CHEMICAL SHIFT VIA REMOTE LIGAND
PROTONATION IN AN IRON(II) COMPLEX
(POSTPRINT)**

Alexandra I. Gaudette, Agnes E. Thorarinsdottir and T. David Harris

Northwestern University

**16 November 2017
Interim Report**

**Distribution Statement A.
Approved for public release: distribution unlimited.**

(STINFO COPY)

**AIR FORCE RESEARCH LABORATORY
MATERIALS AND MANUFACTURING DIRECTORATE
WRIGHT-PATTERSON AIR FORCE BASE, OH 45433-7750
AIR FORCE MATERIEL COMMAND
UNITED STATES AIR FORCE**

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**


1. REPORT DATE (DD-MM-YY) 16 November 2017		2. REPORT TYPE Interim		3. DATES COVERED (From - To) 24 January 2014 – 16 October 2017	
4. TITLE AND SUBTITLE PH-DEPENDENT SPIN STATE POPULATION AND ¹⁹ F NMR CHEMICAL SHIFT VIA REMOTE LIGAND PROTONATION IN AN IRON(II) COMPLEX (POSTPRINT)				5a. CONTRACT NUMBER FA8650-15-D-5518	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 62102F	
6. AUTHOR(S) Alexandra I. Gaudette, Agnes E. Thorarinsdottir, and T. David Harris - Northwestern University				5d. PROJECT NUMBER 4348	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER X0Y5	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Northwestern University 2145 Sheridan Road Evanston, IL 60208				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory Materials and Manufacturing Directorate Wright-Patterson Air Force Base, OH 45433-7750 Air Force Materiel Command United States Air Force				10. SPONSORING/MONITORING AGENCY ACRONYM(S) AFRL/RXAS	
				11. SPONSORING/MONITORING AGENCY REPORT NUMBER(S) AFRL-RX-WP-JA-2017-0501	
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Statement A. Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES PA Case Number: 88ABW-2017-5766; Clearance Date: 16 Nov 2017. This document contains color. Journal article published in Chemical Communications, Vol. 53, 11 Dec 2017. © 2017 The Royal Society of Chemistry. The U.S. Government is joint author of the work and has the right to use, modify, reproduce, release, perform, display, or disclose the work. The final publication is available at http://pubs.rsc.org http://dx.doi.org/10.1039/C7CC08099A					
14. ABSTRACT (Maximum 200 words) An Fe ^{II} complex that features a pH-dependent spin state population, by virtue of a variable ligand protonation state, is described. This behavior leads to a highly pH-dependent ¹⁹ F NMR chemical shift with a sensitivity of 13.9(5) ppm per pH unit at 37 °C, thereby demonstrating the potential utility of the complex as a ¹⁹ F chemical shift-based pH sensor.					
15. SUBJECT TERMS Fe ^{II} ; pH-dependent spin state population; variable ligand protonation state; NMR chemical shift; pH sensor					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT: SAR	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON (Monitor) Claretta Sullivan 19b. TELEPHONE NUMBER (Include Area Code) (937) 255-9148
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			

Cite this: *Chem. Commun.*, 2017, 53, 12962Received 22nd October 2017,
Accepted 10th November 2017

DOI: 10.1039/c7cc08158h

rsc.li/chemcomm

pH-Dependent spin state population and ^{19}F NMR chemical shift *via* remote ligand protonation in an iron(II) complex†

Alexandra I. Gaudette, Agnes E. Thorarinsdottir  and T. David Harris *

An Fe^{II} complex that features a pH-dependent spin state population, by virtue of a variable ligand protonation state, is described. This behavior leads to a highly pH-dependent ^{19}F NMR chemical shift with a sensitivity of 13.9(5) ppm per pH unit at 37 °C, thereby demonstrating the potential utility of the complex as a ^{19}F chemical shift-based pH sensor.

The development of transition metal-based molecules and materials that can be switched between low-spin and high-spin electronic states has constituted a highly active area of research over the past several decades.¹ Indeed, the magnetic bistability of such spin-crossover compounds make them potential candidates for molecular switches² and chemical sensors,^{2,3} as the spin transition can be controlled by a number of external stimuli, such as temperature,⁴ pressure,^{4b,5} and light.^{4b,6} Recently, we and others have begun to explore the potential for spin-switchable molecules as bioresponsive probes for temperature,⁷ anions,⁸ and enzyme activity.⁹ In addition, given the relationship between tissue acidosis and diseases, including cancer¹⁰ and ischemia,¹¹ a compound that undergoes a spin state transition as a function of pH could serve as a valuable tool for pH sensing. Nevertheless, pH-induced spin state switching is rare, and compounds that exhibit such behavior are unsuitable for most biological sensing applications due to $\text{p}K_{\text{a}}$ values far from biological pH¹² or poor stability in water.¹³

One approach toward biological pH sensing is to employ pH-induced changes in the chemical shift of ^{19}F resonances, typically caused by an interconversion between species of different protonation states. Here, the use of ^{19}F magnetic resonance spectroscopy (MRS) offers key advantages over the more commonly employed ^1H MRS, most notably the absence of endogenous fluorine in living systems.¹⁴ Furthermore, the chemical shift of the ^{19}F nucleus is highly sensitive to its local environment, such

that small chemical changes can lead to drastic changes in chemical shift.¹⁵ Indeed, diamagnetic ^{19}F MRS pH probes with $\text{p}K_{\text{a}}$ values near 7 have been developed for *in vivo* applications, where a variation in ^{19}F chemical shift of 12 ppm was observed between the protonated and deprotonated forms.¹⁶ In addition, the sensitivity of ^{19}F MRS probes can be further improved by incorporating paramagnetic metal ions, as the difference in chemical shift between the protonated and deprotonated forms is amplified by the presence of contact (through-bond) and dipolar (through-space) contributions to the chemical shift.^{17,18}

We previously demonstrated that spin-crossover Fe^{II} complexes can facilitate chemical shift-based MR thermometry.^{7b,c} Here, since both the contact and dipolar contributions to the paramagnetic chemical shift scale with $S(S + 1)$,¹⁸ where S is the electronic spin state, thermally-induced spin-crossover from an $S = 0$ ground state to an $S = 2$ excited state afforded a dramatic increase in chemical shift with temperature. Building on these results, we sought to develop a spin-crossover Fe^{II} complex that undergoes a deprotonation-induced spin state change near biological pH for ^{19}F chemical shift-based pH sensing. Herein, we report the synthesis and characterization of an Fe^{II} complex that features a new asymmetric 1,4,7-triazacyclononane (TACN) ligand appended with two 4-hydroxypyridine donors, and demonstrate that pH-induced spin state switching can engender highly sensitive ^{19}F MRS pH probes.

In order to develop an ^{19}F MR probe that undergoes a pH-induced spin state transition, we set out to design a ligand that (1) forms a water-soluble complex with Fe^{II}, (2) features an ^{19}F reporter group, and (3) affords a ligand field that changes dramatically with pH. Toward this end, we selected 4-hydroxy-3,5-dimethyl-2-pyridyl groups as the pH sensing moieties, because the ability of 4-hydroxypyridines to engender significant changes in the electronic structure of transition metal compounds upon protonation or deprotonation has been demonstrated in V,¹⁹ Fe,²⁰ Co,²⁰ Ni,²¹ Ru,²² Pt,²³ Re,²⁴ and Ir-based²⁵ compounds. We thus hypothesized that incorporating hydroxypyridine groups into a spin-crossover complex could create a compound with a pH-sensitive spin state population (see Fig. 1). Along these lines, we targeted the macrocyclic TACN-based ligand H₂L (see Fig. 1, left).

Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208-3113, USA. E-mail: dharris@northwestern.edu

† Electronic supplementary information (ESI) available: Experimental details, crystallographic data, additional magnetic and spectroscopic data, and crystallographic information file (CIF) for 1. CCDC 1574128. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7cc08158h




Fig. 1 Scheme depicting the mechanism for pH sensing with **1**. With increasing pH, the pendent hydroxylpyridine groups are deprotonated, and the resulting weaker ligand field leads to a higher population of the $S = 2$ high-spin state.

The ligand H_2L was prepared following a four-step synthesis outlined in Scheme S1 (ESI[†]). Incorporation of two 4-methoxy-3,5-dimethyl-2-picolyl donors onto a tosylated TACN macrocycle *via* nucleophilic substitution reaction, followed by deprotection with H_2SO_4 and reductive amination with 3-fluoro-6-methyl-2-pyridinecarboxaldehyde, afforded H_2L as a yellow solid. Under anaerobic conditions, reaction of H_2L with one equivalent of $[Fe(H_2O)_6](BF_4)_2$ in MeCN, followed by aqueous workup, gave $[(H_2L)Fe](BF_4)_2$ (**1**) as a yellow-green solid in 52% yield. Subsequent slow cooling of an aqueous solution afforded yellow-green block-shaped single crystals of **1**.

Single-crystal X-ray diffraction analysis for **1** was carried out at 100 K (see Fig. 2 and Table S1, ESI[†]). The compound crystallized in the orthorhombic space group $Pbca$, with one $[(H_2L)Fe]^{2+}$ cation and two $[BF_4]^-$ anions in the asymmetric unit. The two acidic protons on the hydroxylpyridine moieties were located in the Fourier difference map and were freely refined. The coordination environment at the Fe^{II} center in $[(H_2L)Fe]^{2+}$ is highly distorted from octahedral geometry. The Fe–N bond distances range from 2.109(5) to 2.223(5) Å with an average distance of 2.165(5) Å, and the *cis* N–Fe–N angles range from 75.1(2) to 111.3(2)° with an average angle of 90.8(2)°. These structural metrics indicate the presence of high-spin Fe^{II} in the solid state at 100 K. Considering the observation of low-spin Fe^{II} in related compounds at 100 K,^{7c,26} we surmise the high-spin Fe^{II} center in **1** likely results from crystal packing effects.

We next examined the effects of pH on the electronic structure of **1** in aqueous solution. UV-Visible absorption spectra were




Fig. 2 Crystal structure of the complex cation of **1**. Orange, green, red, blue, gray, and white spheres represent Fe, F, O, N, C, and H atoms respectively; non-hydroxyl H atoms are omitted for clarity.

collected for 100 mM phosphate buffer solutions containing 2% (v/v) DMSO between pH 5.39 and 7.94 at 25 °C (see Fig. S1, ESI[†]). At pH 7.94, the spectrum displays an intense absorption band at 271 nm ($\epsilon_{\max} = 27\,000\text{ M}^{-1}\text{ cm}^{-1}$) and three less intense bands at 334 ($\epsilon_{\max} = 2300\text{ M}^{-1}\text{ cm}^{-1}$), 416 ($\epsilon_{\max} = 960\text{ M}^{-1}\text{ cm}^{-1}$), and 453 nm ($\epsilon_{\max} = 1200\text{ M}^{-1}\text{ cm}^{-1}$). We assign these features to π – π^* and metal–ligand charge transfer (MLCT) transitions, respectively, as observed in similar compounds.^{26,27} As the pH is decreased to 5.39, the intensity of the π – π^* absorption decreases ($\epsilon_{\max} = 12\,000\text{ M}^{-1}\text{ cm}^{-1}$), while the intensities of the MLCT bands increase significantly with concomitant shifts to 383 ($\epsilon_{\max} = 2800\text{ M}^{-1}\text{ cm}^{-1}$) and 451 nm ($\epsilon_{\max} = 1600\text{ M}^{-1}\text{ cm}^{-1}$). This increase in MLCT band intensity with decreasing pH suggests that the low-spin state becomes more populated when the ligand is fully protonated.^{7c,28} Furthermore, the presence of a single set of three nearly isosbestic points at 330, 472, and 525 nm indicates an equilibrium between two species, likely $(L^{2-})Fe$ and $[(H_2L)Fe]^{2+}$, and therefore suggests highly similar values of pK_a for the two hydroxyl protons. Such nearly simultaneous double deprotonation in aqueous solution has been reported for similar complexes,^{24,25a} and is further supported by a single sigmoidal inflection point of ϵ vs. pH plots at 271 and 380 nm, providing pK_a values of 5.95(7) and 5.90(2), respectively (see Fig. S2, S3 and Table S2, ESI[†]).

To probe the magnetic properties of **1** and further examine the solution electronic structure as a function of pH, variable-temperature dc magnetic susceptibility data were collected using the Evans method²⁹ in the temperature range 5–60 °C (see Fig. 3, upper). Samples were prepared in 100 mM phosphate buffer solutions with 2% (v/v) DMSO added as a reference. At pH 4.74, the value of $\chi_M T$ increases with increasing temperature, from 1.78(1) $\text{cm}^3\text{ K mol}^{-1}$ at 5 °C to 2.40(1) $\text{cm}^3\text{ K mol}^{-1}$ at 60 °C. This temperature dependence indicates the presence of thermally-induced spin-crossover, with significant population of both an $S = 0$ ground state and $S = 2$ excited state in this temperature range. Assuming a value of $g = 2$ for the $S = 2$ state, insertion into the Curie law gives percentages of high-spin molecules (%HS) of 59% at 5 °C and 80% at 60 °C.³⁰ In contrast, the variable-temperature magnetic susceptibility data for an analogous solution of **1** at pH 7.82 show much smaller temperature dependence. Here, $\chi_M T$ increases only slightly with increasing temperature, from 2.62(1) $\text{cm}^3\text{ K mol}^{-1}$ at 5 °C to 2.70(1) $\text{cm}^3\text{ K mol}^{-1}$ at 60 °C, corresponding to %HS = 87% at 5 °C and 90% at 60 °C. We attribute this shift toward greater population of the $S = 2$ state with increasing pH to a decreased ligand field strength of L^{2-} vs. H_2L , owing to the weaker π acidity of the former, which is consistent with pyridyl substituent effects observed for related spin-crossover iron(II) complexes.^{26,31}

To further probe the pH dependence of the spin state of **1**, dc magnetic susceptibility data were collected on an aqueous solution as a function of pH at 25 and 37 °C (see Fig. 3, lower). At 25 °C, $\chi_M T$ rapidly increases with increasing pH, from a value of $\chi_M T = 2.07(1)\text{ cm}^3\text{ K mol}^{-1}$ at pH 4.74 to $\chi_M T = 2.64(1)\text{ cm}^3\text{ K mol}^{-1}$ at pH 7.82, corresponding to %HS = 69% at pH 4.74 and 88% at pH 7.82. The plot of $\chi_M T$ vs. pH at 37 °C exhibits similar behavior, albeit with less pronounced pH dependence, where $\chi_M T$ increases




Fig. 3 Evans method magnetic susceptibility data for **1** in 100 mM phosphate buffer with 2% (v/v) DMSO reference. Upper: Variable-temperature data at pH 4.74 and 7.82. Lower: Variable-pH data at 25 and 37 °C. Error bars represent standard deviations of the measurements.

from 2.20(1) $\text{cm}^3 \text{K mol}^{-1}$ at pH 4.74 (%HS = 73%) to 2.65(1) $\text{cm}^3 \text{K mol}^{-1}$ at pH 7.82 (%HS = 88%). These results are in accord with the variable-pH behavior observed in the UV-Visible analysis described above, and confirm the pH dependence of the spin state population in **1**.

Finally, we aimed to determine the effect of the pH-dependent spin state population on the ^{19}F NMR chemical shift of **1**. Variable-pH ^{19}F NMR spectra were collected on 2 mM solutions of **1** in 100 mM phosphate buffers with 2% (v/v) DMSO at 25 and 37 °C (see Fig. S4 and S5, ESI†). In the pH range 4.61–7.74, these spectra display a single ^{19}F resonance from the metalated ligand. At pH 4.61, when the hydroxylpyridine groups are predominantly protonated, the ^{19}F resonance appears at 11.89 ppm vs. trifluoroacetic acid (TFA) at 25 °C. As the pH is increased, this resonance shifts downfield, following sigmoidal behavior, to a maximum value of 47.03 ppm at pH 7.74. The data at 37 °C exhibit a similar pH profile, albeit with a slightly smaller pH dependence, where the ^{19}F chemical shift increases from 13.10 to 43.57 ppm between pH 4.61 and 7.74 (see Fig. 4). These data were fit to a Boltzmann sigmoidal function according to eqn (1):

$$\delta = A_2 + (A_1 - A_2)/(1 + \exp((\text{pH} - \text{p}K_a)/dx)) \quad (1)$$

where δ is the observed ^{19}F chemical shift, A_2 is the theoretical chemical shift of the fully deprotonated species, A_1 is the theoretical chemical shift of the fully protonated species, dx is a parameter describing the steepness of the curve, and $\text{p}K_a$ is obtained as the inflection point of the curve. Fits to the data




Fig. 4 Plot of ^{19}F NMR chemical shift vs. pH for **1** in 100 mM phosphate buffer with 2% (v/v) DMSO at 25 and 37 °C. Chemical shift values are referenced to TFA, which was set to 0 ppm. Circles denote experimental data and black lines correspond to sigmoidal fits to the data (eqn (1)).

provided estimated $\text{p}K_a$ values of 6.03(1) at 25 °C and 5.97(1) at 37 °C, which are in excellent agreement with the values obtained from UV-Visible data. Further, rearranging eqn (1) and inserting the values obtained from the sigmoidal fit of each curve (see Table S3, ESI†) gives eqn (2) and (3) for the data at 25 and 37 °C, respectively, which relate chemical shift to pH:

$$\text{pH} = 6.03 + 0.51 \ln(-38.8/(\delta - 48.3) - 1) \quad (2)$$

$$\text{pH} = 5.97 + 0.50 \ln(-33.5/(\delta - 44.5) - 1) \quad (3)$$

In addition, by applying a linear fit to the data between pH 5.25 and 6.95, the change in ^{19}F chemical shift affords a pH sensitivity of 16.1(4) and 13.9(5) ppm per pH unit at 25 and 37 °C, respectively (see Fig. S6, ESI†). While this pH sensitivity may partially stem from electrostatic effects and/or small changes in geometry at Fe^{II} upon deprotonation of the ligand, the dramatic change in ^{19}F chemical shift with pH strongly resembles the variable-pH $\chi_M T$ data discussed above. This similarity suggests that the primary cause of the pH-induced change in ^{19}F chemical shift is an increase in population of the $S = 2$ state, owing to deprotonation of the hydroxylpyridine donors.

The foregoing results demonstrate the potential utility of a pH-induced spin state change for pH sensing using ^{19}F MRS. We show that a change in spin state upon deprotonation of $[(\text{H}_2\text{L})\text{Fe}]^{2+}$ dramatically affects its ^{19}F chemical shift, which can be employed to quantitate pH. Indeed, this complex displays greater pH sensitivity than the current most-sensitive lanthanide-based ^{19}F MRS probes^{17b} and the only reported transition metal-based ^1H MR pH probe.³² To the best of our knowledge, $[(\text{H}_2\text{L})\text{Fe}]^{2+}$ represents the first example of an ^{19}F MRS pH probe that employs a pH-induced spin state change. While the oxygen sensitivity of this compound precludes it from *in vivo* experiments, the principles outlined here could inform the design of chemical shift-based MR pH sensors. Future efforts will focus on developing similar compounds with higher stability under biological conditions.

This research was funded by the Air Force Research Laboratory under agreement no. FA8650-15-5518 and Northwestern

University. The U. S. Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation thereon. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or implied, of AFRL or the U. S. Government. T. D. H. thanks the Alfred P. Sloan Foundation, and A. E. T. acknowledges the Leifur Eiriksson Foundation for funding support.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- (a) P. Gütllich and H. A. Goodwin, in *Spin Crossover in Transition Metal Compounds I*, ed. P. Gütllich and H. A. Goodwin, Springer, Berlin, 2004, pp. 1–47; (b) *Spin-Crossover Materials: Properties and Applications*, ed. M. A. Halcrow, Wiley-VCH, Weinheim, 2013.
- (a) O. Kahn and C. J. Martinez, *Science*, 1998, **279**, 44; (b) K. Senthil Kumar and M. Ruben, *Coord. Chem. Rev.*, 2017, **346**, 176.
- (a) G. J. Halder, C. J. Kepert, B. Moubarak, K. S. Murray and J. D. Cashion, *Science*, 2002, **298**, 1762; (b) E. Coronado, M. Giménez-Marqués, G. M. Espallargas, F. Rey and I. J. Vitórica-Yrezábal, *J. Am. Chem. Soc.*, 2013, **135**, 15986; (c) C. Bartual-Murgui, A. Akou, C. Thibault, G. Molnár, C. Vieu, L. Salmon and A. Bousseksou, *J. Mater. Chem. C*, 2015, **3**, 1277.
- (a) M. A. Halcrow, *Chem. Soc. Rev.*, 2011, **40**, 4119; (b) P. Gütllich, A. B. Gaspar and Y. Garcia, *Beilstein J. Org. Chem.*, 2013, **9**, 342; (c) M. A. Halcrow, *Chem. Lett.*, 2014, **43**, 1178.
- (a) V. Ksenofontov, A. B. Gaspar and P. Gütllich, in *Spin Crossover in Transition Metal Compounds III*, ed. P. Gütllich and H. A. Goodwin, Springer, Berlin, 2004, pp. 23–64; (b) B. Li, R.-J. Wei, J. Tao, R.-B. Huang and L.-S. Zheng, *Inorg. Chem.*, 2010, **49**, 745.
- (a) S. Decurtins, P. Gütllich, C. P. Köhler, H. Spiering and A. Hauser, *Chem. Phys. Lett.*, 1984, **105**, 1; (b) A. Hauser, in *Spin Crossover in Transition Metal Compounds II*, ed. P. Gütllich and H. A. Goodwin, Springer, Berlin, 2004, pp. 155–198; (c) A. Cannizzo, C. J. Milne, C. Consani, W. Gawelda, C. Bressler, F. van Mourik and M. Chergui, *Coord. Chem. Rev.*, 2010, **254**, 2677.
- (a) R. N. Muller, L. Vander Elst and S. Laurent, *J. Am. Chem. Soc.*, 2003, **125**, 8405; (b) I.-R. Jeon, J. G. Park, C. R. Haney and T. D. Harris, *Chem. Sci.*, 2014, **5**, 2461; (c) A. E. Thorarinsdottir, A. I. Gaudette and T. D. Harris, *Chem. Sci.*, 2017, **8**, 2448.
- (a) Z. Ni and M. P. Shores, *J. Am. Chem. Soc.*, 2009, **131**, 32; (b) Z. Ni, A. M. McDaniel and M. P. Shores, *Chem. Sci.*, 2010, **1**, 615; (c) C. M. Klug, A. M. McDaniel, S. R. Fiedler, K. A. Schulte, B. S. Newell and M. P. Shores, *Dalton Trans.*, 2012, **41**, 12577; (d) A. M. McDaniel, C. M. Klug and M. P. Shores, *Eur. J. Inorg. Chem.*, 2013, 943; (e) S. A. Barrett and M. A. Halcrow, *RSC Adv.*, 2014, **4**, 11240.
- (a) V. Stavila, M. Allali, L. Canaple, Y. Stortz, C. Franc, P. Maurin, O. Beuf, O. Dufay, J. Samarut, M. Janier and J. Hasserodt, *New J. Chem.*, 2008, **32**, 428; (b) J. Hasserodt, *New J. Chem.*, 2012, **36**, 1707; (c) F. Touti, P. Maurin and J. Hasserodt, *Angew. Chem., Int. Ed.*, 2013, **52**, 4654; (d) C. Gondrand, F. Touti, E. Godart, Y. Berezanskyy, E. Jeanneau, P. Maurin and J. Hasserodt, *Eur. J. Inorg. Chem.*, 2015, 1376; (e) J. Wang, C. Gondrand, F. Touti and J. Hasserodt, *Dalton Trans.*, 2015, **44**, 15391.
- (a) I. F. Tannock and D. Rotin, *Cancer Res.*, 1989, **49**, 4373; (b) R. J. Gillies, N. Raghunand, G. S. Karczmar and Z. M. Bhujwala, *J. Magn. Reson. Imaging*, 2002, **16**, 430; (c) R. A. Gatenby and R. J. Gillies, *Nat. Rev. Cancer*, 2004, **4**, 891.
- (a) K. Katsura, A. Ekholm, B. Asplund and B. K. Siesjö, *J. Cereb. Blood Flow Metab.*, 1991, **11**, 597; (b) J. J. Lemasters, J. M. Bond, E. Chacon, I. S. Harper, S. H. Kaplan, H. Ohata, D. R. Trollinger, B. Herman and W. E. Cascio, *EXS*, 1996, **76**, 99; (c) T. Kalogeris, C. P. Baines, M. Krenz and R. J. Korthuis, *Int. Rev. Cell Mol. Biol.*, 2012, **298**, 229.
- (a) L. L. Martin, K. S. Hagen, A. Hauser, R. L. Martin and A. M. Sargeson, *J. Chem. Soc., Chem. Commun.*, 1988, 1313; (b) F. Touti, P. Maurin, L. Canaple, O. Beuf and J. Hasserodt, *Inorg. Chem.*, 2012, **51**, 31; (c) R. Nowak, E. A. Prasetyanto, L. De Cola, B. Bojer, R. Siegel, J. Senker, E. Rössler and B. Weber, *Chem. Commun.*, 2017, **53**, 971.
- (a) M. Enamullah, W. Linert, V. Gutmann and R. F. Jameson, *Monatsh. Chem.*, 1994, **125**, 1301; (b) M. Enamullah and W. Linert, *J. Coord. Chem.*, 1995, **35**, 325; (c) Y.-H. Luo, M. Nihei, G.-J. Wen, B.-W. Sun and H. Oshio, *Inorg. Chem.*, 2016, **55**, 8147.
- J. Ruiz-Cabello, B. P. Barnett, P. A. Bottomley and J. W. M. Bulte, *NMR Biomed.*, 2011, **24**, 114.
- W. R. Dolbier, *Guide to Fluorine NMR for Organic Chemists*, John Wiley & Sons, Hoboken, NJ, 2009.
- S. He, R. P. Mason, S. Hunjan, V. D. Mehta, V. Arora, R. Katipally, P. V. Kulkarni and P. P. Antich, *Bioorg. Med. Chem.*, 1998, **6**, 1631.
- (a) A. M. Kenwright, I. Kuprov, E. De Luca, D. Parker, S. U. Pandya, P. K. Senanayake and D. G. Smith, *Chem. Commun.*, 2008, 2514; (b) K. H. Chalmers, E. De Luca, N. H. M. Hogg, A. M. Kenwright, I. Kuprov, D. Parker, M. Botta, J. I. Wilson and A. M. Blamire, *Chem. – Eur. J.*, 2010, **16**, 134.
- I. Bertini and C. Luchinat, *NMR of Paramagnetic Molecules in Biological Systems*, The Benjamin/Cummings Publishing Company, Inc., Menlo Park, CA, 1986.
- L. Yang, A. la Cour, O. P. Anderson and D. C. Crans, *Inorg. Chem.*, 2002, **41**, 6322.
- L. J. Kershaw Cook, F. Tuna and M. A. Halcrow, *Dalton Trans.*, 2013, **42**, 2254.
- Z. Derikvand, M. M. Olmstead, B. Q. Mercado, A. Shokrollahi and M. Shahryari, *Inorg. Chim. Acta*, 2013, **406**, 256.
- (a) S. Klein, W. G. Dougherty, W. S. Kassel, T. J. Dudley and J. J. Paul, *Inorg. Chem.*, 2011, **50**, 2754; (b) D. C. Marelius, S. Bhagan, D. J. Charboneau, K. M. Schroeder, J. M. Kamdar, A. R. McGettigan, B. J. Freeman, C. E. Moore, A. L. Rheingold, A. L. Cooksy, D. K. Smith, J. J. Paul, E. T. Papish and D. B. Grotjahn, *Eur. J. Inorg. Chem.*, 2014, 676.
- H. Zhang, B. Zhang, Y. Li and W. Sun, *Inorg. Chem.*, 2009, **48**, 3617.
- G. F. Manbeck, J. T. Muckerman, D. J. Szalda, Y. Himeda and E. Fujita, *J. Phys. Chem. B*, 2015, **119**, 7457.
- (a) J. DePasquale, I. Nieto, L. E. Reuther, C. J. Herbst-Gervasoni, J. J. Paul, V. Mochalin, M. Zeller, C. M. Thomas, A. W. Addison and E. T. Papish, *Inorg. Chem.*, 2013, **52**, 9175; (b) W.-P. Wu, Y.-J. Xu, S.-W. Chang, J. Deng and Y. Fu, *ChemCatChem*, 2016, **8**, 3375.
- I. Prat, A. Company, T. Corona, T. Parella, X. Ribas and M. Costas, *Inorg. Chem.*, 2013, **52**, 9229.
- (a) I. Bernal, I. M. Jensen, K. B. Jensen, C. J. McKenzie, H. Toftlund and J.-P. Tuchagues, *J. Chem. Soc., Dalton Trans.*, 1995, 3667; (b) V. Ballard, F. Banse, E. Anxolabéhère-Mallart, M. Nierlich and J.-J. Girerd, *Eur. J. Inorg. Chem.*, 2003, 2529.
- W. Linert, M. Konecny and F. Renz, *J. Chem. Soc., Dalton Trans.*, 1994, 1523.
- (a) D. F. Evans, *J. Chem. Soc.*, 1959, 2003; (b) E. M. Schubert, *J. Chem. Educ.*, 1992, **69**, 62.
- O. Kahn, *Molecular Magnetism*, VCH Publishers, Inc., New York, NY, 1993.
- L. J. Kershaw Cook, R. Kulmaczewski, R. Mohammed, S. Dudley, S. A. Barret, M. A. Little, R. J. Deeth and M. A. Halcrow, *Angew. Chem., Int. Ed.*, 2016, **55**, 4327.
- P. B. Tsitovich, J. M. Cox, J. A. Sperryak and J. R. Morrow, *Inorg. Chem.*, 2016, **55**, 12001.