

Award Number: W81XWH-15-C-0070

TITLE: Enhancing mHealth Technology in the PCMH Environment to Activate Chronic Care Patients

PRINCIPAL INVESTIGATOR: Ronald W. Gimbel,

CONTRACTING ORGANIZATION: Clemson University
Clemson, SC 29634

REPORT DATE: Sept 2017

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE		Form Approved OMB No. 0704-0188
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.		
1. REPORT DATE September 2017	2. REPORT TYPE Annual	3. DATES COVERED 17 Aug 2016 - 16 Aug 2017
4. TITLE AND SUBTITLE Enhancing mHealth Technology in the PCMH Environment to Activate Chronic Care Patients		5a. CONTRACT NUMBER W81XWH-15-C-0070
		5b. GRANT NUMBER
		5c. PROGRAM ELEMENT NUMBER -
6. AUTHOR(S) Ronald W. Gimbel, PhD, Principal Investigator E-Mail: rgimbel@clemson.edu		5d. PROJECT NUMBER -
		5e. TASK NUMBER -
		5f. WORK UNIT NUMBER -
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Clemson University 501 Edwards Hall Clemson, SC 29634		8. PERFORMING ORGANIZATION REPORT NUMBER -
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSOR/MONITOR'S ACRONYM(S) -
		11. SPONSOR/MONITOR'S REPORT NUMBER(S) -
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited		
13. SUPPLEMENTARY NOTES -		
14. ABSTRACT - The potential of mobile health (mHealth) technologies in the care of patients with diabetes and other chronic conditions has captured the attention of clinicians and researchers. Efforts to date have incorporated a variety of tools and techniques including web-based portals, tailored behavioral text messaging, remote collection of biometric data, electronic coaching, electronic-based health education, secure e-mail communication between visits, and electronic collection of lifestyle and quality of life surveys. Each of these tools, used alone or in combination, have demonstrated varying degrees of effectiveness. Some of the more promising results have been demonstrated using regular collection of biometric devices, tailored behavioral messaging, secure e-mail communication with clinical teams, and regular reporting of quality of life variables. In this study, we seek to incorporate several of the most promising mHealth capabilities in a patient centered medical home (PCMH) workflow. We aim to address underlying technology need and gaps related to the use of mHealth technology and the activation of patients living with Type-2 diabetes. Stated differently, we enable supporting technologies while seeking to influence patient activation and self-care activities.		

15. SUBJECT TERMS MHCE, Mobile Health Care Environment mHealth, mobile health MHS, Military Health System PAM®, Patient Activation Measure CS-PAM®, Clinician Support for Patient Activation Measure PCMH, patient centered medical home SDSCA, Summary of Diabetes Self-Care Activities SUS, System Usability Scale Type 2 Diabetes User-Centered Design Research TATRC, Telemedicine and Advanced Technology Research Center					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE	Unclassified	113	USAMRMC
Unclassified	Unclassified	Unclassified			19b. TELEPHONE NUMBER <i>(include area code)</i> —

1. INTRODUCTION:

The potential of mobile health (mHealth) technologies in the care of patients with diabetes and other chronic conditions has captured the attention of clinicians and researchers. Efforts to date have incorporated a variety of tools and techniques including web-based portals, tailored behavioral text messaging, remote collection of biometric data, electronic coaching, electronic-based health education, secure e-mail communication between visits, and electronic collection of lifestyle and quality of life surveys. Each of these tools, used alone or in combination, have demonstrated varying degrees of effectiveness. Some of the more promising results have been demonstrated using regular collection of biometric devices, tailored behavioral messaging, secure e-mail communication with clinical teams, and regular reporting of quality of life variables. In this study, we seek to incorporate several of the most promising mHealth capabilities in a patient centered medical home (PCMH) workflow. We aim to address underlying technology need and gaps related to the use of mHealth technology and the activation of patients living with Type-2 diabetes. Stated differently, we enable supporting technologies while seeking to influence patient activation and self-care activities.

2. **KEYWORDS:**

MHCE, Mobile Health Care Environment
mHealth, mobile health
MHS, Military Health System
PAM®, Patient Activation Measure
CS-PAM®, Clinician Support for Patient Activation Measure
PCMH, patient centered medical home
SDSCA, Summary of Diabetes Self-Care Activities
SUS, System Usability Scale
Type 2 Diabetes
User-Centered Design Research
TATRC, Telemedicine and Advanced Technology Research Center

3. ACCOMPLISHMENTS:

- What were the major goals of the project?

FOCUS: Phase I user-centered design research

Major Task 1: Identified potential DM patients to participate in the study **(COMPLETED 10/31/16)** - For Phase 1 of Patient recruitment, we requested use of a *Partial HIPAA Waiver of Authorization for Recruitment* for patient recruitment. Patient recruitment occurred via review of the PCMH clinic schedule, referrals from providers, distributed fliers, and population health databases. We also prescreened potential subject's medical record for inclusion and exclusion criteria to verify eligibility for the study.

Major Task 2: Recruited, randomized, and consented patients to participate in user-centered design research; n=15 per site **(COMPLETED NOV 2016)** - For Phase 1 for the patients, an Informed Consent Document (ICD) and HIPAA authorization was sought in advance from each prospective subject and appropriately documented in accordance with 32 CFR 219.117. The senior research associate or Site PI was available to answer any questions that the patients had in regards to the study. For Phase 1 clinicians, a minimal risk information sheet was given to providers before participating in any study-related procedures. The senior research associate or Site PI was available to answer any questions that the clinician may have had in regards to the study. If the PI was the supervisor for any of the clinicians, the SRA recruited their subjects to prevent any perception of coercion or undue influence.

Phase 1, Patient-Participants: Patient recruitment occurred via review of the PCMH clinic schedule, referrals from providers, distributed fliers, and population health databases. Patients were also prescreened via their medical record for inclusion and exclusion criteria to verify eligibility for the study. The PI verified inclusion and exclusion criteria.

Phase 1, Provider-Participants: Clinicians were invited to participate by word of mouth from the site PI and the SRA. A brief background of the study was given and volunteers who were interested and wanted to support the study met with the SRAs and were asked to review the minimal risk information sheet to be included in the study.

Major Task 3: Conduct 4-5 days of user-centered design research (qualitative) with research participants – each site **(COMPLETED 12/3/16)**

- Phase I travel and research to Nellis AFB was conducted 16-19 NOV 2016. Phase I travel and research to MAMC was conducted 30 NOV – 3 DEC 2016.

Research Steps (Phase I). Provider-participants participated in focus group sessions held in a conference room or other suitable location at both sites during a one hour period with two or more of the research team regarding the use of biomedical devices and technology in patient self-management of diabetes. They were shown a series of mock up (under development) technology devices and user interfaces and asked to provide feedback on their preferences. The session was audio recorded for note-taking purposes.

Major Task 4: Analysis and reporting of user-centered design research **(COMPLETED 12/15/16)**

Subtask 1: Assembled, coded, interpreted and triangulated all data collected in user-centered design research **(COMPLETED 12/15/16)**

Subtask 2: Authored user-centered design assessment report; to be used in infrastructure/design modification prior to Phase II study *(Report + manuscript = major milestone #1)*
(COMPLETED 12/15/16)

Subtask 3: Advised TATRC of user-centered research driven modification recommendations to MHCE; request implementation of recommendations prior to Phase II feasibility study
(COMPLETED 12/15/16)

FOCUS: MHCE modification and readiness to conduct Phase II feasibility study

Major Task 1: TATRC team modified MHCE based on findings from user-centered design research **(COMPLETED 6/30/17)**

Major Task 2: Coordinated all logistical issues and other preparation for Phase II research
(COMPLETED 6/30/17)

Major Task 3: Identify potential DM patients to participate in the study **(Ongoing During Recruiting Process)**

Major Task 4: Recruit, randomize, and consent patients to participate in feasibility study **(Recruiting Process Underway)**

FOCUS: Phase II formal feasibility study

Major Task 1: Launch and conduct a formal 12 month feasibility study of patient activated MCHE **(Ongoing Recruitment Phase)**. **As of 8/14/17, there are a total of 120 participants recruited at both sites with 50 patients recruited in the intervention group and 46 patients recruited in the control group.**

Major Task 2: Per protocol collect and analyze (using appropriate statistical testing) data from MHCE, clinical data from SRAs (each site), and other data sources **(Currently Ongoing – patient information from devices and surveys are populating data)**

What was accomplished under these goals?

What opportunities for training and professional development has the project provided? (Not applicable for current reporting period. Training and professional development will concur during and following Phase II)

a. How were the results disseminated to communities of interest?

2016 AMSUS Annual Continuing Education Meeting - Poster #1 – presented 11/29/16, National Harbor, Maryland - “New Biometric Data Collection, Analysis and Visualization in the DoD’s Mobile Health Care Environment”; Authors: Ron Gimbel, PhD, Clemson University; Jeanette Little, MS, TATRC; Terry Newton, MD, OTSG – USA.

2016 AMSUS Annual Continuing Education Meeting - Poster #2 – presented 11/29/16, National Harbor, Maryland – “Enhancing mHealth Technology in the DoD’s PCMH Environment to Activate Type 2 Diabetes Patients; Authors: Ron Gimbel, PhD; Joel Williams, PhD; Liwei Chen, MD, PhD; Cheryl Dye, PhD; Karen Edwards, MEd; Jeanette Little, MS, Terry Newton, MD

DoD publication clearance was obtained from USA, USAF, TATRC for our first manuscript submitted to JMIR Research Protocols for peer review DEC 2016; JMIR Research Protocols approved manuscript for submission on 12 FEB 2017; JMIR Research Protocols manuscript published on 6 MAR 2017: *Enhancing mHealth Technology in the patient centered medical home environment to activate Type 2 diabetes patients: A multi-site feasibility study protocol*; <http://www.researchprotocols.org/2017/3/e38/>

Poster delivered for American Academy of Health Behavior Scientific Meeting, Tucson, AZ; *Improving Diabetes Self-management through Messages Tailored to Patient Activation Levels*; delivered 19-22 MAR 2017.

Presentation approved for the Military Health System Research Symposium, Kissimmee, FL; *Mobile Health Apps in the Medical Theater of Operation – Exploring the Usability of a Mobile Application for Improving Chronic Disease Self-management in the Patient Centered Medical Home Environment*. AUG 2017.

b. What do you plan to do during the next reporting period to accomplish the goals?

FOCUS: MHCE modification and readiness to conduct Phase II feasibility study

Major Task 4: Recruit, randomize, and consent patients to participate in feasibility study – Recruitment process should be completed by 31OCT2017. We have requested a no-cost extension to complete the study so that our end date for Phase II, along with study wrap-up and reporting should be 31DEC2018.

FOCUS: Phase II formal feasibility study

Major Task 1: Launch and conduct a formal 12 month feasibility study of patient activated MCHE

Major Task 2: Per protocol collect and analyze (using appropriate statistical testing) data from MHCE, clinical data from SRAs (each site), and other data sources

Major Task 3: Author 2-3 manuscripts on patient activated MHCE system design, implementation and on-going use in chronic care self-management (First manuscript has been approved and published; second manuscript is under completion now; future manuscripts will be forthcoming as data is populated)

FOCUS: Phase II study wrap up and reporting

Major Task 1: Complete all data analysis from Phase II study – n/a

Major Task 2: Complete formal reporting and follow-on manuscripts (if any) from Phase II study. Formal DoD report will include specific design and investment recommendations grounded in evidence collected during Phase I and Phase II of the study – n/a

Major Task 3: Formal study closure and IRB closure reports – n/a

4. IMPACT:

- **What was the impact on the development of the principal discipline(s) of the project?** Research benefits include improved understanding of how to advance three joint PCMH principles (coordination of care, improved quality and safety, and enhanced access to care) through the use of mobile technology. We expect to improve the understanding of how to include mHealth technology into the PCMH workflow, as well as exploring how to use mHealth technology in the activation of patients diagnosed with Type 2 diabetes. We also explore how patient complexity and degree of “sickness” may influence whether and how patients use mHealth technologies in self-management of their disease. Finally, we will map patient-entered biomedical data into clinical documentation and a decision support platform useful in chronic care management.
- **What was the impact on other disciplines?** The Clinical Advisory Team, populated by MHS clinicians, academicians, and research associates, established clinical and safety thresholds and provided clinician preferences on optimal data visualization and clinical workflow. To ensure the MHCE thresholds were grounded in evidence, CAT members compared them to those presented in current clinical practice guidelines, systematic reviews, and other peer-reviewed biomedical manuscripts.
- **What was the impact on technology transfer?** (Not applicable for this reporting period)
- **What was the impact on society beyond science and technology?** Phase II research will make an impact on society by improving patient activation through the use of the MHCE which is expected to improve their chronic disease self-management behaviors. When chronic disease self-management behaviors improve, clinical outcomes and a reduction of high-intensity health services utilization will decrease. Our efforts and expected knowledge generated from this study will make a substantial contribution on how to use PAM in health services delivery.

5. **CHANGES/PROBLEMS:**

- **Changes in approach and reasons for change** (Nothing to report)
- **Actual or anticipated problems or delays and actions or plans to resolve them**

Due to a nine-month delayed approval from MAMC IRB (submitted NOV2016 and approved SEPT2017), Phases I and II were not scheduled as planned. Phase II research began in early July, 2017 which has required a no-cost extension for the execution of the forma 12-month study into year three.

TATRC adaptations of the MHCE platform, incorporating recommendations from Phase I research and other technical challenges, have also delayed the start date for Phase II by six months.

- **Changes that had a significant impact on expenditures**

The delay in Phase I and Phase II research activities (and resulting expenditures) have had a significant impact on expenditures, although the study should be completed with the same contract budget as originally set for the contract period. Total budget \$1,230,749.

- **Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents** (Not applicable)
- **Significant changes in use or care of human subjects** (No significant changes)
- **Significant changes in use or care of vertebrate animals.** (Not applicable)
- **Significant changes in use of biohazards and/or select agents** (Not applicable)

6. **PRODUCTS:**

- **Publications, conference papers, and presentations**

- **Journal publications.**

DoD publication clearance was obtained from USA, USAF, TATRC for our first manuscript submitted to JMIR Research Protocols for peer review DEC 2016; JMIR Research Protocols approved manuscript for submission on 12 FEB 2017; JMIR Research Protocols manuscript published on 6 MAR 2017: *Enhancing mHealth Technology in the patient centered medical home environment to activate Type 2 diabetes patients: A multi-site feasibility study protocol*; <http://www.researchprotocols.org/2017/3/e38/>

- **Books or other non-periodical, one-time publications.** (Not applicable)
- **Other publications, conference papers, and presentations.**

2016 AMSUS Annual Continuing Education Meeting - Poster #1 – presented 11/29/16, National Harbor, Maryland - “New Biometric Data Collection, Analysis and Visualization in the DoD’s Mobile Health Care Environment”; Authors: Ron Gimbel, PhD, Clemson University; Jeanette Little, MS, TATRC; Terry Newton, MD, OTSG – USA.

2016 AMSUS Annual Continuing Education Meeting - Poster #2 – presented 11/29/16, National Harbor, Maryland – “Enhancing mHealth Technology in the DoD’s PCMH Environment to Activate

Type 2 Diabetes Patients; Authors: Ron Gimbel, PhD; Joel Williams, PhD; Liwei Chen, MD, PhD; Cheryl Dye, PhD; Karen Edwards, Med.; Jeanette Little, MS, Terry Newton, MD

Poster delivered for American Academy of Health Behavior Scientific Meeting, Tucson, AZ; *Improving Diabetes Self-management through Messages Tailored to Patient Activation levels*; delivered 19-22 MAR 2017.

Presentation approved for the Military Health System Research Symposium, Kissimmee, FL; *Mobile Health Apps in the Medical Theater of Operation – Exploring the Usability of a Mobile Application for Improving Chronic Disease Self-management in the Patient Centered Medical Home Environment*. AUG 2017.

- **Website(s) or other Internet site(s)** – Nothing to report during this reporting period
- **Technologies or techniques**

The Telemedicine and Advanced Technology Research Center (TATRC), in partnership with Clemson University, are expanding the capabilities of the Mobile Health Care Environment (MHCE) to support enhanced communication and patient self-management behaviors in Type 2 diabetes care.

In fiscal year 2017/18 the TATRC-Clemson University accomplished the following:

- Development of initial prototype mCare application interface
 - Pilot testing of the initial prototype mCare application interface
 - User focus group usability testing of the mCare application interface
 - Data analysis of the SUS scoring on the user focus group testing
 - Refinement of the mCare application interface based on pilot and focus group usability testing (PowerPoint slides are attached)
 - Development of the mCare provider portal interface
 - Pilot testing of the mCare provider portal interface
 - Refinement of the mCare provider portal interface (PowerPoint slides are attached)
 - Establishment of normalized fitness tracker data using the Validic API to the MHCE system.
 - Ordering and distribution of the end user equipment to the two study site Research Associates.
 - Development of training manuals and end user guides
 - Training to the Research Associates.
 - Technical Support during the enrollment and initial data collection period (ongoing)
- **Inventions, patent applications, and/or licenses** (Not applicable)
 - **Other Products** (Not applicable)

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

- What individuals have worked on the project?

Name:	<i>Ronald W. Gimbel, PhD</i>
Project Role:	<i>Principal Investigator</i>
<u>Nearest person month worked:</u>	2.4
Contribution to Project:	<i>Dr. Gimbel has performed work as the Principal Investigator.</i>
Name:	<i>Karen Edwards</i>
Project Role:	<i>Research Associate</i>
<u>Nearest person month worked:</u>	12
Contribution to Project:	<i>Karen oversees all aspects of the study along with Dr. Gimbel.</i>
Name:	<i>Jennie Moss</i>
Project Role:	<i>Senior Research Associate</i>
<u>Nearest person month worked:</u>	12
Contribution to Project:	<i>Jennie oversees all aspects of the study located at Nellis AFB.</i>
Name:	<i>Kristy Crawford</i>
Project Role:	<i>Research Associate</i>
<u>Nearest person month worked:</u>	12
Contribution to Project:	<i>Kristy oversees all aspects of the study located at Nellis AFB.</i>

Name:	<i>Marie Rempola Hing</i>
Project Role:	<i>Senior Research Associate</i>
<u>Nearest person month worked:</u>	<u>12</u>
Contribution to Project:	<i>Marie oversees all aspects of the study at Madigan AMC.</i>

- **Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?** Nothing to report for this reporting period
- **What other organizations were involved as partners?**
 - **Organization Name:** TATRC
 - **Location of Organization:** Building 38711, Fort Gordon, Georgia 30905-5650
 - **Partner's contribution to the project**
 - **Financial support** – n/a
 - **In-kind support** – n/a
 - **Facilities** – n/a
 - **Collaboration** – TATRC's staff is working with Clemson University project staff on the project.
 - **Personnel exchanges** – n/a
 - **Other** – n/a

8. SPECIAL REPORTING REQUIREMENTS

- **COLLABORATIVE AWARDS:** No applicable for this reporting period
- **QUAD CHART:** see Appendix

9. APPENDICES:

AMSUS Poster #1

AMSUS Poster #2

American Academy of Health Behavior Scientific Meeting Poster


Presentation Outline approved for the Military Health System Research Symposium

Manuscript copy

Power Point sample slides of the mCare patient product

PowerPoint samples slides of the mCare provider portal interface

Quad Chart thru 7/31/17



fluence patient use and clinical workflow.

Mobile health (mHealth) technology in the care of patients with diabetes has demonstrated varying degrees of effectiveness. In this project we seek to incorporate the most promising capabilities of mHealth technology in a Patient-Centered Medical Home (PCMH) to support the activation of patients with Type 2 diabetes. We anticipate that greater patient activation will lead to improved self-care behaviors and outcomes (clinical and health service).

SAFETY ALGORITHMS

Blood Sugar Clinical thresholds:

Readings above 250 mg/dl:
Contact your doctor if your blood sugar is over 250 mg/dl for 24 hours.

Readings above 300 mg/dl:
If your blood sugar is greater than 300 mg/dl, drink water, be active, and check your blood sugar before meals and at bedtime until it is less than 200. If it does not go down contact your doctor.

Readings less than 100 mg/dl:
If your bedtime blood sugar is less than 100 mg/dl, have a snack such as a glass of milk or a half a sandwich.

Readings less than 70 mg/dl:
If your blood sugar is less than 70 mg/dl, treat with 3 glucose tablets, ½ glass juice, or other snack containing sugar. Recheck blood sugar after 15 minutes. If still less than 70, eat another 3 glucose tablets and recheck after 15 minutes. Repeat until blood sugar is above 70 mg/dl. After your blood sugar is back above 70 mg/dl, eat a small snack if your next meal is more than ½ hour away.

Blood Pressure Safety Thresholds:

For systolic blood pressure above 180: First Send: If you have chest pain, shortness of breath, back pain, numbness/weakness, change in vision or difficulty speaking, call 911 immediately.

Then send: For systolic blood pressure above 180, wait 10 minutes at rest and recheck. If it is still above 180 systolic (the top number) and you do not have any other symptoms, contact your doctor.


For diastolic above 110:
For diastolic above 110 send this first: If you have chest pain, shortness of breath, back pain, numbness/weakness, change in vision or difficulty speaking, call 911 immediately.

Then send this: For Diastolic blood pressure above 110 wait ten minutes at rest and check again. If it is still above 100 diastolic (the bottom number) and you do not have any other symptoms, contact your doctor.


Weight Clinical thresholds:

Weight gain of greater than 5 pounds in one week:
If you note a weight gain of greater than 5 pounds in one week, or you notice increased leg swelling or new or increased shortness of breath, call your doctor.

Activities (SA
was advised
team. Algori
are also inclu



DISCUSSION
Research ben
to advance th
care, improv
care) through
cus on integr
flow, as well



ABSTRACT:

- Telemedicine and Advanced Technology Research Center (TATRC) - Clemson University Partnership
- Expanded the Mobile Health Care Environment (MHCE) system capabilities for Type 2 diabetic self-management behaviors and provider communications
- Research project aimed at enhancing patient activation in the patient centered medical home (PCMH) environment

views, and other biomedical manuscripts.

SAFETY ALGORITHMS

Blood Sugar Clinical thresholds:

Readings above 250 mg/dl:

Contact your doctor if your blood sugar is over 250 mg/dl for 24 hours.

Readings above 300 mg/dl:

If your blood sugar is greater than 300 mg/dl, drink water, be active, and check your blood sugar before meals and at bedtime until it is less than 200. If it does not go down contact your doctor.

Readings less than 100 mg/dl:

If your bedtime blood sugar is less than 100 mg/dl, have a snack such as a glass of milk or a half a sandwich.

Readings less than 70 mg/dl:

If your blood sugar is less than 70 mg/dl, treat with 3 glucose tablets, ½ glass juice, or other snack containing sugar. Recheck blood sugar after 15 minutes. If still less than 70, eat another 3 glucose tablets and recheck after 15 minutes. Repeat until blood sugar is above 70 mg/dl. After your blood sugar is back above 70 mg/dl, eat a small snack if your next meal is more than ½ hour away.

Blood Pressure Safety Thresholds:

For systolic blood pressure above 180: First Send: If you have chest pain, shortness of breath, back pain, numbness/weakness, change in vision or difficulty speaking, call 911 immediately.

Then send: For systolic blood pressure above 180, wait 10 minutes at rest and recheck. If it is still above 180 systolic (the top number) and you do not have any other symptoms, contact your doctor.

For diastolic above 110:

For diastolic above 110 send this first: If you have chest pain, shortness of breath, back pain, numbness/weakness, change in vision or difficulty speaking, call 911 immediately.

Then send this: For Diastolic blood pressure above 110 wait ten minutes at rest and check again. If it is still above 100 diastolic (the bottom number) and you do not have any other symptoms, contact your doctor.

Weight Clinical thresholds:

Weight gain of greater than 5 pounds in one week:

If you note a weight gain of greater than 5 pounds in one week, or you notice increased leg swelling or new or increased shortness of breath, call your doctor.

The project included three major components: data collection, data analysis, and data visualization.

- Primary data collection issues included DoD regulatory requirements, data mapping from peripheral devices, patient entry of free-text data, and agreements with wireless communication providers.







FIGURE 1




METHODS

A multi-stage message development process was guided by patient activation levels and theoretical constructs relevant to those levels.

Step One: Researchers developed messages tailored for each of the four PAM® score levels and five diabetes self-care behaviors measured by the SDSCA: nutrition, physical activity, foot care, smoking cessation, and glucose monitoring.

Step Two: Messages were assessed by two-person teams for content accuracy, reading level, and message appropriateness for PAM® level. Messages were then revised based on this review process. A clinical advisory team consisting of clinicians and researchers reviewed and approved the messages for clinical relevance and content.



Step Three: A pre-test (n=21) of the messages with a similar population as the messages administered. Each person was provided with a random sample of five messages and asked four close-ended questions and one open-ended question about the messages.

Step Four: Researchers reviewed PAM® level specific messages and rated the messages on appropriateness for PAM® level, behavioral theory construct, reading level, and based on this review process.

RESULTS AND DISCUSSION

The multi-stage message development process allowed for triangulation review of appropriate messages. At each step in the process, messages were reviewed for feedback (Stages Two and Four) or by potential study participant feedback (Step Three). The produced messages with a high degree of acceptability by people very similar to the target population. Test participants found the messages to be encouraging, useful, applicable, and of credibility with regard to application of the theoretical constructs within the messages complementary to the counseling activities of health educators and others, and to replace it.”⁷

Exploring the Usability of a Mobile Application for Improving Chronic Disease Self-Management in the Patient Centered Medical Home Environment

Joel E. Williams, MPH, Ph.D.
Associate Professor of Public Health Sciences
joel2@clemson.edu

DISCLAIMER

- This material is based upon work supported by the U.S. Army Medical Research Acquisition Activity under Contract No. W81XWH-15-C-0070. Any opinions, findings and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the U.S. Army Medical Research Acquisition Activity.

Presentation Outline

- Provide an overview of larger study
- Describe Phase 1 research:
 - Background
 - Objectives
 - Methods
 - Findings

Research Group Structure

- Government, Academia, Industry partnership
 - Development Advisory Board
 - Clinical Advisory Team
 - Intervention Team



Acknowledgements – Phase 1 Team

- Sarah F. Griffin, MPH, Ph.D. ¹
- Karen W. Edwards, MEd ¹
- Jennie B. Moss, RN ^{1,2}
- Marie E. Hing, MS ^{1,3}
- Jeanette R. Little, MS ⁴
- Ronald Gimbel, Ph.D. ¹ (Principal Investigator)

¹ Clemson University, Clemson, SC

² Mike O'Callaghan Federal Medical Center, Las Vegas, NV

³ Madigan Army Medical Center, Tacoma, WA)

⁴ US Army Telemedicine and Advanced Technology Research Center, Fort Gordon, GA


Background – Research Study

- Multi-site Military Health System study
 - U.S. Department of Defense W81XWH-15-C-0070 Gimbel (PI)
“Enhancing mHealth Technology in the PCMH Environment to Activate Chronic Care Patients” 08/15 – 01/18
 - Mike O'Callaghan Federal Medical Center (NV)
 - Madigan Army Medical Center (WA)
- Phase 1: Utilized a participatory approach in guiding adaptations of a health system application, called mCare.
- Phase 2: Feasibility study where patients are randomized to biosensors only vs. biosensors + mCare interface + tailored messaging

Background – Biosensors & mCare

- Glucometer: My GlucoHealth
- Blood Pressure Monitor: A&D
- Weight Scale: A&D
- Activity Monitor:
 - Fitbit Charge (wrist)





Menu

Announcements
Health Tips
Safety Alerts

Target Ranges



Alert Grid

Alert URL	Alert Text	Alert Date	View	Action
	Robert Chewning has entered a Blood Glucose > 300 mg/dL	10/06/2016	View	Archive
	Mabel Cooper has entered Blood Glucose > 250 mg/dL for > 24 hrs.	10/06/2016	View	Archive
	Lois Goldstein has entered a Blood Glucose < 100 mg/dL at bedtime	10/06/2016	View	Archive
	Kim Stargell has entered a Blood Glucose < 70 mg/dL	10/06/2016	View	Archive
	Robert Chewning has entered a Systolic BP > 180 mmHG	10/06/2016	View	Archive
	Robert Chewning has entered a Diastolic BP > 110 mmHG	10/06/2016	View	Archive
	Mabel Cooper has had a Weight gain > 5 lbs. in 1 week	10/06/2016	View	Archive
	Lois Goldstein has had No Activity Readings in last 48 hrs.	10/06/2016	View	Archive

Diabetics: Active Patients

Name	Last Sych	Glucose	BP	Activity	Weight	
Robert Chewning	10/06/2016	320 mg/dL	182/112	17,251 steps	235 lbs	View
Mabel Cooper	10/06/2016	306 mg/dL	122/82	5,048 steps	146 lbs	View
Lois Goldstein	10/06/2016	95 mg/dL	118/70	0 steps	163 lbs	View
Kim Stargell	10/06/2016	68 mg/dL	116/78	939 steps	174 lbs	View

Phase 1 Objective & Methods

- Utilize a participatory approach to engage Type 2 diabetics, and the clinicians who treat them, in guiding adaptations of the mCare interface to be used in Phase 2 of this study.

- Mixed methods approach:

Quantitative

- Patient task observations
- Open-ended questions during usability testing to capture Patient thoughts

Quantitative

- Patient + Clinician feedback
- Patient task completion notes

Methods – Design

- Science Panel on Interactive Communication and Health (Robinson et al., 1998)
- International Organization for Standardization (ISO) 9241-11 Usability framework : usability definitions and evaluation metrics
- Georgsson & Staggers (2016) model
- Mixed methods evaluation (QUAN + QUAL)

Methods – Context of Use (Patients)

- **Users:** Describe relevant characteristics of the users.
- **Tasks:** Specific activities (device interactions) that will be required of users.
- **Equipment:** Hardware, software and peripheral devices (My GlucoHealth glucometer, A&D blood pressure monitor, A&D weight scale, Fitbit Charge).
- **Environment:** Relevant characteristics of the testing environment including: technical (WiFi), and physical (testing space).

Methods – Usability Metrics (Patients)

- **Effectiveness:** Extent to which the user can achieve a goal with accuracy and completeness.
 - *(1) the degree of task completion AND*
 - *(2) total number of errors per task*
- **Efficiency:** Level of effort and resource usage which is required by the user in order to achieve a goal in relation to accuracy and completeness.
 - *Timed each individual task and compute average time for each task across users.*
- **Satisfaction:** Extent to which users are free from discomfort, and their attitudes towards the use of the product.
 - *Single Ease Question (SEQ – Sauro & Dumas, 2009)*
 - *Open-ended questions (developed by Phase 1 Team)*
 - *System Usability Scale (SUS – Sauro & Lewis, 2012)*

Methods – Usability Tasks (Patients)

- Login and System navigation
- Goal setting
- Specific tasks for each biosensor:
 - Blood pressure (manual entry, Bluetooth sync, graphs)
 - Blood sugar (manual entry, graphs)
 - Weight scale (manual entry, Bluetooth sync, graphs)
 - Fitbit (graph interpretation only)

Methods – Usability (Clinicians)

- Focus groups
 - Physicians and Clinical Staff
 - Recorded
 - Field notes
- Feedback on:
 - Perceived usability for patient
 - Workflow issues / concerns
 - Message content - system alerts, patient reminders
 - Backend portal (doctor's view)

Findings – Patients

- All diabetics indicated that mCare would “help them manage their diabetes” and give their healthcare provider a “better report of their health.” As expected, patients rated navigation tasks as less difficult and peripheral device tasks, e.g., syncing, as more difficult.
- Diabetics committed fewer errors with basic navigation tasks and more with peripheral devices. Further, a higher proportion of diabetics made errors with external device tasks.
- Diabetics and clinicians alike suggested minor changes regarding the look and function of the application, e.g., adding more color and contrast, making buttons larger.

White (n=14, 70%) African-American (n=2, 10%) Native Hawaiian/Pacific Islander (n=2, 10%) Asian (n=2, 10%)

Male (n=14, 70%) Female (n=6, 30%)

Findings – Patients

- Comfort with using apps or technology (5 point scale)
 - *Means, All: 3.3 (Madigan: 2.8 , Nellis: 3.8)*
- Task completion times, *Range: 3.29 sec - 249.45 sec*
- Single Ease Use Question (7 point scale)
 - *Range of Means: 2.8 – 6.6*
- System Usability Scale (SUS)
 - SUS overall, *Mean (SD): 83.8 (14.9) = A+*
 - Usability Sub Factor, *Mean (SD): 86 (13.4) = A+*
 - Learnability Sub Factor, *Mean (SD): 75 (27.5) = A-*

Findings – Clinicians

- Overall, Clinicians were pleased with the mCare system and optimistic about both the backend portal (“doctor’s view”) and application utility for patients.
- Clinicians also had additional suggestions specifically related to alerts, e.g., parameters for alerts sent to patients, color coding alerts for ease of clinician review.
- Patient and Clinician suggestions were reviewed and incorporated as adaptations by the technology team as allowed by system constraints.

19 Physicians (2 DO, 17 MD) ; 1-19 years experience

14 Clinical staff (FNP, nurses, disease managers, pharmacists) ; 4-39 years experience

TATRC Recommendations

- 29 specific recommendations were made to TATRC
 - Only 4 could NOT be completed
 - additional colors (limited colors available)
 - remove signal and refresh buttons (part of the base system)
 - alert icons (limited icon choices available)
 - automatic syncing of devices (system cannot do this)
 - >86% of recommended changes WERE made
 - Usability testing is CRITICAL in understanding the needs of end users and will provide a more meaningful interface for Phase 2 participants

Major Recommendations/Changes

Patients

- increase size of icons
- separate glucose graphs (no overlay)
- increase font size
- allow past dates for manual entry

Clinicians

- change default blood glucose entry to unclassified
- define "after meal" glucose as ≥ 120 minutes post meal
- simplify safety alerts (with Clinical Advisory Team)
- add icons beside safety alerts
- add patient target ranges on graphs

References

- Gimbel R, Shi L, Williams JE, Dye CJ, Chen L, Crawford P, Shry EA, Griffin SF, Jones KO, Sherrill WW, Truong K, Little JR, Edwards KW, Hing M, Moss JB. Enhancing mHealth Technology in the Patient-Centered Medical Home Environment to Activate Patients With Type 2 Diabetes: A Multisite Feasibility Study Protocol. *JMIR Res Protoc* 2017;6(3):e38. DOI: 10.2196/resprot.6993
- Mattias Georgsson & Nancy Staggers. (2016). Quantifying usability: An evaluation of a diabetes mHealth system on effectiveness, efficiency, and satisfaction metrics with associated user characteristics. *J Am Med Inform Assoc*, 23, 5-11. doi:10.1093/jamia/ocv099.
- Robinson TN, Patrick K, Eng TR, Gufstason D. (1998). An evidence-based approach to interactive health communication: a challenge to medicine in the information age. *JAMA*, 280(14):1264-1269.
- Sauro, J., & Dumas, J. S. (2009). Comparison of three one-question, post-task usability questionnaires. In *Proceedings of CHI 2009* (pp. 1599–1608). Boston, MA: ACM.
- Sauro, J., & Lewis, J. R (2012). Quantifying the user experience: Practical statistics for user research. Morgan Kaufmann, Waltham MA, USA.

Protocol

Enhancing mHealth Technology in the Patient-Centered Medical Home Environment to Activate Patients With Type 2 Diabetes: A Multisite Feasibility Study Protocol

Ronald Gimbel¹, PhD; Lu Shi¹, PhD; Joel E Williams¹, MPH, PhD; Cheryl J Dye¹, MA, PhD; Liwei Chen¹, MD, PhD; Paul Crawford², MD; Eric A Shry³, MD; Sarah F Griffin¹, MPH, PhD; Karyn O Jones¹, MA, PhD; Windsor W Sherrill¹, MBA, MHA, PhD; Khoa Truong¹, MA, MPhil, PhD; Jeanette R Little⁴, MS; Karen W Edwards¹, MEd; Marie Hing¹, MS; Jennie B Moss¹, BSN, MS

¹Department of Public Health Sciences, Clemson University, Clemson, SC, United States

²Nellis Family Medicine Residency Program, Mike O'Callaghan Federal Hospital, Las Vegas, NV, United States

³Madigan Army Medical Center, Tacoma, WA, United States

⁴MHIC Laboratory Lead, Telemedicine & Advanced Technology Research Center, U.S. Army Medical Research & Materials Command, Fort Gordon, GA, United States

Corresponding Author:

Ronald Gimbel, PhD

Department of Public Health Sciences

Clemson University

501 Edwards Hall

Clemson, SC, 29634-0745

United States

Phone: 1 864 656 1969

Fax: 1 864 656 6227

Email: rgimbel@clemson.edu

Abstract

Background: The potential of mHealth technologies in the care of patients with diabetes and other chronic conditions has captured the attention of clinicians and researchers. Efforts to date have incorporated a variety of tools and techniques, including Web-based portals, short message service (SMS) text messaging, remote collection of biometric data, electronic coaching, electronic-based health education, secure email communication between visits, and electronic collection of lifestyle and quality-of-life surveys. Each of these tools, used alone or in combination, have demonstrated varying degrees of effectiveness. Some of the more promising results have been demonstrated using regular collection of biometric devices, SMS text messaging, secure email communication with clinical teams, and regular reporting of quality-of-life variables. In this study, we seek to incorporate several of the most promising mHealth capabilities in a patient-centered medical home (PCMH) workflow.

Objective: We aim to address underlying technology needs and gaps related to the use of mHealth technology and the activation of patients living with type 2 diabetes. Stated differently, we enable supporting technologies while seeking to influence patient activation and self-care activities.

Methods: This is a multisite phased study, conducted within the US Military Health System, that includes a user-centered design phase and a PCMH-based feasibility trial. In phase 1, we will assess both patient and provider preferences regarding the enhancement of the enabling technology capabilities for type 2 diabetes chronic care management. Phase 2 research will be a single-blinded 12-month feasibility study that incorporates randomization principles. Phase 2 research will seek to improve patient activation and self-care activities through the use of the Mobile Health Care Environment with tailored behavioral messaging. The primary outcome measure is the Patient Activation Measure scores. Secondary outcome measures are Summary of Diabetes Self-care Activities Measure scores, clinical measures, comorbid conditions, health services resource consumption, and technology system usage statistics.

Results: We have completed phase 1 data collection. Formal analysis of phase 1 data has not been completed. We have obtained institutional review board approval and began phase 1 research in late fall 2016.

Conclusions: The study hypotheses suggest that patients can, and will, improve their activation in chronic care management. Improved activation should translate into improved diabetes self-care. Expected benefits of this research to the scientific community and health care services include improved understanding of how to leverage mHealth technology to activate patients living with type 2 diabetes in self-management behaviors. The research will shed light on implementation strategies in integrating mHealth into the clinical workflow of the PCMH setting.

Trial Registration: ClinicalTrials.gov NCT02949037. <https://clinicaltrials.gov/ct2/show/NCT02949037>. (Archived by WebCite at <http://www.webcitation.org/6oRyDzqei>)

(*JMIR Res Protoc* 2017;6(3):e38) doi:[10.2196/resprot.6993](https://doi.org/10.2196/resprot.6993)

KEYWORDS

mHealth; diabetes mellitus; patient activation; patient-centered medical home; patient centered care; eHealth; health information

Introduction

Diabetes mellitus is a chronic disease with high rates of disability, impaired quality of life, and premature death [1-4]. The prevalence of type 2 diabetes is increasing at an alarming rate in the United States; in 2013, the estimated number of patients was between 20 million and 27 million, or about 7% to 10% of the adult population [2,3]. Research suggests that, if current trends continue, diabetes will be diagnosed in 1 in 3 adults in the United States by 2050 [4,5]. Diabetes is the leading cause of blindness, nontraumatic amputations, and adult renal failure, and reduces life expectancy by 5-10 years [2]. The individual symptom burden (eg, chronic pain, neuropathy, depression, and physical disability) is substantial and significantly increases in the older adult population [1]. In the United States, an average individual with diabetes incurs medical expenditures of about US \$13,700 a year, of which about US \$7900 is attributable to diabetes [4]. This represents an expenditure about 2.3 times greater than that for a diabetes-free individual [4].

Numerous primary care-based efforts have been aimed at reducing both the disease burden on individuals and the cost of diabetes care. A contemporary strategy is the management of patients with diabetes within the context of the patient-centered medical home (PCMH) setting. A key PCMH principle is the appropriate use of information technology to support optimal patient care, performance measurement, patient education, and enhanced communication [6]. Several case studies from various US health systems show the benefit of the PCMH model to improved diabetes care [7]. There is published evidence on the positive impact of PCMH-based care in psychosocial outcomes of patients with diabetes [8].

The potential of mHealth technologies in the care of patients with diabetes and other chronic conditions has captured the attention of clinicians and researchers. Efforts to date have incorporated a variety of tools and techniques, including Web-based portals [9-11], short message service (SMS) text messaging [9,12-14], remote collection of biometric data [12,15], electronic coaching [14], electronic-based health education [13], secure email communication between visits [16-18], electronic collection of lifestyle and quality-of-life surveys, and personal health records (PHRs). Each of these tools, used alone or in combination, has demonstrated varying degrees of effectiveness. Some of the more promising results have been demonstrated using regular collection of biometric

devices (eg, glucometers, activity monitors) [12], SMS text messaging [12-14], secure email communication with clinicians and clinical teams [9,16,17], and regular reporting of quality-of-life variables aligned with decision support. In this study, we seek to incorporate many of the most promising mHealth capabilities in a PCMH workflow led by a clinical advisory team. We aim to address underlying technology needs and gaps related to the use of mHealth technologies and the activation of patients with type 2 diabetes.

The Concept of Patient Activation

Self-management for patients with type 2 diabetes and other chronic conditions includes following complex treatment regimens, monitoring chronic conditions, and making lifestyle changes [19-22]. The chronic care model suggests that activated patients are better able to function in the role of self-manager [21,23]. An activated patient has the motivation, confidence, and skills necessary to enact behavioral changes and make health-related decisions [24-27]. These patients ask questions and collaborate with their health care provider [19,26-28]. Research shows that activated patients have more positive clinical outcomes, are more likely to receive preventive care, and have lower health care-related costs [24,26,29].

A recommended strategy in patient activation is the concept of “preactivating” patients prior to clinical encounters [20,30]. The concept incorporates active targeted communication and follow-up from the health care team [30]. Interventions to include educational programs [31], care coaching [32], and motivational interviewing [33] have been attempted to improve patient activation with varied success [34]. However, these efforts have infrequently been tailored to potential intrinsic differences in how the patients approach their disease. Theoretically, research suggests that patient activation can be increased [19,35-37]. Conceptualizing activation as a dynamic variable allows researchers to target this motivating factor that can potentially influence health behaviors [21,24,38,39].

Previous Research on Patient Portals, Personal Health Records, Patient Activation, and Improved Outcomes

Federal legislation and movement toward patient centeredness in the United States has fueled interest in providing patients with access to their health information, enhanced communication with clinical environments, and greater emphasis on self-care [40-43]. Early research on portal and PHR use and patient activation provided mixed results. Several studies reported a positive significant relationship between use of portals and

PHRs and activation of patients [41,43-45], while other studies did not realize a significant finding [40,46,47]. The design of these published studies prevented any in-depth inquiry into why (or not) portal and PHR use influenced patient activation. Their authors posited a variety of possible factors, including the target patient population [44], time since severe diagnosis or symptoms and episodes [46,47], and patient age (activation being higher in adults than in children) [45]. One study suggested that tailoring a portal or PHR intervention to the patient activation level may optimize intervention efficiency [43].

Early research on increased activation and improved clinical outcomes using patient portal and PHR-based interventions has also provided mixed results. Several studies demonstrated a relationship between increased patient activation and improved intermediate clinical outcomes (eg, hypertension, smoking, body mass index, and glycated hemoglobin [HbA_{1c}]) [48], while a major study did not record a significant finding regarding the same outcomes [42]. It is noteworthy that these early studies did not provide substantial detail on design issues related to the portal or PHR, or whether the intervention included behavioral reinforcement.

User-Centered Design

Design science will inform our development and testing [49,50]. User-centered design will guide development, following participatory design methods to understand more specifically how patients experience diabetes on a daily basis, what clinicians need to know from patients, and how to create a shared communication system for better decision making [51]. Consistent with the guidelines set forth by the Science Panel on Interactive Communication and Health [52], our evaluation design will incorporate the 3 elements of formative, process, and outcome evaluation. Methods include (1) clinician focus groups and in-depth patient interviews to define key knowledge variables that are personally and clinically relevant, (2) iterative usability testing with patients, and (3) iterative observations of the system in clinical settings [53].

Military Health System: An Overview

The US Military Health System (MHS) is a large integrated health system that cares for about 9.39 million beneficiaries through its TRICARE insurance product and its substantial direct care system consisting of tertiary facilities, community hospitals, and clinics globally. Nearly 35% of its beneficiary pool are active duty members and their dependents, with a larger population (about 56%) being retirees and their beneficiaries [54]. The MHS direct care system is robust. Facilities are accredited by the Joint Commission (formerly the Joint Commission on Accreditation of Healthcare Organizations), and the MHS operates a dedicated educational infrastructure to support medical and nursing education programs [54]. The MHS has a connecting health information technology infrastructure to support clinical care and clinical operations.

Hypotheses

Hypothesis 1: User-centered design will allow developers to create a patient-centered interactive and tailored mobile technology for use in the PCMH setting.

Hypothesis 2: The use of interactive and tailored mobile technology, the Mobile Health Care Environment (MHCE), employed in the PCMH setting will increase the activation of patients with chronic type 2 diabetes.

Hypothesis 3: The use of interactive and tailored mobile technology in a PCMH setting will increase diabetes self-care activities.

Hypothesis 4: Patients who engage at a higher rate with the interactive and tailored mobile technology in a PCMH setting will realize greater improvement in clinical measures.

The primary goal of the research is to enhance patient activation levels and improve self-management of type 2 diabetes through the use of the MHCE in the PCMH setting. While there are published studies aimed at improving the activation and care of patients with diabetes in the United States, to our knowledge, no study has sought to enhance care of patients with diabetes using a fully comprehensive and adaptable MHCE-like system. We seek to demonstrate improvement in patient activation measured by the Patient Activation Measure (PAM) instrument [21,55]. We believe that, in improving their activation, patients will also realize an improvement in diabetes self-care activities measured by their Summary of Diabetes Self-Care Activities (SDSCA) [56] scores.

Methods

Trial Design

This is a multisite, phased study conducted within the MHS that includes a user-centered design phase and a PCMH-based feasibility trial. In phase 1, we will assess both patient and provider preferences regarding the enhancement of the MHCE technology capabilities for type 2 diabetes chronic care management. The phase 2 research will be a single-blinded (patients only) 12-month feasibility study that will incorporate randomization principles. We will employ a 1:1 allocation ratio between intervention and control.

Inclusion and Exclusion Criteria

Inclusion criteria for patient participation in phase 1 or 2 research are the following: (1) men and women aged 18 years or older, (2) able to understand and read English, (3) enrolled for primary care to one of the target PCMH sites, and (4) having a diagnosis of type 2 diabetes. Additionally, with respect to phase 2 patients, we will seek to recruit a maximum of 120 (per site), with a distribution of patients with PAM levels 1-4, a sample representative of the patients enrolled in the PCMH. We did not derive the 120 per site recruitment numbers from power calculations, but deemed them to be sufficient. Finally, participants for phase 2 must be available for a 12-month study.

Inclusion criteria for clinician participation in phase 1 or 2 research are the following: (1) being a physician, physician assistant, nurse practitioner, or nurse employed at the target site, and (2) providing care for patients with type 2 diabetes.

Exclusion criteria for patient participation in phase 1 or 2 research are the following: (1) pregnant women, (2) non-English-speaking patients, (3) receiving hospice care, (4) having active cancer and receiving treatment with chemotherapy

or radiation therapy, (5) taking warfarin, (6) recipient of gastric bypass or similar procedure, (7) having a diagnosis of uncontrolled hypothyroidism, (8) having known Cushing syndrome, (9) being treated with oral steroids, (10) having known liver disease, (11) having a current diagnosis of cognitive impairments that would interfere with use of technology, (12) having congestive heart failure, in New York Heart Association functional class 3 or 4, and (13) unable to use a mobile device due to cognitive or physical impairments during initial screening. We exclude pregnant women because they require careful monitoring due to potential medical complications for the woman and unborn child. While some mHealth studies seek to include additional exclusions based on age, educational level, or technical literacy, our research team rejected adding any additional exclusion criteria beyond the 13 listed above. We purposely seek the “average” patient with type 2 diabetes in the target population. Feedback from our clinician investigators and research staff at our clinical sites is encouraging that these patients will be capable of using the intervention technology.

Exclusion criteria for clinician participation in phase 1 or 2 research are the following: (1) not affiliated with the target site, and (2) not providing care for patients with type 2 diabetes.

Participant Enrollment

We will recruit patients via review of the PCMH clinic schedule, referrals from providers, distributed posters and fliers, and population health databases. Potential participants will be prescreened through verification of the inclusion and exclusion criteria based on a medical record review. Interested participants will be scheduled for a screening visit with study staff to provide informed consent and be administered the PAM instrument. Patients' PAM scores will place them in a stratified group, where they will be randomly allocated.

Clinicians practicing in the respective PCMH sites will be invited to participate by word of mouth from the site's principal investigator; this is a convenience sample. The clinician participants who would like to participate in the study will meet with the senior research associate to review the minimal-risk information sheet to be included in the study. For phase 2, clinicians will sign an informed consent form. The clinician participants will not be blinded in the study, nor allocated to intervention or control groups.

Setting and Site Selection

We seek to purposefully assess the MHCE implementation for diabetes care in 2 distinctly different PCMH environments and geographic locations. The risks of attracting very different populations are mitigated by rather comprehensive inclusion and exclusion criteria, which will ensure similarity regarding patient acuity. The patient base includes those on active duty,


retirees, and dependents who have typically spent years in the military and have been stationed at various locations. Both of the selected facilities are federal facilities and operated by the MHS.

Madigan Army Medical Center, the US Army's second largest military treatment facility located in Tacoma, Washington, is a tertiary facility with a level II trauma center and robust graduate medical education programs. They serve a patient base of approximately 118,000 patients; about 7500 (or >6%) are living with type 2 diabetes. Of the diabetes population, about 15% are active duty members or their dependents, and about 85% are retirees and their dependents. Over half of the patients with diabetes are 57-76 years of age. The study location within the medical center is a PCMH managed by the Department of Internal Medicine. There are approximately 14,300 patients enrolled in this PCMH supported by a staff of 77 (12 staff physicians; 8 residents) responsible for their care.

Mike O'Callaghan Federal Medical Center is a federal facility in the greater Las Vegas, Nevada area, that serves approximately 47,000 patients; about 4500 (or >9%) are living with type 2 diabetes. Of the diabetes population, about 4% are active duty members or their dependents, and about 96% are retirees and their dependents. Over 72% of the patients with diabetes are in their 60s or older. The study location within the medical center is a PCMH managed by the Department of Family Medicine. There are approximately 7500 patients enrolled in this PCMH supported by a staff of 62 (9 staff physicians; 26 residents) responsible for their care.

Description of the Mobile Health Care Environment

The US Department of Defense (DoD) MHCE system is a secure health information system designed to support health services delivery and mHealth. The MHCE meets all physical and information security mandates, as prescribed by federal law and DoD regulation, for the protection of personal health information and personally identifiable information. The MHCE was developed by the DoD Telemedicine and Advanced Technology Research Center as a platform to support mHealth. Its first major application was to support patient engagement for wounded warriors rehabilitating in their communities. In the study, soldiers on average responded to $\geq 60\%$ of weekly questionnaires related to behavioral health challenges, posttraumatic stress, or traumatic brain injury [57]. Our study is Telemedicine and Advanced Technology Research Center's second major application. The MHCE is designed to remotely support patients by sending automated reminders, announcements, wellness tips, alerts, and status questionnaires. Figure 1 is a visual example of the graphical user interface that patients will see when accessing the MHCE. In this study, we enhance MHCE capabilities in several ways.

Figure 1. Mobile Health Care Environment home screen (patient view). BP: blood pressure.

Intervention Overview

Our intervention is based on an enhanced MHCE in several ways. First, we add the capability to include collection and visualization of data from Bluetooth-enabled medical devices.

This includes mapping data from device output into the MHCE, developing data visualization appropriate for mHealth and clinical care (eg, graphing outcomes, temporal trend patterns), migrating data in an analysis cell, and developing decision-support algorithms that signal safety alerts and need

for behavioral reinforcement. Devices used in this study include a scale, glucometer, blood pressure reader, and activity monitor. Second, we expand the capacity of the MHCE analysis cell to manage large amounts of data and to conduct both routine reports and research applications. Third, we add patient activation and associated measurement instruments for capturing baseline and ongoing changes to patient activation. Fourth, we expand the MHCE messaging platform that research associates, and later clinical support staff, can use to send tailored behavioral messaging to patients in an effort to influence greater activation and reinforce positive behavior.

The MHCE can be accessed by mobile phones or tablets that use either an IOS or Android platform. The MHCE requires Internet access for patients to sync data from devices (addressed above) to the MHCE backend portal, to receive tailored behavioral messages, or to use other functions. During the study, patients will additionally receive SMS messages with hyperlinks to a separate secure information system platform used for administration and analysis of PAM and SDSCA instruments. MHCE activity, or lack thereof, will be monitored by senior research associates, who can prompt patients via tailored behavioral messages or direct contact.

Tailored Behavioral Messaging

A primary component of the MHCE system is tailored behavioral messaging. Tailored behavioral messages are more likely than generic messages to facilitate health behavior change when they are aligned with individuals' beliefs, lifestyle, demographics, social norms, or interests [58-60].

In this study, the research team has developed behavioral messages tailored for each of the 4 PAM score levels; in total we have developed 360 messages. The messages fall within 9 functional areas common to diabetes care: nutrition, home monitoring, physical activity, blood pressure, foot care, medications, smoking, glucose control, and general behavioral reinforcement. The messages are consistent with general concepts and goals of self-management behaviors consistent with the DoD-Veterans Affairs clinical practice guideline for type 2 diabetes and the SDSCA survey instrument.

Since different PAM levels require different strategies, we addressed varying needs through a combination of applied constructs. Specifically, level 1 messages must address the emotional state of feeling overwhelmed and passive with an emphasis on the importance of taking action. To address the needs of PAM level 1 patients, we use constructs from social

networks and social support theory [61], specifically that of emotional support that emphasizes expressions of empathy and caring. We encourage a sense of hope by expressing the belief and expectation that the message recipient can change his or her situation and overcome difficulties. Constructs from the transtheoretical model [62] such as visioning, dramatic relief, self-reevaluation, and environmental reevaluation also guided level 1 message development.

PAM level 2 messages build knowledge and self-efficacy to engage in a behavior and focus on ways to take small steps that don't require much in-depth knowledge. Self-efficacy and the confidence a person feels about performing a particular activity was a primary construct used to develop these messages with a focus on one of the main strategies to build self-efficacy, that of taking small steps that are likely to result in performance accomplishment. Outcome expectations, or the anticipatory outcomes of a behavior, stated in ways that would likely appeal to the expectancies or values a person places on the outcome, was also an important construct [63].

PAM level 3 messages assume some knowledge and focus more on building self-management skills such as goal setting and self-monitoring. For messages in this level, we used transtheoretical model [62] constructs relevant to the preparation and action stages of behavior change.

PAM level 4 messages about staying the course and avoiding relapse when stressed were grounded in the transtheoretical model constructs guiding processes used in the maintenance stage of change. Also used in level 4 message development were strategies developed in a relapse prevention model [64] such as identifying high-risk situations for relapse and the development of specific coping strategies for those situations.

In phase 2 of our study, tailored behavioral messages will be sent to each intervention group participant, via the MHCE accessed through their mobile device, based on both senior research associate-initiated and algorithm-automated schedules and thresholds developed according to PAM level, SDSCA responses, and agreed-upon general rotation. Figure 2 offers examples. The senior research associates will use the MHCE backend portal control panel for manual rotational scheduling of messages to be delivered 3 days per week (typically Monday, Wednesday, and Friday) within the MHCE system. Participant responses to the SDSCA may trigger additional messaging if their clinical readings from biomedical devices exceed established safety thresholds.

Figure 2. Example of tailored health messaging.

Scenario: A 60-year-old woman who is a PAM level 2 on the activation scale, meaning that she is becoming aware of her need for self-management but still struggling and is capable of setting only simple goals. She may indicate on the SDSCA that she is not completing 150 minutes of physical activity each week. We know from her clinical data that her BMI is 35. She might receive messages like some of these:

"Diabetes is a tough disease, but by taking small steps, you can learn to control the disease rather than let it control you."

"Physical activity is very important to controlling diabetes, and you can increase your activity little by little by adding an extra 5 minutes of walking here and there throughout the day."

"When you park your car at work, park at the end of the parking lot so that you walk a little further to the building."

"Get up from your desk each hour to walk down the hall for a drink of water."

At lunch, walk 5 minutes before sitting down to eat."

"At the end of the day, walk across the parking lot back to your car."

"By making small changes, you can accumulate about 20-25 extra minutes of physical activity per day!"

"If you increase your physical activity, you will feel better, have more energy, and very likely lose weight."

"Think about what it will be like to be your "best self" - taking on the day with more energy and feeling in control!"

Phase 1 User-Centered Design Study Flow

In phase 1 we will evaluate and gain feedback from patients with diabetes regarding MHCE app navigation, use of external devices, ease of use, and satisfaction. We will collect baseline research participant data to include basic demographic data and clinical measures following verification of informed consent. One researcher-facilitator will lead individual participants through usability testing and the additional researcher-observer will document observations. During a facilitator-provided demonstration of the MHCE, the facilitator will ask each participant to concurrently navigate to each component of the MHCE system via a mini tablet device under their control. For each task, we will ask 3 open-ended questions to evaluate task-specific user satisfaction regarding the look and layout of

the app, how the app functions, and any specific issues that are confusing. Next, the facilitator will give a brief demonstration of the external devices that will be used in the study: a blood pressure monitor, a glucometer, a digital precision weight scale, and a Fitbit Charge wireless activity and sleep wristband. For each device, we will ask participants to (1) manually upload data, (2) sync each device with the app, and (3) interpret graphs. While it would be preferable to observe the MHCE in the context of where the patient would actually use the system, financial limitations prohibit such expanded usability observation research.

Research staff will evaluate usability by applying definitions and usability evaluation metrics guided by the International Organization for Standardization's 9241-11 usability framework

and mHealth usability research [65]. Specific metrics to evaluate usability are effectiveness, efficiency, and satisfaction. We will evaluate effectiveness via task completion and error coding. We will assess timed task completion as a task being completed with ease, being completed with minor mistakes, or not completed. Errors will be coded using a codebook developed by the phase 1 team. The observer will also note when users commit errors they cannot solve or commit errors that prevent further progress. We will use the Single Ease Question to evaluate informant satisfaction immediately after performing each task [66]. The System Usability Scale (SUS) will evaluate overall informant satisfaction with the MHCE [67].

We will also assess provider preferences in phase 1 using focus groups of clinicians and nurses recruited from the 2 study sites. Two trained qualitative researchers will facilitate the focus groups. We will take field notes during the focus groups and audio record each session to ensure accuracy of the field notes. The facilitators will use a semistructured interview guide to elicit clinician and nurse feedback about the MHCE. After briefly demonstrating the app, facilitators will ask 6 broad questions (with probes), developed by the phase 1 team in conjunction with study coinvestigators. These questions are designed to elicit feedback from participants regarding app design, alerts (general), wording of alerts, perceived usefulness to patients for promoting self-management, clinical usefulness and workflow, and backend portal data summaries. We will probe specific issues related to clinical usefulness of the MHCE in the context of the clinical workflow of the PCMH environment.

A 4-member team will complete a thematically organized data analysis of the clinician and nurse feedback using an inductive narrative approach [68-70]. We will begin with an analysis of

field notes from 1 randomly selected provider and 1 nurse to create an initial codebook. We will expand the codebook as we continue to code field notes. The analysis team will divide the coding duties so that each transcript is coded by 2 independent coders [71]. The team will meet during the coding process to address consensus, update the coding structure, and revisit any previously coded field notes that need to be reviewed again based on these updates. Codes will be applied to the transcripts using Atlas.ti software version 7.5.10. Codes drawn from the interview guide will serve as the organizing framework for analysis. As new themes emerge, we will expand the narrative.

Phase 2 Controlled Study: Patient Enrollment and Study Flow

For phase 2 we will recruit 240 patients (120 per site), with half assigned as a control group. Eligible patients will be first assigned to 4 strata according to their PAM score. After all patients are identified and assigned into the strata, simple randomizations will be performed within each stratum to assign patients to either the MHCE or usual care groups. Patients will be randomly allocated to either the control or the intervention group based on their PAM scores.

We will modify the MHCE system between phase 1 and phase 2 research, incorporating phase 1 observations and optimizing system usability at the patient level. We will collect baseline research participant data, including basic demographic data and clinical measures, following verification of informed consent.

MHCE Intervention Versus Usual Care

Patients in both the intervention and usual care (control) groups will receive a device package as outlined in [Textbox 1](#). These devices will collect and record biometric data. All patients will be trained in using biomedical devices and peripheral equipment.

Textbox 1. Patient device package (intervention and control groups).

- Activity monitor (Bluetooth and cloud enabled)
- Scale (Bluetooth enabled)
- Blood pressure cuff (Bluetooth enabled)
- Glucometer (Bluetooth enabled)

For the patient groups allocated to the intervention, their devices will be mapped to the MHCE system accessible from the patients' mobile phone or an iPad mini tablet device. Data from their biomedical devices will be visually presented in the MHCE with trend and scalable options ([Figure 3](#)).

Safety algorithms will be mapped to these clinical data to alert the patient and, depending on the measure, the clinical team when readings exceed established thresholds. The intervention groups will also have full access to and will receive the tailored behavioral messaging outlined above. At time of study

enrollment, we will provide a tablet device to patients who are fully eligible to participate, are allocated to the intervention group, but do not have a mobile phone (with iOS or Android operating system).

In both the intervention and control groups, the patients' clinician and PCMH support team will be notified of the patients' enrollment in the study. The intervention patients will be encouraged to regularly use the MHCE system as a tool to improve their diabetes self-care.

Figure 3. Example of visualization of patient data. BP: blood pressure.



Initial Outcome Measures for Patient Component

Primary outcome measures are PAM scores. Secondary outcome measures in the study are (1) SDSCA responses, (2) clinical measures (Textbox 2), (3) comorbid conditions (eg, uncontrolled

plasma glucose, hypertension, hyperlipidemia, stroke, eye disease, coronary heart disease), (4) SUS survey scores, (5) MHCE usage statistics, and (6) health services utilization measures.

Textbox 2. Clinical measures in phase 2.


- Glycated hemoglobin (HbA_{1c})
- Low-density lipoprotein
- High-density lipoprotein
- Height and weight
- Abdominal circumference
- Systolic blood pressure
- Diastolic blood pressure

Patient Activation Measure Instrument

The self-reported PAM survey is associated with self-management behaviors, medication adherence, patient satisfaction, and quality of life [55,72]. Within a diabetes-specific population, PAM is not related to knowledge regarding HbA_{1c} (the standard measure of average blood glucose level [73]), but is associated with better glycemic control [74]. Interventions, including educational programs [31], care coaching [32], and motivational interviewing [33], have been

attempted to improve this activation with varied success. Specifically, patient activation can be increased with targeted, patient-centered, repeated messaging [19]. The PAM is a valid, reliable, unidimensional, probabilistic Guttman-like scale that was validated over a decade ago [21] and is a standard tool to measure patient activation. We will administer the PAM at screening visits in phases 1 and 2, and electronically every 3 months during phase 2 for both the intervention and control groups. Figure 4 outlines the 4 PAM levels.

Figure 4. The 4 levels within the Patient Activation Measure (PAM) survey.



Summary of Diabetes Self-Care Activities Instrument

The SDSCA instrument is a brief self-report instrument for measuring levels of self-management across different components of the diabetes regimen [56]. The SDSCA includes 11 core items associated with diabetes self-care. The SDSCA has been successfully used in numerous diabetes studies both within and outside the United States [56,75-78]. The SDSCA has been validated and is considered a standard instrument in diabetes care for measuring self-care activities, with its validation and reliability published nearly two decades ago [56]. We will administer the SDSCA at the intake visit for phase 2, and electronically every 2 weeks during phase 2 for both the intervention and control groups.

Clinical Measures

We will collect clinical measures (Textbox 2) from patients at intake during phase 1 research. We will collect and compare changes in patient clinical measures for both groups in phase 2 at 3 points: intake, midpoint (month 6), and conclusion (month 12). For patients assigned to the MHCE intervention group, the MHCE system will also record weight, systolic blood pressure, diastolic blood pressure, and blood glucose values to the MHCE module on a regular basis via Wi-Fi or Bluetooth-enabled peripheral equipment.

Clinician Support for Patient Activation Measure

The Clinician Support for Patient Activation Measure (CS-PAM) instrument measures clinician beliefs about patient self-management behavior. The CS-PAM has been a valid and reliable instrument in use since 2010 [25]. The CS-PAM score indicates an individual clinician's overall level of endorsement or belief about the importance of patient self-management, as well as beliefs about the importance of specific patient competency categories [25].

In phase 2, we will measure clinician support for patient self-management by the CS-PAM. PCMH clinicians (ie, physicians, nurse practitioners, and physician assistants) in this study will take the CS-PAM at 3 points in the study: beginning, midpoint (month 6), and conclusion (month 12).

System Usability Scale Survey

The SUS survey is a 10-item Likert-like scaled survey used to convey a subjective assessment of system usability. The instrument was developed over 15 years ago and is used to measure the usability of websites. The SUS was validated on several occasions, with perhaps the largest validation study (including 10 years' worth of data) conducted in 2008 [79]. In this study we will substitute the term "MHCE system" for the term "website" in the instrument. We will conduct the SUS survey at the conclusion of the encounter for phase 1 patients, and at midpoint (months 5-6) and study conclusion (months 11-12) for phase 2 patients in the intervention group.

MHCE Usage Statistics

Our technology enablement partners will embed counters (invisible to patients) that track usage of MHCE components. These counters will export usage data to our research analysis database. Summary statistics and trends will be analyzed with comparison.

Comorbid Conditions

We will assess and document comorbid conditions (eg, hypertension, hyperlipidemia) among both the control and intervention groups during prescreening of eligibility, at intake, at study midpoint, and at study conclusion. While not primary outcome measures, any change over time and whether the number and type of comorbid conditions influence patient use of MHCE will be assessed.

Data Analysis Strategy

We will conduct the primary analyses for phase 2 using an intent-to-treat approach. Study participants will be retained in their original assignment groups after the random allocation in the analysis. Achievement of randomization will be evaluated through the comparison of baseline key variables between the MHCE intervention group and the control group. We will also compare baseline key characteristics between eligible patients who participate in the study and those who do not participate to examine the potential for bias.

To test hypotheses 2 and 3, that patients who participate in MHCE will have higher PAM, SDSCA, and SUS scores and improved selected clinical outcomes and comorbid conditions than their counterparts in usual care, we will use multivariate regression models (logistic regression if the outcome is a binary variable and linear regression if the outcome is a continuous variable) with the intervention assignment as the primary independent variable. Stratified analyses will be conducted (eg, sex, race, and initial PAM score).

The primary comparison will be outcomes at 12 months. Additional analyses will use longitudinal analysis models using a generalized estimating equation, which will include outcomes at both 6 and 12 months.

To test hypothesis 4, that patients who engage at a higher rate with the interactive and tailored mobile technology in MHCE will realize greater improvement in clinical measures (eg, HbA_{1c} values; [Textbox 2](#)), we will use multivariate linear regression models. Clinical outcomes will be the dependent variables and will be tested separately. The main independent variable will be MHCE usage. We will examine the association between the dependent variable and MHCE usage by using the generalized estimate equation with adjustment of potential confounders (eg, age, sex, race, duration of disease, use or nonuse of insulin).

Trial Status

At the time of publication, we have completed phase 1 data collection. Formal analysis of phase 1 data has not been completed. Institutional review board approval (study and site implementation) has been obtained and phase 1 research commenced in late fall 2016.

Results

The hypotheses of the study suggest that patients can, and will, improve their activation in chronic care self-management. Improved activation should translate into improved diabetes self-care. While not powered in this study, improved self-management activities should lead to fewer emergency situations (and trips to the emergency department), weight loss (in many cases), improved blood pressure, and improved clinical measures. Cumulatively, the gains should translate into improved quality of life if our hypotheses are supported.

This study has been approved by the institutional review boards of Clemson University (protocol #IRB2015-234) and the Madigan Army Medical Center (representing both DoD sites; reference #216073). Study personnel will follow protocol with all informed consent mandates directed by the institutional review boards; informed consent in this study includes both patients and clinicians or key clinical staff. This trial was registered with ClinicalTrials.gov (NCT02949037) on October 31, 2016.

Discussion

Expected benefits of this research and development effort to the scientific community and health care services include improved understanding of how to advance 3 joint PCMH principles (ie, better coordination of care, improved quality and safety, and enhanced access to care) through the use of mobile technology and improved understanding of how to include mHealth technology in the clinical workflow of the PCMH health services model, as well as improved understanding of how to use mHealth technology to activate patients with a diagnosis of type 2 diabetes in disease self-management behaviors. We also expect to improve understanding of how patient complexity and degree of “sickness” may influence patient use or nonuse of mHealth technologies in self-management of their disease, and to explore how to map patient-entered biomedical data onto clinical documentation and a decision-support platform useful in chronic care management.

Our study design is not immune from potential threats to validity. Patients allocated to the control arm will be issued the same peripheral devices as the intervention group and, while they may not achieve the same degree of activation, they may realize improvement if they use the equipment being issued to them. Though this behavioral mechanism could benefit patients in the control group, a strong activation change in the control arm could conceal the behavioral benefit of our intervention when we compare patient behavior from the 2 arms.

We are aware that we did not conduct a power calculation for sample size, since this project was funded as a feasibility study, not a randomized controlled trial. Thus, sample size estimates are neither required nor appropriate. We additionally recognize that a formal randomized controlled trial would be preferred to our current design. A follow-on randomized controlled trial is our goal once we have collected sufficient data and have a better understanding of how patients will use this chronic care health

information technology system. At that point we will properly power the study. legitimately be able to predict the intervention effect and

Acknowledgments

The authors appreciate and would like to acknowledge the contributions of Terry J Newton, MD (US Army Office of the Surgeon General), who provided feedback on the systemwide implications of the research efforts and offered general guidance. We also acknowledge Ms Holly Pavliscsak (US Army Telemedicine and Advanced Technology Research Center), who helped shape early technology development of our mHealth intervention.

Funding: The study was extramurally funded, via the Broad Agency Announcement, by the US Army Medical Research Acquisition Activity; contract # W81XWH-15-C-0070. The funder did not influence the design of the study or strategies related to its collection, analysis, or interpretation of data.

Federal disclaimer: The views expressed are those of the authors and do not reflect the official policy of the Department of the Army, the Department of the Air Force, the Department of Defense, or the US federal government.

Authors' Contributions

RG initially conceptualized the study, and both RG and LS wrote the initial draft of the manuscript. PC, EAS, RG, WWS, KWE, MH, and JBM assisted in the setting narrative, enrollment strategies, and institutional review board-related issues. The technical aspects of the intervention and MHCE description were authored by JRL and RG. The biostatistics and data analysis strategy were developed by LC and KT. The patient activation and behavioral messaging component, including examples, was developed by CJD, JEW, SFG, and KOJ. The user-centered design component was researched and authored by JEW and SFG. The outcome measures component was conceptualized and authored by PC, EAS, KT, LS, CJD, JEW, LC, JBM, MH, WWS, and KWE. The clinical components, including inclusion and exclusion criteria, were developed by PC, EAS, CD, WWS, JBM, MH, KWE, and RG. The MHS review and component were authored by JBM, MH, KWE, and RG. All authors read, contributed to, critically reviewed, and approved the final manuscript.

Conflicts of Interest

None declared.

References

1. Sudore RL, Karter AJ, Huang ES, Moffet HH, Laiteerapong N, Schenker Y, et al. Symptom burden of adults with type 2 diabetes across the disease course: diabetes & aging study. *J Gen Intern Med* 2012 Dec;27(12):1674-1681 [[FREE Full text](#)] [doi: [10.1007/s11606-012-2132-3](#)] [Medline: [22854982](#)]
2. Vigersky RA. An overview of management issues in adult patients with type 2 diabetes mellitus. *J Diabetes Sci Technol* 2011 Mar 01;5(2):245-250 [[FREE Full text](#)] [Medline: [21527089](#)]
3. Scollan-Koliopoulos M, Bleich D, Rapp KJ, Wong P, Hofmann CJ, Raghuwanshi M. Health-related quality of life, disease severity, and anticipated trajectory of diabetes. *Diabetes Educ* 2013;39(1):83-91. [doi: [10.1177/0145721712467697](#)] [Medline: [23174664](#)]
4. American Diabetes Association. Economic costs of diabetes in the U.S. in 2012. *Diabetes Care* 2013 Apr;36(4):1033-1046 [[FREE Full text](#)] [doi: [10.2337/dc12-2625](#)] [Medline: [23468086](#)]
5. Boyle JP, Thompson TJ, Gregg EW, Barker LE, Williamson DF. Projection of the year 2050 burden of diabetes in the US adult population: dynamic modeling of incidence, mortality, and prediabetes prevalence. *Popul Health Metr* 2010 Oct 22;8:29 [[FREE Full text](#)] [doi: [10.1186/1478-7954-8-29](#)] [Medline: [20969750](#)]
6. Patient-Centered Primary Care Collaborative. Joint principles of the patient-centered medical home. 2007 Jan 01. URL: <https://www.pcpcc.org/about/medical-home> [accessed 2017-03-02] [[WebCite Cache ID 6ofFoxX10](#)]
7. Nielsen M, Langner B, Zema C, Hacker T, Grundy P. Benefits of Implementing the Primary Care Patient-Centered Medical Home: A Review of Cost & Quality Results. Washington, DC: Patient-Centered Primary Care Collaborative; 2012:41.
8. Jortberg BT, Miller BF, Gabbay RA, Sparling K, Dickinson WP. Patient-centered medical home: how it affects psychosocial outcomes for diabetes. *Curr Diab Rep* 2012 Dec;12(6):721-728. [doi: [10.1007/s11892-012-0316-1](#)] [Medline: [22961115](#)]
9. Wade-Vuturo AE, Mayberry LS, Osborn CY. Secure messaging and diabetes management: experiences and perspectives of patient portal users. *J Am Med Inform Assoc* 2013 May 1;20(3):519-525 [[FREE Full text](#)] [doi: [10.1136/amiainl-2012-001253](#)] [Medline: [23242764](#)]
10. Yu CH, Bahniwal R, Laupacis A, Leung E, Orr MS, Straus SE. Systematic review and evaluation of web-accessible tools for management of diabetes and related cardiovascular risk factors by patients and healthcare providers. *J Am Med Inform Assoc* 2012;19(4):514-522 [[FREE Full text](#)] [doi: [10.1136/amiainl-2011-000307](#)] [Medline: [22215057](#)]
11. Osborn CY, Mayberry LS, Mulvaney SA, Hess R. Patient web portals to improve diabetes outcomes: a systematic review. *Curr Diab Rep* 2010 Dec;10(6):422-435 [[FREE Full text](#)] [doi: [10.1007/s11892-010-0151-1](#)] [Medline: [20890688](#)]

12. Faridi Z, Liberti L, Shuval K, Northrup V, Ali A, Katz DL. Evaluating the impact of mobile telephone technology on type 2 diabetic patients' self-management: the NICHE pilot study. *J Eval Clin Pract* 2008 Jun;14(3):465-469. [doi: [10.1111/j.1365-2753.2007.00881.x](https://doi.org/10.1111/j.1365-2753.2007.00881.x)] [Medline: [18373577](#)]
13. Arora S, Peters AL, Agy C, Menchine M. A mobile health intervention for inner city patients with poorly controlled diabetes: proof-of-concept of the TExT-MED program. *Diabetes Technol Ther* 2012 Jun;14(6):492-496. [doi: [10.1089/dia.2011.0252](https://doi.org/10.1089/dia.2011.0252)] [Medline: [22524591](#)]
14. Quinn CC, Shardell MD, Terrin ML, Barr EA, Ballew SH, Gruber-Baldini AL. Cluster-randomized trial of a mobile phone personalized behavioral intervention for blood glucose control. *Diabetes Care* 2011 Sep;34(9):1934-1942 [[FREE Full text](#)] [doi: [10.2337/dc11-0366](https://doi.org/10.2337/dc11-0366)] [Medline: [21788632](#)]
15. Newton KH, Wiltshire EJ, Elley CR. Pedometers and text messaging to increase physical activity: randomized controlled trial of adolescents with type 1 diabetes. *Diabetes Care* 2009 May;32(5):813-815 [[FREE Full text](#)] [doi: [10.2337/dc08-1974](https://doi.org/10.2337/dc08-1974)] [Medline: [19228863](#)]
16. Harris LT, Haneuse SJ, Martin DP, Ralston JD. Diabetes quality of care and outpatient utilization associated with electronic patient-provider messaging: a cross-sectional analysis. *Diabetes Care* 2009 Jul;32(7):1182-1187 [[FREE Full text](#)] [doi: [10.2337/dc08-1771](https://doi.org/10.2337/dc08-1771)] [Medline: [19366959](#)]
17. Bredfeldt CE, Compton-Phillips AL, Snyder MH. Effects of between visit physician-patient communication on Diabetes Recognition Program scores. *Int J Qual Health Care* 2011 Dec;23(6):664-673 [[FREE Full text](#)] [doi: [10.1093/intqhc/mzr061](https://doi.org/10.1093/intqhc/mzr061)] [Medline: [21937586](#)]
18. Lyles CR, Grothaus L, Reid RJ, Sarkar U, Ralston JD. Communication about diabetes risk factors during between-visit encounters. *Am J Manag Care* 2012 Dec;18(12):807-815 [[FREE Full text](#)] [Medline: [23286610](#)]
19. Ledford CJW, Ledford CC, Childress MA. Extending Physician ReACH: influencing patient activation and behavior through multichannel physician communication. *Patient Educ Couns* 2013 Apr;91(1):72-78. [doi: [10.1016/j.pec.2012.11.011](https://doi.org/10.1016/j.pec.2012.11.011)] [Medline: [23219484](#)]
20. Ledford CJW, Ledford CC, Childress MA. Exploring patient activation in the clinic: measurement from three perspectives. *Health Educ Behav* 2013 Jun;40(3):339-345. [doi: [10.1177/1090198112455173](https://doi.org/10.1177/1090198112455173)] [Medline: [22984210](#)]
21. Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the Patient Activation Measure (PAM): conceptualizing and measuring activation in patients and consumers. *Health Serv Res* 2004 Aug;39(4 Pt 1):1005-1026 [[FREE Full text](#)] [doi: [10.1111/j.1475-6773.2004.00269.x](https://doi.org/10.1111/j.1475-6773.2004.00269.x)] [Medline: [15230939](#)]
22. Ledford CJW. Exploring the interaction of patient activation and message design variables: message frame and presentation mode influence on the walking behavior of patients with type 2 diabetes. *J Health Psychol* 2012 Oct;17(7):989-1000. [doi: [10.1177/1359105311429204](https://doi.org/10.1177/1359105311429204)] [Medline: [22253330](#)]
23. Wagner EH, Glasgow RE, Davis C, Bonomi AE, Provost L, McCulloch D, et al. Quality improvement in chronic illness care: a collaborative approach. *Jt Comm J Qual Improv* 2001 Feb;27(2):63-80. [Medline: [11221012](#)]
24. Greene J, Hibbard JH. Why does patient activation matter? An examination of the relationships between patient activation and health-related outcomes. *J Gen Intern Med* 2012 May;27(5):520-526 [[FREE Full text](#)] [doi: [10.1007/s11606-011-1931-2](https://doi.org/10.1007/s11606-011-1931-2)] [Medline: [22127797](#)]
25. Hibbard JH, Collins PA, Mahoney E, Baker LH. The development and testing of a measure assessing clinician beliefs about patient self-management. *Health Expect* 2010 Mar;13(1):65-72. [doi: [10.1111/j.1369-7625.2009.00571.x](https://doi.org/10.1111/j.1369-7625.2009.00571.x)] [Medline: [19906211](#)]
26. Cortes DE, Mulvaney-Day N, Fortuna L, Reinfeld S, Alegría M. Patient-provider communication: understanding the role of patient activation for Latinos in mental health treatment. *Health Educ Behav* 2009 Feb;36(1):138-154 [[FREE Full text](#)] [doi: [10.1177/1090198108314618](https://doi.org/10.1177/1090198108314618)] [Medline: [18413668](#)]
27. Griffin SJ, Kinmonth A, Veltman MW, Gillard S, Grant J, Stewart M. Effect on health-related outcomes of interventions to alter the interaction between patients and practitioners: a systematic review of trials. *Ann Fam Med* 2004;2(6):595-608 [[FREE Full text](#)] [doi: [10.1370/afm.142](https://doi.org/10.1370/afm.142)] [Medline: [15576546](#)]
28. Hibbard JH, Mahoney ER, Stock R, Tusler M. Do increases in patient activation result in improved self-management behaviors? *Health Serv Res* 2007 Aug;42(4):1443-1463 [[FREE Full text](#)] [doi: [10.1111/j.1475-6773.2006.00669.x](https://doi.org/10.1111/j.1475-6773.2006.00669.x)] [Medline: [17610432](#)]
29. Greene J, Hibbard JH, Sacks R, Overton V, Parrotta CD. When patient activation levels change, health outcomes and costs change, too. *Health Aff (Millwood)* 2015 Mar 1;34(3):431-437. [doi: [10.1377/hlthaff.2014.0452](https://doi.org/10.1377/hlthaff.2014.0452)] [Medline: [25732493](#)]
30. Bodenheimer T. Helping patients improve their health-related behaviors: what system changes do we need? *Dis Manag* 2005 Oct;8(5):319-330. [doi: [10.1089/dis.2005.8.319](https://doi.org/10.1089/dis.2005.8.319)] [Medline: [16212517](#)]
31. Li T, Wu HM, Wang F, Huang CQ, Yang M, Dong BR, et al. Education programmes for people with diabetic kidney disease. *Cochrane Database Syst Rev* 2011 Jun 15(6):CD007374. [doi: [10.1002/14651858.CD007374.pub2](https://doi.org/10.1002/14651858.CD007374.pub2)] [Medline: [21678365](#)]
32. Lin W, Chien H, Willis G, O'Connell E, Rennie KS, Bottella HM, et al. The effect of a telephone-based health coaching disease management program on Medicaid members with chronic conditions. *Med Care* 2012 Jan;50(1):91-98. [doi: [10.1097/MLR.0b013e31822dcedf](https://doi.org/10.1097/MLR.0b013e31822dcedf)] [Medline: [21993059](#)]

33. Linden A, Butterworth SW, Prochaska JO. Motivational interviewing-based health coaching as a chronic care intervention. *J Eval Clin Pract* 2010 Feb;16(1):166-174. [doi: [10.1111/j.1365-2753.2009.01300.x](https://doi.org/10.1111/j.1365-2753.2009.01300.x)] [Medline: [20367828](#)]
34. Ledford CJW, Canzona MR, Cafferty LA, Hodge JA. Mobile application as a prenatal education and engagement tool: a randomized controlled pilot. *Patient Educ Couns* 2016 Apr;99(4):578-582. [doi: [10.1016/j.pec.2015.11.006](https://doi.org/10.1016/j.pec.2015.11.006)] [Medline: [26610389](#)]
35. Deen D, Lu W, Rothstein D, Santana L, Gold MR. Asking questions: the effect of a brief intervention in community health centers on patient activation. *Patient Educ Couns* 2011 Aug;84(2):257-260. [doi: [10.1016/j.pec.2010.07.026](https://doi.org/10.1016/j.pec.2010.07.026)] [Medline: [20800414](#)]
36. Deen D, Lu WH, Weintraub MR, Maranda MJ, Elshafey S, Gold MR. The impact of different modalities for activating patients in a community health center setting. *Patient Educ Couns* 2012 Oct;89(1):178-183. [doi: [10.1016/j.pec.2012.04.012](https://doi.org/10.1016/j.pec.2012.04.012)] [Medline: [22683294](#)]
37. Lu W, Deen D, Rothstein D, Santana L, Gold MR. Activating community health center patients in developing question-formulation skills: a qualitative study. *Health Educ Behav* 2011 Dec;38(6):637-645. [doi: [10.1177/1090198110393337](https://doi.org/10.1177/1090198110393337)] [Medline: [21558464](#)]
38. Fowles JB, Terry P, Xi M, Hibbard J, Bloom CT, Harvey L. Measuring self-management of patients' and employees' health: further validation of the Patient Activation Measure (PAM) based on its relation to employee characteristics. *Patient Educ Couns* 2009 Oct;77(1):116-122. [doi: [10.1016/j.pec.2009.02.018](https://doi.org/10.1016/j.pec.2009.02.018)] [Medline: [19356881](#)]
39. Coleman EA, Parry C, Chalmers S, Min S. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med* 2006 Sep 25;166(17):1822-1828. [doi: [10.1001/archinte.166.17.1822](https://doi.org/10.1001/archinte.166.17.1822)] [Medline: [17000937](#)]
40. Toscos T, Daley C, Heral L, Doshi R, Chen Y, Eckert GJ, et al. Impact of electronic personal health record use on engagement and intermediate health outcomes among cardiac patients: a quasi-experimental study. *J Am Med Inform Assoc* 2016 Jan;23(1):119-128. [doi: [10.1093/jamia/ocv164](https://doi.org/10.1093/jamia/ocv164)] [Medline: [26912538](#)]
41. Crouch PB, Rose CD, Johnson M, Janson SL. A pilot study to evaluate the magnitude of association of the use of electronic personal health records with patient activation and empowerment in HIV-infected veterans. *PeerJ* 2015;3:e852 [FREE Full text] [doi: [10.7717/peerj.852](https://doi.org/10.7717/peerj.852)] [Medline: [25802815](#)]
42. Wagner PJ, Dias J, Howard S, Kintziger KW, Hudson MF, Seol Y, et al. Personal health records and hypertension control: a randomized trial. *J Am Med Inform Assoc* 2012 Aug;19(4):626-634 [FREE Full text] [doi: [10.1136/amiainl-2011-000349](https://doi.org/10.1136/amiainl-2011-000349)] [Medline: [22234404](#)]
43. Solomon M, Wagner SL, Goes J. Effects of a Web-based intervention for adults with chronic conditions on patient activation: online randomized controlled trial. *J Med Internet Res* 2012;14(1):e32 [FREE Full text] [doi: [10.2196/jmir.1924](https://doi.org/10.2196/jmir.1924)] [Medline: [22353433](#)]
44. Nagykalai Z, Aspy CB, Chou A, Mold JW. Impact of a wellness portal on the delivery of patient-centered preventive care. *J Am Board Fam Med* 2012 Apr;25(2):158-167 [FREE Full text] [doi: [10.3122/jabfm.2012.02.110130](https://doi.org/10.3122/jabfm.2012.02.110130)] [Medline: [22403196](#)]
45. Viau KS, Jones JL, Murtaugh MA, Gren LH, Stanford JB, Bilder DA. Phone-based motivational interviewing to increase self-efficacy in individuals with phenylketonuria. *Mol Genet Metab Rep* 2016 Mar;6:27-33 [FREE Full text] [doi: [10.1016/j.ymgmr.2016.01.002](https://doi.org/10.1016/j.ymgmr.2016.01.002)] [Medline: [27014576](#)]
46. Riippa I, Linna M, Rönkkö I, Kröger V. Use of an electronic patient portal among the chronically ill: an observational study. *J Med Internet Res* 2014 Dec 08;16(12):e275 [FREE Full text] [doi: [10.2196/jmir.3722](https://doi.org/10.2196/jmir.3722)] [Medline: [25488754](#)]
47. Riippa I, Linna M, Rönkkö I. The effect of a patient portal with electronic messaging on patient activation among chronically ill patients: controlled before-and-after study. *J Med Internet Res* 2014;16(11):e257 [FREE Full text] [doi: [10.2196/jmir.3462](https://doi.org/10.2196/jmir.3462)] [Medline: [25413368](#)]
48. Kim JY, Wineinger NE, Steinhubl SR. The influence of wireless self-monitoring program on the relationship between patient activation and health behaviors, medication adherence, and blood pressure levels in hypertensive patients: a substudy of a randomized controlled trial. *J Med Internet Res* 2016 Jun 22;18(6):e116 [FREE Full text] [doi: [10.2196/jmir.5429](https://doi.org/10.2196/jmir.5429)] [Medline: [27334418](#)]
49. Hevner A, March S, Park J, Ram S. Design science in information systems research. *MIS Q* 2004;28(1):75-105.
50. Peffers K, Tuunanen T, Rothenberger M, Chatterjee S. A design science research methodology for information systems research. *J Manage Inform Systems* 2007;24(3):45-77.
51. Hesse BW, Shneiderman B. eHealth research from the user's perspective. *Am J Prev Med* 2007 May;32(5 Suppl):S97-103 [FREE Full text] [doi: [10.1016/j.amepre.2007.01.019](https://doi.org/10.1016/j.amepre.2007.01.019)] [Medline: [17466825](#)]
52. Robinson TN, Patrick K, Eng TR, Gustafson D. An evidence-based approach to interactive health communication: a challenge to medicine in the information age. Science Panel on Interactive Communication and Health. *JAMA* 1998 Oct 14;280(14):1264-1269. [Medline: [9786378](#)]
53. Neuhauser L, Kreps G. Participatory design and artificial intelligence: strategies to improve health communication for diverse audiences. 2011 Presented at: AAAI Spring Symposium Series; March 21-23, 2011; Palo Alto, CA, USA.
54. Gimbel RW, Pangaro L, Barbour G. America's "undiscovered" laboratory for health services research. *Med Care* 2010 Aug;48(8):751-756. [doi: [10.1097/MLR.0b013e3181e35be8](https://doi.org/10.1097/MLR.0b013e3181e35be8)] [Medline: [20613659](#)]

55. Harvey L, Fowles JB, Xi M, Terry P. When activation changes, what else changes? The relationship between change in patient activation measure (PAM) and employees' health status and health behaviors. *Patient Educ Couns* 2012 Aug;88(2):338-343. [doi: [10.1016/j.pec.2012.02.005](https://doi.org/10.1016/j.pec.2012.02.005)] [Medline: [22459636](https://pubmed.ncbi.nlm.nih.gov/22459636/)]
56. Toobert DJ, Hampson SE, Glasgow RE. The summary of diabetes self-care activities measure: results from 7 studies and a revised scale. *Diabetes Care* 2000 Jul;23(7):943-950 [FREE Full text] [Medline: [10895844](https://pubmed.ncbi.nlm.nih.gov/10895844/)]
57. Pavliscsak H, Little JR, Poropatich RK, McVeigh FL, Tong J, Tillman JS, et al. Assessment of patient engagement with a mobile application among service members in transition. *J Am Med Inform Assoc* 2016 Jan;23(1):110-118. [doi: [10.1093/jamia/ocv121](https://doi.org/10.1093/jamia/ocv121)] [Medline: [26363048](https://pubmed.ncbi.nlm.nih.gov/26363048/)]
58. Head KJ, Noar SM, Iannarino NT, Grant HN. Efficacy of text messaging-based interventions for health promotion: a meta-analysis. *Soc Sci Med* 2013 Nov;97:41-48. [doi: [10.1016/j.socscimed.2013.08.003](https://doi.org/10.1016/j.socscimed.2013.08.003)] [Medline: [24161087](https://pubmed.ncbi.nlm.nih.gov/24161087/)]
59. Nelson LA, Mayberry LS, Wallston K, Kripalani S, Bergner EM, Osborn CY. Development and usability of REACH: a tailored theory-based text messaging intervention for disadvantaged adults with type-2 diabetes. *JMIR Hum Factors* 2016 Sep 08;3(2):e23 [FREE Full text] [doi: [10.2196/humanfactors.6029](https://doi.org/10.2196/humanfactors.6029)] [Medline: [27609738](https://pubmed.ncbi.nlm.nih.gov/27609738/)]
60. Hibbard JH, Greene J, Tusler M. Improving the outcomes of disease management by tailoring care to the patient's level of activation. *Am J Manag Care* 2009 Jun;15(6):353-360 [FREE Full text] [Medline: [19514801](https://pubmed.ncbi.nlm.nih.gov/19514801/)]
61. Israel BA, Rounds KA. Social networks and social support: a synthesis for health educators. *Adv Health Educ* 1987;12:65-80.
62. Prochaska JO, Velicer WF, Rossi JS, Goldstein MG, Marcus BH, Rakowski W, et al. Stages of change and decisional balance for 12 problem behaviors. *Health Psychol* 1994 Jan;13(1):39-46. [Medline: [8168470](https://pubmed.ncbi.nlm.nih.gov/8168470/)]
63. Bandura A. *Social Foundations of Thought and Action*. Englewood Cliffs, NJ: Prentice-Hall; 1986:544.
64. Marlatt G. Relapse prevention: theoretical rationale/overview of the model. In: Marlatt GA, Gordon JR. editors. *Relapse Prevention*. New York, NY: Guilford Press; 1985.
65. Georgsson M, Staggers N. Quantifying usability: an evaluation of a diabetes mHealth system on effectiveness, efficiency, and satisfaction metrics with associated user characteristics. *J Am Med Inform Assoc* 2016 Jan;23(1):5-11 [FREE Full text] [doi: [10.1093/jamia/ocv099](https://doi.org/10.1093/jamia/ocv099)] [Medline: [26377990](https://pubmed.ncbi.nlm.nih.gov/26377990/)]
66. Sauro J, Dumas J. Comparison of three one-question, post-task usability questionnaires. In: *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. 2009 Presented at: CHI '2009; April 4-9, 2009; Boston, MA, USA. [doi: [10.1145/1518701.1518946](https://doi.org/10.1145/1518701.1518946)]
67. Sauro J, Lewis J. *Quantifying the User Experience: Practical Statistics for User Research*. Waltham, MA: Morgan Kaufmann; 2012.
68. Kidd PS, Parshall MB. Getting the focus and the group: enhancing analytical rigor in focus group research. *Qual Health Res* 2000 May;10(3):293-308. [Medline: [10947477](https://pubmed.ncbi.nlm.nih.gov/10947477/)]
69. Thorn S. Data analysis in qualitative research. *Evid Based Nurs* 2000;3:68-70. [doi: [10.1136/ebn.3.3.68](https://doi.org/10.1136/ebn.3.3.68)]
70. Thomas DR. A general inductive approach for analyzing qualitative evaluation data. *Am J Eval* 2006 Jun 01;27(2):237-246. [doi: [10.1177/1098214005283748](https://doi.org/10.1177/1098214005283748)]
71. Miles M, Huberman A. *Qualitative Data Analysis*. 2nd edition. Thousand Oaks, CA: Sage Publications; 1994.
72. Mosen DM, Schmittiel J, Hibbard J, Sobel D, Remmers C, Bellows J. Is patient activation associated with outcomes of care for adults with chronic conditions? *J Ambul Care Manage* 2007 Mar;30(1):21-29. [Medline: [17170635](https://pubmed.ncbi.nlm.nih.gov/17170635/)]
73. Rask KJ, Ziemer DC, Kohler SA, Hawley JN, Arinde FJ, Barnes CS. Patient activation is associated with healthy behaviors and ease in managing diabetes in an indigent population. *Diabetes Educ* 2009;35(4):622-630. [doi: [10.1177/0145721709335004](https://doi.org/10.1177/0145721709335004)] [Medline: [19419972](https://pubmed.ncbi.nlm.nih.gov/19419972/)]
74. Rogvi S, Tapager I, Almdal TP, Schiøtz ML, Willaing I. Patient factors and glycaemic control--associations and explanatory power. *Diabet Med* 2012 Oct;29(10):e382-e389. [doi: [10.1111/j.1464-5491.2012.03703.x](https://doi.org/10.1111/j.1464-5491.2012.03703.x)] [Medline: [22540962](https://pubmed.ncbi.nlm.nih.gov/22540962/)]
75. Gonzalez JS, Safren SA, Delahanty LM, Cagliero E, Wexler DJ, Meigs JB, et al. Symptoms of depression prospectively predict poorer self-care in patients with type 2 diabetes. *Diabet Med* 2008 Sep;25(9):1102-1107 [FREE Full text] [doi: [10.1111/j.1464-5491.2008.02535.x](https://doi.org/10.1111/j.1464-5491.2008.02535.x)] [Medline: [19183315](https://pubmed.ncbi.nlm.nih.gov/19183315/)]
76. Watkins YJ, Quinn LT, Ruggiero L, Quinn MT, Choi Y. Spiritual and religious beliefs and practices and social support's relationship to diabetes self-care activities in African Americans. *Diabetes Educ* 2013 Feb;39(2):231-239 [FREE Full text] [doi: [10.1177/0145721713475843](https://doi.org/10.1177/0145721713475843)] [Medline: [23411653](https://pubmed.ncbi.nlm.nih.gov/23411653/)]
77. Primožič S, Tavčar R, Avbelj M, Dernovšek MZ, Oblak MR. Specific cognitive abilities are associated with diabetes self-management behavior among patients with type 2 diabetes. *Diabetes Res Clin Pract* 2012 Jan;95(1):48-54. [doi: [10.1016/j.diabres.2011.09.004](https://doi.org/10.1016/j.diabres.2011.09.004)] [Medline: [21963107](https://pubmed.ncbi.nlm.nih.gov/21963107/)]
78. Cohen HW, Shmukler C, Ullman R, Rivera CM, Walker EA. Measurements of medication adherence in diabetic patients with poorly controlled HbA(1c). *Diabet Med* 2010 Feb;27(2):210-216 [FREE Full text] [doi: [10.1111/j.1464-5491.2009.02898.x](https://doi.org/10.1111/j.1464-5491.2009.02898.x)] [Medline: [20546266](https://pubmed.ncbi.nlm.nih.gov/20546266/)]
79. Bangor A, Kortum PT, Miller JT. An empirical evaluation of the System Usability Scale. *Int J Hum Comput Interact* 2008 Jul 30;24(6):574-594. [doi: [10.1080/10447310802205776](https://doi.org/10.1080/10447310802205776)]

Abbreviations

CS-PAM: Clinician Support Patient Activation Measure

DoD: Department of Defense

HbA1c: glycated hemoglobin

MHCE: Mobile Health Care Environment

MHS: Military Health System

PAM: Patient Activation Measure

PCMH: patient-centered medical home

PHR: personal health record

SDSCA: Summary of Diabetes Self-Care Activities

SMS: short message service

SUS: System Usability Scale

Edited by A McDougall; submitted 20.11.16; peer-reviewed by K Pal, G Graffigna; comments to author 03.01.17; revised version received 16.01.17; accepted 12.02.17; published 06.03.17

Please cite as:

Gimbel R, Shi L, Williams JE, Dye CJ, Chen L, Crawford P, Shry EA, Griffin SF, Jones KO, Sherrill WW, Truong K, Little JR, Edwards KW, Hing M, Moss JB

Enhancing mHealth Technology in the Patient-Centered Medical Home Environment to Activate Patients With Type 2 Diabetes: A Multisite Feasibility Study Protocol

JMIR Res Protoc 2017;6(3):e38


URL: <http://www.researchprotocols.org/2017/3/e38/>

doi: [10.2196/resprot.6993](https://doi.org/10.2196/resprot.6993)

PMID: [28264792](https://pubmed.ncbi.nlm.nih.gov/28264792/)

©Ronald Gimbel, Lu Shi, Joel E Williams, Cheryl J Dye, Liwei Chen, Paul Crawford, Eric A Shry, Sarah F Griffin, Karyn O Jones, Windsor W Sherrill, Khoa Truong, Jeanette R Little, Karen W Edwards, Marie Hing, Jennie B Moss. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 06.03.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.

Patient Presentation



FaceTime



Calendar



Photos



Camera



Contacts



Clock



Maps



Videos



Photo Booth



Notes



Reminders



iTunes Store



App Store



iBooks



Settings



Fitbit



iGluco



Heka BP



MedM Health



mCare Plus



Messages




Safari





Mail



Music











U.S. ARMY



OK



[mCare Access/Accounts Help](#)



Checking connection, Please wait...

Main menu

**CLINICAL DISCLAIMER**

If you think you may be experiencing a medical emergency, call 911. If you need medical advice, please contact your doctor.

The information in this secure mobile app is provided as a wellness information resource only, and is not to be used or relied on for any diagnostic or treatment purposes. This information should never be used as a substitute for professional diagnosis and treatment. Please consult your health care provider, before making any healthcare decisions or for guidance about a specific medical condition.

Accept 



mCare

connecting patients via their personal cell phones



Close menu



menu



Change Pin



Alerts



Questionnaires



Contacts



Announcements



Appointments



Reminders



Tips



Diabetes



Logout

**are**

their personal cell phones





Palliative Care Process





Health Tip



Palliative Care Process



Palliative Care is a comprehensive team approach that entails pain and symptom management, emotional support and counseling, and advanced care planning.

Close menu



menu



Change Pin



Alerts



Questionnaires



Contacts



Announcements



Appointments



Reminders



Tips



Diabetes



Logout



are

their personal cell phones





John Smith





Contact

Name:

John Smith

Phone Number:

[1234567890](tel:1234567890)

Email:

JohnSmith@mCareSample.com

Role:

PL

Update



mCare

connecting patients via their personal cell phones





Menu

Announcements
Health Tips
Safety Alerts

Target Ranges

Menu

Devices



Settings




mCare




My Devices

Last 7 Days

Blood Glucose



Blood Pressure



Physical Activity



Summary




Glucose



BP



Ac



Settings

Goals



Units





Goals

Blood Glucose

Low	81	mg/dL
High	180	mg/dL

Blood Pressure

Systolic Low	90	mmHg
Systolic High	120	mmHg
Diastolic Low	60	mmHg
Diastolic High	80	mmHg

Activity

Daily	10000	steps
-------	------------------------------------	-------

Weight

	150	lbs
--	----------------------------------	-----

Cancel

Save





Alerts

Health Tips

Weekly Measuring

When you are ready to make measuring glucose part of your daily life, make a plan for measuring your glucose regularly for a week. Write down your plan and be specific about when, where, and how you will measure your glucose.

Today



1:28 PM

500

After Meal

If your blood sugar is greater than 300 mg/dL, drink water, be active, and check your blood sugar before meals and at bedtime until it is less than 200.

If it does **NOT** go down, contact your doctor.



1:28 PM

500

After Meal

Contact your doctor if your blood sugar is **over** 250 mg/dL for 24 hours.



1:28 PM

500

After Meal

Contact your doctor if your blood sugar is **over** 250 mg/dL for 24 hours.



1:27 PM

200/110

IMPORTANT: If you have chest pain, shortness of breath, back pain, numbness/weakness, change in vision or difficulty speaking, call 911 immediately!

Otherwise, for systolic blood pressure **above** 181, wait ten minutes at rest and recheck. If your systolic reading (the top number) is still above 181 and you do not have any other symptoms, contact your doctor.



1:27 PM

200/110/110

If your pulse is greater than 100 beats per minute at REST, contact your healthcare provider.




1:18 PM


If you need assistance with your F¹⁰⁰ monitor, please contact your assigned mCare Research Associate or the mCare Help Desk staff for assistance.

Your assigned points of contact can be found in the Contacts section.










Add new value option available on 30 day, 7 day, or day view.





Add Reading



Blood Glucose Reading

1

Please Select Type Of Reading

All Readings

Current Reading

2

Take Reading



Insert Strip

.....

Place Droplet of
Blood on Strip

.....



Eject Strip

3

Sync Reading



Add From My Device



Manually Input

Cancel

iPad1:45 PM89%

Reading Details

Blood Glucose Reading

1464114164214

226

mg/dL

Taken at: 1:45 pm on May 3, 2017

Blood Glucose Reading Classification

Before Meal	After Meal	Bedtime
Fasting	Unclassified	

Enter Notes Here...

Cancel

Delete


Save



Reading Details



Blood Glucose Reading



Taken at: 1:45 pm on May 3, 2017

Blood Glucose Reading Classification

Before Meal

After Meal

Bedtime




Please select a Reading Classification for your Blood Glucose reading


Enter Notes Here...

Cancel

Delete

Save





iPad 2:53 PM 75%

Enter Reading Values

Blood Glucose Reading

Enter Reading

mg/dL

Entry time:

Wed, May 3, 2017 2:53 PM

Blood Glucose Reading Classification

Before Meal	After Meal	Bedtime
Fasting	Unclassified	

Time Since Meal


120 minutes


Enter Notes Here...

Cancel

Delete

Save







Add Reading



Add Blood Pressure Reading:



Add From My Device



Tap To Add Manually

Cancel



Summary



Glucose



BP



Activity



Weight



Add Reading



Blood Pressure Reading

- 1 Put on the blood pressure cuff

*Ensure cuff is 3 to 5cm above the left elbow
- 2 Connect Device



Add From My Device


- 3 Press **Start** on device





Manually Input

Cancel



Reading Details



Blood Pressure Reading



65  bpm

Taken at: 1:47 pm on May 3, 2017

Enter Notes Here...

Cancel


Delete

Save



DAY 7 DAY 30 DAY

From 28-Sep-2016 To 04-Oct-2016



Today		
8:27 AM	121/80 mmHg	>
Yesterday		
9:47 AM	142/90 mmHg	>
28-Sep-2016		
10:27 AM	113/75 mmHg	>



Add Reading



Add Blood Pressure Reading:



Add From My Device



Tap To Add Manually



Cancel



Summary



Glucose



BP



Activity



Weight



Enter Reading Values



Blood Pressure Reading

Enter
Systolic

mmHg

Enter
Diastolic

mmHg

Enter
Pulse

bpm

Entry time: Fri, May 12, 2017 10:50 AM

Enter Notes Here...

Cancel

Delete

Save




My Devices




Last 7 Days

Blood Glucose



Blood Pressure



Physical Activity

↓ More



Summary



Glucose




BP



Activity



Weight





Register

In order for mCare to use your FitBit data, you will have to register your user account.
Please click the link below to register.

Register Device





Fitbit

CONNECTED ✓

REFRESH ↻ DISCONNECT ⌵

[MORE DETAILS](#)



Garmin Connect

CONNECT +

Garmin designs, manufactures and markets GPS navigation, communication and sonar products.

[MORE DETAILS](#)




Jawbone UP

CONNECT +

UP is a wristband and app that tracks how you sleep, move and eat.

[MORE DETAILS](#)





Add Reading



Add Weight Reading:



Add From My Device



Tap To Add Manually



Cancel



Summary



Glucose



BP



Activity



Weight

iPad 3:54 PM 56%

Add Reading

Weight Reading

Enter Weight

155 lbs

Entered at: 3:54 pm on October 5, 2016

Enter Notes Here...

Cancel Delete Save

1 2 3 4 5 6 7 8 9 0

- / : ; () \$ & @ return

#+= undo . , ? ! ' " #+=

ABC ABC



Add Reading



Add Weight Reading:



Add From My Device



Tap To Add Manually

Cancel



Summary



Glucose



BP



Activity



Weight



Add Reading



Weight Reading

1

Connect Device



Add From My Device

2

Stand on weight scale



Manually Input


Cancel



Reading Details



Weight Reading



Taken at: 1:35 pm on May 15, 2017

Enter Notes Here...

Cancel

Delete

Save

Provider Back-end Portal

Log-In Page

Log in

You will be unable to login without a created account by an mCare Personnel. Please click [here](#) to contact the help desk to request an account.

User Name

marietesting

Password


.....

Log in

[Forgot your password?](#)

[Contact](#) | [Facebook](#) | [FAQ](#) | [About](#) | [Help](#)

Home Page



Hello marietesting!

[Log off](#)

Alert Grid

Classification	Type	Alert Text	Alert Date	View	Action
		m mcare has entered a Diastolic BP > 160	05/19/2017	View	Archive
		m mcare has entered a Systolic BP > 200	05/19/2017	View	Archive
		m mcare has entered a resting Pulse < 50 bpm	05/19/2017	View	Archive

1 - 3 of 3 items

2.

Diabetes: Active Patients

Name	Last Sync	Glucose	BP	Pulse	Activity	Weight	View
ziggy d razak	03/16/2017	80 mg/dL	110/80	80	5176 steps	190 lbs	View
zaheir Razak	05/18/2017	120 mg/dL	170/80	210	6761 steps	168 lbs	View
Stanford B Ventura	04/10/2017	278 mg/dL	109/77	75	N/A	129 lbs	View
Mabel R Vasquez	05/02/2017	93 mg/dL	107/72	58	9372 steps	141 lbs	View
Stewart Ventura	05/05/2017	20 mg/dL	130/86	81	N/A	128 lbs	View
Leila D Goff	05/05/2017	233 mg/dL	N/A	N/A	N/A	N/A	View
Jennifer Perkins	04/19/2017	N/A	N/A	N/A	N/A	N/A	View
Nathan A Montgomery	05/16/2017	45 mg/dL	129/200	200	N/A	N/A	View

1.

Any Values in RED are out of

Patient's Weekly Summary Page



Hello marietesting!

Log off

Diabetic Dashboard: Mabel Vasquez

Summary Blood Glucose Blood Pressure Activity Weight

Last 7 Days Readings

From: 5/23/2017



To: 5/30/2017



Export to PDF

1.

Glucose




Date Of Reading

■ After Meal ■ Before Meal ■ Bedtime ■ Fasting ■ Unclassified

Blood Pressure



Blood Glucose-Monthly Patient Readings



Blood Pressure

Summary Blood Glucose Blood Pressure Activity Weight


Day7 Day30 Day

From: 5/2/2017To: 5/16/2017

Export to PDF

3.

4.



1.


2. —> ■ Systolic ■ Diastolic ■ BPM

5.

Export to PDF

Date ▾	Time ▾	Blood Pressure Reading	Pulse	Notes
05/04/2017	10:56 AM	145 / 120 mmHg	148 bpm	
05/04/2017	10:44 AM	130 / 86 mmHg	81 bpm	
05/02/2017	7:08 AM	180 / 160 mmHg	120 bpm	

Activity-Monthly View





Diabetic Dashboard: Mabel Vasquez

Summary Blood Glucose Blood Pressure Activity **Weight**

Day 7 Day 30 Day

From: 4/2/2017 To: 5/28/2017



Date ▼	Time ▼	Weight Reading	Notes
04/14/2017	6:19 AM	139.4 lbs	
04/14/2017	6:17 AM	139 lbs	
04/13/2017	8:40 AM	137.6 lbs	
04/13/2017	8:39 AM	137.6 lbs	
04/05/2017	6:13 AM	141.2 lbs	after breakfast
04/05/2017	6:12 AM	141.2 lbs	
04/05/2017	6:11 AM	140.8 lbs	
04/04/2017	5:58 AM	140.6 lbs	

Enhancing mHealth Technology in the PCMH environment to Activate Chronic Care Patients

ERMS/Log Number: 14210004

Award Number: W81XWH-15C-0070



PI: Ronald W. Gimbel, PhD

Org: Clemson University


Award Amount: \$1.22M

Study/Product Aim(s)

- Integrate mHealth technology into the clinical workflow of the PCMH model;
- Activate Type 2 diabetes patients in disease self-management behaviors through novel use of mHealth technology;
- Advance understanding of how patient complexity and degree of "sickness" may influence mHealth technology use in disease self-management; and
- Map patient-initiated biomedical data into clinical documentation and a decision support useful in chronic care management.

Approach

A multi-site phased feasibility study, conducted in the PCMH environment of Nellis AFB and Madigan AMC. Includes user-centered design research and feasibility testing followed by 12 month clinical trial. Study is partnership between TATRC (technology developers), Clemson University (researchers), and MTFs.



Timeline and Cost

Activities	CY	15	16	17	18
CRADA & IRB approvals					
User-centered design & testing					
Clinical trial at 2 MTFs					
Data analysis & reporting					
Estimated Budget (\$K)		\$151	\$480	\$518	\$81

Goals/Milestones

CY15 Goal –Regulatory & administrative approvals (2 months)

- ☐ Clemson IRB and CRADA approval obtained
- ☐ Madigan IRB approval pending
- ☐ Research team training conducted; Clemson RA, Madigan AMC and Nellis AFB SRAs hired; Clinical and Developmental Advisory Teams established

CY16 Goals – Phase I user-centered design and feasibility testing

- ☐ User-centered design research
- ☐ Obtain Madigan IRB approval
- ☐ Feasibility testing with patients
- ☐ Modify mHealth technology incorporating lessons learned

CY17 Goals – Phase II clinical trials at 2 MTFs

- ☐ Randomize patients
- ☐ Formal clinical trials of enhanced mHealth technology

CY18 Goal – Data analysis and reporting (4 months)

- ☐ Analyze data (including time series data)
- ☐ Formal reports and manuscripts

Budget Expenditure to Date

Projected Expenditure: \$1,230,749

Actual Expenditure: \$747,888.40 at 7/31/17

Updated: 15 MAY 2017