

AWARD NUMBER: W81XWH-11-1-0657

TITLE: Treatment-Based Classification versus Usual Care for Management of Low Back Pain

PRINCIPAL INVESTIGATOR: MAJ Daniel Rhon

RECIPIENT: Geneva Foundation
Tacoma, WA 98402

REPORT DATE: October 2017

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release; distribution is unlimited.

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE October 2017		2. REPORT TYPE Final		3. DATES COVERED 8.1.2011 – 7.31.2017	
4. TITLE AND SUBTITLE Treatment-Based Classification versus Usual Care for Management of Low Back Pain				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-11-1-0657	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) MAJ Daniel Rhon – daniel_rhon@baylor.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Geneva Foundation 917 Pacific Avenue, Suite 600 Tacoma, WA 98402				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT PURPOSE: The primary purpose of this study was to compare two management strategies for Soldiers with low back pain. Both received self-management education which as termed "usual care" and one group received immediate physical therapy. The secondary purpose was to compare downstream healthcare utilization associated with the two management strategies. DESIGN: A randomized controlled trial SCOPE: Active duty Soldiers with low back pain – 1-year follow-up period MAJOR FINDINGS: Improvements favored the early physical therapy group, but only short term (4 weeks). At 1 year, although both groups were significantly better, neither was superior. The influence of the self-management education component may have been underestimated, and likely better than "usual care". In addition, while the Oswestry Disability Index is considered the best outcome measures for low back pain, it may not be sensitive enough to capture meaningful change in military service members. Most service members did not qualify for the study as their disability score was too low, despite having disabling back pain. *This was a Career Development Grant with dual focus on mentoring/developing new scientists.					
15. SUBJECT TERMS Low back pain, military readiness, primary care, physical therapy					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Unclassified	18. NUMBER OF PAGES 21	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
1. Introduction	4
2. Keywords	4
3. Overall Project Summary	4
4. Key Research Accomplishments	5
5. Conclusion	6
6. Publications, Abstracts, and Presentations	6
7. Inventions, Patents and Licenses	6
8. Reportable Outcomes	6
9. Other Achievements	8
10. References	8
11. Appendices	8

1. INTRODUCTION:

Low back pain is the most significant contributor to lost work days related to injury in the entire U.S. Armed Forces. The detrimental impact on combat readiness of low back pain cannot be understated, as back problems are the number one cause of evacuation from Iraq and Afghanistan, making it one of the largest causes of attrition in Soldiers in combat. Back pain in the military is a predictor of back pain after the military, making this a relevant issue to the civilian healthcare systems within the U.S. This randomized clinical trial seeks additional evidence to determine if early physical therapy access using a treatment based classification algorithm will result in greater improvements in function and quality of life and decreased healthcare utilization over 1 year as compared to a stepped “usual care” strategy in 120 active duty Soldiers presenting with low back pain.

2. **KEYWORDS:** low back pain, primary care, clinical guidelines, early intervention

3. OVERALL PROJECT SUMMARY:

SOW Tasks:

Initial Tasks (months 1-7): Coordinate IRB approval, investigator participation and subject recruitment in conjunction with ongoing standard of care for patients at MAMC healthcare clinics.

Task #	Task Title	Status
a.	Submit study protocol to MAMC IRB (months 0-2)	Submitted successfully June 2011
b.	Educate participating clinicians in the 2 treatment algorithms (months 2-4)	Research assistant was hired and completed training. Two meetings with staff at the medical clinic where enrollment will occur too place (December 2011 and in January 2012)
c.	Receive approval for study by MAMC IRB (month 6)	Completed August 2011
d.	Receive approval for study by CIRO (months 6-11)	Completed November 2011
e.	Establish administrative support for enrolling subjects (months 5-8)	Had a face to face meeting with Dr. Fritz and Dr. Cleland at a conference in November 2011 and February 2012. We walked through the entire methods. Subject folders have been created for data collection.
f.	Trial registered with clinicaltrials.gov (months 8-9) http://clinicaltrials.gov/ct2/show/NCT01556581	Trial registered in March 2012

Aim 1: Compare the effectiveness of two primary care management strategies for patients with a recent onset of combat-related LBP.

Task 1a (months 11-22): Enrollment into study (119 subjects). Active duty Soldiers with low back pain are randomized into one of two primary care management strategies (usual care stepped approach or early referral to physical therapy for treatment-based classification approach). Subjects are consented, baseline measures taken, randomization to treatment group occurs, and then allocated intervention is given.

Task #	Task Title	Status
1a	Recruitment and enrollment of subjects began in March 2012	We finished up enrollment at the at all sites, with over 400 subjects screened and

		119 enrolled.
	Other notes:	Met with Dr. Julie Fritz twice during this year, once while simultaneously attending a conference, and also a site-visit to UoU.

Task 1b (months 12-34): Track outcomes at 4 weeks, 12 weeks, and 1 year after initial enrollment.

Method of tracking outcomes: Follow-up re-assessments will be performed 4 weeks, 12 weeks and 1-year after the baseline examination. Follow-up assessments will be performed by a Research Assistant blinded to the patient's treatment group assignment. Subjects will be called 2 weeks prior to their projected follow-up date and scheduled a follow-up appointment. Subjects will arrive at their appointment and fill out the appropriate outcome measures. The data from the outcomes will be placed in a patient folder with only their subject ID for identification. Data will then be entered into a protected spreadsheet as described in the protocol.

Task #	Task Title	Status
1b	Follow-up of Subjects began	We have completed the 1-year follow-up for all subjects.
1c	Analysis of data to compare effectiveness	complete

Aim 2: Compare the subsequent healthcare utilization associated with two management strategies for patients with a recent onset combat-related LBP.

STATUS: Healthcare utilization was pulled from the Military Health System Data Repository (MDR) for all subjects enrolled in the study. Data was cleaned, abstracted, and merged with the self-reported outcomes. The data was then utilized for analysis of downstream healthcare utilization and costs between groups.

Aim 3: Compare any differences in psychosocial factors between success and failures within both groups of treatments.

STATUS: Analysis complete.

Challenges:

The primary challenge to this project was regarding recruitment. When we first planned the study, we had counted on normal sick call processes to capture patients seeking care for low back pain. Shortly after starting, a new sick call process went into place where patients could book appointments online rather than having to come in first thing in the morning. This meant that a much broader footprint was needed in primary care to capture all potential subjects coming in for low back pain, and the study had not been budgeted to accommodate this.

Plan: Findings have been disseminated at scientific conferences, and the primary manuscript is being completed for submission to a scientific journal. New follow-on projects to continue evaluating optimal care pathways for low back pain are in progress, stemmed from this initial project.

4. KEY RESEARCH ACCOMPLISHMENTS:

- The value of an optimal self-management program has been raised again, in light of these findings. There is need for improved self-management tools, that can be delivered outside of traditional settings.
- The need for improved outcome measures of disability, pain, and function specifically for service members has been raised.
- Early physical therapy was still more effective than usual care in the short-term (4 weeks).

5. **CONCLUSION:** There was no difference between usual care and early physical therapy after 1 year. Physical therapy provided greater improvement in disability after 4 weeks. As both groups improved, the impact of the education may have been underestimated. Patients in the physical therapy group utilized greater back pain-related healthcare resources, but overall healthcare costs did not differ compared to usual care. However, we may have underestimated the effects of a focused self-management program that addressed psychosocial beliefs and risk factors. All patients received this, which is a much better intervention than “usual care”. This needs further investigation in further prospective trials.

6. **PUBLICATIONS, ABSTRACTS, AND PRESENTATIONS:**

- (1) Lay Press: None
- (2) Peer-Reviewed Scientific Journals:
Rhon D, Fritz J. COMParative Early Treatment Effectiveness between physical therapy and usual care for low back pain (COMPETE): study protocol for a randomized controlled trial. *Trials* 2015;16:423.
PMID: 26399603
- (3) Invited Articles: None
- (4) Abstracts:
(Presentation of primary results)
Rhon DI, Fritz JM. *Effectiveness and downstream healthcare utilization for patients that received early physical therapy versus usual care for low back pain: a randomized clinical trial.*
- a. Presentations made during the last year

Rhon DI, Fritz JM. *Effectiveness and downstream healthcare utilization for patients that received early physical therapy versus usual care for low back pain: a randomized clinical trial.* Podium, International Low Back and Neck Forum, Oslo, Norway, September 2017. (publication pending)

7. **INVENTIONS, PATENTS AND LICENSES:** N/A

8. **REPORTABLE OUTCOMES:**




Figure 1. Primary outcome measure (Oswestry Disability Index). No significant between group difference out at 1 year (even after adjusting for baseline score, age, prior history of back pain, and psychosocial variables).




Figure 2. Pain rating. No significant between group difference out at 1 year (even after adjusting for baseline score, age, prior history of back pain, and psychosocial variables).




Figure 3. Downstream total healthcare costs and specifically low back pain related costs between groups. Significantly higher costs in the early PT group (likely related to PT visits).

There was no difference between usual care and early PT after 1 year. PT provided greater improvement in disability after 4 weeks. As both groups improved, the impact of the education may have been underestimated. Patients in the PT group utilized greater back pain-related healthcare resources, but overall healthcare costs did not differ compared to UC.

- 9. OTHER ACHIEVEMENTS:** As the primary focus of this award was to serve as a conduit for research career enhancement, it was a tremendous success on that account. This award created an excellent opportunity for MAJ Rhon, as an early career researcher, to develop skills and receive excellent mentoring from a seasoned and leading researcher in the field of back pain. The grant allowed for a postdoctoral project through the University of Utah. It also was instrumental in providing experience and pilot data for larger back pain related studies. The most recent, was a \$6.5 million grant awarded by a collaboratory between NIH, DoD, and the VA, to fund this same team as Co-Primary Investigators (Dr. Dan Rhon and Dr. Julie Fritz).

Story Link: <https://www.nih.gov/news-events/news-releases/federal-agencies-partner-military-veteran-pain-management-research>

10. REFERENCES:


1. Cohen SP, Gallagher RM, Davis SA, Griffith SR, Carragee EJ. Spine-area pain in military personnel: a review of epidemiology, etiology, diagnosis, and treatment. *Spine J* 2012;12(9):833–42.
2. Ernat J, Knox J, Orchowski J, Owens B. Incidence and risk factors for acute low back pain in active duty infantry. *Mil Med* 2012;177(11):1348–51.
3. Mattila VM, Kyröläinen H, Santtila M, Pihlajamäki H. Low back pain during military service predicts low back pain later in life. *PLoS One* 2017;12(3):e0173568.
4. Rhon D, Fritz J. COMParative Early Treatment Effectiveness between physical therapy and usual care for low back pain (COMPETE): study protocol for a randomized controlled trial. *Trials* 2015;16(1):423.

11. APPENDICES:

Appendix A: Protocol Publication

Appendix B: Abstract Presentation to International Low Back Pain Forum

Appendix C: Quad Chart



COMParative Early Treatment Effectiveness between physical therapy and usual care for low back pain (COMPETE): study protocol for a randomized controlled trial

Rhon and Fritz

STUDY PROTOCOL

Open Access



COMParative Early Treatment Effectiveness between physical therapy and usual care for low back pain (COMPETE): study protocol for a randomized controlled trial

Daniel Rhon^{1*} and Julie Fritz²

Abstract

Background: Low back pain is among the leading causes of medical visits and lost duty days among members of the United States Armed Forces and represents the highest 5-year risk of permanent disability in the US Army. For certain elements of care, the timing may be just as important as the type of care. The purpose of this study is to assess the impact of the timing of access to a physical therapist by patients with low back pain, by looking at outcomes and low back pain-related healthcare utilization over a 1-year period.

Methods/Design: This trial will be a two-arm pragmatic randomized clinical trial occurring at two different clinical sites in the Military Health System. We will assess outcomes and related downstream costs for patients who access physical therapy at the primary care level compared to those that receive usual care only. There will be 220 consecutive patients randomized to receive care in either group (early physical therapy or usual care only) for the first 4 weeks, and these patients will then be allowed to receive any additional care dictated by their primary care provider for the following year. The primary outcome measure is the Oswestry Disability Index. Secondary outcome measures are the Global Rating of Change, Patient Satisfaction and 1-year healthcare utilization. Follow-ups will occur at 4 weeks, 3 months and 1 year.

Discussion: This trial takes a pragmatic approach to delivering care by enabling a usual care environment for managing low back pain, while also allowing immediate access to physical therapy. After the initial intervention, the patient's primary provider can continue to manage the patient as he/she normally would in practice. The Military Health System Data Repository will capture all low back pain-related healthcare utilization that occurs in order to allow for a comparison between groups. Analysis from retrospective cohorts has shown improved outcomes and decreased costs for patients that received early versus late physical therapy, but this has yet to be shown in prospective trials.

Trial registration: ClinicalTrials.gov NCT01556581 initially on 14 March 2012.

Background

Low back pain is a priority healthcare condition due to its high prevalence and the economic healthcare burden it imposes on society. In fact, the World Health Organization has listed it in the top 10 conditions with the highest disease burden on society. The problem is even worse in the military where the stress and strain of combat add to the prevalence of this injury, making it

one of the largest causes of attrition in soldiers both in combat and in the garrison [1–4]. Over the last decade, low back pain has been among the leading causes of medical visits and lost duty days among members of the US Armed Forces [5, 6] and represents the highest 5-year risk of permanent disability in the US Army [7]. This makes it an important target for clinically oriented research to improve management and reduce costs.

Physical therapy is a treatment option for low back pain. Although utilization of physical therapy is relatively low, 7 to 16 % of those that seek care do so for low back pain, [8] and this utilization is slowly increasing [9, 10].

* Correspondence: daniel.i.rhon.mil@mail.mil

¹Center for the Intrepid, Brooke Army Medical Center, 3551 Roger Brooke Drive, Fort Sam Houston, TX 78234, USA

Full list of author information is available at the end of the article

Despite the increasing utilization of physical therapy and associated costs, there is surprisingly little information on the effectiveness of physical therapy as an early management strategy for patients with low back pain. Recent evidence suggests that the timing of physical therapy may be one of the more important variables related to prognosis because it is associated with improved healthcare costs downstream and decreased future healthcare utilization. Two recent studies found that early physical therapy (<14 days) was associated with significantly less healthcare utilization and costs for the 18- to 24-month period past the episode of back pain [11, 12]. A separate study compared an initial management strategy of either advanced imaging or physical therapy for low back pain and found that choosing advanced imaging over physical therapy as the initial strategy resulted in significantly greater overall 1-year healthcare costs related to back pain (average of \$4,793 more per person) [13]. This means that initial care decisions made by healthcare providers about low back pain, such as early referral to physical therapy, can have a substantial impact on cost and long-term outcomes. The timing of interventions, such as physical therapy, is an area that needs further inquiry, and has the potential to influence referral recommendations and treatment approaches for low back pain. While the value of early physical therapy for low back pain has been shown in cohort studies, it has not been evaluated prospectively in a clinical trial.

The purpose of this study is to assess the impact of early physical therapy in patients seeking care for low back pain. We will assess outcomes and related downstream costs for patients seen in primary care that either are referred for physical therapy immediately or are managed via usual care pathways.

Methods/Design

Study design

This study will be a prospective multicenter randomized clinical trial with two arms, comparing the effectiveness of two strategies for managing patients with a recent onset of low back pain. One strategy will be usual care (UC) based on a stepped care approach. If patients do not improve with initial care options, then the next level of care is provided, *etcetera* (for example, nonsteroidal anti-inflammatory drugs (NSAIDs) and rest, progression to advanced imaging and/or specialty care referral). The other strategy will involve immediate Physical Therapy (PT). The study was designed following the standard protocol items for randomized trials (SPIRIT) guidelines for planning clinical trials [14]. The study will examine functional outcomes and subsequent 1-year healthcare utilization in each group. The primary difference between the strategies is the management in the first 4 weeks. The UC group will be managed with stepped care

during this period, receiving advice, education, activity limitation guidance and medications if needed but no early physical therapy. The PT group will receive all the same care, but will also receive eight sessions of physical therapy guided by a treatment-based classification (TBC) approach during the first 4 weeks. The differences in care between the two interventions will occur primarily in the first 4 weeks. The approach will be pragmatic in order to compare the effects of the two management strategies under realistic clinical circumstances. Therefore, we will not attempt to balance provider contact time between groups over the first 4 weeks or control further treatment decisions made after the first 4 weeks. Assessors will be blinded to the treatment group.

Subject selection

Subjects will be a sample of convenience made up of male or female service members on active duty between the ages of 18 and 60 years (or emancipated minors), seen at two large military medical center. Subjects will be given the option to enroll in the study in consecutive order, as they report for medical care at the primary care clinics and meet the inclusion criteria. Women comprise 15 % of all soldiers, and therefore, we expect our sample to have 15 to 20 female subjects accordingly.

Hypothesis

The following hypotheses will be tested:

1. Early intervention strategies utilizing physical therapy will result in greater improvements in function and disability in the long-term compared to a stepped usual care approach.
2. Early intervention strategies utilizing physical therapy will result in decreased use of other medical resources (medications, imaging, additional healthcare) in the long-term (1 year) compared to a stepped usual care approach.
3. Baseline psychosocial factors will be associated with changes in clinical outcomes within each treatment group.

Recruitment and enrollment

Subjects will be recruited from two large U.S. Army Medical Centers from the regular pool of patients seeking medical care for low back pain. These patients will be approached and asked if they want to participate in the study. If they agree, they will then be consented and screened. Consent will be obtained from each participant. Ethical review and approval has been provided by the Western Region Medical Center Institutional Review Board at Madigan Army Medical Center.

Randomization

Randomization will be conducted using a blocked randomization list (block size 2 or 4) generated prior to the study. Randomization assignments will be sealed in opaque envelopes prior to beginning the study. Opening the randomization envelope will occur after completion of the baseline examination and the advice and education intervention, to avoid bias in the delivery of this aspect of the intervention. Patients in both groups will be instructed to follow-up with their primary care provider as needed. Patients in the PT group will be scheduled to begin physical therapy within 3 days.

We will record reasons for a subject dropping out of the study during any stage of the trial, and we will record all reasons for nonparticipation in the study to enable our ability to calculate an overall participation rate. The proposed flow of subjects through the study is depicted in Fig. 1.

Treatment procedures

Those that meet the eligibility criteria will go through a baseline assessment and fill out the baseline surveys and self-report measures. All subjects will be evaluated and medically screened for red flags or other medical issues that would make them ineligible to receive physical therapy. After the baseline examination and all baseline measures are complete, all patients will receive


a 20-minute educational session consistent with current evidence on self-management strategies (Additional file 1: Appendix 1). This will be considered the usual case. Every subject enrolled in the study will be given the option to receive a written excusal or modification of activity for up to 30 days and a 10-day prescription of NSAIDs and/or muscle relaxers if there are no medical contraindications (history of allergies to medications, GI upset, ulcers, GI pathology or renal pathology). Patients often receive this medication for back pain in primary care, and we do not want to encourage them to seek additional care outside the study in order to receive these medications if they have a strong desire for them, particularly if they are randomized to receive immediate physical therapy. Therefore, a reasonable dose will be offered to all subjects but will not be mandatory. Finally, the patient will be randomized either to receive only usual care or to see a physical therapist immediately for care.

Usual Care treatment group

Patients in the usual care (UC) group will receive the initial management outlined above. Consistent with a stepped care approach supported by practice guidelines, [15] no additional intervention will be provided to these patients, and they will be encouraged to follow-up for additional consultation on an as-needed basis if they feel they need additional care. At that time, consistent with practice guidelines, decisions on further treatments and/or referrals will be made by the patient's primary care provider based on clinical judgment. All aspects of treatment provided to the UC group are part of the standard of care.

Physical Therapy (PT) treatment group

Patients in the PT group will receive the same initial management described above, and will then be referred to physical therapy within 48 hours. A maximum delay of 2 days should be inconsequential as typical recovery for back pain is 6 weeks, with a plateau at around 12 weeks [16]. The physical therapy treatment will be determined using a treatment-based classification approach, which has acceptable reliability [17]. This approach attempts to match a patient with an appropriate treatment based on their clinical presentation and relevant impairments. The three primary initial treatment approaches will use manual therapy (Additional file 2: Appendix 2), core strengthening (Additional file 2: Appendix 3), or an extension-oriented treatment approach (Additional file 2: Appendix 4). Typical of clinical practice within participating physical therapy clinics, the treating physical therapist will be allowed to modify the initial treatment strategy after 2 weeks of treatment if the patient demonstrates either substantial clinical improvement or worsening of symptoms. Substantial



clinical improvement or worsening will be based on a repeat administration of the Oswestry Disability Index and patient self-report. Substantial improvement will be defined first by patient report and then verified objectively by administration of an Oswestry with improvement (that is, decrease in pain) of 50 % or more. Worsening will be identified first by patient report, and then defined as any deterioration (that is, increase) on a repeat Oswestry score. If either substantial improvement or deterioration is observed, the treating physical therapist may modify the initial treatment strategy. In other words, patients will also be progressed through the treatment program based on the clinical judgment of the treating physical therapist. Although they will start in one treatment group, they can progress to treatment in another group based on the directions of the physical therapist. Using the example above, if a patient has back pain with radicular leg symptoms that are successfully treated with the extension-oriented treatment approach after only five sessions and shows a subsequent improvement of 50 % on the Oswestry Disability Index, then the physical therapist may decide that the patient would benefit from core stabilization exercises and transition to this approach for the last three sessions, if impairments remain. All aspects of treatment provided to the PT group are part of the standard of care.

Patients in this group may receive a maximum of eight sessions over the 4-week treatment period (2 sessions per week). This is a typical treatment dosage for physical therapy for low back pain. At the end of this treatment period, patients are dismissed from care. If they feel that they need any further care, they are instructed to follow up with their primary care manager. All patients in this group will also be given an exercise program via handout (Additional file 3: Appendix 5) to perform at home.

Data collection

The baseline examination will include a physical examination and completion of the self-report questionnaires listed here. The primary outcome variable is the Oswestry Disability Index. Secondary outcome variables include numeric pain rating, global rating of change, additional healthcare utilization and lost work/training time.

Physical examination

A clinician will conduct a physical examination composed of the examination variables required to identify the patient's treatment-based cluster. These variables include questions related to the distribution and duration of the patient's symptoms, neural tension tests, measures of spinal range of motion including the effects of motion on the patient's symptoms, assessment of spinal mobility and tests of trunk muscle function.

Self-report questionnaires

After baseline assessment, the follow-up re-assessments will be performed at 4 weeks, 12 weeks and 1 year after the baseline examination. The assessor will be blinded to the patient's treatment group assignment.

Oswestry Disability Index

The Oswestry Disability Questionnaire (OSW) was originally described by Fairbank et al. [18] as a condition-specific measure of functional status for patients with LBP. The OSW is a 10-item scale with higher numbers indicating greater disability. We will use the modified version that replaces the sex life item with an employment/homemaking item due to poor compliance with the former [19, 20]. The OSW is widely used in research on non-operative management of patients with LBP [21]. Our previous research has found the modified OSW to be used in this study to have high levels of test-retest reliability among stable patients (ICC = 0.90), good construct validity and responsiveness to change for patients with acute LBP, with a minimum clinically important difference (MCID) of six points for patients with acute LBP receiving physical therapy [19].

Numeric Pain Rating Scale

A 0 to 10 numeric pain rating scale (NPRS; '0' indicating no pain, and '10' indicating the worst imaginable pain) will be used to assess LBP intensity. Numeric pain scales are known to have excellent test-retest reliability [20]. Our previous research has found the NPRS to be responsive to change, with a minimum clinically important difference of two points among patients with acute LBP receiving physical therapy [22].

Fear Avoidance Belief Questionnaire

The Fear Avoidance Beliefs Questionnaire (FABQ) [23] will be used to measure patients' beliefs about how physical activity and work may affect their LBP and perceived risk for re-injury. The FABQ contains two subscales: a seven-item work subscale (FABQW) and a four-item physical activity subscale (FABQPA). Test-retest reliability of the FABQ subscales is high [23, 24], and validity is supported by associations with disability and work loss in patients with acute and chronic LBP [25, 26]. Heightened fear-avoidance beliefs have been shown to be a risk factor for the development of chronic LBP following an acute episode [27, 28].

Pain Catastrophizing Scale

The Pain Catastrophizing Scale (PCS) is a 13-item patient-report scale developed to measure the extent to which people catastrophize in response to pain [29]. Each item is scored from 0 ('not at all') to 4 ('all the time'). The PCS is reported as a total score, with higher

scores indicating greater catastrophizing, and the scale is composed of three sub-scales: Rumination (four items, for example, 'When I am in pain, I keep thinking about how badly I want the pain to stop'); Magnification (three items, for example, 'When I am in pain, I become afraid that the pain will get worse'); and Helplessness (six items, for example, 'When I am in pain, I feel I can't go on'). The PCS has been shown to have high levels of internal consistency and construct validity [29, 30]. Pain catastrophizing has also been found to play a role in the transition from acute to chronic LBP [27].

Epworth Sleepiness Scale

The Epworth Sleepiness Scale (ESS) assessed the likelihood of falling asleep in eight scenarios commonly encountered in daily life [31]. The likelihood is ranked on a 0 to 3 scale, with 0 indicating that the individual would never doze and 3 indicating a high chance of dozing. It is one of the most commonly used self-assessment tools for measuring sleepiness [32]. Although not validated specifically in a population with low back pain, it has been shown to be a valid and reliable tool [33] for assessing sleep in other settings and populations [32, 34–37]. This may have a significant role in affecting outcomes of patients with low back pain as several preliminary studies have shown a correlation between chronic back pain and quality of sleep [38, 39]. Sleep deprivation is commonly seen in the active duty population with a high operational tempo.

Global Rating of Change

The patient will complete a global rating of change scale (GRC) at each follow-up. A 15-point scale is used as described by Jaeschke and colleagues [40], and this scale requires the patient to rate the degree of change in his or her condition from the beginning of treatment to the present. The midpoint of the scale is no change (0). Ratings from -1 to -7 represent varying degrees of a worsening of the patient's condition, whereas rating from +1 to +7 represent varying degrees of improvement.

Patient satisfaction Patient satisfaction with the care received for their LBP will be measured using a 10-item instrument that has been validated and found capable of distinguishing among three different dimensions of satisfaction (caring, information and treatment effectiveness) among patients with LBP attending primary care [41].

Pain body diagram

A body diagram will be completed to identify the location and nature of symptoms [42]. Body diagrams can be used to reliably categorize the most distal extent of a patient's symptoms [42, 43].

Profile data

A profile is the official instrument used to restrict physical activities and work duties due to medical conditions. It is the only official document that allows a soldier to miss work or training. It provides a list of the activities that cannot be performed along with a duration of time that the restriction is in effect. It is what will be used to track work-loss days. This is recorded in number of days.

Healthcare utilization

Finally, healthcare utilization data will be collected from the Military Health System Data Repository (MDR) database and will be confirmed via electronic medical record review. Healthcare utilization data will be used to determine any subsequent medical utilization related to low back pain and the economic impact of those injuries. These data will also allow a comparison of costs between the two treatment approaches used in this study.

Data analysis

The data analysis for hypothesis number 1 will use separate analyses of covariance (ANCOVA) to compare the mean change in the primary (OSW) and secondary (NPRS, FABQ, PCS and GRC) outcomes from baseline to each follow-up assessment between groups while controlling for the baseline level of the outcome being analyzed to account for regression to the mean. We will assess the influence of baseline psychosocial measures (PCS and FABQ) on outcomes in both groups by introducing the scores as covariates into the ANCOVA in order to address hypothesis #3. For hypothesis #2, we will use chi-square tests and calculate odds ratios with confidence intervals for the specific healthcare utilization frequently sought by patients with LBP (MRI, injections, work loss as measured by the DA3329 limited duty profile).

We will make every effort to achieve complete data collection. For data that are missing, we will use multiple imputation (MI). Under the MI approach, baseline and follow-up factors beyond the variable being analyzed can be incorporated into the imputation model to account for dependence of the missing data mechanism on other measured factors, such as predictors of patient characteristics and prior scores. We will apply the method of data augmentation using Markov Chain Monte Carlo (MCMC) to generate imputed values. Four steps are involved: a) preparing the dataset for MI, which includes identifying all outcome variables likely to be involved in later planned or unplanned analyses, as well as likely predictors of missingness itself, evaluating distributions for normality and outliers and taking appropriate transformations to normality as needed; b) carrying out the MI, where variance and covariance estimates based on

observed data are used to iteratively estimate maximum likelihood values for all participants, with multiple replacement values for each missing data point then being drawn randomly from the posterior distribution and perturbed with error; c) carrying out analyses identically on all versions of the imputed data; and d) combining tests and parameter estimates and adjusting for between-imputation variance to yield the final statistical results.

We will conduct all analyses based on intention-to-treat principles. This means that we will analyze all patients in the groups to which they were randomized regardless of compliance with the treatment protocols or follow-up assessments. Noncompliance with follow-up assessments will be handled using the MI approach outlined above. Per-protocol sensitivity analyses may be conducted if a high degree of treatment noncompliance is observed.

Sample size estimation

Back pain and lower extremity injuries make up 44 % of all injuries for which soldiers seek care. Patients with back pain are seen on a daily basis in this healthcare system. Assuming at least 85 % of the patients complete the OSW at 12 weeks, enrollment of 110 subjects per group (total sample size 220 subjects) will provide at least 84 % power to detect a difference of seven points change in OSW at 12 weeks, assuming a standard deviation in the change in OSW of 16 points (treatment effect = 43.8 % of one standard deviation). The MCID for the OSW has been estimated at 6 points [19]. Previous work with patients with acute LBP indicate that these estimates of anticipated effect size and standard deviation are realistic, [44–46] and would be consistent with detecting an effect that is at least slightly in excess of the threshold for seeing significant change.

The natural history of recovery from back pain and our experience in prior studies suggest that differences tend to become less pronounced at 1 year than at 12 weeks. We believe 12 weeks is the appropriate time frame for the primary endpoint because it reflects the potential impact of treatment, and clinically, if patients are not experiencing improvement at this time point, they seek additional treatments. Thus, we are using this endpoint for the purposes of our power analysis rather than 1 year. The 4-week assessment will allow us to evaluate immediate post-treatment changes, and the 1-year assessment permits us to evaluate the potential long-term effects.

Inclusion criteria

The inclusion criteria are as follows:

1. Military personnel on active duty and eligible for healthcare at a military treatment facility;

2. A primary complaint of low back pain defined as symptoms of pain and/or numbness between the 12th rib and buttocks with or without symptoms into the leg(s), which, in the opinion of the provider, are originating from tissues of the lumbar region;
3. Duration of current episode of low back pain $\leq$ 90 days;
4. Age 18 to 60 years (or emancipated minors on active duty); and
5. Available for the following 4 weeks to complete a regimen of treatment.

Exclusion Criteria

Exclusion criteria are as follows:

1. Oswestry Disability Index <20 % (prevent a ceiling effect).
2. History of receiving any medical care for this episode of low back pain within the last 3 months.
3. Prior surgery to the thoraco-lumbar spine or pelvis.
4. This episode of back pain is due to a traumatic fracture.
5. Pending a medical or physical evaluation board or discharge process, pending any litigation related to the condition, or planning on getting out of the military within the next 9 months.
6. Any “red flags” that would indicate a potentially serious condition or other significant disease process. These could include but not limited to cauda equina syndrome, large or rapidly progressing neurological deficit, fracture, cancer, ankylosing spondylitis, or other systemic disease.
7. Current episode occurred because of a motor vehicle accident.
8. Currently pregnant (or history of pregnancy in the previous 6 months).

Discussion

There are challenges to implementing a clinical trial such as this. One of these is the nature of treatments that may potentially not be made available to patients, particularly those randomized to early physical therapy. Unless red flag screening indicates otherwise, imaging will not be ordered for patients. While this follows published clinical guidelines, and although outcomes are not shown to improve (and in some cases worsen) when advanced imaging is ordered, studies show that patient satisfaction is associated with higher rates of ordering of imaging [47]. Another challenge is the implementation of what is traditionally a specialty care service (physical therapy), but now is used at the primary care level. Physical therapy is typically a sequential service following primary care management, even though it is only utilized

for 7-16 % of patients that present to primary care for back pain [11, 12].

It is feasible for patients with a new episode of LBP to receive physical therapy as part of an early management strategy [8]. However, physical therapy also contributes substantially to the healthcare costs associated with managing LBP [48, 49]. Along with the increasing utilization of physical therapy for LBP, are the corresponding costs that are also increasing [9, 10]. In the last 10 years, physical therapists in the US Army are now providing medical care further up on the front lines with the infantry soldiers as an organic asset to their unit. These therapists are in a prime position to facilitate early access to physical therapy interventions and have done so successfully in a combat-deployed setting [50, 51]. Considering that about 60 % of patients report recurrent symptoms, and 20 % report persistent disabling symptoms following an initial LBP episode [52, 53], research examining the most effective initial management strategies and the potential role of physical therapy is clearly needed. Suboptimal effectiveness of primary care management for patients with an acute episode of LBP can also lead to use of increasingly invasive and costly interventions, including injections, narcotic pain medications and surgical procedures [54–56]. In addition to the significant fourfold increase in prescription pain medication use in soldiers, other significant military healthcare costs are associated with the diagnostic management of these soldiers. In the current combat environment, advanced diagnostic imaging is not readily available [51]. Tests that would routinely be ordered in regular hospitals/clinics, when ordered in a combat setting, often require an evacuation out of the theater, incurring significant additional evacuation costs and time loss from duty. Especially for low back pain, many of these tests do not offer a valid prognosis or provide guidance for treatment and likely lead to an increase in overall medical costs [20, 57]. Reducing trends toward increased utilization and costs associated with prolonged episodes of LBP will require more effective early management strategies [58]. In a combat deployed setting, effective management strategies have the potential to decrease significantly the number of unnecessary evacuations out of the combat theater. The initial primary care encounter may therefore provide the best window of opportunity to improve outcomes and reduce costs related to low back pain.

Trial status

The study is still actively recruiting subjects at the time of submission.

Additional files

Additional file 1: Appendix 1. Self-management Handout. (PDF 5448 kb)

Additional file 2: Appendix 2. Treatment-based Classification Approach - Manual Therapy. **Appendix 3.** Treatment-based Classification Approach - Core Strengthening. **Appendix 4.** Treatment-based Classification Approach - Extension Oriented Treatment Approach. (PDF 792 kb)
Additional file 3: Appendix 5. Home Exercise Program. (PDF 2337 kb)

Abbreviations

FABQ: Fear Avoidance Belief Questionnaire; GRC: Global Rating of Change; LBP: low back pain; MDR: Military Health System Data Repository; MI: multiple imputation; MCID: minimum clinically important difference; MCMC: Markov Chain Monte Carlo; NPRS: Numerical Pain Rating Scale; OSW: Oswestry Disability Index; PCS: Pain Catastrophizing Scale; PT: physical therapy; SPIRIT: standard protocol items for randomized trials; TBC: treatment-based classification; UC: usual care.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

JF conceived the study. JF and DR derived the design of the study. All authors provided input into creating the protocol and making it applicable to the military setting. JF provided the statistical approach. DR drafted the manuscript. All authors read and approved the final manuscript.

Authors' information

JF is a Professor, Department of Physical Therapy and Associate Dean for Research at the University of Utah and served as a postdoc mentor to DR during this project. DR works as a physical therapist in the military health system.

Acknowledgements

This project was funded by the Career Development Award through the Peer Reviewed Orthopedic Research Program, part of the Congressionally Directed Medical Research Program: Award W81XWH-11-1-0657 (PRORP-CDA).

Disclaimer

This research was supported by the Department of Defense Peer Reviewed Orthopedic Research Program under award number (W81XWH-11-1-0657). Views and opinions of, and endorsements by the author(s) do not reflect those of Madigan Army Medical Center, Brooke Army Medical Center, the US Army or the Department of Defense.

Author details

¹Center for the Intrepid, Brooke Army Medical Center, 3551 Roger Brooke Drive, Fort Sam Houston, TX 78234, USA. ²Department of Physical Therapy, College of Health, University of Utah, 520 Wakara Way, Salt Lake City, Utah 84108, USA.

Received: 12 May 2015 Accepted: 14 September 2015

Published online: 23 September 2015

References

- Cohen SP, Brown C, Kurihara C, Plunkett A, Nguyen C, Strassels SA. Diagnoses and factors associated with medical evacuation and return to duty for service members participating in Operation Iraqi Freedom or Operation Enduring Freedom: a prospective cohort study. *Lancet*. 2010;375:301–9.
- Harman DR, Hooper TI, Gackstetter GD. Aeromedical evacuations from Operation Iraqi Freedom: a descriptive study. *Mil Med*. 2005;170:521–7.
- Cohen SP, Griffith S, Larkin TM, Villena F, Larkin R. Presentation, diagnoses, mechanisms of injury, and treatment of soldiers injured in Operation Iraqi Freedom: an epidemiological study conducted at two military pain management centers. *Anesth Analg*. 2005;101(4):1098–103.
- Cohen SP, Brown C, Kurihara C, Plunkett A, Nguyen C, Strassels SA. Diagnoses and factors associated with medical evacuation and return to duty among nonmilitary personnel participating in military operations in Iraq and Afghanistan. *CMAJ*. 2011;183:e289–295.
- Ernat J, Knox J, Orchowski J, Owens B. Incidence and risk factors for acute low back pain in active duty infantry. *Mil Med*. 2012;177:1348–51.

6. Center AFHS. Absolute and Relative Morbidity Burdens Attributable to Various Illness and Injuries. U.S. Armed Forces. Medical Surveillance Monthly Report MSMR. 2007;14:18–23.
7. Lincoln AE, Smith GS, Amoroso PJ, Bell NS. The natural history and risk factors of musculoskeletal conditions resulting in disability among US Army personnel. *Work*. 2002;18:99–113.
8. Freburger JK, Carey TS, Holmes GM. Physician referrals to physical therapists for the treatment of spine disorders. *Spine J*. 2005;5:530–41.
9. Feuerstein M, Marcus SC, Huang GD. National trends in nonoperative care for nonspecific back pain. *Spine J*. 2004;4:56–63.
10. Martin BI, Deyo RA, Mirza SK, Turner JA, Comstock BA, Hollingworth W, et al. Expenditures and health status among adults with back and neck problems. *JAMA*. 2008;299:656–64.
11. Childs JD, Fritz JM, Wu SS, Flynn TW, Wainner RS, Robertson EK, et al. Implications of early and guideline adherent physical therapy for low back pain on utilization and costs. *BMC Health Serv Res*. 2015;15:150.
12. Fritz JM, Childs JD, Wainner RS, Flynn TW. Primary care referral of patients with low back pain to physical therapy: impact on future health care utilization and costs. *Spine*. 2012;37:2114–21.
13. Fritz JM, Brennan GP, Hunter SJ. Physical therapy or advanced imaging as first management strategy following a new consultation for low back pain in primary care: associations with future health care utilization and charges. *Health Services Res*. 2015. doi:10.1111/1475-6773.12301.
14. Chan AW, Tetzlaff JM, Altman DG, Laupacis A, Gotzsche PC, Krleza-Jeric K, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med*. 2013;158:200–7.
15. van Tulder MW, Koes B, Seitsalo S, Malmivaara A. Outcome of invasive treatment modalities on back pain and sciatica: an evidence-based review. *Eur Spine J*. 2006;15 Suppl 1:82–92.
16. Artus M, van der Windt D, Jordan KP, Croft PR. The clinical course of low back pain: a meta-analysis comparing outcomes in randomised clinical trials (RCTs) and observational studies. *BMC Musculoskelet Disord*. 2014;15:68.
17. Henry SM, Fritz JM, Trombley AR, Bunn JY. Reliability of a treatment-based classification system for subgrouping people with low back pain. *J Orthop Sports Phys Ther*. 2012;42:797–805.
18. Fairbank JC, Couper J, Davies JB, O'Brien JP. The Oswestry low back pain disability questionnaire. *Physiotherapy*. 1980;66:271–3.
19. Fritz JM, Irrgang JJ. A comparison of a modified Oswestry Low Back Pain Disability Questionnaire and the Quebec Back Pain Disability Scale. *Phys Ther*. 2001;81:776–88.
20. Jensen MP, Turner JA, Romano JM. What is the maximum number of levels needed in pain intensity measurement? *Pain*. 1994;58:387–92.
21. Deyo RA, Battie M, Beurskens AJ, Bombardier C, Croft P, Koes B, et al. Outcome measures for low back pain research. A proposal for standardized use. *Spine*. 1998;23:2003–13.
22. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine*. 2005;30:1331–4.
23. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 1993;52:157–68.
24. Jacob T, Baras M, Zeev A, Epstein L. Low back pain: reliability of a set of pain measurement tools. *Arch Phys Med Rehabil*. 2001;82:735–42.
25. Crombez G, Vlaeyen JW, Heuts PH, Lysens R. Pain-related fear is more disabling than pain itself: evidence on the role of pain-related fear in chronic back pain disability. *Pain*. 1999;80:329–39.
26. Fritz JM, George SZ, Delitto A. The role of fear-avoidance beliefs in acute low back pain: relationships with current and future disability and work status. *Pain*. 2001;94:7–15.
27. Buer N, Linton SJ. Fear-avoidance beliefs and catastrophizing: occurrence and risk factor in back pain and ADL in the general population. *Pain*. 2002;99:485–91.
28. Fritz JM, George SZ. Identifying psychosocial variables in patients with acute work-related low back pain: the importance of fear-avoidance beliefs. *Phys Ther*. 2002;82:973–83.
29. Osman A, Barrios FX, Kopper BA, Hauptmann W, Jones J, O'Neill E. Factor structure, reliability, and validity of the Pain Catastrophizing Scale. *J Behav Med*. 1997;20:589–605.
30. Osman A, Barrios FX, Gutierrez PM, Kopper BA, Merrifield T, Grittmann L. The Pain Catastrophizing Scale: further psychometric evaluation with adult samples. *J Behav Med*. 2000;23:351–65.
31. Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep*. 1991;14:540–5.
32. Gander PH, Marshall NS, Harris R, Reid P. The Epworth Sleepiness Scale: influence of age, ethnicity, and socioeconomic deprivation. *Epworth Sleepiness scores of adults in New Zealand*. *Sleep*. 2005;28:249–53.
33. Johns MW. Reliability and factor analysis of the Epworth Sleepiness Scale. *Sleep*. 1992;15:376–81.
34. Lundt L. Use of the Epworth Sleepiness Scale to evaluate the symptom of excessive sleepiness in major depressive disorder. *Gen Hosp Psychiatry*. 2005;27:146–8.
35. Bausmer U, Gouweris H, Selivanova O, Goepel B, Mann W. Correlation of the Epworth Sleepiness Scale with respiratory sleep parameters in patients with sleep-related breathing disorders and upper airway pathology. *Eur Arch Otorhinolaryngol*. 2010;267:1645–8.
36. Chedraui P, Perez-Lopez FR, Mendoza M, Leimberg ML, Martinez MA, Vallarino V, et al. Factors related to increased daytime sleepiness during the menopausal transition as evaluated by the Epworth sleepiness scale. *Maturitas*. 2010;65:75–80.
37. Feng J, He QY, Zhang XL, Chen BY. Epworth Sleepiness Scale may be an indicator for blood pressure profile and prevalence of coronary artery disease and cerebrovascular disease in patients with obstructive sleep apnea. *Sleep Breath*. 2012;16:31–40.
38. O'Donoghue GM, Fox N, Heneghan C, Hurley DA. Objective and subjective assessment of sleep in chronic low back pain patients compared with healthy age and gender matched controls: a pilot study. *BMC Musculoskelet Disord*. 2009;10:122.
39. Marty M, Rozenberg S, Duplan B, Thomas P, Duquesnoy B, Allaert F. Quality of sleep in patients with chronic low back pain: a case-control study. *Eur Spine J*. 2008;17:839–44.
40. Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertain the minimal clinically important difference. *Control Clin Trials*. 1989;10:407–15.
41. Cherkin D, Deyo RA, Berg AO. Evaluation of a physician education intervention to improve primary care for low-back pain. II Impact on patients. *Spine*. 1991;16:1173–8.
42. Chan CW, Goldman S, Ilstrup DM, Kunselman AR, O'Neill PI. The pain drawing and Waddell's nonorganic physical signs in chronic low-back pain. *Spine*. 1993;18:1717–22.
43. Beattie PF, Meyers SP, Stratford P, Millard RW, Hollenberg GM. Associations between patient report of symptoms and anatomic impairment visible on lumbar magnetic resonance imaging. *Spine*. 2000;25:819–28.
44. Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med*. 2004;141:920–8.
45. Fritz JM, Delitto A, Erhard RE. Comparison of classification-based physical therapy with therapy based on clinical practice guidelines for patients with acute low back pain: a randomized clinical trial. *Spine*. 2003;28:1363–71.
46. Brennan GP, Fritz JM, Hunter SJ, Thackeray A, Delitto A, Erhard RE. Identifying subgroups of patients with acute/subacute "nonspecific" low back pain: results of a randomized clinical trial. *Spine*. 2006;31:623–31.
47. Chou R, Qaseem A, Owens DK, Shekelle P. Diagnostic imaging for low back pain: advice for high-value health care from the American College of Physicians. *Ann Intern Med*. 2011;154:181–9.
48. Kominski GF, Heslin KC, Morgenstern H, Hurwitz EL, Harber PL. Economic evaluation of four treatments for low-back pain: results from a randomized controlled trial. *Med Care*. 2005;43:428–35.
49. Dagenais S, Caro J, Haldeman S. A systematic review of low back pain cost of illness studies in the United States and internationally. *Spine J*. 2008;8:8–20.
50. Rhon DI. A physical therapist experience, observation, and practice with an infantry brigade combat team in support of Operation Iraqi Freedom. *Mil Med*. 2010;175:442–7.
51. Rhon DI, Gill N, Teyhen D, Scherer M, Goffar S. Clinician perception of the impact of deployed physical therapists as physician extenders in a combat environment. *Mil Med*. 2010;175:305–12.
52. Hestbaek L, Leboeuf-Yde C, Manniche C. Low back pain: what is the long-term course? A review of studies of general patient populations. *Eur Spine J*. 2003;12:149–65.
53. Dunn KM, Jordan K, Croft PR. Characterizing the course of low back pain: a latent class analysis. *Am J Epidemiol*. 2006;163:754–61.
54. Deyo RA, Mirza SK. Trends and variations in the use of spine surgery. *Clin Orthop Relat Res*. 2006;443:139–46.

55. Luo X, Pietrobon R, Curtis LH, Hey LA. Prescription of nonsteroidal anti-inflammatory drugs and muscle relaxants for back pain in the United States. *Spine*. 2004;29:531–7.
56. Friedly J, Chan L, Deyo R. Increases in lumbosacral injections in the Medicare population: 1994 to 2001. *Spine*. 2007;32:1754–60.
57. Carragee E, Alamin T, Cheng I, Franklin T, van den Haak E, Hurwitz E. Are first-time episodes of serious LBP associated with new MRI findings? *Spine J*. 2006;6:624–35.
58. Wasiak R, Kim J, Pransky G. Work disability and costs caused by recurrence of low back pain: longer and more costly than in first episodes. *Spine*. 2006;31:219–25.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Effectiveness and downstream healthcare utilization for patients that received early physical therapy versus usual care for low back pain: a randomized clinical trial

MAJ Daniel Rhon, Brooke Army Medical Center
Dr. Julie Fritz, University of Utah

Background: Low back pain is one of the primary musculoskeletal complaints that military service members seek care for, contributing to the 25 million injury-related limited duty days each year. Early physical therapy interventions have the potential to improve outcomes, and reduce downstream healthcare utilization and costs.

Methods: Active duty service members in the US Military Health System seeking care for low back pain from a general practitioner were recruited. After completing the visit with their GP, patients attended a 30-minute self-management class with strong focus on the psychosocial aspects of low back pain. Then patients were randomized to usual care (follow up with GP as needed) versus immediate start in a 3-week physical therapy program, and followed out to 1 year. The primary outcome was the Oswestry Disability Index at 1 year, and secondary outcomes included the numeric pain rating scale, global rating of change, and 1-year healthcare utilization and costs. An ANCOVA was used to compare differences between groups, adjusting for baseline score, age, pain catastrophizing, and fear avoidance beliefs. Significance was set at 0.05. The trial received ethics approval from the US Army Western Region Institutional Review Board, and was registered on [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01556581): NCT01556581

Results: 119 patients (mean age 27.2 years; mean bmi 27.8 kg/m²; 15.1% female) enrolled in the study (61 randomized to usual care; 58 to early PT). Patients in both groups had a significant improvement of approximately 50% on the Oswestry at 1 year compared to baseline, but no difference between groups. There was a significant difference in oswestry scores (mean change 5.3, 95CI 0.41 to 10.1; p=0.021) favoring the early PT group at 4 weeks, but no differences between groups at 12 weeks and 1 year. Total mean healthcare costs over the 1-year were about the same in each group (UC = \$3454, SD 375; PT = \$3138, SD 364), but the portion related specifically to low back pain was lower in the usual care group (\$1060; SD \$1513) compared to early PT (\$2016, SD \$1590).

Discussion and conclusions: While short term improvements at 4 weeks favored the early physical therapy group, at one year there was no difference between interventions. Patients in the PT group also utilized greater back pain-related healthcare resources during the 1 year follow-up. The study was underpowered, and it is possible that a Type II error could have prevented the detection of significant effect. As patients in both groups showed significant improvement at 1 year, it is possible that the influence of the 30-minute self-management class was underestimated.

Disclaimer: The view(s) expressed herein are those of the author(s) and do not reflect the official policy or position of Brooke Army Medical Center, the U.S. Army Medical Department, the U.S. Army Office of the Surgeon General, the Department of the Army, Department of Defense, or the U.S. Government.

Early Treatment-Based Classification versus Usual Care for Management of Low Back Pain (OR100037) PRORP: Career Dev Award



PI: MAJ Daniel Rhon Org: Brooke Army Medical Center/The Geneva Foundation

Problem, Hypothesis and Military Relevance

Problem: MSK injuries are a primary source of disability in the U.S. Military, accounting for \$548 million annually in direct costs. Low back pain is the leading single cause.

Hypothesis: The overall hypothesis guiding the study is that the additional initial treatment expense incurred by early implementation will result in superior short-term clinical effectiveness, and will be more cost-effective in the long-term due to reduced healthcare utilization.

Military Relevance: Over the last decade, LBP is among the leading cause of medical visits and lost duty days among members of the US Armed Forces.



Proposed Solution

- The objective is to compare the effectiveness and subsequent healthcare use associated with early physical therapy access compared with to a usual stepped care approach, looking at overall 1-year outcomes, overall healthcare utilization in each group, and differences in psychosocial beliefs between responders & non-responders.
- The hypothesis is that early physical therapy will be more cost-effective in the long term due to reduced healthcare utilization.

Timeline and Total Cost (direct and indirect)

Activities	1	2	3	4	5	6
IRB approval, training of personnel & med staff, recruitment initiation						
Ongoing recruitment to reach target goal of 220 subjects						
1-YR Follow-up complete						
Healthcare Utilization Analysis						
Estimated Budget (260.8K)	58.1k	72.2k	57.4k	48.5k	24.5k	

Early Treatment-Based Classification versus Usual Care for Management of Low Back Pain (OR100037) PRORP: Career Dev Award



PI: MAJ Daniel Rhon

Org: Madigan Army Medical Center/The Geneva Foundation

Award Information Log Number/Contract Number: W81XWH-11-1-0657 Period of Performance: 8/1/2011-7/31/2016 Award Amount: \$260,884 GOR: Miriam Darnell, Ph.D. Collaborators: Dr. Julie Fritz, PT, Ph.D. University of Utah	Problem Areas • Many problems early related to recruitment (leadership changes and changes in command support, sequestration affecting focus of the MTF, changes in procedures for LBP patients to seek care, etc). However, we adapted and modified our approach (several times) and were able to finish recruitment! • Funding for our PTA/RA was supposed to expire summer of 2014, but we have a no-cost extension approved for an extra year. The Site PIs (unfunded) are now helping manage the follow-up process.
Key Research Accomplishments • 119 subjects were enrolled and managed in our trial. • All subjects were followed for 1 year after enrollment • Analyses of primary outcome measures and healthcare utilization between the 2 groups has been completed.	Next Steps • Poor adherence to clinical practice guidelines for low back pain in the MHS continues to be a problem, leading to overutilization of healthcare resources and now tied with poor outcomes (use more resources = worse outcomes). • Validation of the STarTBack risk screening tool for LBP in the military is a crucial next step as it can help solve this problem. It is a simple and valid decision-making algorithm for appropriate referral/management of civilian patients with LBP (who should go early to PT and who can wait)