



NATIONAL DEFENSE RESEARCH INSTITUTE

Effects of Medication-Assisted Treatment (MAT) for Opioid Use Disorder on Functional Outcomes

A Systematic Review

Margaret A. Maglione, Laura Raaen, Christine Chen, Gulrez Shah Azhar, Nima Shahidinia, Mimi Shen, Ervant J. Maksabedian Hernandez, Roberta M. Shanman, Susanne Hempel

For more information on this publication, visit www.rand.org/t/RR2108

Published by the RAND Corporation, Santa Monica, Calif.

© Copyright 2018 RAND Corporation

RAND® is a registered trademark.

Limited Print and Electronic Distribution Rights

This document and trademark(s) contained herein are protected by law. This representation of RAND intellectual property is provided for noncommercial use only. Unauthorized posting of this publication online is prohibited. Permission is given to duplicate this document for personal use only, as long as it is unaltered and complete. Permission is required from RAND to reproduce, or reuse in another form, any of its research documents for commercial use. For information on reprint and linking permissions, please visit www.rand.org/pubs/permissions.

The RAND Corporation is a research organization that develops solutions to public policy challenges to help make communities throughout the world safer and more secure, healthier and more prosperous. RAND is nonprofit, nonpartisan, and committed to the public interest.

RAND's publications do not necessarily reflect the opinions of its research clients and sponsors.

Support RAND

Make a tax-deductible charitable contribution at
www.rand.org/giving/contribute

www.rand.org

Preface

Over the past two decades, the U.S. Department of Defense (DoD) has invested unparalleled resources into developing effective treatments for military-related psychological health conditions. Systematic reviews are a key component in the knowledge translation process and function to translate the available research into evidence-based health care guidelines that promote optimal clinical care. Although a few government agencies, including the U.S. Department of Veterans Affairs and the Agency for Healthcare Research and Quality, have established evidence synthesis centers, there is no similar center within DoD that focuses exclusively on psychological health issues. Thus, the Southern California Evidence-based Practice Center, housed at the RAND Corporation, has been awarded a three-year contract to synthesize research on psychological health interventions important to military populations. This document is a systematic review performed during year two of this three-year project. The review will be of interest to military health policymakers and practitioners who oversee or implement Medication Assisted Treatment for opioid use disorder.

None of the authors has any conflict of interest to declare.

This research is sponsored by the Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury and conducted within the Forces and Resources Policy Center of the RAND National Defense Research Institute, a federally funded research and development center sponsored by the Office of the Secretary of Defense, the Joint Staff, the Unified Combatant Commands, the Navy, the Marine Corps, the defense agencies, and the defense Intelligence Community. For more information on the RAND Forces and Resources Policy Center, see www.rand.org/nsrd/ndri/centers/frp.html or contact the director (contact information is provided on the webpage).

Abstract

At the request of the Defense Centers of Excellence (DCoE) for Psychological Health and Traumatic Brain Injury, this systematic review (PROSPERO 2017: CRD 42017058608) synthesizes evidence on the effects of medication-assisted treatment (MAT) on functional outcomes among patients with opioid use disorder (OUD). Functional outcomes included cognitive (e.g., memory), occupational (e.g., return to work), physical (e.g., fatigue), behavioral/social (e.g., family function), and neurological (e.g., balance) function.

We searched five databases from inception to January 2017 and bibliographies of systematic reviews to identify English-language controlled trials, case control studies, and cohort comparisons of one or more groups; cross-sectional studies were excluded. Two independent reviewers screened identified literature, abstracted study-level information, and assessed the quality of included studies. Meta-analyses used the Hartung-Knapp method for random-effects models. The quality of evidence (QoE) was assessed using the Grades of Recommendation, Assessment, Development and Evaluation (GRADE) approach.

A comprehensive search and 1,327 full-text publication screenings yielded 27 randomized controlled trials (RCTs) and ten observational studies meeting inclusion criteria. The studies reported highly diverse functional outcome measures. No RCT was rated as having low risk of bias, but several methodologically sound observational studies were identified. The statistical power to detect differences in functional outcomes was unclear in the majority of studies.

Regarding cognitive outcomes, a large observational study found that MAT users had twice the risk of injurious traffic accidents as nonusers (low QoE). Two studies reported that MAT users performed significantly worse in working memory and cognitive speed than matched controls with no history of substance use disorder (SUD) or opioid use (very low QoE). Regarding occupational outcomes, patients with OUD on MAT showed no differences from those treated without MAT. One cohort study found that fewer persons with OUD who were treated with buprenorphine reported fatigue than did persons with OUD who were untreated; other physical function outcomes either showed mixed evidence or no difference. One study showed patients taking buprenorphine or methadone scored worse in aggressive responding than did controls with no history of SUD (very low QoE); studies of other behavioral/social function outcomes either showed mixed evidence or no statistically significant difference.

Contents

Preface	iii
Abstract	iv
Figures and Tables	vii
Summary	viii
Acknowledgments	xv
Abbreviations	xvi
1. Introduction	1
2. Methods	4
Literature Search	4
Eligibility Criteria	4
Inclusion Screening	5
Data Extraction	5
Risk of Bias	6
Data Synthesis	7
Quality of Evidence	8
3. Results	9
Search Results	9
Description of Included Studies	10
Risk of Bias for Individual Included Studies	12
Key Question 1: What Are the Effects of MAT (Using Buprenorphine, Buprenorphine Plus Naloxone, Methadone, or Naltrexone) for OUD on Functional Outcomes Compared to Wait-List, Placebo, Treatment Without Medication, Any Other Comparator, or Each Other (e.g., Buprenorphine Versus Naltrexone)?	19
Key Question 1a: Does the Effect Vary by Type of Medication?	31
Key Question 1b: Does the Effect Vary by Route of Administration?	37
Key Question 1c: Does the Effect Vary by Length of Treatment, Follow-Up Time, or Later Cessation of MAT?	40
Key Question 1d: Does the Effect Vary by Treatment Modality (e.g., Methadone Clinic Versus Prescription Medication Taken at Home)?	43
4. Discussion	45
Summary of Findings	45
Prior Systematic Reviews	62
Strengths and Limitations	63
Implications for Future Research and Practice	64
Appendix A. Food and Drug Administration Warnings Table	66
Appendix B. Search Methodology	67
Appendix C. Risk of Bias Instruments	71

Appendix D. Evidence Table	77
Appendix E. Excluded Publications	114
Publications That Do Not Meet Inclusion Criteria (with Reasons for Exclusion).....	114
References	248

Figures and Tables

Figures

Figure 3.1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Article Flow Diagram	10
Figure 3.2. Insomnia: MAT (Buprenorphine, Naltrexone) Versus Placebo	23
Figure 3.3. Illegal Activity: Methadone Versus Passive Control.....	25
Figure 3.4. Percentage Arrested or Incarcerated: MAT Versus No MAT.....	27
Figure 3.5. Percentage Arrested or Incarcerated: Methadone Versus No MAT	28
Figure 3.6. Percentage Arrested or Incarcerated: Naltrexone Versus No MAT	28
Figure 3.7. Verbal Memory: Buprenorphine Versus Methadone.....	31
Figure 3.8. Attention: Buprenorphine Versus Methadone.....	32
Figure 3.9. Fatigue: Buprenorphine Versus Methadone.....	34
Figure 3.10. Insomnia: Buprenorphine Versus Methadone	35

Tables

Table 3.1. Risk of Bias for Controlled Trials.....	14
Table 3.2. Risk of Bias for Observational Studies	17
Table 4.1. Summary of Findings	47

Summary

Introduction

In response to the growing epidemic of opioid misuse, federal agencies in the United States were directed to improve access to medication-assisted treatment (MAT). MAT is the use of approved medications combined with counseling, other behavioral therapies, and patient monitoring to treat opioid use disorder (OUD). Medications approved in the United States for MAT for OUD include methadone, buprenorphine, Suboxone (a combination of buprenorphine and naloxone), and naltrexone.

The Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury commissioned a systematic review of the effects of MAT for OUD on functional outcomes. These outcomes include cognitive (e.g., memory), physical (e.g., fatigue), occupational (e.g., return to work), behavioral/social (e.g., family function), and neurological (e.g., balance) function. Such outcomes are important in determining whether active-duty service members can be deployed.

Key Questions

This review was guided by the following key question (KQ) and subquestions:

1. What are the effects of MAT (using buprenorphine, buprenorphine plus naloxone, methadone, or naltrexone) for OUD on functional outcomes compared with wait-list, placebo, treatment without medication, any other comparator, or each other (e.g., buprenorphine versus naltrexone)?
 - a. Do the effects vary by type of medication?
 - b. Do the effects vary by route of administration (e.g., oral versus injection versus implant)?
 - c. Do the effects vary by length of treatment, follow-up time, or later cessation of MAT?
 - d. Do the effects vary by treatment modality (e.g., methadone clinic versus prescription medication taken at home)?

Methods

To answer the KQs, we searched PubMed, PsycINFO, EMBASE, Cumulative Index to Nursing and Allied Health Literature, Cochrane CENTRAL, and Cochrane Database of Systematic Reviews from inception to January 2017, as well as bibliographies of existing systematic reviews to identify reports of English-language controlled trials, case-control studies, and cohort studies that compared two or more groups and reported baseline and follow-up data on functional outcomes. Cross-sectional studies were excluded, as were studies of MAT

medications not approved for use for OUD in the United States. We included two types of studies: (1) those that compared MAT-treated patients with OUD to persons with OUD who were not treated with MAT (i.e., they received another treatment, placebo, treatment as usual, or no treatment) and (2) those that compared MAT-treated patients with OUD to matched controls with no history of substance use disorder (SUD).

Two independent reviewers screened 6,292 identified citations using predetermined eligibility criteria. Because functional outcomes are most often reported as secondary outcomes in studies of substance abuse treatment, we retrieved full-text copies of all studies that assessed the efficacy of MAT for OUD and therefore potentially could meet our inclusion criteria and combed the results sections for relevant outcomes. Reviewers abstracted prespecified study-level information and assessed each included study's risk of bias (ROB); all abstracted data were checked by the project lead for accuracy.

Meta-analyses were conducted using the Hartung-Knapp method for random-effects models when sufficient data were available. Continuous outcomes were expressed as standard mean differences (SMDs), and categorical outcomes were expressed as relative ratios (RRs) together with the 95-percent confidence intervals (CIs). The overall quality of evidence (QoE) was assessed using the Grades of Recommendation, Assessment, Development, and Evaluation approach, and we differentiated high, moderate, low, and very low confidence in summary results (conclusions).

Results

Despite a comprehensive literature search that identified 1,327 publications that were scrutinized as full text, only 37 studies (27 randomized controlled trials [RCTs] and ten observational studies reported in 41 articles) met inclusion criteria. No RCT was rated as having low ROB, primarily because of lack of participant blinding, high attrition rate, and lack of reporting of the method of randomization and allocation concealment. Several observational studies with low ROB were identified.

The studies reported on a large number of highly diverse functional outcome measures, including verbal memory, attention, insomnia, fatigue, and criminal activity. Functional measures were primary outcomes in only six of the RCTs; it is unclear if the other trials, which were statistically powered to detect differences in illicit use of opioids or treatment retention, had adequate power to detect differences in functional effects.

KQ 1: Effects of MAT on Functional Outcomes

We found that although MAT patients performed significantly better on some functional outcomes than persons with OUD who did not receive MAT, they performed worse on several cognitive measures than did matched “healthy” controls with no history of SUD or opioid use.

Because of the limited number of studies identified and the moderate-to-high ROB of many of them, QoE is low or very low for all evidence statements.

Cognitive Function

Individuals with OUD using MAT versus healthy controls with no history of SUD.

A very large observational study conducted in France found that persons prescribed MAT had twice the risk of injurious traffic accidents than nonusers (low QoE). A cohort study measuring working memory found that OUD patients using either buprenorphine or methadone scored significantly worse than did matched controls with no history of SUD or opioid use (very low QoE). Another cohort study measured cognitive speed and found that OUD patients who used either buprenorphine or methadone scored significantly worse than did controls with no history of SUD or opioid use (very low QoE).

Two studies reported that MAT patients performed no worse than healthy controls in verbal memory tasks. One cohort study reported a difference in verbal memory favoring methadone patients, while another cohort study found no difference between methadone patients and healthy controls (very low QoE). Both studies found no difference between OUD patients taking buprenorphine and healthy controls (low QoE).

Two cohort studies showed no significant difference in attention between MAT-treated OUD patients and healthy controls with no history of SUD or opioid use (low QoE).

Individuals with OUD who were treated with MAT versus individuals with OUD not treated with MAT.

No studies that met our inclusion criteria compared cognitive outcomes between persons with OUD who were treated with MAT to persons with OUD who were not treated with MAT medications.

Occupational Function

Individuals with OUD who were treated with MAT versus individuals with OUD not treated with MAT.

Three RCTs and two observational studies that measured employment outcomes found no difference between MAT patients and persons with OUD treated for substance abuse without MAT (very low QoE).

Physical Function

Individuals with OUD who were treated with MAT versus controls with no history of SUD.

We identified no studies that reported physical function that met our inclusion criteria.

Individuals with OUD who were treated with MAT versus individuals with OUD not treated with MAT.

In one cohort study, a significantly lower percentage of patients with OUD who received buprenorphine reported fatigue than did untreated persons with OUD, while there was no difference in the rate of fatigue between persons with OUD who received methadone and these controls (very low QoE).

A meta-analysis of four RCTs found no difference in the percentage reporting insomnia between participants receiving MAT and those receiving a placebo (RR 1.02; CI 0.61, 1.71; moderate QoE). A methadone cohort study also reported no difference in insomnia (very low QoE).

One RCT comparing methadone versus a non-MAT intervention reported no difference in mean ASI medical scores. A cohort study comparing buprenorphine treatment with syringe exchange also reported no difference for this measure (very low QoE).

Behavioral/Social Function

Individuals with OUD who were treated with MAT versus controls with no history of SUD.

One small cohort study reported aggression outcomes; an RCT that randomized OUD patients to either buprenorphine or methadone found that individuals receiving either treatment scored significantly worse on aggressive responding than did controls with no history of SUD (very low QoE).

Individuals with OUD who were treated with MAT versus individuals with OUD not treated with MAT.

Studies reporting on crime (classified as a dysfunctional outcome) showed mixed results. Two RCTs reported that OUD patients on methadone spent fewer days engaged in criminal activity than those randomized to a placebo or wait-list (SMD -0.57; CI -1.00, -0.13; low QoE). However, a meta-analysis of RCTs that reported the percentage arrested or incarcerated found no statistically significant difference between patients randomized to MAT and those not randomized to MAT (RR 0.75; CI 0.46, 1.23; low QoE). An RCT and a cohort study that used a scale assessing illegal activities reported significantly better scores for OUD patients treated with MAT than for participants provided with psychosocially enhanced detox or syringe exchange. Another RCT and another cohort study reported no statistically significant difference for the mean number of charges or the mean number of arrests between MAT-treated patients and those not receiving treatment (all very low QoE).

One RCT reported no significant differences in mean family or psychiatric function scores between MAT and placebo groups; a cohort study also reported no statistically significant differences in a psychiatric function score (very low QoE).

Neurological Function

No studies that compared OUD patients who received MAT to those without MAT or that compared OUD patients who received MAT to controls with no history of SUD reported on neurological outcomes (e.g., hyporeflexia, balance, coordination).

KQ 1a: Effects by Type of Medication

Sixteen studies compared the effects of different MAT medications; of these, ten compared buprenorphine to methadone.

A meta-analysis of three RCTs that compared the effects of buprenorphine treatment on fatigue to that of methadone showed a significantly lower prevalence of fatigue in buprenorphine patients than in methadone patients (RR 0.62; CI 0.41, 0.95; moderate QoE). In absolute terms, 52 fewer buprenorphine patients per 1,000 reported fatigue compared with methadone patients.

Three RCTs that focused on cognitive function compared the effects of buprenorphine to methadone; no statistically significant differences in memory, cognitive speed and flexibility, attention, or vision were reported between treatments, with the exception of a small study where buprenorphine patients performed better than did methadone patients in vision tracking (low QoE for memory and attention because of high ROB and imprecision; very low QoE for cognitive speed, cognitive flexibility, and vision because of high ROB, imprecision, and lack of replication).

A meta-analysis of three RCTs reporting on insomnia found no statistical difference between buprenorphine and methadone groups (low QoE). Two RCTs reporting on pain perception found no significant difference between buprenorphine and methadone groups (low QoE). The only RCT of Suboxone versus methadone that reported functional outcomes found no significant difference in pain-rating scores at six months (very low QoE). One RCT of methadone versus naltrexone found no significant difference in the mean number of days patients engaged in illegal activity (very low QoE). A large nationally representative observational study found no differences in the increase in the proportion of participants who were employed and the number of arrests in the past 30 days among patients who received extended-release naltrexone administered by injection, oral naltrexone, Suboxone, or psychosocial treatment without medication (very low QoE).

KQ 1b: Effects by Route of Administration

We identified three RCTs that directly compared routes of administration of the same medication that reported on functional outcomes (physical and social function). None reported significant differences on these outcomes.

One RCT found no difference in risk of insomnia between oral Suboxone and Suboxone implant, and one RCT found no difference in effect on social function between methadone administered orally and by injection (very low QoE). Two RCTs found no difference in effects on mental health or physical health between methadone administered orally and that administered by injection (very low QoE). One large nationally representative observational study reported no differences in percentage employed or arrested during treatment among patients receiving Suboxone, oral naltrexone, injection naltrexone, or non-MAT treatment (very low QoE).

KQ 1c: Effects by Length of Treatment, Follow-Up, and Later Cessation

Three studies followed MAT patients longitudinally but none found an interaction effect of intervention by time on the reported results (ASI family component and psychiatric component) (very low QoE).

One RCT and one cohort study reported no statistically significant effect of length of treatment by treatment group on memory, attention, medical score, or legal score (very low QoE). A meta-regression across studies found no indication that RCTs with longer follow-up differed in effects on sleep (insomnia) or legal outcomes (arrests).

We identified only one study that addressed the question of how and when functional outcome effects change after cessation of MAT. The study compared MAT patients who remained in treatment to former patients no longer on MAT. Current MAT patients were significantly more likely to be not working in the past 30 days than former patients. The difference between current and former MAT treatment groups in terms of the percentage arrested in the past 30 days was not statistically significant (very low QoE).

KQ 1d: Effects by Treatment Modality

In the United States, methadone is traditionally dispensed under supervision at a methadone clinic, while buprenorphine is prescribed by a physician and can be distributed at a pharmacy. These two drugs are compared in KQ 1a. The results of studies that compared different modalities using the same medication are presented in the next section.

One cohort study with low ROB reported fewer nondrug-related crimes were committed by OUD patients prescribed methadone from a general practitioner's office than by those dispensed methadone at a traditional methadone clinic (very low QoE).

One RCT reported no differences in ASI psychiatric, legal, employment, or medical scores between a group take-home methadone program with twice a week distribution versus twice per month; however, the statistical power to detect differences in these functional outcomes is unknown (very low QoE).

Discussion

Some studies found significant effects in favor of MAT regarding the amount of criminal activity or legal status compared with persons with OUD not receiving MAT. In several studies, MAT patients performed significantly worse than matched controls with no history of SUD or opioid use on measures of aggression, working memory, and cognitive speed. However, it is unclear if the observed differences are because of MAT or because of long-term use of opioids in general. Although healthy controls are usually matched to patients on demographic and other characteristics, these individuals clearly differ in substance abuse history and may differ in unreported psychological, psychiatric, and family history characteristics that contribute to poor function. Quality of evidence for most outcomes was low or very low.

It is unclear in many instances if participants met standards required for military deployment. No studies were conducted on active-duty service members or reported performance on specific occupational tasks. No studies reported the current or former occupations of participants, and applicability of the outcome measures to successful military deployment was not discussed in any study.

Conclusions

Making clinical and policy recommendations is beyond the scope of the systematic review; the goal of this report was to summarize, synthesize, and assess the quality of the existing evidence. Despite an exhaustive and systematic search, the small number of studies that report on outcomes of interest and the weaknesses in the body of evidence prevent any strong conclusions about the effects of MAT on functional outcomes or differences in effects among medication types, route of administration, treatment modality, or length of treatment.

Acknowledgments

This research is sponsored by the Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury (DCoE). We thank Fuad Issa at DCoE, Melissa Fraine and Jennifer Bodart at the Defense Health Agency, and Thomas Concannon at RAND for their helpful feedback on the study protocol. We thank Thomas Concannon, William Sauve at Greenbrook TMS, and Marija Kelber and Marjorie Campbell of the Deployment Health Clinical Center for their helpful comments on the report. In addition, we thank Bradley Belsher, project monitor at DCoE overseeing this project, for his ongoing support of the project.

Abbreviations

AIDS	acquired immune deficiency syndrome
CI	confidence interval
CNS	central nervous system
DCoE	Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury
DoD	U.S. Department of Defense
DR2	decision and reaction test
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition
FDA	U.S. Food and Drug Administration
GRADE	Grades of Recommendation, Assessment, Development and Evaluation
HR	hazard ration
ITT	intention-to-treat
KEEP	Key Extended Entry Program
KQ	key question
MAP	Maudsley Addiction Profile
MAT	Medication-Assisted Treatment (for opioid use disorder)
MCQ	Memory Complaint Questionnaire
MD	mean difference
MMT	methadone maintenance therapy
NR	not reported
OMT	opioid maintenance treatment
OR	odds ratio
OD	opioid use disorder
PASAT	Paced Auditory Serial Addition Task
PSAP	Point Subtraction Aggression Paradigm
PVT	peripheral vision test
QOE	quality of evidence
RCT	randomized controlled trial
RD	risk difference
RE	random effect
ROB	risk of bias
RR	relative ratio
RST3	reactive stress tolerance test
RWT	Regensburg Word Fluency Test
SAMHSA	U.S. Substance Abuse and Mental Health Services Administration

SD	standard deviation
SEP	syringe-exchange program
SF-36	36-Item Short-Form Survey
SMD	standard mean difference
SUD	substance use disorder
TAP	Test for Attentional Performance
TAU	treatment as usual
TMT-A	Trail-Making Test A
VA	U.S. Department of Veterans Affairs
VLMT	Verbal Learning and Memory Test
WMS-III	Wechsler Memory Scale, Third Edition
XR-NTX	extended-release naltrexone

1. Introduction

Opioids are a class of drugs that includes the illicit drug heroin as well as the prescription pain relievers oxycodone, hydrocodone, codeine, morphine, and fentanyl (National Institute on Drug Abuse, undated). The misuse of these drugs has increased tremendously during the past 15 years. In 2014, 586,000 Americans had a substance use disorder (SUD) involving heroin, and 1.9 million had an SUD involving a prescription pain reliever (Substance Abuse and Mental Health Services Administration and Center for Behavioral Health Statistics and Quality, 2015). The same year, more than 28,000 people died of opioid overdose, an increase of more than 400 percent since 1999 (Rudd et al., 2016). According to a retrospective cohort analysis of data from 2008 through 2010 (Jones, 2013), 80 percent of new heroin users in the United States started out misusing prescription painkillers.

A study of almost 2,600 U.S. soldiers that had returned from deployment to Iraq or Afghanistan found that 44 percent had chronic pain and 15.1 percent regularly used opioids (Toblin et al., 2014). These rates are much higher than the estimates of 26 and 4 percent, respectively, for the general civilian population (Toblin et al., 2014). Drug overdose deaths more than doubled among active-duty personnel between 2006 and 2011 (Headquarters, Department of the Army, 2012). Prescription medications (most commonly oxycodone) were involved in 68 percent of those deaths.

Medication-Assisted Treatment (MAT) is the use of medications combined with counseling, other behavioral therapies, and patient monitoring to treat opioid use disorder. Medications approved in the United States for MAT include methadone, buprenorphine, a combination of buprenorphine and naloxone marketed as Suboxone, and naltrexone.

In response to the growing epidemic of opioid misuse, former President Barack Obama issued an Executive Memorandum in October 2015 (The White House, 2015) directing all federal agencies to improve access to MAT. Federal agencies that directly provide health care services, contract to provide health care services, reimburse for health care services, or facilitate access to health benefits were directed to identify any barriers individuals with opioid use disorder would encounter in accessing MAT and to submit an action plan to address the barriers. The U.S. Department of Defense (DoD), which provides health care services through TRICARE, is one such federal agency. The current DoD/U.S. Department of Veterans Affairs (VA) Clinical Practice Guideline for the Management of Substance Use Disorders strongly recommends the use of opioid agonists for MAT, either methadone in an Opioid Treatment Program or Suboxone through physician office-based opioid treatment (VA, 2015). In pregnant women, buprenorphine alone is recommended. The guideline strongly recommends extended-release injectable naltrexone, an opioid antagonist, for patients for whom treatment with an opioid agonist is contraindicated.

The DoD/VA recommendations were based on a systematic review of SUD treatment focused on effects on consumption of opioids and other drugs, adherence to treatment, and adverse events. The recommendations were also based on the balance of desirable and undesirable medication effects, values and preferences of patients, and DoD mission readiness; however, “functional outcomes” that may be related to occupational performance were not addressed by the systematic review. Neurocognitive ability is a key determinant of functional outcomes. Memory, reaction time, cognitive processing speed, and vigilance are measures of neurocognitive ability that affect problem solving, skill acquisition, occupational performance, and social function. Such outcomes are important in determining whether active-duty service members can be deployed. Currently, active-duty service members taking methadone for MAT are ineligible for deployment.

Medications used in MAT are described in the next paragraph. Appendix A shows the U.S. Food and Drug Administration (FDA) warnings regarding adverse events.

Methadone was developed in the 1940s and first introduced as a treatment for heroin addicts in the United States in the 1960s (Mattick et al., 2009). The FDA regulated the use of methadone for opioid dependent patients in 1972 (Ball and Ross, 1991). Methadone is administered orally, most often as a liquid. Diskette, powder, and tablet forms are also available (World Health Organization, 2009). Methadone must be prescribed by a physician and administered in a specialized clinic. Patients are typically initiated at low dosages, with the first dose between 10 and 20 milligrams (mg). The FDA limits the first dose to 30 mg for the first treatment day (World Health Organization, 2009). After, no regulations limit the maximum daily dose of methadone. However, doses typically range from 60 mg to 120 mg (American Society of Addiction Medicine, 2015). Recommended duration of treatment is at least one year, although treatment may be maintained for many years. Supplemental psychosocial support may be recommended depending on the individual’s situation.

Buprenorphine for opioid addiction was approved by the Drug Addiction Act of 2000 and by the FDA in 2002 (U.S. Department of Health and Human Services, 2005). Buprenorphine is a partial agonist opioid drug receptor modulator that can be prescribed alone or in conjunction with naloxone, an antagonist of the mu-opioid receptor. The combination of the two drugs is marketed as Suboxone. Buprenorphine can be prescribed only by physicians who have received specialized training or are certified in addiction medicine or psychiatry. No specific requirements regulate pharmacies that fill and dispense buprenorphine (U.S. Department of Health and Human Services, 2015); however, it is recommended that the first dose be administered in a setting where medical staff can observe the patient (McNicholas, 2004). Buprenorphine is most often administered either as a pill or sublingually. When administered with naloxone, the ratio of buprenorphine to naloxone is 4 to 1 (U.S. Department of Health and Human Services, revised 2012). The American Society for Addiction Medicine recommends that heroin patients receive their first buprenorphine dose within six to 12 hours after their last heroin dose and other opioid-dependent patients receive their first dose 24 to 72 hours after their last opioid dose. Duration of

buprenorphine maintenance treatment can range from a few months to many years. A buprenorphine implant was approved by the FDA in 2016.

Naltrexone was developed in 1963 and approved in 1984 by the FDA for the treatment of opioid dependence (U.S. Department of Health and Human Services, 2009). Naltrexone is an opioid antagonist, which means that patients undergoing naltrexone treatment should abstain from all opioids from seven to ten days before initiation through the completion of treatment. Extended-release naltrexone (XR-NTX) can be ordered by a physician, physician assistant, or nurse practitioner (U.S. Department of Health and Human Services, 2015). In the United States, naltrexone can be administered orally or intramuscularly. For oral administration, dosages are usually 50 mg per day or 100 mg three times per week supplemented with a 150 mg dose once per week for oral naltrexone. For intramuscular administration, a dose of 380 mg is typically administered every month (American Society of Addiction Medicine, 2015). Duration of treatment depends on the patient's symptoms and condition (American Society of Addiction Medicine, 2015). Supplemental treatment—such as counseling, therapy, or social support programs—is recommended during treatment (U.S. Department of Health and Human Services, 2015). Naltrexone is also available in an implant form in Australia and some European countries; however, it is not approved by the FDA for use in the United States.

This systematic review aims to synthesize estimates of the effects of MAT for opioid use disorder (OUD) on functional outcomes including cognitive (e.g., memory), occupational (e.g., return to work), physical (e.g., fatigue), behavioral/social (e.g., family function), and neurological (e.g., balance) functions.

We assess whether effects differ by type of medication, route of administration (e.g., injection or oral), length of treatment, and modality.

The following question and subquestions guide this systematic review:

1. What are the effects of MAT (using buprenorphine, buprenorphine plus naloxone, methadone, or naltrexone) on functional outcomes compared with wait-list, placebo, treatment without medication, any other comparator, or one another (e.g., buprenorphine versus naltrexone)?
 - a. Do the effects vary by type of medication?
 - b. Do the effects vary by route of administration (e.g., oral versus injection versus implant)?
 - c. Do the effects vary by length of treatment, follow-up time, or later cessation of MAT?
 - d. Do the effects vary by treatment modality (e.g., methadone clinic versus prescription medication taken at home)?

2. Methods

Literature Search

We searched PubMed, PsycINFO, EMBASE, Cumulative Index to Nursing and Allied Health Literature, Cochrane CENTRAL, and Cochrane Database of Systematic Reviews for English-language studies of the effects of MAT for OUD. The search strategy was developed by a senior librarian in RAND's Knowledge Services that was informed by search results of prior feasibility scans conducted for this project and existing systematic reviews on the topic. The search string is presented in Appendix B. Because functional outcomes are often reported as secondary outcomes (substance use measures are frequently primary outcomes), we did not restrict the literature search to citations referencing functional outcomes. Instead, we retrieved and screened full texts of trials and observational studies of the MAT interventions to determine whether relevant outcomes were reported in the publication. Prior systematic reviews on MAT were reference-mined for possibly relevant studies.

Eligibility Criteria

The inclusion and exclusion criteria are summarized in the following PICOTSS (participants, interventions, comparators, outcomes, timing, settings, and study design) framework.

- Participants
 - Studies of male and female participants, 18 years of age or older, were eligible for inclusion. Studies of pregnant women were excluded.
- Interventions
 - Studies evaluating methadone, buprenorphine, buprenorphine plus naloxone (Suboxone), or naltrexone for MAT for OUD were eligible, regardless of route of administration. Studies evaluating MAT interventions not approved in the United States, such as slow-release morphine or heroin, were excluded. Studies of opioid detox only, without maintenance, were excluded.
- Comparators
 - Studies that compared MAT with treatment of OUD without medications, with wait-list control, with no-treatment, or with other active treatments were included, as were head-to-head studies of MAT comparing medications, settings, and route of administration. Studies that compared MAT patients to matched controls with no history of SUD or opioid use were also eligible.

- Outcomes
 - Studies were required to report functional outcomes, including cognitive processing (e.g., memory, reaction time, attention, vigilance), occupational function (e.g., return to work), physical function, behavioral/social function (criminal activity, arrests, family function), or neurological function. Studies were required to assess outcomes in all participants or screen for the presence or absence of events in all included participants; studies reporting individual adverse events only for selected patients (e.g., reasons for dropping out of the study) were excluded.
- Timing
 - Studies could involve any treatment duration referred to as “maintenance,” and any follow-up period was eligible.
- Settings
 - Studies were limited to outpatient settings (i.e., methadone clinic or doctor’s office). Studies conducted in hospitals (inpatient), and residential rehabilitation facilities were excluded. MAT studies that began in prison and followed patients after release, through an outpatient phase, were eligible.
- Study design
 - Studies were limited to controlled trials, with or without random assignment, and observational studies (cohort or case-control) that compare two or more groups and report baseline and follow-up measures. Cross-sectional studies were excluded.

Inclusion Screening

Following a pilot session to ensure similar interpretation of the inclusion and exclusion criteria, two reviewers independently screened titles and abstracts of retrieved citations. Citations judged as potentially eligible by one or both reviewers were obtained as full text.

Full-text publications were screened against inclusion and exclusion criteria by two independent reviewers; disagreements were resolved through discussion within the review team with the project lead making the final decision. Reasons for exclusion at each stage were recorded in an electronic database.

Data Extraction

Data-collection forms were designed by the project lead with input from the project team and tested on three randomly selected studies. Forms were modified accordingly, and a final pilot on a random selection of studies was conducted to ensure agreement of interpretation. Two reviewers independently abstracted categorical study-level data using database software designed for systematic reviews, and discrepancies were resolved through discussion at weekly review team meetings. Qualitative information was abstracted by one reviewer and checked by the project lead. Two reviewers abstracted all outcome data; to ensure quality, the data extraction

accuracy was checked by the project lead and a statistician.

Information extracted from individual studies included the following:

- study ID and year
- participants: gender, age, race/ethnicity, years of opioid use, and population description
- interventions: medication used, dosage, route of administration, and any behavioral approach (e.g., cognitive behavioral therapy, substance abuse counseling, 12 step)
- comparators: type and description of comparator (e.g., wait-list control, treatment without medication, substance-abuse counseling plus placebo, and matched controls with no history of SUD, or head-to-head trials of MAT types/dosages)
- outcomes: functional domain, method of measurement, metric of data expression (e.g., means, proportions) and corresponding results (e.g., effect estimate, precision), and functional adverse events (e.g., insomnia) for all follow-ups
- timing: time-points of outcome assessment and duration of intervention
- setting: country and clinical setting/treatment modality where medication was administered
- study design: inclusion and exclusion criteria, sample size, and items relevant to risk of bias (ROB) assessment.

When two or more publications appeared to be reporting on the same study, participant descriptions were compared to identify studies with multiple publications and companion papers. All publications that contributed unique data to an included study were included for data extraction. All analyses were conducted at the study level; publications reporting on the same outcomes for the same participants entered the analyses only once, so participants were not counted multiple times in the findings. The review is based on data published in peer-reviewed scientific journals and did not include conference abstracts or dissertations.

Outcome data were based on intention-to-treat (ITT) analyses reported in the included studies. In the absence of ITT data, we used the number randomized as the denominator for count data and the number of participants at follow-up for means and proportions.

Risk of Bias

Reviewers extracted information on selection bias, performance bias, detection bias, attrition bias, and reporting bias of the included studies. The first five included studies were rated in dual; kappa for the ROB for these five studies was 0.82. The remaining studies were reviewed by one researcher and then reviewed for accuracy by the project leader. For controlled trials, we used the Cochrane Risk of Bias tool (Higgins and Green, 2011), which assesses the following sources of bias: random sequence generation (selection bias), allocation concealment (selection bias), blinding of participants and providers (performance bias), blinding of outcome assessors (detection bias), completeness of reporting outcome data (attrition bias), and selective outcome reporting (reporting bias). For the observational studies, we examined the representativeness of the MAT patients, the baseline similarity of the compared groups, efforts to match groups or use consecutive patients, how outcomes were obtained (official records, direct observation, or self-

report), and whether potential confounders were adjusted for. The full ROB instruments are included as Appendix C.

To determine study-limitation evaluations for the quality of evidence assessment and for potential sensitivity analyses, the project lead categorized each study as having overall low, moderate, or high ROB. We used the following methodology, based on the most important items from the Higgins and Green handbook:

- *Low ROB*: Comparable groups are initially assembled and maintained throughout the study with at least 80-percent follow-up at one year; reliable, valid measurement is used and applied equally to all groups; interventions are clearly described; all important outcomes are considered; appropriate attention is given to confounders in analysis; ITT analysis is used for trials.
- *Moderate ROB*: One or more of the following issues is found in the study: Some, but not major, differences between groups exist at follow-up; measurement instruments are acceptable but not ideal, although are generally applied equally; some, but not all, important outcomes are considered; some, but not all, potential confounders are accounted for in analyses. ITT analysis is used for trials.
- *High ROB*: One or more of the following “fatal flaws” is found in the study: Initially assembled groups are not comparable or maintained throughout the study, unreliable or invalid measurements are used or applied unequally across groups, key confounders are given little to no attention in analyses, and ITT analysis is not used.

Data Synthesis

The purpose of the systematic review was to synthesize the evidence of MAT for opioid use disorder on functional outcomes. We differentiated cognitive function, occupational function, physical function, behavioral/social function, and neurological function.

Effects in studies using categorical variables were summarized as relative ratios (RRs) together with their 95-percent confidence intervals (CIs) comparing treatment and control group results. Effects estimates for studies using continuous variables were computed as mean differences (MDs) between the treatment and comparison groups. Where more than one study was available that reported on the outcome of interest and studies did not use the same assessment scale, we computed standard mean differences (SMDs) to allow comparisons across studies that used different measures.

When sufficient data were available and clinical diversity was acceptable, we conducted meta-analysis to pool results across included randomized controlled trials (RCTs) for the outcomes of interest. To avoid population heterogeneity, we conducted separate analyses comparing (1) persons with OUD on MAT to persons with OUD not on MAT and (2) MAT patients to controls with no history of SUD or opioid use. We used the Hartung-Knapp-Sidik-Jonkman method for our random-effects meta-analysis (Hartung, 1999; Hartung and Knapp, 2001; Sidik and Jonkman, 2006). This approach is preferred when the number of studies to be pooled is small and when there is evidence of heterogeneity (IntHout, Ioannidis, Borm, 2014). It

has been shown that the error rates are more robust than with the DerSimonian and Laird method (Sánchez-Meca and Marin-Martínez, 2008). We report the I-squared statistic as a measure of between-study heterogeneity. Meta-analyses used the longest follow-up point where studies reported on multiple data points (data for all follow-ups are displayed in the evidence table in Appendix D).

Quality of Evidence

The quality of evidence was assessed for major outcomes using the Grades of Recommendation, Assessment, Development, and Evaluation (GRADE) approach (Balslem et al., 2011). Namely, the body of evidence was assessed based on the following dimensions: study limitations (low, medium, or high), consistency (consistent, inconsistent, or unknown), directness (direct or indirect), and precision (precise or imprecise) (Egger et al., 1997).

Per the GRADE system, if RCTs are identified that respond to a particular question, the quality of evidence is initially rated as high and then downgraded when results are primarily based on studies with substantial limitations; when results are inconsistent across individual studies or in the presence of substantial heterogeneity in pooled analyses; when the result is based on only a single study without replication in an independent research study; when conclusions are based on indirect evidence (e.g., effects based on meta-regressions across studies in the absence of head-to-head comparisons); and when pooled results are imprecise estimates of the treatment effect (CIs are wide, spanning effect sizes with different clinical conclusions).

We differentiated four levels of the quality of evidence:

- *High* indicates that the review authors are very confident that the effect estimate lies close to the true effect for a given outcome, as the body of evidence has few or no deficiencies. As such, the reviewers believe the findings are stable. That is, further research is very unlikely to change confidence in the effect estimate.
- *Moderate* indicates that the review authors are moderately confident that the effect estimate lies close to the true effect for a given outcome, as the body of evidence has some deficiencies. As such, the reviewers believe that the findings are likely to be stable, but further research may change confidence in the effect estimate and may even change the estimate.
- *Low* indicates that the review authors have limited confidence that the effect estimate lies close to the true effect for a given outcome, as the body of evidence has major or numerous (or both) deficiencies. As such, the reviewers believe that additional evidence is needed before concluding either that the findings are stable or that the effect estimate lies close to the true effect.
- *Very low* indicates that the review authors have very little confidence that the effect estimate lies close to the true effect for a given outcome, as the body of evidence has very major deficiencies. As such, the true effect is likely to be substantially different from the estimated effect; thus, any estimate of effect is very uncertain.


3. Results

Search Results

A total of 6,292 citations were identified for review through electronic literature searches and reference-mining of relevant systematic reviews (see Figure 3.1). Two trained reviewers excluded 4,965 citations before full-text review. The primary reason for exclusion was that the study did not assess MAT ($n = 4,425$) for OUD; these articles may have studied the use of MAT for other purposes (e.g., treatment of alcohol abuse, detoxification). Studies were also excluded if they studied a MAT medication not approved by the FDA ($n = 21$) or used study designs ($n = 492$) outside the scope of this review. Medications included in this review were limited to methadone, buprenorphine, Suboxone, and naltrexone. Study designs in this review included controlled trials and observational studies with at least one comparison group that reported baseline and follow-up data. Cross-sectional studies were excluded. Additionally, 23 duplicate articles were excluded, and four articles could not be retrieved using their citations.

Full-text review was also conducted by two independent reviewers. Based on the full-text review of 1,327 articles, 1,290 articles were excluded and 37 articles were included in the final analysis. Articles were excluded if they reported on MAT studies using a medication outside the scope of this review ($n = 160$), only measured the effects of a co-intervention such as counseling ($n = 241$), did not report any functional outcomes ($n = 307$), or used out-of-scope study designs ($n = 415$). Fifty-one articles reported on populations beyond the scope of this review (e.g., adolescents and pregnant women), 42 articles were excluded for setting (inpatient or residential), 13 were not published in English, 54 were conference abstracts, one was a dissertation, and two were excluded for other reasons. Appendix E lists excluded publications with reasons for exclusion. Four articles reported follow-up data on the 37 included studies; these data were abstracted and included in our analyses.

Figure 3.1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) Article Flow Diagram



Description of Included Studies

Design

Twenty-seven controlled trials, four case-control studies, and six cohort comparisons reporting baseline and follow-up data were included. Sample size for controlled trials ranged from 32 to 585 participants enrolled. Observational studies ranged from an open-label

nonrandomized study that enrolled 38 participants to a case-control study of all injurious traffic accidents in a French government database ($n = 72,685$).

Setting

The vast majority of studies were conducted in North America (19) or Europe (16). One study was conducted in Australia and another in Asia.

Fourteen studies were conducted at methadone clinics, nine were conducted at other substance abuse treatment centers, three were conducted at VA facilities, and seven were conducted at physician offices. There were six studies of “take-home” medications, administered either as an implant or as take-home methadone. In seven studies, MAT was initiated in prison, and patients were followed up after release. These numbers are not mutually exclusive, as head-to-head studies of MAT could be conducted in more than one setting; for example, methadone might be distributed at a methadone clinic, while in the same study, buprenorphine might be distributed at a physician’s office located in an outpatient clinic. Setting was unclear or not reported in eight studies; several of these studies assessed methadone, which can be assumed to have been administered under supervision at a methadone clinic.

Participants

The mean age of participants ranged from 24.4 (standard deviation [SD] 3.6) years to 44.4 (SD 9.2) years. Four studies enrolled only men, one study enrolled only women, and four studies did not report gender composition. The remaining studies ranged from 6.5 to 46.3 percent women. Racial composition was infrequently reported. Mean length of pre-intervention opioid use ranged from 2.0 to 17.7 years in controlled trials; the majority of trials required *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (DSM-IV) diagnosis for eligibility.

Interventions

The 37 studies included a total of 58 MAT arms: 34 methadone arms, 16 buprenorphine arms, five Suboxone arms, and seven naltrexone arms. The numbers do not add to 58, as two studies had MAT arms that grouped two medications together: one observational study of auto accidents in France grouped drivers taking prescription methadone or buprenorphine together, while an observational study of cognitive ability grouped patients on Suboxone with those on buprenorphine alone. Route of administration was oral (e.g., liquid methadone or sublingual buprenorphine) in 44 arms; three additional arms used injectable methadone; and five arms studied buprenorphine, Suboxone, or naltrexone administered via an implant that continuously supplied the medication for at least one month. Route of administration was not reported for six arms.

In controlled trials, the length of treatment at follow-up ranged from one to 36 months; the mean was 7.2 months (SD 7.0 months).

Comparators

Eleven of the 27 controlled trials were head-to-head trials of MAT that did not include a non-MAT comparison group. In addition, two of the ten observational studies did not have a non-MAT comparison. Six studies had a placebo group, six had a group that received counseling or psychosocial treatment only, three had matched, untreated persons with OUD, and three studies compared MAT patients to a matched control group with no history of SUD or opioid use. The following comparison groups were each included in one study: a syringe-exchange program (SEP), residential therapeutic community, and wait-list. One observational study assessed whether persons prescribed methadone or buprenorphine had a higher frequency of injurious traffic accidents than persons not on those medications. Another cohort study compared patients who had dropped out of methadone treatment to those who had stayed enrolled for several years. Finally, one study followed opioid-dependent prisoners who were randomized to either methadone or no treatment, after their release into the community.

Risk of Bias for Individual Included Studies

Controlled Trials

Table 3.1 shows ROB items and overall ratings for controlled trials. None of the controlled trials obtained an overall rating of low ROB, which required participant and researcher blinding, acceptable attrition rate, and intention-to-treat analysis. Nineteen trials were judged to have moderate ROB; the eight other trials were rated as having high ROB.

Random sequence generation. Eleven trials had unclear selection bias because they did not report their random sequence generation method; one trial reported an unsatisfactory method. Fifteen other trials reported adequate random sequence generation methods (e.g., computerized random generator).

Allocation concealment. ROB is low if the participants and investigators enrolling participants could not foresee assignment. Fourteen trials did not report an allocation concealment method. The allocation concealment method was adequate in 11 trials; in two trials, the method was inadequate.

Blinding of participants and providers. Participant blinding of substance abuse interventions can be difficult, especially if the comparator group is randomized to a waiting list or treatment as usual. Nineteen trials did not blind participants. Eight trials used identical placebos, and participants were blinded.

Blinding of outcome assessors. Three trials had unclear risk of detection bias because they did not report whether outcome assessors were blind to participant intervention conditions. Fourteen trials explicitly indicated that the outcome assessors were blind to intervention assignment.

Outcome data. Trials were judged to have attrition bias if more than 20 percent were lost to follow-up at one year or less or more than 30 percent were lost to follow-up at more than one year. Twelve trials had low risk of attrition bias, 13 had high risk, and two had unclear attrition. Only 12 trials used ITT analysis; for two trials, use of ITT was unclear.

Selective outcome reporting. No trials were judged to be high risk because of selective outcome reporting. Functional outcomes were usually secondary outcomes; the objective of most trials was to assess the effect of the intervention on illicit opioid use.

Other. The baseline demographic and substance use characteristics of comparison groups were statistically similar in the vast majority of trials (25). One trial did not report these statistics, while one trial reported significant differences at baseline.

Seven trials reported inadequate compliance with medication or differences in compliance between groups. Eleven trials did not report on compliance.

Table 3.1. Risk of Bias for Controlled Trials

Reference	Was Allocation Sequence (Randomization Method) Adequately Generated?	Was ALLOCATION Adequately Concealed (Prior to Assignment)?	Were Participants Adequately BLINDED?	Were OUTCOME ASSESSORS Adequately BLINDED?	Incomplete Outcome Data (ATTRITION BIAS) Because of Amount, Nature, or Handling of Incomplete Outcome Data	Is There Evidence of SELECTIVE OUTCOME REPORTING Bias (Yes/No)?	INTENTION-TO-TREAT Analysis? (Yes/No)	Group SIMILARITY AT BASELINE	Was There Incomplete COMPLIANCE with Interventions Across Groups?	Additional Bias: Bias Because of Problems Not Covered Elsewhere in the Table	Overall ROB
Bale et al., 1980	No	No	No	Yes	No	No	No	Unclear	Unclear	No	High
Compton et al., 2012	Unclear	No	No	No	Yes	No	No	Yes	No	No	High
Cornish et al., 1997	Unclear	Unclear	No	Unclear	Yes	No	Yes	Yes	Yes	No	Moderate
Coviello et al., 2010	Yes	Unclear	No	Yes	No	No	Yes	No	Unclear	No	Moderate
Dole et al., 1969	Yes	Unclear	No	No	No	No	No	Yes	Unclear	Yes	High
Fudala et al., 2003	Yes	Yes	Yes	Yes	No	No	No	Yes	No	No	Moderate
Gerra et al., 2007	Unclear	Unclear	No	No	Unclear	No	No	Yes	Unclear	No	High
Giacomuzzi et al., 2006	Unclear	Unclear	No	No	No	No	Unclear	Yes	No	No	High
Kinlock et al., 2009 (also reported in Gordon, 2008)	Unclear	Unclear	No	No	No	No	Yes	Yes	No	No	Moderate
Lee et al., 2016	Yes	Unclear	No	No	No	No	Yes	Yes	Unclear	No	Moderate
Ling et al., 2010	Unclear	Yes	Yes	Yes	Yes	No	Yes	Yes	No	No	Moderate
Lobmaier et al., 2010	Yes	Yes	No	No	Yes	No	Yes	Yes	Yes	Yes	High
Magura et al., 2009	Yes	Yes	Yes	Yes	Yes	No	No	Yes	Yes	No	Moderate
Mattick et al., 2003	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	No	Moderate

Reference	Was Allocation Sequence (Randomization Method) Adequately Generated?	Was ALLOCATION Adequately Concealed (Prior to Assignment)?	Were Participants Adequately BLINDED?	Were OUTCOME ASSESSORS Adequately BLINDED?	Incomplete Outcome Data (ATTRITION BIAS) Because of Amount, Nature, or Handling of Incomplete Outcome Data	Is There Evidence of SELECTIVE OUTCOME REPORTING Bias (Yes/No)?	INTENTION-TO-TREAT Analysis? (Yes/No)	Group SIMILARITY AT BASELINE	Was There Incomplete COMPLIANCE with Interventions Across Groups?	Additional Bias: Bias Because of Problems Not Covered Elsewhere in the Table	Overall ROB
Metrebian et al., 2015 (also reported in Byford et al., 2013)	Yes	Yes	No	Yes	No	No	Yes	Yes	Unclear	No	Moderate
Neri et al., 2005	Yes	Unclear	No	No	No	No	No	Yes	No	No	Moderate
Neumann et al., 2013	Yes	Unclear	No	Yes	Yes	No	No	Yes	Yes	No	High
Newman and Whitehill, 1979	Unclear	Unclear	Yes	Yes	Yes	No	Yes	Yes	Unclear	No	Moderate
Rosenthal et al., 2013	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	No	Moderate
Schwartz, 2009 (also reported in Schwartz, 2007)	Yes	Unclear	No	No	No	No	No	Yes	Unclear	No	Moderate
Sees et al., 2000	Yes	Yes	No	Yes	No	No	Yes	Yes	Yes	No	Moderate
Senay et al., 1993	Unclear	Unclear	No	Unclear	Unclear	No	No	Yes	Unclear	No	Moderate
Soyka et al., 2005	Unclear	Unclear	No	No	Yes	No	No	Yes	Unclear	No	High
Soyka et al., 2008	Unclear	Unclear	No	Yes	Yes	No	Unclear	Yes	Unclear	Yes	High
Strain et al., 1993	Unclear	Yes	Yes	Unclear	Yes	No	No	Yes	Yes	No	Moderate
Strang et al., 2000	Yes	Yes	No	Yes	No	No	No	Yes	No	No	Moderate
Tiihonen et al., 2012	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	No	No	Moderate

Observational Studies

Table 3.2 shows ROB items and overall ratings for observational studies. Four studies were judged as having moderate ROB, four were judged as having low ROB, and two were judged as having high ROB.

Participants. Five studies included participants who were truly representative of the average MAT patient in the community; these studies often took place in countries where all MAT patients are enrolled in a registry. Two studies included participants who were somewhat representative of the average MAT patient in the community, and two studies included a selected group of MAT users (e.g., volunteers). One study did not describe the derivation of the subject cohort. All studies either controlled for important factors such as demographic characteristics and substance abuse history in their analyses or, in the case of cohort comparisons, reported no significant differences in participant characteristics at baseline.

Outcome data. To be categorized as low ROB, observational studies were required to report less than 20 percent loss to follow-up at one year, less than 30 percent loss at one to five years, less than 40 percent loss at six to ten years, and less than 50 percent loss at 11 to 18 years. The attrition rate was inadequate in six studies. Three observational studies had 100 percent follow-up; these studies used administrative program data, legal records, or national registries and used record linkage to obtain outcome data. Five other studies used self-report of outcomes, while the other two used blind assessment of the participants.

Table 3.2. Risk of Bias for Observational Studies

Reference	Representativeness of the Exposed Cohort	Selection of the Nonexposed Cohort (e.g., Healthy Controls or Opioid-Dependent Participants Receiving a Different Medication or Not Receiving MAT)	Ascertainment of Exposure	Demonstration That Outcome of Interest Was Not Present at Start of Study	Comparability of Cohorts on the Basis of the Design or Analysis	Assessment of Outcome	Was Follow-Up Long Enough for Outcome to Occur?	Adequacy of Follow-Up Cohorts	Overall ROB
Aalto et al., 2011	Truly representative of the average MAT patient in the community	Drawn from a different source	Secure record	Yes	Controls for demographics, controls for additional factors	Self-report	Yes	Dropout rate not acceptable	High
Corsenac et al., 2012	Somewhat representative of the average MAT patient in the community	Drawn from the same community as the exposed cohort	Secure record	Yes	Controls for demographics, controls for additional factors	Record link	Yes	Complete follow-up—all subjects accounted for	Low
Coviello et al., 2011	Truly representative of the average MAT patient in the community	Drawn from the same community as the exposed cohort	Secure record, structured interview	Yes	Controls for demographics, controls for additional factors	Self-report	Yes	Subjects lost to follow-up unlikely to introduce bias	Low
Crits-Christoph et al., 2015	Selected group of users (e.g., volunteers)	Drawn from the same community as the exposed cohort	Secure record	Yes	Controls for demographics	Record link	Yes	Follow-up rate not acceptable	Moderate
Farrell-MacDonald et al., 2014	Selected group of users (e.g., volunteers)	Drawn from the same community as the exposed cohort	Secure record	Yes	Controls for demographics	Record link	Yes	Complete follow-up—all subjects accounted for	Low

Reference	Representativeness of the Exposed Cohort	Selection of the Nonexposed Cohort (e.g., Healthy Controls or Opioid-Dependent Participants Receiving a Different Medication or Not Receiving MAT)	Ascertainment of Exposure	Demonstration That Outcome of Interest Was Not Present at Start of Study	Comparability of Cohorts on the Basis of the Design or Analysis	Assessment of Outcome	Was Follow-Up Long Enough for Outcome to Occur?	Adequacy of Follow-Up Cohorts	Overall ROB
Giacomuzzi et al., 2005	No description of the derivation of the cohort	No description of the derivation of the nonexposed cohort	Secure record, structured interview	Yes	No differences in population characteristics	Independent blind assessment	Yes	Follow-up rate not acceptable those lost	High
Giacomuzzi et al., 2003	Somewhat representative of the average MAT patient in the community	Drawn from the same community as the exposed cohort	Secure record	Yes	Controls for additional factors	Self-report	Yes	Follow-up rate not acceptable	Moderate
Gossop et al., 1999	Truly representative of the average MAT patient in the community	Not applicable	Secure record	Yes	Controls for demographics	Self-report	Yes	Follow-up rate not acceptable	Moderate
Rapeli et al., 2009	Truly representative of the average MAT patient in the community	Drawn from a different source	Secure record	Yes	Controls for additional factors	Independent blind assessment	Yes	Follow-up rate not acceptable	Moderate
Reijneveld and Plomp, 1993	Truly representative of the average MAT patient in the community	Drawn from the same community as the exposed cohort	Secure record	Yes	No differences in population characteristics	Self-report	Yes	Subjects lost to follow-up unlikely to introduce bias	Low

Key Question 1: What Are the Effects of MAT (Using Buprenorphine, Buprenorphine Plus Naloxone, Methadone, or Naltrexone) for OUD on Functional Outcomes Compared to Wait-List, Placebo, Treatment Without Medication, Any Other Comparator, or Each Other (e.g., Buprenorphine Versus Naltrexone)?

The results are organized by the following functional effect outcome categories: cognitive, occupational, physical, behavioral/social, and neurological. Within these categories for KQ 1, we report comparisons of MAT patients to matched controls with no history of SUD separately from comparisons of persons with OUD on MAT to persons with OUD who do not receive MAT. Results from RCTs are reported first, followed by results from observational studies. Meta-analyses are documented in forest plots.

Results for comparisons between two MAT drugs such as methadone and buprenorphine are addressed in KQ 1a.

Cognitive Function

Individuals with OUD Who Were Treated with MAT Versus Controls with No History of SUD

We found two studies (Soyka et al., 2008; Rapeli et al., 2009) that reported on a battery of cognitive tests and one study (Corsenac et al., 2012) that reported on driving. The cognitive test results are documented in detail in the evidence table (Appendix D). Both studies reported on memory and attention.

Memory

Two studies measured verbal memory. Soyka et al. (2008) randomized persons with OUD to either oral methadone or sublingual buprenorphine. At eight to ten weeks, the patients completed the Verbal Learning and Memory Test (VLMT). Rapeli et al. (2009) likewise randomized persons with OUD to either oral methadone or sublingual buprenorphine; at six to nine months, patients completed a list-learning task from the Memory for Persons Data. Both studies compared patient results to those of matched controls with no history of SUD or opioid use. We converted the individual results to SMDs to facilitate the comparison across studies. Soyka et al. (2008) reported an SMD in verbal memory of 0.81 (CI 0.25, 1.36) in favor of patients on methadone while Rapeli et al. (2009) reported no difference between the groups (SMD 0.00; CI -0.74, 0.74). For patients on buprenorphine, no statistically significant differences were observed: Soyka et al. (2008) reported SMD 0.43 (CI -0.11, 0.98) and Rapeli et al. (2009) SMD -0.57 (CI -0.30, 0.16).

Rapeli also measured working memory using the Letter-Number Sequencing task from the Wechsler Memory Scale, Third Edition (WMS-III). At six to nine months, buprenorphine

patients scored significantly worse than controls (MD -2.8; CI -4.80, -0.80) as did methadone patients (MD -3.1; CI -5.08, -1.12).

Attention

The two studies previously described also measured attention. Soyka et al. (2008) reported the d2 Test of Attention at eight to ten weeks, while Rapeli reported results for the Test for Attentional Performance (TAP) at six to nine months in a separate publication from the memory results. The studies reported no statistically significant differences between patients on methadone and controls with no history of SUD or opioid use (Rapeli et al. [2009] SMD -0.38; CI -1.13, 0.37; Soyka et al. [2008] SMD -0.18; CI -0.72, 0.36). Both studies reported that buprenorphine patients scored lower than controls (Rapeli et al. [2009] SMD -0.63; CI -1.42, 0.16; Soyka et al. [2008] SMD -0.55; CI -1.10, 0.01) but the differences were not statistically significant.

Cognitive Speed

Soyka et al. (2008) measured cognitive and perceptual motor speed using the Trail Making Test of MAT treatment; lower scores indicate faster speed and better performance. At eight to ten weeks, patients on both buprenorphine (MD 5.5; CI 0.32, 10.68) and methadone (MD 5.9; CI 1.49, 10.31) scored significantly worse than controls with no history of SUD or opioid use.

Driving

One study that reported on driving ability met the inclusion criteria. Using traffic accident data from police reports and the national police database of injurious crashes, Corsenac et al. (2012) conducted a case-control study to investigate the association between the use of buprenorphine or methadone and the risk of being responsible for a road traffic crash ($n = 72,685$). Data on reimbursed medicines dispensed within six months before the crash were obtained by linking included drivers to the national health care insurance database using their national ID, gender, and date of birth. The authors found users of methadone or buprenorphine to be at a higher risk of injurious road traffic crashes. Analyses of odds ratios (ORs), which adjusted for age, gender, socioeconomic category, region, location, time of day, month, vehicle type, alcohol level, injury severity, concomitant levels 2 and 3 medicine exposure (for highest levels of risk), and long-term chronic diseases found use of buprenorphine or methadone to be associated with a two-fold increase in risk for injurious road traffic crashes (OR 2.02; CI 1.40, 2.91).

Individuals with OUD Who Were Treated with MAT Versus Individuals with OUD Not Treated with MAT

No studies that met our inclusion criteria compared cognitive outcomes between MAT patients and persons with OUD who were not treated with MAT medications.

Occupational Function

Individuals with OUD Who Were Treated with MAT Versus Controls with No History of SUD

No studies that met our inclusion criteria compared occupational function of MAT patients to controls with no history of SUD or opioid use.

Individuals with OUD Who Were Treated with MAT Versus Individuals with OUD Not Treated with MAT

Employment

Three RCTs that reported on occupational functioning compared patients on MAT to those enrolled in non-MAT interventions. Sees et al. (2000) compared a psychosocially enriched methadone maintenance program to a 180-day detoxification program. At 12 months, among 134 participants completing the study, they found no statistically significant difference (MD -0.01 ; CI $-0.10, 0.08$) in the Addiction Severity Index (ASI) employment score between methadone MAT and detox-only groups. At 12 months, Kinlock et al. (2009) found no significant difference in the number of days employed in the past 30 among male former prisoners randomized to either methadone MAT or passive referral upon release from a Baltimore corrections facility ($n = 204$). Finally, Coviello et al. (2010) reported the difference in percentage of criminal justice clients ($n = 111$) employed six months after release between a group on naltrexone and a group receiving standard psychological treatment without medication. They found 66 percent and 52 percent employment rates for the two groups, respectively; this difference was not statistically significant.

We also identified two observational studies that reported employment outcomes. Using the U.S. Substance Abuse and Mental Health Services Administration (SAMHSA) Treatment Episode Data Set, Crits-Christoph et al. (2015) conducted a retrospective analysis of opioid misusers who were under community supervision by a state correctional agency (i.e., on parole or probation) who received outpatient substance abuse treatment ($n = 873$). The authors compared participants who received XR-NTX, oral naltrexone, Suboxone, or psychosocial treatment without medication. Median length of treatment for patients on XR-NTX was 97 days, compared with 63 days for those on oral naltrexone ($p = 0.13$). Patients on Suboxone stayed 69 days ($p = 0.06$ compared with XR-NTX) and those who received treatment without MAT stayed 63 days ($p = 0.005$). Controlling for group differences using propensity scores based on several intake variables (e.g., severity), we found the difference among the four groups in the outcome “increase in the proportion of subjects who were employed (from intake to discharge)” to not be significant. In the second observational study, Reijneveld and Plomp (1993) conducted a small cohort comparison ($n = 38$) of patients who left the methadone maintenance system in the Netherlands with those who stayed five years. The authors discovered no differences in terms of “time spent on job” and “paid job.” However, more patients who stayed in treatment spent “sixteen hours or more with some legal aim per week.” The authors concede that their study was not designed to investigate the causes behind these differences.

Physical Function

Individuals with OUD Who Were Treated with MAT Versus Controls with No History of SUD

No studies that met our inclusion criteria compared physical function of MAT patients to controls with no history of SUD or opioid use.

Individuals with OUD Who Were Treated with MAT Versus Individuals with OUD Not Treated with MAT

For the physical function domain, we identified studies reporting on fatigue, insomnia, and an indicator of general medical problems. Results are described in the next section.

Fatigue


We identified one RCT that compared naltrexone to a placebo (Tiihonen et al., 2012). The randomized double-blind controlled trial of 100 drug-dependent patients observed no significant differences in fatigue between patients treated with naltrexone compared to those treated with a placebo implant (RR 1.00; CI 0.06, 15.55) at a ten-week follow-up.

A cohort study compared fatigue in 120 patients randomized with methadone or buprenorphine after six-month treatment to that of a group of untreated persons with OUD evaluated upon admission (Giacomuzzi et al., 2005). Fewer methadone patients reported fatigue than did the untreated opioid users (RR 0.78; CI 0.56, 1.09); however, the difference was not statistically significant. In the same study, buprenorphine patients were significantly less likely to report fatigue than the untreated group (RR 0.47; CI 0.29 to 0.76).

Insomnia

Four randomized controlled trials of MAT versus placebo reported the prevalence of insomnia among participants. Fudala et al. (2003), Ling et al. (2010), and Rosenthal et al. (2013) studied buprenorphine while Tiihonen et al. (2012) studied naltrexone. The studies conducted a follow-up at between one and six months ($n = 645$). Figure 3.2 shows the pooled results.

Figure 3.2. Insomnia: MAT (Buprenorphine, Naltrexone) Versus Placebo



NOTE: RE = random effect.

Across all available studies, there was no difference in those who reported insomnia between the MAT patients and the patients receiving a placebo (RR 1.02; CI 0.61, 1.71; 4 RCTs; I^2 18 percent). Little between-study heterogeneity was detected.

One additional study compared MAT patients with untreated persons to OUD evaluated upon admission. One hundred twenty opioid-dependent patients (Giacomuzzi et al., 2005) were randomized to either buprenorphine or methadone. At a six-month follow-up, fewer methadone and buprenorphine patients than control patients reported insomnia, but the difference was not significant (methadone: RR 0.72; CI 0.47, 1.08; buprenorphine: RR 0.72; CI 0.47, 1.08).

Overall Medical Problems

One RCT and one observational study reported the ASI medical score. One study used the European version of the index, while the other used the fifth U.S. revision. We converted results to SMDs to facilitate the comparison across studies.

Sees et al. (2000) randomized patients with OUD to either a methadone maintenance program or detoxification-only program. At 12 months, among 134 participants completing the study, they found no statistically significant difference (SMD -0.20; CI -0.54, 0.15) in the ASI medical score.

Aalto et al. (2011) conducted an observational study comparing the effectiveness of buprenorphine maintenance treatment with syringe exchange, using a matched set of patients in an SEP. Among the 60 enrolled participants, they found no statistically significant difference in mean ASI medical score at 12 months (SMD -0.57; CI -1.29, 0.15).

Behavioral/Social Function

Individuals with OUD Who Were Treated with MAT Versus Controls with No History of SUD

Aggression was the only behavioral/social function outcome reported in studies that compared MAT patients to controls with no history of SUD or opioid use.

Aggression

One study reported on aggression. Gerra et al. (2007) examined differences in aggression among patients randomized to either oral methadone or sublingual buprenorphine and a control group with no history of SUD or opioid use (total $n = 45$). Using the Point Subtraction Aggression Paradigm (PSAP) software program, the authors measured “aggressive responding,” “escape responding,” and “point-maintained responses” as metrics for aggression at three-month follow-up. More aggression is indicated by higher scores in the first two measures and a lower score for “point-maintained responses.” Compared with controls, the methadone group scored significantly higher in aggressive responding (MD 230.79; CI 188.79, 272.79) and escape responding (MD 82.79; CI -1, 166.58) and lower in point-maintained responses (MD -2256; CI -2509.48, -200.52). The buprenorphine group also scored significantly higher in aggressive responding (MD 226.97; CI 187.54, 266.4) and lower in point-maintained responses (MD -579.66; CI -786.67, -372.65). The difference in escape responding was not statistically significant (MD 39.89; CI -64.35, 144.129).

Individuals with OUD Who Were Treated with MAT Versus Individuals with OUD Not Treated with MAT

Several studies that compared individuals with OUD who were treated to MAT and those who were not treated reported on different aspects of social or behavioral function, including family functioning, psychological function, and criminal activity. Criminal activity, as a component of antisocial behavior, was assessed in multiple studies but operationalized in a variety of different ways (e.g., the number of arrests, ASI legal score).

Family Functioning

In the study by Sees et al. (2000), which randomized 179 participants to either methadone maintenance or a “psychologically enriched” detoxification program, the ASI was used to measure family function (a lower score indicates progress). At a one-year follow-up, the difference in scores between groups was not statistically significant (MD -0.01; CI -0.04, 0.02).

Psychological Function

Two studies (including one RCT) reported on effects on the ASI psychiatric score but each used different versions, and scores were converted to SMDs.


Sees et al. (2000) randomized patients with OUD to either a methadone maintenance program or detoxification-only program. The RCT reported no differences in the ASI psychiatric score between the two groups (SMD -0.20; CI -0.55, 0.14) at 12 months.

Aalto et al. (2011) conducted an observational study comparing the effectiveness of buprenorphine maintenance treatment with an SEP. Among the 60 enrolled participants, there was no difference between groups in mean ASI psychiatric score at 12 months (SMD 0.04; CI -0.67, 0.75).

Amount of Illegal Activity

Two RCTs of methadone that measured the amount of illegal activity among participants were identified. Schwartz et al. (2007) randomized 319 subjects with OUD to either oral methadone (mean 80 mg) or wait-list; at ten months, they reported the number of days subjects engaged in illegal activity in the past 30 days. Gordon et al. (2008) randomized 141 former prisoners to oral methadone (target 60 mg daily) or a passive referral to methadone and at six months reported the number of days subjects participated in crime. Figure 3.3 shows the pooled results across the two RCTs ($n = 460$).

Figure 3.3. Illegal Activity: Methadone Versus Passive Control



Methadone patients spent significantly less time engaged in crime than did control patients (SMD -0.57; CI -1.00, -0.13; 2 RCTs; I^2 74 percent). This analysis detected substantial between-study heterogeneity. While the direction of effects was similar, point estimates varied somewhat, but given the small number of studies, we were unable to investigate sources of systematic differences between the studies.

Additional studies also reported on crime outcomes, but reported data, study designs, outcomes, or comparator did not allow a combined analysis. A double-blind RCT of 95 opioid-dependent participants studied the effects of two methadone doses (20 mg and 50 mg) compared with a placebo group (Strain et al., 1993). At a five-month follow-up, the mean number of days of illegal activity in the past 30 was not significantly different between the 20 mg methadone

group and the placebo group (MD 3.00 days; CI -2.34, 8.34). However, the mean number of days of illegal activity in the past 30 was significantly lower in the 50 mg methadone group compared with the placebo group (MD -2.50 days; CI -7.63, -2.63). The number of crimes committed was also significantly lower in the 50 mg methadone group (MD -9.23) and 20 mg methadone group (MD -4.0) compared with the placebo group.


One RCT (Coviello et al., 2010) included 111 opioid-abusing criminal offenders, randomized to receive oral naltrexone with standard psychological treatment or standard psychological treatment alone (control). No significant difference was observed between the naltrexone group and control group in the average number of charges (MD -0.30) or convictions per month (MD 0.00) (SD not reported).

One additional, large cohort study (Crits-Christoph et al., 2015) also reported the number of days engaged in criminal activity. This study followed participants who received XR-NTX, oral naltrexone, Suboxone, or psychosocial treatment without medication (total $n = 2,882$) and found no significant differences across treatment groups in mean number of arrests 30 days after discharge. The authors suggested that their study had a short observation period and outcomes such as arrests should be studied over longer treatment durations and study periods.

Percentage of Participants Arrested, Incarcerated, or Engaging in Illegal Activity

Six RCTs of MAT versus no MAT reported the percentage of participants arrested or incarcerated. Comparators included passive referral (Kinlock et al., 2009), wait-list (Schwartz, 2009), behavioral treatment without MAT (Lee et al., 2016; Bale et al., 1980), or placebo (Dole et al., 1969; Cornish et al., 1997). Three studies reported a follow-up at six months, while the other three reported this measure at one year. Pooled results are shown in Figure 3.4 ($n = 1,302$).


Figure 3.4. Percentage Arrested or Incarcerated: MAT Versus No MAT



The difference in arrest rates between MAT patients and those not receiving MAT was not statistically significant (RR 0.75; CI 0.46, 1.23; 6 RCTs; I^2 85 percent). Substantial heterogeneity among studies was detected; thus, we conducted a sensitivity analysis. Excluding a methadone study from 1969 that reported the largest effect (Dole et al., 1969) did not substantially reduce heterogeneity (I^2 83 percent), but excluding all studies reporting on data before 1980 did reduce heterogeneity (I^2 66 percent); however, the effect estimate also did not change substantially and still did not show a statistically significant difference between studies (RR 0.81; CI 0.50, 1.30; 4 RCTs).

Four trials included in the above meta-analysis compared patients receiving methadone to participants who did not receive MAT (Kinlock et al., 2009; Schwartz, 2009; Bale et al., 1980; Dole et al., 1969). Pooled results are shown in Figure 3.5 ($n = 943$).


Figure 3.5. Percentage Arrested or Incarcerated: Methadone Versus No MAT



No difference was seen in the arrest rates between methadone patients and those not receiving MAT (RR 0.80; CI 0.33, 1.91; 4 RCTs; I^2 89 percent). Considerable heterogeneity among studies was detected in this analysis.

Two trials compared patients who received naltrexone to participants who did not receive MAT (Lee et al., 2016; Cornish et al., 1997). Pooled results ($n = 359$) are shown in Figure 3.6.

Figure 3.6. Percentage Arrested or Incarcerated: Naltrexone Versus No MAT



NOTE: CI numbers are obscured because the point estimates reported in these two studies were substantially different.

No significant difference in arrest/incarceration rates was observed between the naltrexone patients and participants not receiving MAT (RR 0.64; CI 0.02, 19.77; 2 RCTs; I^2 49 percent). Moderate between-study heterogeneity was detected, and the wide CI indicated that the point estimates reported in the two individual studies were substantially different, suggesting that the weighted average is not a good representation of the effect. (This explains why the CI numbers are obscured in the figure.) While both studies favored the MAT intervention, only one reported a statistically significant effect.

Three observational studies also reported the percentage of individuals arrested, incarcerated, or engaging in illegal activity. Based on retrospective administrative data on women offenders admitted to the Correctional Service of Canada's opioid maintenance treatment (OMT) program between 2003 and 2008 ($n = 137$), Farrell-MacDonald et al. (2014) assessed the effect of OMT on postrelease criminal reoffending and correctional readmission. Analysis by Cox proportional hazard modeling showed that patients on continued methadone maintenance therapy (MMT) had a 65-percent lower risk of returning to custody than the other two groups of terminated treatment post-release (MMT-T) and non-MMT (MMT-N) controls. Coviello et al. (2011) compared outcomes among 230 opioid-dependent patients who received active methadone maintenance or passive referral to a methadone clinic. At a six-month follow-up, the difference in the percentage of each group engaged in illegal activity in the past 30 days was not significant (RR 0.89; CI 0.45, 1.75). Reijneveld and Plomp (1993) compared outcomes of patients who left the Netherlands' methadone maintenance system to those who stayed five years ($n = 38$). Defining criminal problems as "caught at least one time by police in past half year" and "[a]t least some illegal activity/past week," they found no statistically significant differences between the two cohorts in terms of these criminal problems.

Legal Status

Two studies (Aalto et al., 2011; Sees et al., 2000) included ASI legal score as an outcome; this composite score considers self-reported frequency of illegal activity, illegal income, and feelings regarding severity of legal issues. Both studies found significant differences between MAT patients and the control group. The first study ($n = 60$) compared a matched cohort of patients enrolled in either buprenorphine maintenance treatment or an SEP and found the buprenorphine group had significantly better outcomes (i.e., lower ASI legal scores) at three, six, and 12 months (12-month MD -0.25 ; CI -0.39 , -0.11). The second study randomized 179 participants to either methadone maintenance or a "psychologically enriched" detoxification program: At a one-year follow-up, the methadone group had a significantly lower (better) mean ASI legal score (MD 0.08; CI 0.02, 0.14).

Neurological Function

No studies reported outcomes that could be categorized as neurological (e.g., hyperreflexia, balance, coordination).

Summary

With respect to cognitive function, two studies found that MAT patients performed as well or better on verbal memory tasks as healthy controls with no history of SUD or opioid use. Two studies showed no significant difference in attention between MAT patients and matched healthy controls; however, one of these studies measured working memory and found that both buprenorphine and methadone patients scored significantly worse than controls. One of these two studies also measured cognitive speed and found that both buprenorphine and methadone patients scored significantly worse than controls with no history of SUD or opioid use.

Regarding occupational function, three RCTs and two observational studies found no significant differences between MAT patients and persons with OUD treated without medication. Our meta-analysis of four RCTs found no significant difference between subjects receiving MAT and those receiving placebo in the percentage reporting insomnia.

Regarding physical function, MAT was not associated with fatigue in one cohort study and one RCT. In one RCT, subjects completed the ASI family and psychiatric components; the authors reported no significant difference in mean score between MAT and placebo groups.

Regarding social or behavioral function (including family functioning, psychological function, and criminal activity), one small study reported aggression outcomes: Both buprenorphine and methadone users scored significantly worse than matched controls with no history of SUD or opioid use on aggressive responding. In addition, a very large observational study conducted in France found that MAT (either buprenorphine or methadone) users had twice the risk of injurious traffic accidents than nonusers. Most studies that measured crime reported that persons with OUD randomized to MAT committed fewer crimes than those randomized to placebo or passive control (e.g., wait-list). Likewise, two studies reporting the ASI legal component found MAT patients had significantly better scores than persons with OUD provided with psychosocially enhanced detox or syringe exchange. However, our meta-analyses of RCTs that reported the percentage arrested or incarcerated found no significant difference between patients randomized to MAT or no MAT.

Key Question 1a: Does the Effect Vary by Type of Medication?

This section compares effects by type of medication. Evidence stems from studies comparing patients taking methadone to patients taking buprenorphine, a study comparing Suboxone (buprenorphine and naloxone) to methadone, a study comparing buprenorphine alone with Suboxone, and a study comparing the naltrexone implant with oral methadone. Results are reported by functional outcome category, first presenting results from studies that randomly assigned participants, followed by evidence from observational studies.


Cognitive Function

We identified three RCTs (Soyka et al., 2008; Rapeli et al., 2009, Soyka et al., 2005) that reported on a battery of cognitive tests, comparing groups taking different MATs. The cognitive test results are documented in detail in the evidence table (Appendix D). Results on memory, attention, and vision are presented in detail in the next sections.

Memory

Two head-to-head RCTs measured verbal memory. Soyka et al. (2008) randomized persons with OUD to either sublingual buprenorphine or oral methadone. At eight to ten weeks, patients completed the VLMT. Rapeli et al. (2009) likewise randomized persons with OUD to either sublingual buprenorphine or oral methadone; at six to nine months, patients completed a list learning task from the Memory for Persons Data. Figure 3.7 shows the pooled results for patients on buprenorphine compared with patients on methadone.

Figure 3.7. Verbal Memory: Buprenorphine Versus Methadone



NOTE: The numbers are somewhat obscured because the wide CI of the pooled result indicated that the effect estimates were substantially different.


Pooled analysis ($n = 87$) indicates that the difference in effects on verbal memory was not statistically significantly different between the groups (SMD -0.34 ; CI $-2.44, 1.76$; 2 RCTs; I^2 9 percent). The analysis detected very little between-study heterogeneity, given that only two small studies were included. However, the wide CI of the pooled result indicated that the effect estimates were substantially different. (This is why the CI numbers are somewhat obscured in the figure.) Still, both individual studies reported the same direction of effects.

Rapeli et al. (2009) also measured working memory using the Letter-Number Sequencing task from the WMS-III. At six to nine months, the difference in mean score between the buprenorphine and methadone groups was not significant (MD 0.30; CI $-1.43, 2.03$). Similarly, in the Rapeli et al. study, scores on the Paced Auditory Serial Addition Task (PASAT), which also measures working memory, were not significantly different at six to nine months (MD 1.6, CI $-4.94, 8.14$).

Attention

The two trials just described also measured attention. Soyka et al. (2008) reported on the d2 Test of Attention at eight to ten weeks, while Rapeli et al. (2009) reported results for the TAP at six to nine months in an article published two years after publication of the memory results. One additional trial that randomized 46 patients to either buprenorphine or methadone measured effects on attention (Soyka et al., 2005). Figure 3.8 shows the pooled results for patients on buprenorphine compared with patients on methadone ($n = 131$).

Figure 3.8. Attention: Buprenorphine Versus Methadone



Pooled analysis indicates that the difference in attention across all available studies was not statistically significant (SMD -0.12; CI -0.79, 0.51; 3 RCTs; I^2 14 percent). The analysis detected very little heterogeneity among studies.

Vision

Patients enrolled in the RCT by Soyka et al. (2005) also completed a peripheral vision test as part of the Act and React Test System. No difference in mean reaction time (seconds) was observed between buprenorphine and methadone patients at eight to ten weeks (MD 0.00, CI -0.23, 0.23). Buprenorphine patients scored significantly better than methadone patients in visual tracking performance (MD -1.10; CI -2.20, 0.00). No statistically significant difference in visual perception was observed, as measured by correct answers on a tachistoscope test (MD -0.20; CI -2.58, 2.18).

Cognitive Speed

Soyka et al. (2005) also assessed cognitive and perceptual motor speed using the Trail Making Test at eight to ten weeks of MAT treatment. No significant differences were observed between patients on buprenorphine versus methadone (MD -0.4; CI -5.98, 5.18).

Cognitive Flexibility

Soyka et al. (2005) also reported no statically significant difference in mean Regensburger Word Fluency Test scores between the buprenorphine and methadone treatment groups (MD 0.90; CI -3.57, 5.37). This test is a commonly used measure of cognitive flexibility.

Occupational Function

No head-to-head trials comparing different medications reported occupational outcomes. One large observational study reported on employment.

Employment

As noted previously, using the SAMHSA Treatment Episode Data Set, Crits-Christoph et al. (2015) compared patients ($n = 873$) who received XR-NTX, oral naltrexone, Suboxone, or psychosocial treatment without medication. Controlling for group differences using propensity scores based on several intake variables (e.g., severity), the difference among the four groups in the outcome “increase in the proportion of subjects who were employed (from intake to discharge)” was not significant.


Physical Function

Within the physical function domain, we identified head-to-head studies that reported on fatigue, insomnia, pain, and nausea.

Fatigue

Three RCTs ($n = 547$) comparing buprenorphine with methadone reported the percentage of subjects who experienced fatigue or somnolence during the trial (Mattick et al., 2003; Neri et al., 2005; Giacomuzzi et al., 2006). Follow-up times ranged from 13 weeks to one year. As shown in Figure 3.9, fewer buprenorphine patients than methadone patients reported fatigue across studies; the effect was statistically significant (RR 0.62; CI 0.41, 0.95; 3 RCTs; I^2 5 percent).

Figure 3.9. Fatigue: Buprenorphine Versus Methadone




The analysis detected very little between-study heterogeneity.

Insomnia

The three RCTs mentioned immediately above also reported the number of patients experiencing insomnia (Mattick et al., 2003; Neri et al., 2005; Giacomuzzi et al., 2006).

Figure 3.10. Insomnia: Buprenorphine Versus Methadone



The difference in prevalence between the buprenorphine and methadone groups was not statistically significant (RR 1.11; CI 0.70, 1.75; 3 RCTs; I^2 6 percent). The analysis detected very little heterogeneity among the studies.

Pain

Two RCTs of buprenorphine versus methadone reported pain outcomes. Compton et al. (2012) randomized 82 patients with OUD to either sublingual buprenorphine or oral methadone. Pain response was measured at baseline (treatment entry) and at 12 to 18 weeks. Hyperalgesia was present among OUD patients at baseline and did not improve significantly over the course of treatment. No difference was seen in pain detection (measured in both volts and seconds to detection) or pain tolerance (also measured in both volts and seconds to detection) between the buprenorphine and methadone groups (details provided in the evidence table in Appendix D).

Mattick et al. (2003) also conducted an RCT that compared sublingual buprenorphine and oral methadone ($n = 405$ enrolled, 216 completed). The percentage of patients reporting pain during the trial did not differ between the groups (RR 0.95; CI 0.59, 1.53).

One RCT that compared Suboxone to methadone also reported on pain. Neumann et al. (2013) reported patient Numeric Pain Rating Scale (range 1 to 10, lower scores better) scores at six months; the difference between the groups was not statistically significant (MD -0.96 ; CI -3.54 , 1.63).

Finally, Fudala et al. (2003) randomized patients with OUD to either Suboxone or buprenorphine alone. The difference in the percentage of patients who reported experiencing pain during the trial at four weeks was not statistically significant (RR 1.20; CI 0.71, 2.04).

Nausea

Neri et al. (2005) randomized 62 patients with OUD to sublingual buprenorphine or oral methadone. At a one-year follow-up, the percentage of patients reporting nausea during the trial did not differ between groups (RR 1.01; CI 0.31, 3.33).

Behavioral/Social Function

One head-to-head study of different medications reported on aggression, and several studies reported on crime.

Aggression

One study reported on aggression. Gerra et al. (2007) compared aggression between patients randomized to oral methadone and sublingual buprenorphine (total $n = 30$). Using the PSAP software program, the study measured “aggressive responding,” “escape responding,” and “point-maintained responses” as metrics for aggression at a three-month follow-up. More aggression is indicated by higher scores in the first two measures and a lower score for “point-maintained responses.” The differences between the two groups in aggressive responding (MD –3.82; CI –41.76, 34.12) and escape responding (MD –42.9; CI –132.96, 47.16) were not statistically significant. The buprenorphine group scored significantly better than the methadone group in point-maintained responses (MD 1676.34; CI 1456.18, 1896.50).

Criminal Activity

An open-label RCT compared a naltrexone implant to oral methadone among 46 heroin-dependent criminal offenders. At a six-month follow-up, mean days of illegal activity per month did not differ between the naltrexone implant group and the methadone control group (MD 0.50; CI –9.98 to 10.98).

The large observational study by Crits-Christoph et al. (2015) found no difference in number of arrests in the prior 30 days, at treatment discharge, among patients who received XR-NTX, oral naltrexone, Suboxone, or psychosocial treatment without medication using the SAMHSA Treatment Episode Data Set. Group differences were adjusted using propensity scores based on several intake variables (e.g., severity). As stated earlier, the authors suggested that their study had a short observation period and outcomes such as arrests should be studied over longer periods.

Percentage of Participants Arrested or Incarcerated

A randomized controlled trial of 116 heroin-dependent inmates compared the effects of Suboxone and oral methadone in a correctional setting. Participants transferred to community MAT treatment used the same medications after release. The percentage of participants who were arrested in the three months following release was not significantly different between

participants receiving Suboxone and methadone (MD -0.02 ; CI $-0.33, 0.29$). Likewise, the difference in reincarceration rates was not significant (RR 0.80 ; CI $0.53, 1.20$).

Neurological Function

Only one head-to-head study of different medications reported on a neurological function outcome.

Stress Tolerance

Soyka et al. (2005) randomized 62 patients to either sublingual buprenorphine or oral methadone. Patients completed the Reactive Stress Tolerance test at eight to ten weeks. The differences in mean score for phase 1 (MD 0.5 ; CI $-2.0, 3.0$), phase 2 (MD -2.4 ; CI $-14.28, 9.48$), and phase 3 (MD 2.6 , CI $-7.91, 13.11$) were not statistically significant.

Summary

We did not find statistically robust evidence that treatment effects systematically vary by medication for any of the function domains or specific outcomes except fatigue which was reported by a smaller percentage of buprenorphine than methadone patients across three RCTs. Three RCTs that focused on cognitive function reported no statistically significant differences in memory, cognitive speed or flexibility, attention, or vision between buprenorphine and methadone, with the exception of a small study where buprenorphine patients performed better in vision tracking. Regarding aggression, one RCT reported the buprenorphine group scored significantly better than the methadone group in “point-maintained responses,” but there was no difference in two other measures of aggression.

A few studies that reported on employment and neurological function using diverse measures reported no differences among medications. We identified only one RCT of naltrexone that reported functional outcomes; compared with methadone, no difference was seen in the mean number of days patients engaged in illegal activity.

Key Question 1b: Does the Effect Vary by Route of Administration?

In this section, we describe head-to-head studies that compared two or more routes of administration for the same medication. Standard of care for buprenorphine is sublingual administration, while methadone is generally administered as syrup for oral consumption. Results for comparisons of buprenorphine versus methadone are reported in the previous section (KQ 1a), which compared two or more different medication types.

We identified three RCTs (Metrebian et al., 2015; Strang et al., 2000; Rosenthal et al., 2013) reporting on effect variation by route of administration. Metrebian et al. randomized 127 patients to either injectable or oral methadone; both were administered under supervision. Strang et al.

randomized 37 patients to either injectable or oral methadone. Rosenthal et al. randomized 287 patients to buprenorphine implants, placebo implants, or Suboxone.

The RCTs reported on physical function and social function but used different outcomes and measures.

Physical Function

Physical Health

Reporting on the physical functioning component of the 36-Item Short-Form Survey (SF-36) at six months, Metrebian et al. (2015) found no statistically significant difference between injectable and oral methadone (MD 7.40; CI –4.21, 19.01). Strang et al. (2000) found no statistically significant difference in Maudsley Addiction Profile physical health scores (MD –0.80; CI –5.10, 3.50) for injectable versus oral methadone at six months.

Insomnia

Rosenthal et al. (2013) found no statistically significant difference in the percentage of participants reporting insomnia between the oral Suboxone and buprenorphine implant groups (RR 0.52; CI 0.24, 1.11) at six months.

Behavioral/Social Function

Mental Health

Regarding the mental health component of the SF-36, Metrebian et al. (2015) found no statistically significant difference between the oral and injectable methadone groups at six months (MD –2.16; CI –8.94, 4.62). Strang et al. (2000) found no statistically significant differences on the Maudsley Addiction Profile mental health score between the injectable methadone group and the oral group (MD –0.30; CI –1.01, 0.41) at six months.

SF-36 Social Functioning Component

Metrebian et al. (2015) reported no statistically significant difference between the oral versus injectable methadone groups (MD –5.20; CI –16.57, 6.17) at six months on the social functioning component of the SF-36.

Crime

Metrebian et al. (2014) found no statistically significant difference between the oral and the injectable methadone groups in the percentage of participants engaging in criminal activity in the previous month (RR 0.90; CI 0.40, 2.04) at six months. The small RCT by Strang et al. (2000) found no statistically significant difference between the two groups (MD –3.60; CI –10.02, 2.82) in the mean number of days patients reported committing shoplifting, robbery, burglary, and fraud at six months.

Summary

Overall, we did not find any indication that treatment effects systematically vary by route of administration. However, only a few studies that can answer this question have been published to date. In addition, all identified studies reported on relatively small samples and it is unclear whether studies had sufficient statistical power to detect effects in functional outcomes. Finally, the existing studies report on unique operationalizations of functional effects, and we found no measure that was used in more than one study.

Key Question 1c: Does the Effect Vary by Length of Treatment, Follow-Up Time, or Later Cessation of MAT?

We identified two observational studies (Rapeli et al., 2011; Aalto et al., 2011) and one RCT (Schwartz et al., 2009) that reported outcomes at multiple follow-up times and used these outcomes to assess variations in effect by length of treatment. In addition, one observational study (Reijneveld et al., 1993) was identified that compared patients who stayed in methadone maintenance for five years to patients who ended treatment before five years. The MDs between groups for each measure and each follow-up time are displayed in the evidence table in Appendix D, and results are summarized narratively in this section.

In addition, we conducted meta-regression analyses across studies. We added the length of follow-up to the meta-analysis model to determine if the length of follow-up systematically affected the differences between patients on MAT and patients not receiving MAT treatment. Given the small number of studies reporting on the same outcome, this indirect analysis was possible for only two outcomes (insomnia and percentage arrested or incarcerated).

Cognitive Function

One RCT reported cognitive function at multiple follow-up times. Rapeli et al. (2011) randomized persons with OUD to either oral methadone or sublingual buprenorphine. Fourteen buprenorphine and 12 methadone patients were administered a battery of cognitive tests at five to eight months and 11 to 16 months. Scores were compared with those of 14 matched healthy nondrug-using controls. Specific domains are described in the next sections.

Memory

Rapeli et al. (2011) assessed working memory using the Letter-Number Sequencing task from the WMS-III and the PASAT, while verbal memory was assessed using the Logical Memory score from the WMS-III. The authors found no significant differences between groups, time effect, or group by time interaction for the verbal memory measure.

Attention

Rapeli et al. (2011) assessed attention using two scores from the Attentional Performance Test: Go-NoGo reaction time and Go-NoGo errors. No significant time or group by time interaction effects were seen in these measures.

Occupational Function

Employment

Reijneveld et al. (1993) compared a random sample of patients enrolled in a methadone treatment program in the Netherlands for at least five years ($n = 21$) with a random sample of

those who ended treatment in less than five years ($n = 17$). Patients who were no longer on MAT were significantly less likely to report spending no time working in the past month (RR 0.54; CI 0.29, 1.00).

Physical Function

Insomnia

Our meta-regression analysis of RCTs reporting insomnia found no indication that studies with longer follow-ups reported stronger or weaker treatment effects ($p = 0.20$).

Overall Medical

Aalto et al. (2011) compared ASI medical component scores between patients on buprenorphine and matched controls in an SEP at three, six, and 12 months ($n = 60$). The reported p values for the tests of interaction between time and group were not significant ($p = 0.47$).

Behavioral/Social Function

Mental Health

The cohort study reported by Aalto et al. (2011) also compared ASI psychiatric component scores. The p value for the test of interaction between time (three, six, and 12 months) and group was significant ($p < 0.001$). However, results must be interpreted with caution because they appear to be primarily associated with baseline differences between groups. In addition, the SEP had a high dropout rate (from 30 patients to 11 at the one-year follow-up, compared with 30 to 25 in the buprenorphine group).

Family Function

Aalto et al. (2011) reported the ASI family/relationships component. P values for the tests of interaction between time (three, six, and 12 months) and group were significant ($p = 0.05$). This result appears to be associated primarily with a significant difference in improvement in the buprenorphine group compared with the control group at three months of treatment and no between-group differences at six or 12 months.

Crime

Schwartz et al. (2009) randomized 319 subjects with OUD to either methadone or wait-list. The proportion of each group who had been arrested did not differ significantly over six, 12, or 24 months (six-month RR 0.78, CI 0.48, 1.26; 12-month RR 1.03, CI 0.70, 1.50; 24-month RR 0.86, CI 0.66, 1.12).

Aalto et al. (2011) also compared ASI legal component scores between buprenorphine patients and matched controls in an SEP. *P* values for the tests of interaction between time (three, six, and 12 months) and group were not significant ($p = 0.31$).

The Reijneveld study compared patients enrolled in methadone treatment for at least five years to those who ended treatment earlier reported no difference in the percentage reporting engaging in illegal activity in the prior week (RR 0.62; CI 0.06, 6.25).

A meta-regression analysis found no association between longer follow-ups and differences between groups for the outcome percentage arrested or incarcerated ($p = 0.65$).

Summary

Of the three studies that followed MAT patients longitudinally, one found an effect of time and group on the ASI family and psychiatric components, but results are difficult to interpret. One study found that methadone patients who remained in treatment for five years had worse employment outcomes than patients who left treatment earlier. The studies had very small sample sizes and did not control for other factors possibly associated with outcomes, and no other study was identified that reported on the same characteristic of interest. A meta-regression found no indication that studies with longer follow-ups reported stronger treatment effects. In sum, we did not find robust evidence that the treatment effect systematically varies by length of treatment, follow-up time, or later cessation of MAT.

Key Question 1d: Does the Effect Vary by Treatment Modality (e.g., Methadone Clinic Versus Prescription Medication Taken at Home)?

In this section, we describe head-to-head studies comparing two or more treatment modalities that use the same medication. In the United States, methadone is traditionally dispensed under supervision at a methadone clinic, whereas buprenorphine is prescribed by a physician and can be distributed at a pharmacy. Comparisons between these two drugs are described in KQ 1a.

We identified two studies (Senay et al., 1993; Gossop et al., 1999) that compared effects of treatment modalities. Senay randomized 130 patients to take-home methadone distributed either two times per week or twice per month. Gossop compared a cohort of patients ($n = 452$) dispensed daily methadone either at a methadone clinic or prescribed by a general practitioner. Supervision (to be provided at retail pharmacies) was prescribed by only 14 percent of general practitioners.

Behavioral/Social Function

Addiction Severity Index Psychiatric Score

The RCT by Senay et al. (1993) reported no significant differences between groups receiving take-home methadone on different schedules at six months (SMD 0.02; CI -0.01, 0.04).

Crime

The cohort study by Gossop et al. (1999) reported on crimes (excluding drug possession and drug selling). At the six-month follow-up, the authors found a statistically significant difference favoring patients prescribed methadone from a general practitioner's office compared with patients who were dispensed the treatment at a traditional methadone clinic (MD -3.80; CI -7.06, -0.54).

The RCT by Senay et al. (1993) reported six-month ASI legal component scores and reported no difference in outcome for patients receiving take-home methadone on different schedules (MD 0.00; CI -0.02, 0.02).

Occupational Function

Senay et al. (1993) also found no difference in ASI employment scores between groups receiving take-home methadone on two different schedules (MD 0.00; CI -0.11, 0.11).

Physical Function

At six months, Senay reported no statistically significant differences in mean ASI medical scores between patient groups on the two different take-home methadone schedules (MD -0.01; CI -0.04, 0.02).

Summary

Only two studies that reported functional effects compared one treatment modality to another. One RCT reported no differences in ASI psychiatric, legal, employment, or medical scores between groups on take-home methadone distributed two times per week versus twice per month. One cohort study reported fewer nondrug-related crimes committed by patients prescribed methadone from a general practitioner's office compared with patients who were dispensed methadone at a traditional methadone clinic.

4. Discussion

This chapter begins with a summary of findings organized by KQs. KQs are organized by functional area, and within functional area, by comparison group. We then compare our findings with those of prior systematic reviews on the topic, describe the strengths and limitations of the research, and discuss the implications of our findings.

Summary of Findings

Despite an exhaustive search of more than 1,300 publications that were scrutinized as full text, only 37 studies met inclusion criteria. Because functional outcomes are most often reported as secondary outcomes in studies of substance abuse treatment, we retrieved full copies of all MAT studies of OUD patients that potentially would meet our inclusion criteria for study design and combed the results sections for relevant outcomes: 27 RCTs and ten observational studies reported in 41 publications were included in this review. No RCT was rated as low ROB, but several methodologically sound observational studies were identified. The statistical power to detect effects in functional outcomes was unclear in the majority of studies. The studies reported highly diverse functional outcome measures, with the exception of verbal memory, attention, insomnia, fatigue, and criminal activity.

Regarding function of MAT patients compared with “healthy” controls with no history of SUD or opioid use, a large observational study found that MAT users had twice the risk of injurious traffic accidents of nonusers. We identified one study that measured working memory and another that reported cognitive speed: MAT users performed significantly worse than “healthy” controls with no history of SUD or opioid use in these studies. Based on two studies, it appears that MAT users do not perform worse on verbal memory tasks than healthy controls. One study showed that patients taking buprenorphine or methadone scored higher in aggressive responding than healthy controls.

Evidence was mixed when MAT patients were compared with persons with OUD who were not on MAT. One cohort comparison found that fewer buprenorphine patients reported fatigue than did persons with OUD who did not receive MAT; other physical and behavioral/social function outcomes had mixed findings or showed no differences.

We found little statistically robust evidence that treatment effects systematically vary by medication. A comparison across RCTs found a significantly lower prevalence of fatigue in buprenorphine patients compared with methadone patients. Direct comparisons of functional effects by route of administration, length of treatment, and treatment modality were scarce and reported mixed results. A meta-regression found no indication that longer follow-up periods are associated with effect sizes for insomnia or percentage arrested.

In sum, weaknesses in the body of evidence prevent any strong conclusions about the effects of MAT on functional outcomes or differences among medication types, treatment modalities, or length of treatment. Rigorous studies of functional effects could be designed and funded to strengthen the body of literature.

Detailed findings are described in this chapter and shown in Table 4.1, along with the quality of evidence rating for each outcome. For each outcome, the table displays the number and type of studies, relative and absolute effects, quality of evidence criteria, and the GRADE category for all outcomes of interest.

Table 4.1. Summary of Findings

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Key Question 1. Effects of MAT on functional outcomes												
<i>MAT versus healthy nondrug-using controls</i>												
Cognitive function: verbal memory—methadone versus healthy nondrug-using controls	2 cohorts	High	Inconsistent	Direct	Imprecise	N/A	1.6 (1.6) 16.3 (3.1)	3.1 (2) 14.2 (3.1)	1 favored methadone, SMD 0.81 (0.25, 1.36); 1 no statistically significant difference, SMD 0.00 (−0.74, 0.74).	N/A	N/A	Very low
Cognitive function: verbal memory—buprenorphine versus healthy nondrug-using controls	2 cohorts	High	Consistent	Direct	Imprecise	N/A	1.6 (1.6) 16.3 (3.1)	2.6 (2.7) 14.1 (3.3)	No statistically significant difference, SMD 0.43 (−0.11, 0.98), SMD −0.57 (−0.30, 0.16)	N/A	N/A	Low
Cognitive function: attention—methadone versus healthy nondrug-using controls	2 cohorts	High	Consistent	Direct	Imprecise	N/A	0.2 (0.4) 481.6 (67.4)	0.5 (1.0) 468.4 (74.7)	No statistically significant difference, SMD −0.38 (−1.13, 0.37), SMD −0.18 (−0.72, 0.36)	N/A	N/A	Low
Cognitive function: attention—buprenorphine versus healthy nondrug-using controls	2 cohorts	High	Consistent	Direct	Imprecise	N/A	0.2 (0.4) 481.6 (67.4)	0.6 (0.8) 439.2 (83.4)	No statistically significant difference, SMD −0.63 (−1.42, 0.16); SMD −0.55 (−1.10, 0.01)	N/A	N/A	Low
Cognitive function: working memory—methadone versus healthy non-drug-using controls	1 cohort	High	Not replicated	Direct	Imprecise	N/A	11.6 (2.9)	8.6 (2.1)	Favors controls, SMD	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
									-1.14 (-1.94, -0.34)			
Cognitive function: working memory—buprenorphine versus healthy nondrug-using controls	1 cohort	High	Not replicated	Direct	Imprecise	N/A	11.6 (2.9)	9.2 (2.3)	Favors controls, SMD -0.89 (-1.64, -0.14)	N/A	N/A	Very low
Cognitive function: cognitive speed—methadone versus healthy nondrug-using controls	1 cohort	High	Not replicated	Direct	Imprecise	N/A	24.3 (6.9)	30.2 (8.6)	Favors controls, SMD 0.74 (0.16, 1.33)	N/A	N/A	Very low
Cognitive function: cognitive speed—buprenorphine versus healthy nondrug-using controls	1 cohort	High	Not replicated	Direct	Imprecise	N/A	24.3 (6.9)	29.8 (10.5)	Favors controls, SMD 0.61 (0.02, 1.21)	N/A	N/A	Very low
Cognitive function: driving accidents—methadone and buprenorphine versus nondrug users	1 case control	Low	Not replicated	Direct	Precise	N/A	196 total	72,489 total	MAT users more likely in accidents, OR 2.02 (1.40, 2.91)	Not calculable	Not calculable	Low
Behavioral/social function: aggression—methadone versus healthy nondrug-using controls	1 cohort	High	Not replicated	Direct	Precise	N/A	112.36 (60.57)	343.15 (56.74)	Methadone higher in aggressive responses, SMD 3.83 (2.62, 5.03)	N/A	N/A	Very low
Behavioral/social function: aggression—buprenorphine versus healthy nondrug-using controls	1 cohort	High	Not replicated	Direct	Precise	N/A	112.36 (60.57)	339.33 (48.99)	Buprenorphine higher in aggressive responses, SMD 4.01 (2.77, 5.25)	N/A	N/A	Very low
MAT versus no MAT												
Physical function: fatigue—methadone versus untreated opioid users	1 cohort	High	Not replicated	Direct	Imprecise	N/A	76.8/120	20/40	No statistically significant difference, RR 0.78 (0.56, 1.09)	640	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Physical function: fatigue—buprenorphine versus untreated opioid users	1 cohort	High	Not replicated	Direct	Imprecise	N/A	76.8/120	12/40	Favors buprenorphine, RR 0.47 (0.29, 0.76)	640	–339.2 (–454.4, –153.6)	Very low
Physical function: insomnia—MAT (buprenorphine or naltrexone) versus placebo	4 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	39.24/266	57.05/375	No statistically significant difference, RR 1.02 (0.61, 1.71)	146.5	N/A	Moderate
Physical function: insomnia—methadone versus untreated opioid users	1 cohort	High	Not replicated	Direct	Imprecise	N/A	67.2/120	16/40	No statistically significant difference, RR 0.72 (0.47 to 1.08)	560	N/A	Very low
Physical function: ASI medical score—methadone versus psychosocial + detox	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.20 (0.31)	0.14 (0.292)	No statistically significant difference, SMD –0.20 (–0.54, 0.15)	N/A	N/A	Very low
Physical function: ASI medical score—buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	Imprecise	N/A	0.47 (0.28)	0.29 (0.32)	No statistically significant difference SMD –0.57 (–1.29, 0.15)	N/A	N/A	Very low
Behavioral/social function: ASI psych function—methadone versus psychosocial + detox	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.15 (0.189)	0.11 (0.205)	No statistically significant difference, SMD –0.20 (–0.55, 0.14)	N/A	N/A	Very low
Behavioral/social function: ASI psych function—buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	Imprecise	N/A	0.33 (0.22)	0.34 (0.27)	No statistically significant difference, SMD 0.04 (–0.67, 0.75)	N/A	N/A	Very low
Behavioral/social function: ASI family function—methadone versus psychosocial + detox	1 RCT	Low	Not replicated	Direct	Precise	N/A	0.15 (0.113)	0.14 (0.086)	No statistically significant difference, SMD –0.01 (–0.04, 0.02)	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Behavioral/social function: crime—days of illegal activity, methadone versus wait-list or passive referral	2 RCTs	Low	Consistent results, but substantial heterogeneity	Direct	Precise		7.3 (9.64) 56.5 (45)	2.1 (9.45) 28.5 (45)	RCTs favored methadone, SMD -0.57 (-1.00, -0.13)	N/A	N/A	Low
Behavioral/social function: crime—mean number of charges, month, naltrexone versus treatment as usual	1 RCT	Low	Not replicated	Direct	Not reported	N/A	0.5 (not reported [NR])	0.2 (NR)	No statistically significant difference, MD 0.00 (SD not reported)	N/A	N/A	Very low
Behavioral/social function: crime—percentage arrested or convicted, MAT (methadone, naltrexone) versus no MAT	6 RCTs	Moderate	Inconsistent	Direct	Imprecise	Not calculable	225.05/521	190.49/527	No statistically significant difference, RR 0.75 (0.46, 1.23)	500	N/A	Low
Behavioral/social function: crime—percentage arrested or convicted, methadone versus no MAT	4 RCTs	Moderate	Consistent	Direct	Imprecise	Not calculable	170.58/349	146.61/340	No statistically significant difference, RR 0.80 (0.33, 1.94)	500	N/A	Low
Behavioral/social function: crime—percentage arrested or convicted, naltrexone versus no MAT	2 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	54.57/172	35.04/187	No statistically significant difference, RR 0.64 (0.02, 19.77)	425	N/A	Low
Behavioral/social function: crime—mean number of arrests 30 days after discharge, XR-NTX, oral naltrexone, or Suboxone versus psychosocial treatment alone	1 cohort	Low	Not replicated	Direct	Not reported	N/A	38/677	6.56/196	No statistically significant difference, RRs not reported	56	N/A	Very low
Behavioral/social function: ASI legal—methadone versus psychosocial + detox	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.05 (0.13)	0.13 (0.193)	Favors methadone, SMD 0.50 (0.15, 0.85)	N/A	N/A	Very low
Behavioral/social function: ASI legal—buprenorphine versus syringe exchange	1 cohort	Low	Not replicated	Direct	Imprecise	N/A	0.38 (0.21)	0.13 (0.19)	Favors buprenorphine, SMD -1.25 (-2.01, -0.48)	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Occupational function: employment, percentage employed, naltrexone versus treatment as usual	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	16.64/32	20.46/31	No statistically significant difference, RR 1.21 (0.80 to 1.84)	520	N/A	Very low
Occupational function: employment, ASI score—methadone versus psychosocial + detox	1 RCT	Low	Not replicated	Direct	Precise	N/A	0.77 (0.262)	076 (0.251)	No statistically significant difference, SMD -0.04 (-0.38, 0.3)	N/A	N/A	Very low
Occupational function: employment, number of days working in past 30, methadone versus passive referral	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	12 (10.3)	8.5 (10.5)	No statistically significant difference (-7.01, 0.01)	N/A	N/A	Very low
Occupational function: employment—five years of methadone versus < five years	1 cohort	High	Not replicated	Direct	Unclear	N/A	6.97/17	15.96/21	No statistically significant difference	760	N/A	Very low
Occupational function: employment—XR-NTX, oral naltrexone, or Suboxone versus psychosocial treatment alone	1 cohort	High	Not replicated	Direct	Unclear	N/A	153/677	37.32/196	No statistically significant difference	226	N/A	Very low
Key Question 1a. Do the effects differ by drug?												
<i>Buprenorphine versus methadone</i>												
Cognitive function: verbal memory	2 RCTs	High	Consistent	Direct	Imprecise	Not calculable	3.1 (2.0) 14.9 (0.2)	2.6 (2.7) 14.6 (0.7)	No statistically significant difference, SMD -0.34 (-2.44, 1.76)	N/A	N/A	Low
Cognitive function: attention	3 RCTs	High	Consistent	Direct	Imprecise	Not calculable	466.5 (72.1) 468.4 (74.7) 0.5 (1.0)	474.2 (54.3) 439.2 (83.4) 0.6 (0.8)	No statistically significant difference, SMD -0.12 (-0.76, 0.52)	N/A	N/A	Low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Cognitive function: vision, reaction time	1 RCT	High	Not replicated	Direct	Imprecise	N/A	1.2 (0.4)	1.2 (0.4)	No statistically significant difference, SMD 0 (−0.58, 0.58)	N/A	N/A	Very low
Cognitive function: vision, tracking performance	1 RCT	High	Not replicated	Direct	Imprecise	N/A	4.8 (2.4)	3.7 (1.3)	No statistically significant difference, SMD −0.55 (−1.14, 0.04)	N/A	N/A	Very low
Cognitive function: visual perception	1 RCT	High	Not replicated	Direct	Imprecise	N/A	30.7 (4)	30.5 (4.2)	No statistically significant difference, SMD −0.05 (−0.63, 0.53)	N/A	N/A	Very low
Physical function: insomnia	3 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	43.2/271	45.96/255	No statistically significant difference, RR 1.11 (0.70, 1.75)	241	N/A	Low
Physical function: fatigue	3 RCTs	Low	Consistent	Direct	Precise	Not calculable	42.18/ 271	24.6/255	Favors buprenorphine RR 0.62 (0.41, 0.95)	138	−52.44 (−81.42, −6.9)	Moderate
Physical function: pain	2 RCTs	High	Consistent	Direct	Imprecise	Not calculable	30.3/202	26.88/192	No statistically significant difference Detection (seconds) SMD −0.2 (−1.16, 0.76); (volts) SMD −0.06 (−1.02, 0.9) Tolerance (seconds) SMD 0.42 (−0.55, 1.39); (volts)	N/A	N/A	Low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
									SMD 0.26 (−0.7, 1.22)			
Physical function: nausea	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	5/29	4/23	No statistically significant difference, RR 1.01 (0.31, 3.33)	172	N/A	Very low
Behavioral/social: aggression	1 RCT	High	Not replicated	Direct	Imprecise	N/A	343.15 (56.73)	339.33 (48.99)	No statistically significant difference, SMD −0.07 (−0.79, 0.65)	N/A	N/A	Very low
Neurological: stress tolerance	1 RCT	High	Not replicated	Direct	Imprecise	N/A	2.8 (3.6)	3.3 (4.9)	No statistically significant difference Phase 1 SMD 0.12 (−0.46, 0.69) Phase 2 SMD −0.11 (−0.69, 0.46) Phase 3 SMD 0.14 (−0.44, 0.72)	N/A	N/A	Very low
Suboxone versus buprenorphine												
Physical function: pain	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	24/107	19/103	No statistically significant difference, RR 1.22 (0.71, 2.08)	184	N/A	Very low
Suboxone versus methadone												
Physical function: pain	1 RCT	High	Not replicated	Direct	Imprecise	N/A	6.1 (0.87)	5.2 (0.99)	No statistically significant difference, SMD −0.28 (−1.05, 0.50)	N/A	N/A	Very low
Behavioral/social: crime	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.71 (0.77)	0.69 (0.95)	No statistically significant	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
									difference, SMD -0.02 (-0.39, 0.34)			
Naltrexone versus methadone												
Behavioral/social: crime	1 RCT	High	Not replicated	Direct	Imprecise	N/A	14.4 (13.11)	14.9 (12.34)	No statistically significant difference, SMD 0.04 (-0.78, 0.86)	N/A	N/A	Very low
XR-NTX, oral naltrexone, Suboxone, versus psychosocial treatment												
Behavioral/social: employment	1 observational study	Low	Not replicated	Direct	Not reported	N/A	153/677	37.32/196	No statistically significant difference	226	N/A	Very low
Behavioral/social: crime	1 observational study	Low	Not replicated	Direct	Not reported	N/A	379.12/677	65.6/196	No statistically significant difference	56	N/A	Very low
Key Question 1b. Do the effects vary by route of administration?												
Methadone: injectable versus oral												
Physical function: SF-36 physical score	2 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	75.9 (28.7) 14.2 (6.5)	83.3 (23.2) 13.4 (6)	No statistically significant difference, SMD 0.28 (-0.16, 0.73) SMD -0.13 (-0.81, 0.56)	N/A	N/A	Low
Behavioral/social: SF-36 mental health score	2 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	38 (17.25) 1.6 (1.3)	35.84 (12.81) 1.3 (0.6)	No statistically significant difference SMD -0.14 (-0.58, 0.3) SMD -0.30 (-0.99, 0.39)	N/A	N/A	Low
Behavioral/social: SF-36 social score	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	70.1 (25.4)	64.9 (26.1)	No statistically significant difference,	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
									SMD -0.2 (-0.64, 0.24)			
Behavioral/social: crime	2 RCTs	Low	Consistent	Direct	Imprecise	Not calculable	9/38 5.8 (10.8)	9/42 2.2 (7.3)	No statistically significant difference, SMD -0.39 (-1.08, 0.3)	237	N/A	Low
Suboxone: implant versus sublingual												
Physical function: insomnia	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	9./114	17.62/119	No statistically significant difference, RR 0.52 (0.24, 1.11)	130	N/A	Very low
Key Question 1c. Do the effects vary by length of treatment, follow-up time, or later cessation of MAT?												
Length of treatment												
Cognitive function: memory, methadone versus buprenorphine versus healthy matched controls	1 RCT	High	Not replicated	Direct	N/A	N/A	N/A	N/A	No significant effect	N/A	N/A	Very low
Cognitive function: attention, methadone versus buprenorphine versus healthy matched controls	1 RCT	High	Not replicated	Direct	N/A	N/A	N/A	N/A	No significant effect	N/A	N/A	Very low
Physical: medical ASI score, buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	N/A	N/A	N/A	N/A	No significant effect $p = 0.47$	N/A	N/A	Very low
Behavioral/social function: ASI psychiatric score, buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	N/A	N/A	N/A	N/A	Time by group interaction ($p < 0.001$), see text	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Behavioral/social function: ASI family function score, buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	N/A	N/A	N/A	N/A	Time by group interaction ($p = 0.05$), see text	N/A	N/A	Very low
Behavioral/social function: ASI legal score, buprenorphine versus syringe exchange	1 cohort	High	Not replicated	Direct	N/A	N/A	N/A	N/A	No significant effect, $p = 0.31$	N/A	N/A	Very low
Behavioral/social function: percentage arrested, methadone versus wait-list	1 RCT	Moderate	Not replicated	Direct	N/A	N/A	N/A	N/A	No significant effect	N/A	N/A	Very low
Physical function: insomnia, meta-regression	4 RCTs	Low	Consistent	Indirect	Imprecise	Not calculable	N/A	N/A	No systematic effect, $p = 0.20$	N/A	N/A	Low
Behavioral/social: percentage arrested, meta-regression	6 RCTs	Moderate	Inconsistent	Indirect	Imprecise	Not calculable	N/A	N/A	No systematic effect, $p = 0.65$	N/A	N/A	Very low
Cessation of MAT												
Occupational: percentage employed	1 cohort	High	Not replicated	Direct	Imprecise	N/A	15.96/21	6.97/17	Favors ceased MAT, RR 0.54 (0.29, 1.00)	760	-349.6, (-539.6, 0)	Very low
Behavioral/social: percentage arrested	1 cohort	High	Not replicated	Direct	Imprecise	N/A	2.1/21	1.02/17	No significant difference, RR 0.62 (0.06, 6.25)	100	N/A	Very low
Key Question 1d. Do the effects vary by treatment modality (e.g., methadone clinic versus prescription medication taken at home)?												
Twice per month versus take-home methadone: 2 or 3 times per week												
Physical function: ASI medical	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.025 (0.069)	0.015 (0.052)	No statistically significant difference, SMD -0.17 (-0.57, 0.23)	N/A	N/A	Very low
Behavioral/social: ASI psychological	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.021 (0.065)	0.036 (0.086)	No statistically significant difference, SMD 0.19 (-0.21, 0.58)	N/A	N/A	Very low

Outcome	Number of Studies, Design	Risk of Bias	Consistency	Directness	Precision	Publication Bias	Absolute Risk, Control: Score (SD) or n/N	Absolute Risk, Intervention Group: Score (SD) or n/N	Relative Effect-Direction/ Magnitude (95% CI)	Control Risk (per 1,000)	Absolute Effect (95% CI)	GRADE
Behavioral/social: ASI legal	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.013 (0.057)	0.014 (0.049)	No statistically significant difference, SMD 0.02 (-0.38, 0.42)	N/A	N/A	Very low
Occupational: ASI employment	1 RCT	Low	Not replicated	Direct	Imprecise	N/A	0.072 (0.273)	0.069 (0.275)	No statistically significant difference, SMD -0.01 (-0.41, 0.39)	N/A	N/A	Very low
<i>Methadone: methadone clinic versus general practitioner's (GP's) office</i>												
Behavioral/social: number of nondrug crimes	1 cohort	Low	Not replicated	Direct	Imprecise	N/A	2.4 (8)	6.2 (22.4)	Favors GP office, MD 3.8 (0.54, 7.06)	N/A	N/A	Very low

NOTE: ASI = Addiction Severity Index; CI = confidence interval; N/A = not applicable; RCT = randomized controlled trial, RR = relative risk; SD = standard deviation; SMD = standardized mean difference. For dichotomous variables, n = number of persons experiencing an event (e.g., arrest), while N = sample size.

Key Question 1: Effect of MAT on Functional Outcomes

Of the five functional outcome categories, we identified at least one study in four of the categories: cognitive, physical, behavioral/social, and occupational. Outcomes reported in more than one study were verbal memory, attention, insomnia, fatigue, and criminal activity. Several functional outcomes were reported in only one study; if this was the case, we rated the quality of evidence as very low unless the study was of extremely high quality and reported precise results.

MAT Patients Versus “Healthy” Controls

Two studies randomized patients to either methadone or buprenorphine and compared their performance on a battery of cognitive tests to that of healthy controls with no history of SUD or opioid use. One of these studies found a significant difference in verbal memory favoring methadone patients compared with controls but no difference for buprenorphine. The other found no difference when healthy controls were compared with users of either medication. Quality of evidence is very low for methadone versus controls for verbal memory; we downgraded by an additional level because only one of the studies reported statistically significant results. No differences in attention were observed between the methadone or buprenorphine groups and controls. One of these studies also measured working memory and found that both buprenorphine and methadone patients scored significantly worse than controls. The other study measured cognitive speed and found that both MAT groups scored significantly worse than controls. Quality of evidence for both working memory and cognitive speed was rated very low due to high risk of study bias and lack of replication. Regarding driving, a very large observational study with low ROB found that MAT users (buprenorphine or methadone) had twice the risk of traffic accidents as nonusers, controlling for other important factors. Still, lack of RCT data led us to rate the quality of evidence for this outcome as low.

Regarding behavioral/social function, one small study reported aggression outcomes; patients randomized to buprenorphine or methadone scored significantly worse than controls with no history of OUD on aggressive responding. Quality of evidence is very low.

MAT Patients Versus Persons with OUD Not on MAT

No studies of MAT versus no MAT reported cognitive or behavioral function.

Regarding physical function, our meta-analysis of four RCTs found no significant difference in the percentage reporting insomnia between participants receiving MAT (buprenorphine, naltrexone) and those receiving placebo. Quality of evidence for insomnia was moderate because the RCTs had low ROB and results were consistent but imprecise. In one cohort study, a significantly lower percentage of buprenorphine patients reported fatigue than did persons with OUD who did not receive MAT, while the rate of fatigue did not differ between methadone patients and controls. Quality of evidence for the outcome of fatigue for methadone treatment is very low given the lack of RCT evidence, lack of replication, and high ROB. One RCT of

methadone versus a non-MAT intervention reported no difference in ASI medical scores; quality of evidence was very low because of imprecision and lack of replication. A cohort study that compared buprenorphine to syringe exchange also reported no difference in this score.

Regarding behavioral/social function, one RCT found no significant differences in mean family or psychiatric score between MAT and placebo groups; quality of evidence was rated very low for these outcomes due to lack of replication. Two RCTs with low ROB that measured crime reported that patients on methadone spent fewer days engaged in criminal activity than those randomized to placebo or passive control (e.g., wait-list); results were precise, but quality of evidence was low due to heterogeneity. Likewise, two studies that reported legal outcomes found MAT patients had significantly better scores than subjects provided with psychosocially enhanced detox or syringe exchange; quality of evidence was very low for methadone and buprenorphine because only one study reported this outcome for each drug. In contrast, our meta-analyses of RCTs that reported the percentage arrested or incarcerated found no significant difference between patients randomized to MAT or no MAT; quality of evidence for percentage arrested was low for methadone and naltrexone due to heterogeneity and imprecision.

Three RCTs and two observational studies reported no significant differences in employment outcomes between MAT patients and persons with OUD treated without MAT.

In sum, we identified several RCTs, and this robust study design has the potential to allow strong conclusions to be drawn. However, no RCT was rated as having low ROB, usually because of lack of participant blinding, high attrition, or a combination of both. Given the difficulties in blinding head-to-head studies of medications that are clearly different in appearance (methadone is usually administered as a syrup, while buprenorphine is sublingual) and the nature of SUD, these challenges were anticipated. Some studies compared MAT patients to matched controls with no history of SUD, whereas other studies compared MAT patients to persons with OUD who received a placebo or a non-MAT intervention or were assigned to a passive control condition. Several of the studies that compared MAT patients to persons with OUD who did not receive MAT reported significant positive effects of MAT on functional outcomes. However, in several studies, MAT patients performed significantly worse than matched controls with no history of SUD. Because of the limited number and quality of the studies, the quality of evidence supporting significant differences is low or very low.

Key Question 1a: Effects by Type of Medication

Sixteen studies compared different MAT medication types. Ten compared buprenorphine, a relatively new drug, to methadone, which was first implemented widely in the 1970s. Other studies compared buprenorphine to Suboxone, Suboxone to methadone, and methadone to naltrexone. Only cognitive function (memory, attention), physical function (insomnia, fatigue, pain), and behavioral (crime) outcomes were reported in more than one study.

We found little statistically robust evidence that treatment effects systematically vary by medication. Three RCTs that focused on cognitive function compared buprenorphine to

methadone; no statistically significant differences in memory, cognitive speed and flexibility, attention, or vision were reported, with the exception of a small study in which buprenorphine patients performed better in vision tracking than those on methadone. Quality of evidence for memory and attention was low due to high ROB and imprecision. Quality of evidence for cognitive speed, cognitive flexibility, and vision was very low due to high ROB, imprecision, and lack of replication.

Regarding physical function, a meta-analysis of three RCTs showed a significantly lower prevalence of fatigue in buprenorphine patients than in methadone patients. In absolute terms, 52 fewer buprenorphine patients than methadone patients per 1,000 reported fatigue; quality of evidence for fatigue was moderate, as the trials had low ROB, and results were consistent and precise. Our meta-analysis of three RCTs that reported insomnia found no statistical difference between the buprenorphine and methadone groups; the insomnia analysis included the same trials, but quality of evidence was low because the results were imprecise. Two RCTs that reported pain found no significant difference between buprenorphine and methadone groups: Quality of evidence was low due to high ROB and imprecision.

The only RCT of Suboxone versus methadone that reported functional outcomes found no significant difference in pain-rating scores at six months. The only RCT of methadone versus naltrexone that reported functional outcomes found no significant difference in the mean number of days patients engaged in illegal activity. Finally, a large observational study of a nationally representative sample found no difference in the outcomes “increase in the proportion of subjects who were employed (from intake to discharge)” and “number of arrests in the past 30 days (at treatment discharge),” among patients who received XR-NTX, oral naltrexone, Suboxone, or psychosocial treatment without medication. Group differences were adjusted using propensity scores based on several intake variables (e.g., severity). Despite the low ROB of this observational study, quality of evidence was very low due to lack of replication and lack of RCT data.

Key Question 1b: Effects by Route of Administration

Despite the considerable literature available on MAT, we identified only three RCTs that directly compared routes of administration and reported on functional outcomes. Specifically, the identified studies reported on physical and social function, but they used different outcomes and measures. None reported significant differences on these outcomes. Quality of evidence was very low for no difference in risk of insomnia between oral Suboxone and Suboxone implant; evidence was downgraded due to lack of replication. Quality of evidence is low for no difference between the effects of oral and injected methadone on mental health or physical health due to imprecision and small sample size. Quality of evidence was very low for no difference in effect between oral and injected methadone on social function because of lack of replication, imprecision, and small sample size. One large observational study on a nationally representative sample reported no differences in percentage employed or arrested during treatment among

patients receiving Suboxone, oral naltrexone, injection naltrexone, or non-MAT treatment; quality of evidence was very low due to lack of replication and lack of RCT data.

Key Question 1c: Effect by Length of Treatment, Follow-Up, and Later Cessation

Among three studies that followed MAT patients longitudinally, effect of time by group was significant in only one cohort study for two outcomes: ASI family component and psychiatric component. That study had high ROB because of an unacceptable drop-out rate. We judged the quality of evidence for an effect of treatment length by group on family and psychiatric function to be very low, as the only identified study has a high ROB, is not an RCT, and has not been replicated independently. Likewise, we rated the quality of evidence as very low for an effect of length of treatment by group on memory, attention, and legal issues because of ROB and lack of replication; each outcome was reported in one study that reported multiple follow-ups.

Our meta-regression analyses found no indication that RCTs with longer follow-ups reported larger (or smaller) differences between groups for the outcomes insomnia or percentage arrested. Quality of evidence for insomnia is low because although the four included RCTs had low ROB, the small number of studies has low statistical power to detect differences. Similarly, the quality of evidence for percentage arrested was very low given the small number of studies and overall moderate ROB.

We identified only one study (Reijneveld and Plomp, 1993) that contributed to the question of how and when functional outcome effects change after cessation of MAT. This small cohort study compared MAT patients who remained in MAT treatment at five years to patients who left treatment prior to five years. Current MAT patients were significantly more likely to be out of work than were former patients. The percentage arrested in the past 30 days was not significantly different. Quality of evidence for these outcomes was very low because of small sample size ($n = 38$), imprecision, lack of replication, and lack of RCT data.

Key Question 1d: Effects by Treatment Modality

Only two studies were identified that compared treatment modalities. One RCT, with low ROB, reported no differences in ASI psychiatric, legal, employment, or medical scores between groups on take-home methadone distributed two times per week versus twice per month. However, given the unclear statistical power to detect differences in these functional outcomes and the lack of replication of the results, the quality of evidence was judged to be very low. One cohort study with low ROB reported a lower rate of nondrug-related crimes by patients prescribed methadone from a general practitioner's office compared with patients dispensed methadone at a traditional methadone clinic. Quality of evidence was rated very low due to lack of replication and lack of RCT evidence.

Prior Systematic Reviews

We identified four recent systematic reviews relevant to this project; all focused on cognitive function. All used less-restrictive inclusion criteria than our review: The majority of the included individual studies used cross-sectional designs with no follow-up. The results reported in these publications echo our findings that MAT patients may perform worse than controls without a history of SUD or opioid use, although quality of evidence in our review was rated low or very low. Prior reviews reported mixed results regarding comparison of MAT patients with former opioid users.

Baldacchino et al. (2017) conducted several meta-analyses to synthesize data from 23 studies. Outcomes of interest included cognitive flexibility, attention and information processing, short-term memory, long-term memory, and impulsivity. Two studies were longitudinal in design, and 21 were cross-sectional, observational studies. Of these, 21 studies compared chronic methadone patients to “healthy participant populations,” and seven studies compared chronic methadone patients to abstinent former opioid users. The author conducted meta-analyses on impulsivity, cognitive flexibility, short-term memory, long-term memory, and attention; methadone patients scored significantly worse than controls for all outcomes. Standardized effect sizes were large, ranging from 0.41 to 0.89. When current methadone patients were compared with abstinent former opioid users, nonsignificant differences in effect size were observed for impulsivity, cognitive flexibility, and attention. For short- and long-term memory, better performance was found among former opioid users than among chronic methadone patients. In sum, this review suggests that some neuropsychological functional domains may be negatively affected by chronic methadone use. However, it is unclear if the observed differences were due to methadone use or long-term use of opioids in general.

Strand et al. (2013) examined the effects of methadone and buprenorphine maintenance on driving ability, including 54 experimental studies with a control group and five epidemiological studies with risk analysis. The authors included the following populations: persons with OUD receiving MAT with methadone or buprenorphine, persons with OUD not receiving MAT, and controls without a significant history of drug use. Meta-analysis was not conducted. The epidemiological studies had mixed findings: The largest analyzed a national database using case-control and case-crossover analyses. The case-control analysis revealed a higher risk of causing a motor vehicle accident among patients who received methadone and/or buprenorphine the day of the accident (OR 2.9; CI 1.51, 3.16). However, the case-crossover analysis did not find a significant relationship between methadone and/or buprenorphine exposure and a motor vehicle accident (OR 1.26; CI 0.93, 1.70). Among the experimental studies, patients who received methadone showed signs of impairment compared with controls in 127 of 407 tests across 28 studies, and patients receiving buprenorphine showed signs of impairment compared with controls in 22 of 83 tests across seven studies. Across the eight studies that directly compared methadone to buprenorphine, patients taking buprenorphine showed significantly less

impairment than those taking methadone in 13 of 76 tests. This review concluded that buprenorphine and methadone can impair driving abilities in healthy controls and may impair some functions in maintenance patients. Our systematic review found only one study that assessed the ability to drive an automobile. This observational study of injurious traffic accidents in France found statistically significant results; controlling for many important factors, patients taking methadone or buprenorphine were twice as likely to be involved in injurious accidents as individuals not on MAT.

Wang, Woules, and Russell (2013) synthesized the results from 35 studies, including 22 cross-sectional, five longitudinal, and eight RCTs on the effects of methadone maintenance on cognitive function compared with controls without a history of SUD. Meta-analysis was not conducted. Cognitive impairments among methadone patients included decreased performance on memory (eight studies), attention (two studies), psychomotor speed (four studies), decisionmaking (two studies), emotional interpretation (two studies), and verbal function (two studies) when compared with controls. However, the authors noted that two studies observed no significant differences, three studies observed small differences, and two studies found improved reaction times among methadone patients.

Finally, Biernacki et al. (2016) synthesized results from 22 studies that compared current or former opioid users to controls who did not have a significant history of drug abuse and were not currently using illicit drugs. This review categorized MAT patients as current opioid users and did not differentiate them from current users of illicit drugs. The design of included studies was not reported. Fifteen studies measured decisionmaking among current opioid users; in a meta-analysis, performance on decisionmaking measures was significantly worse in current users than in controls with no history of SUD. Among current opioid users, no significant association was found between the duration of opioid use and the size of the effect on decisionmaking based on results from 11 studies. Performance on decisionmaking measures was significantly worse among former opioid users when compared with controls with no history of SUD but not significantly different when compared between former and current users.

In sum, these recent reviews indicate that MAT patients perform worse in cognitive function than healthy controls with no history of SUD or opioid use. However, it is unclear if the observed differences are due to MAT or long-term use of opioids in general. Although controls are usually matched to patients on demographic and other characteristics, they clearly differ in substance abuse history and may differ in unreported psychological, psychiatric, and family history characteristics that might contribute to poor function.

Strengths and Limitations

This review has several strengths: an *a priori* research design, duplicate study selection and data abstraction of study information, a comprehensive search of electronic databases, ROB assessments, and use of comprehensive quality of evidence assessments to formulate review

conclusions. To not miss relevant studies, we screened 1,327 full text articles for functional outcomes.

However, our review has several limitations. First, very few studies of MAT reported functional outcomes; 37 MAT studies that met our study design criteria reported cognitive, physical, occupational behavioral/social, or neurological outcomes; while 307 studies of MAT were excluded for no functional outcomes. This lack of relevant data was not entirely unexpected, as studies of interventions for SUD tend to focus on reduction or cessation of substance use, treatment retention, and harm reduction. Functional outcomes were secondary outcomes in most included studies. Functional measures were primary outcomes in only six controlled trials; it is unclear if the other trials, which were statistically powered to detect differences in illicit use of opioids or treatment retention, had adequate power to detect differences in function.

The small number of studies reporting specific functional outcome measures limited our ability to conduct meta-analyses across medication types, comparators, settings, and routes of administration. Only six studies of naltrexone reported functional outcomes; most evidence is based on trials of methadone, buprenorphine, or buprenorphine plus naloxone (marketed as Suboxone). Furthermore, the included controlled trials had moderate to high ROB, primarily because of lack of participant blinding, high attrition rates, and failure to report the method of randomization and allocation. We did not contact individual study authors: Results and quality ratings reported in the review are based on published data. Finally, although we calculated the I-squared statistic to assess heterogeneity among studies included in our meta-analyses, some undetected heterogeneity may exist. The I-squared statistic is dependent on statistical power, which is primarily influenced by the number of studies and secondarily by the size of the studies; our meta-analyses included a small number of studies, and study sample size was often small compared with typical studies of medications and health care interventions.

Implications for Future Research and Practice

Making clinical and policy recommendations is beyond the scope of the systematic review; the goal of this report was to summarize, synthesize, and assess the quality of the existing evidence. Weaknesses in the body of evidence prevent any strong conclusions about the effects of MAT on functional outcomes or differences among medication types, treatment modalities, or length of treatment. Some studies that compared MAT patients to persons with OUD who did not receive MAT reported significant beneficial effects. However, this finding does not imply that performance meets the standards required for military deployment.

One RCT reported that the mean number of days of illegal activity in the past 30 days was not significantly different between the 20 mg methadone group and the placebo group but was significantly lower in the 50 mg methadone group. The number of crimes committed was significantly lower in both methadone groups compared with the placebo group, with the 50 mg

group reporting significantly fewer crimes than the 20 mg group (MD -9.23 versus -4.0). These results suggest a possible dose-response relationship that could be explored through future research, as dosage issues were beyond the scope of this project.

Importantly, in several studies, MAT patients performed significantly worse than matched controls with no history of SUD or OUD on measures of aggression, working memory, and cognitive speed. Unfortunately, no studies were conducted on active-duty service members or reported performance on specific occupational tasks. No studies reported the current or former occupations of participants, and applicability of the outcome measures to successful military deployment was not discussed in any study.

The original GRADE system (Atkins et al., 2004), developed to support clinical practice guidelines, recommended four factors to be considered when making a recommendation: (1) the trade-offs, taking into account the estimated size of the effect for the main outcomes, the confidence limits around those estimates, and the relative value placed on each outcome; (2) the quality of the evidence; (3) translation of the evidence into practice in a specific setting, taking into consideration important factors that could be expected to modify the size of the expected effects, such as availability of necessary expertise; and (4) uncertainty about baseline risk for the population of interest (to accurately balance benefits and harms). We rated the quality of evidence according to GRADE criteria; quality is low or very low for all outcomes. If data on the ability of MAT patients to perform typical activities of deployed service members in diverse occupations are needed, rigorous studies of applicable tasks could be designed and funded. RCTs are the highest standard of evidence; however, trials of this nature would be difficult to conduct. The time needed to obtain results that reflect real-world MAT programs might be prohibitive, given DoD's decisionmaking time line. It is possible that secondary analyses of existing administrative or health databases could be conducted quickly; however, we are aware of no sources of MAT data that collect information on detailed task performance. If decisions on deployment must be made without conducting further studies, expert opinion, political will, feasibility, troop morale, and other factors should be considered.

Appendix A. Food and Drug Administration Warnings Table

FDA Warnings	Buprenorphine^a	Buprenorphine with Naloxone (Suboxone)^a	Methadone^b	Naltrexone^c
Respiratory	Respiratory depression	Respiratory depression	Respiratory depression	—
Cardiac	—	—	Arrhythmias, cardiac conduction effects, including QT interval prolongation and torsades de pointes	—
Hepatic	Cytolysis hepatitis and hepatitis with jaundice	Cytolysis hepatitis and hepatitis with jaundice	—	Hepatocellular injury in excessive doses
Central nervous system (CNS)	CNS depression	CNS depression	Interactions with CNS depressants Interactions with alcohol and drugs of abuse	—
Others	Allergic reactions Dependence Opiate withdrawal effects	Allergic reactions Dependence Opiate withdrawal effects	Interactions with alcohol and drugs of abuse Head injury and increased intracranial pressure Acute abdominal conditions Hypotensive effect Dependence	Eosinophilic pneumonia Injection-site reactions

^a "Patient Information Leaflet, Patient Information: Suboxone, Subutex," undated.

^b Roxane Laboratories, undated.

^c "Vivitrol," undated.

Appendix B. Search Methodology

DATABASE SEARCHED AND TIME PERIOD COVERED

PubMed: January 1, 1970–January 16, 2017

LANGUAGE

English

SEARCH STRATEGY

((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR “Opioid-Related Disorders”[Mesh]
AND
“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone
AND
(humans[MESH] OR ((inprocess[sb] OR publisher[sb] OR pubmednotmedline [sb]) NOT (mice[ti] OR mouse[ti] OR rats[ti] OR rat[ti] OR dogs[ti])))

DATABASE SEARCHED AND TIME PERIOD COVERED

PubMed: January 1, 1970–December 31, 2000

LANGUAGE

English

SEARCH STRATEGY

“Opioid-Related Disorders”[Mesh] OR narcotic* OR opiate* OR opioid* OR heroin OR morphine
AND
misuse or abus* or addict* OR habit* OR withdraw*
AND
random* OR randomized controlled trial[pt] OR randomized controlled trials OR rct* OR blind* OR double-blind* OR single-blind*

DATABASE SEARCHED AND TIME PERIOD COVERED

PsycINFO: January 1, 1970–January 31, 2017

LANGUAGE

English

SEARCH STRATEGY

TI (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR SU (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR AB (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone)
AND

TI ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR SU ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR AB ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*))

AND

TI (randomi* OR “systematic review”) OR SU (randomi* OR “systematic review”) OR AB (randomi* OR “systematic review”)

DATABASE SEARCHED AND TIME PERIOD COVERED

PsycINFO: January 1, 1970–January 16, 2017

LANGUAGE

English

SEARCH STRATEGY

[TI (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR SU (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR AB (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR DE “Narcotic Antagonists” OR DE “Nalorphine” OR DE “Naloxone” OR DE “Naltrexone”

AND

DE “Opiates” OR DE “Codeine” OR DE “Heroin” OR DE “Morphine”

AND

DE “Drug Addiction” OR DE “Addiction” OR DE “Drug Dependency” OR DE “Heroin Addiction” OR DE “Drug Abuse” OR DE “Drug Overdoses” OR DE “Drug Withdrawal” OR DE “Intravenous Drug Usage” OR DE “Substance Use Disorder”

OR

TI (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR SU (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR AB (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR DE “Narcotic Antagonists” OR DE “Nalorphine” OR DE “Naloxone” OR DE “Naltrexone”

AND

TI (narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*) OR SU ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR AB ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*))

AND

TI (control* OR case-control*) OR SU (control* OR case-control*) OR AB (control* OR case-control*)

OR

TI (random* OR rct*) OR SU (random* OR rct*) OR AB (random* OR rct*)

AND

Narrow by Population: - human

DATABASE SEARCHED AND TIME PERIOD COVERED

CINAHL: January 1, 1970–December 31, 2016

LANGUAGE

English

SEARCH STRATEGY

TI (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR AB (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone) OR SU (“medication assisted treatment” OR “medication-assisted treatment” OR buprenorphine OR methadone OR naltrexone)

AND

TI ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR AB ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*)) OR SU ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse or abus* or addict* OR habit* OR withdraw*))

AND

TI (randomi* OR “systematic review”) OR SU (randomi* OR “systematic review”) OR AB (randomi* OR “systematic review”)

OR

TI (control* OR case-control) OR AB (control* OR case-control) OR SU (control* OR case-control)

DATABASE SEARCHED AND TIME PERIOD COVERED

Cochrane Database of Systematic Reviews, Other Reviews, CENTRAL: January 1, 2000–October 31, 2016

SEARCH STRATEGY

(narcotic* or opiate* or opioid* or heroin or morphine) and (misuse or abus* or addict* or habit* or withdraw*):ti,ab,kw (Word variations have been searched) OR MeSH descriptor: [Opioid-Related Disorders] explode all trees

AND

“medication assisted treatment” or “medication-assisted treatment” or buprenorphine or methadone or naltrexone:ti,ab,kw (Word variations have been searched)

DATABASE SEARCHED AND TIME PERIOD COVERED

Cochrane CENTRAL: January 1, 1970–December 31, 2000

SEARCH STRATEGY:

(narcotic* or opiate* or opioid* or heroin or morphine) and (misuse or abus* or addict* or habit* or withdraw*):ti,ab,kw (Word variations have been searched) OR MeSH descriptor: [Opioid-Related Disorders] explode all trees

AND

“medication assisted treatment” or “medication-assisted treatment” or buprenorphine or methadone or naltrexone:ti,ab,kw (Word variations have been searched)

DATABASE SEARCHED AND TIME PERIOD COVERED

EMBASE: From inception–February 1, 2017

LANGUAGE

English

SEARCH STRATEGY

‘narcotic dependence’/exp OR ((narcotic* OR opiate* OR opioid* OR heroin OR morphine) AND (misuse OR abus* OR addict* OR habit* OR withdraw* OR depend*))

AND

‘medication assisted treatment’ OR ‘medication-assisted treatment’ OR buprenorphine OR methadone OR naltrexone

AND

random* OR rct* OR blind* OR ‘double blind*’ OR ‘single blind*’ OR systematic OR ‘meta analy*’ OR ‘double blind procedure’/de OR ‘randomized controlled trial’/de OR ‘randomized controlled trial (topic)’/de OR ‘systematic review’/de

Appendix C. Risk of Bias Instruments

QUALITY ASSESSMENT: CONTROLLED TRIALS

Was the allocation sequence (randomization method) adequately generated?

There is a LOW RISK OF BIAS if the investigators describe a random component in the sequence generation process such as: referring to a random number table, using a computer random number generator, coin tossing, shuffling cards or envelopes, throwing dice, drawing of lots.

There is a HIGH RISK OF BIAS if the investigators describe a non-random component in the sequence generation process, such as: sequence generated by odd or even date of birth, date (or day) of admission, hospital or clinic record number; or allocation by judgement of the clinician, preference of the participant, results of a laboratory test or a series of tests, or availability of the intervention.

If the trial was not randomized, please check “high risk”

IF HIGH RISK OF BIAS, EXPLAIN IN NOTES.

Low risk

High risk notes



Was ALLOCATION adequately concealed (prior to assignment)?

There is a LOW RISK OF BIAS if the participants and investigators enrolling participants could not foresee assignment because one of the following, or an equivalent method, was used to conceal allocation: central allocation (including telephone, web-based and pharmacy-controlled randomization); sequentially numbered drug containers of identical appearance; or sequentially numbered, opaque, sealed envelopes.

There is a HIGH RISK OF BIAS if participants or investigators enrolling participants could possibly foresee assignments and thus introduce selection bias, such as allocation based on: using an open random allocation schedule (e.g. a list of random numbers); assignment envelopes were used without appropriate safeguards (e.g. if envelopes were unsealed or non-opaque or not sequentially numbered);

alternation or rotation; date of birth; case record number; or other explicitly unconcealed procedures.

IF HIGH RISK OF BIAS, EXPLAIN IN NOTES.

[Clear Response](#)

High risk notes

Were participants adequately BLINDED?

There is a LOW RISK OF BIAS if blinding of participants was ensured and it was unlikely that the blinding could have been broken; or if there was no blinding or incomplete blinding, but the review authors judge that the outcome is not likely to be influenced by lack of blinding.

- ☐ Low risk
- ☐ High risk
- ☐ Unclear

[Clear Response](#)

High risk notes

Were OUTCOME ASSESSORS adequately BLINDED?

There is LOW RISK OF BIAS if the blinding of the outcome assessment was ensured and it was unlikely that the blinding could have been broken.

Please make sure you assess the blinding of whoever measured the FUNCTIONAL OUTCOME, rather than outcomes such as drug use.

- ☐ Low risk
- ☐ High risk
- ☐ Unclear

[Clear Response](#)

High risk notes

Incomplete outcome data (ATTRITION BIAS) due to amount, nature or handling of incomplete outcome data

There is a LOW RISK OF BIAS if there were no missing outcome data; reasons for missing outcome data were unlikely to be related to the true outcome;

missing outcome data were balanced in numbers, with similar reasons for missing data across groups

(****The percentage of withdrawals and drop-outs should not exceed 20% for short-term follow-up [≤ 1 year] and 30% for long-term follow-up [> 1 year]****).

IF HIGH RISK OF BIAS, EXPLAIN IN NOTES. Again, please assess for FUNCTIONAL outcome, rather than other outcomes.


Participants may provide urine for drug testing, but not show up for functional tests or provide data on functional outcomes at follow-up.

- ☐ Low risk
- ☐ High risk

☐ Unclear

[Clear Response](#)

High risk notes



Is there evidence of SELECTIVE OUTCOME REPORTING bias (Yes/No)?

Are all outcomes in the Methods section (all pre-specified outcomes) reported, were all components of composite outcomes reported?

DESCRIBE ISSUES IN NOTES.


☐ Yes

☐ No

☐ Unclear

[Clear Response](#)

Notes



INTENTION-TO-TREAT analysis? (Yes/No)

Intention to treat (ITT) analysis means all patients who were enrolled and randomly allocated to treatment are included in the analysis and are analyzed in the groups to which they were randomized. Everyone who is randomized in the trial is considered to be part of the trial regardless of whether he or she completes the trial.

YES if they state ITT and methods used were actually ITT, or ****all**** participants were analyzed in the group to which they were allocated by randomization (no cross-over).

IF NO ITT, EXPLAIN IN NOTES.


☐ Yes

☐ No

☐ Unclear

[Clear Response](#)

Notes



Group SIMILARITY AT BASELINE

There is LOW RISK OF BIAS if groups are similar at baseline for demographic and other factors ("Table 1").
Also LOW risk of bias if any baseline differences were adjusted for in all relevant analyses.
IF HIGH RISK OF BIAS, EXPLAIN IN NOTES.

- ☐ Low risk
- ☐ High risk
- ☐ Unclear

[Clear Response](#)

Notes

Was there incomplete COMPLIANCE with interventions across groups?

There is LOW RISK OF BIAS if compliance with the interventions was acceptable ($\geq 80\%$ across intervention duration), based on the reported actual compliance during or at the end of the intervention.

There is HIGH RISK OF BIAS if compliance was low ($< 80\%$) during or at the end of the intervention.

There is UNCLEAR RISK OF BIAS if these data were not reported.

- ☐ Low risk
- ☐ High risk
- ☐ Unclear

[Clear Response](#)

Notes

Additional Bias: Bias due to problems not covered elsewhere in the table

- ☐ Yes
- ☐ No

[Clear Response](#)

Notes

QUALITY ASSESSMENT: OBSERVATIONAL STUDIES

Selection

1) Representativeness of the exposed cohort

- ☐ Truly representative of the average MAT patient in the community
- ☐ Somewhat representative of the average MAT patient in the community
- ☐ Selected group of users, e.g. nurses, volunteers
- ☐ No description of the derivation of the cohort

2) Selection of the non-exposed cohort (e.g. "healthy controls" or opioid dependent participants receiving a different medication or not receiving MAT)

- ☐ Drawn from the same community as the exposed cohort
- ☐ Drawn from a different source
- ☐ No description of the derivation of the non-exposed cohort
- ☐ Not applicable

3) Ascertainment of exposure

- ☐ Secure record (e.g. medical records, participation in a specific treatment program)
- ☐ Structured interview
- ☐ Written self-report
- ☐ No description

4) Demonstration that outcome of interest was not present at start of study (if relevant, which will almost never be the case) or author's statement that a valid outcome measure was chosen (e.g. a validated instrument measuring depression, anxiety, insomnia, etc).

- ☐ Yes
- ☐ No

Comparability

1) Comparability of cohorts on the basis of the design or analysis

If the authors describe factors for which they adjusted or noted that cohorts were matched on important factors and listed the factors, count that as a "yes"

- ☐ Study controls for demographic information, if drug users not in treatment -- controls for severity of drug use
- ☐ Study controls for any additional factor (This criteria could be modified to indicate specific control for a second important factor)

Outcome

1) Assessment of outcome

- ☐ Independent blind assessment

☐ Record linkage

☐ Self-report

☐ No description

2) Was follow-up long enough for outcome to occur (e.g. if the authors say they chose a particular follow-up time, definitely select "yes"; otherwise use your own judgment).

☐ Yes (select an adequate follow-up period for outcome of interest)

☐ No

3) Adequacy of follow-up cohorts

☐ Complete follow-up -- all subjects accounted for

☐ Subjects lost to follow up unlikely to introduce bias - small number lost - >80% retention for ≤ 1 year followup; >30% loss for 1-5 years followup; >40% loss for 6-10 years followup; >50% loss for 11-18 years followup; or description provided of those lost

☐ Follow-up rate not acceptable (according to levels described above) and no description of those lost

☐ No statement

Appendix D. Evidence Table

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Aalto et al., 2011 Location: Europe Study design: case-control	Number enrolled: 60 Number completed: 54 Mean age: buprenorphine: 26.7 (4.9), SEP: 26.5 (4.6) Percentage female: 20 Race/ethnicity: NR Years of opioid use: buprenorphine: 3.6 (1.7) SEP: 4.2 (1.8) Inclusion criteria: The inclusion criteria for the buprenorphine program were opioid dependence under the International Classification of Diseases–10 criteria and previous participation in at least one detoxification treatment. Another 30 opioid-dependent patients, who reported mainly using buprenorphine and who participated in the SEP, were recruited for the study in Helsinki between October 2005 and May 2006. Exclusion criteria: NR	Intervention: buprenorphine Setting: other substance abuse treatment Route of administration: oral Duration (months): 23 Non-MAT comparator: 1. SEP	Physical : ASI medical score, 12 months, buprenorphine (outpatient versus control—matched controls in an SEP, MD: −0.18, 95% CI: (−0.39, 0.03) Behavioral/social : ASI legal score, 12 months, buprenorphine—outpatient versus control—matched controls in an SEP, MD: −0.25, 95% CI: (−0.39, −0.11) ASI psychiatric score, 12 months, buprenorphine—outpatient versus control—matched controls in an SEP, MD: 0.01, 95% CI: (−0.16, 0.18)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Bale et al., 1980 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 585 Number completed: 545 Mean age: >25 years = 63% Percentage female: 0 Race/ethnicity: White: 39%, Black: 41%, Hispanic: 12% Years of opioid use: NR Inclusion criteria: Subject must have been using narcotics daily just prior to admission. He or she must not have been recently discharged from a longer-term treatment program at the hospital (because the nature of the discharge could preclude his readmission to a longer-term program). Exclusion criteria: Major psychiatric problems. Some patients were later found to be ineligible for VA treatment because of an “undesirable” military discharge or because they had falsified their names or records to gain entry to the detoxification ward. These patients were excluded from the study at the point when such information was discovered. Finally, patients who had felony charges pending were excluded.	Intervention: methadone Setting: methadone clinic Route of administration: oral Duration (months): NR Non-MAT comparator: 1. The family is a residential therapeutic community with 20 to 35 residents, including five to seven heroin addicts. The program is strongly committed to abstinence from all psychoactive substances and uses group confrontation and support (often called “attack” or “Synanon style” therapy). 2. Quadrants, a therapeutic community with 20 to 30 residents. All of the program residents have been addicted to heroin. Is less committed to complete abstinence, and some prescribed medications (but not methadone) are used. Uses limited group confrontation in combination with other therapeutic techniques. 3. Satori, a therapeutic community with 15 to 20 residents, all have been addicted to heroin. Psychoactive medications, including methadone, are occasionally prescribed. Emphasis on historical material and reconstruction therapy, attack style of confrontation is not employed.	Behavioral/social : Arrests during year (proportion of group), 1 year, all subjects (combined) in residential program versus outpatient MMT program, RR: 0.94, 95% CI: (0.68, 1.28) Percentage participants reincarcerated, 1 year, all subjects (combined) in residential program versus outpatient MMT program, RR: 1.24, 95% CI: (0.52, 2.95)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Compton et al., 2012 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 103 Number completed: 51 Mean age: control: 30.14 (11), buprenorphine: 33.41 (9) methadone: 34.55 (12) Percentage female: 35.9 Race/ethnicity: White: 79.6%, Black: 5.8%, Asian: 1.9%, other: 12.6% Years of opioid use: NR Inclusion criteria: At least 18 years of age; in good physical health; agreeable to and capable of signing an informed consent; no existing conditions that would affect sensitivity to cold (e.g., Raynaud's disease, urticaria); no neuropathology that would affect pain responses (e.g., peripheral neuropathy, neuropathic pain); and no cardiovascular conditions that could put participants at risk for blood pressure increases. Noncontrol group participants were seeking opioid maintenance therapy for the treatment of a DSM-IV diagnosed heroin dependence disorder. Exclusion criteria: A known sensitivity to buprenorphine or methadone; dependence on alcohol, benzodiazepines, or other drugs of abuse (except nicotine); any acute medical condition that would make participation medically hazardous; acute psychosis, severe depression, or in need of acute inpatient treatment/suicidal; taken levo-alpha-acetylmethadol, methadone or naltrexone within 30 days of enrolling in the study; discontinued participation in an opiate-substitution (i.e., methadone, levo-alpha-acetylmethadol) treatment program within 30 days of enrolling in the study; or any pending legal action that could prohibit sustained participation. For the control group, a current or past history of substance abuse, current use of analgesic medication, or being a nursing or pregnant female excluded participation.	Intervention: methadone Setting: other substance abuse treatment Route of administration: NR Duration (months): 4.5 Intervention 2: buprenorphine Setting: other substance abuse treatment Route of administration: NR Duration (months): 4.5 Non-MAT Comparator: 1. drug-free controls	Physical : Pain detection—volts, 18 weeks, buprenorphine versus methadone, MD: -0.94, 95% CI: (-5.16, 3.28) Pain tolerance—volts, 18 weeks, buprenorphine versus methadone, MD: 3.61, 95% CI: (-2.13, 9.35) Pain detection—volts, 18 weeks, buprenorphine versus control, MD: 2.09, 95% CI: (-4.7, 8.88) Pain tolerance—volts, 18 weeks, buprenorphine versus control, MD: -9.17, 95% CI: (-19.42, 1.08) Pain detection—volts, 18 weeks, buprenorphine versus methadone, MD: -0.84, 95% CI: (-12.1, 10.42) Pain tolerance—volts, 18 weeks, buprenorphine versus methadone, MD: 4.04, 95% CI: (-9.44, 17.52) Pain detection time (seconds), 18 weeks, buprenorphine versus control, MD: -3.63, 95% CI: (-6.79, -0.47) Pain tolerance—time (seconds), 18 weeks, buprenorphine versus control, MD: -25.35, 95% CI: (-38.87, -11.83) Pain detection time (seconds), 18 weeks, methadone versus control, MD: -2.69, 95% CI: (-7.32, 1.94) Pain tolerance—time (seconds), 18 weeks, methadone versus control, MD: -28.96, 95% CI: (-42.83, -15.09) Pain detection time (seconds), 18 weeks, methadone versus control, MD: 2.93, 95% CI: (-8.05, 13.91) Pain tolerance—time (seconds), 18 weeks, methadone versus control, MD: -13.21, 95% CI: (-27.82, 1.4)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Cornish et al., 1997 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 51 Number completed: 24 Mean age: 39 Percentage female: 10 Race/ethnicity: White: 24%, Black: 62%, Hispanic: 14% Years of opioid use: NR Inclusion criteria: Federal probationers or parolees with a history of opioid addiction Exclusion criteria: NR	Intervention: naltrexone Setting: an office within the probation department Route of administration: oral Duration(months): 6 Non-MAT comparator: 1. Subjects were required to attend three orientation and counseling sessions per week during the first 2 weeks of the study. These sessions were conducted by the research technician and focused on obtaining drug use and treatment histories, acquired immune deficiency syndrome (AIDS) education and risk reduction, and orientation to the research protocol. In addition, the subjects saw their parole/probation officer (PO) and provided a monitored urine specimen and breathalyzer reading twice weekly. From week 3 through week 24, control subjects saw their PO twice weekly and one of the two visits was randomly selected for the subject to provide a monitored urine specimen and breathalyzer reading.	Behavioral/social: Percentage participants reincarcerated, 6 months, naltrexone versus placebo, RR: 0.46, 95% CI: (0.23, 0.94)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Corsenac et al., 2012 Location: Europe Study design: case control	Number enrolled: 72,685 Number completed: 72,685 Mean age: NR Percentage female: 31.5 Race/ethnicity: NR Years of opioid use: not applicable Inclusion criteria: Drivers were included through their national ID, gender, and date of birth, as extracted from police reports. An application, based on optical character recognition, was developed to automatically extract from the image files the date of the crash, an individual's national ID, gender, and date of birth. Data on reimbursed medicines dispensed within six months before the crash were obtained by linking included drivers to the national health care insurance database using their national ID, gender, and date of birth. Exclusion criteria: not applicable	Intervention: methadone, buprenorphine Setting: not applicable, study of traffic accident data Route of administration: not applicable Duration (months): not applicable Non-MAT comparator: 1. General population. Collected from police reports and databases through ID extraction.	Cognitive: Responsible for traffic accidents, buprenorphine, or methadone versus nondrug user, 3 years, OR: 2.02, 95% CI: (1.40, 2.91)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Coviello et al., 2010 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 111 Number completed: 34 Mean age: 33.5 Percentage female: 18 Race/ethnicity: White: 47%, Black: 26%, Hispanic: 27% Years of opioid use: Heroin: 7.7 Inclusion criteria: Subjects were eligible for participation if they: (1) signed an informed consent form agreeing to randomization to one of the two treatment groups; (2) were between the ages of 18 and 55; (3) had a diagnosis of opioid dependence based on DSM-IV criteria and a structured psychiatric interview; (4) were in good general health as determined by a complete physical examination and laboratory tests; (5) had been assigned to probation/parole for a minimum of 6 months; and (6) had a negative result for urinary opioids and reported being at least 3 days opioid-free prior to randomization. Exclusion criteria: Subjects were excluded if they had: (1) current severe alcohol dependence that required medical supervision for alcohol withdrawal symptoms; (2) current psychosis, dementia, mental retardation, or history of schizophrenia; (3) clinically significant abnormalities in hematology, chemistry, or urinalysis; (4) clinically significant cardiovascular, neurological, hepatic, renal, pulmonary, metabolic, endocrine, or gastrointestinal disorders; (5) a diagnosis of chronic pain disorder; or (6) taken an opioid antagonist within the prior six months. Female subjects who were pregnant or lactating, or women of childbearing potential who were not using birth control were also excluded.	Intervention: naltrexone Setting: university research offices Route of administration: oral Duration (months): 6 Non-MAT comparator: 1. TAU (treatment as usual): psychological treatment. Participants in the TAU group received 6 months of psychosocial treatment at one of several community-based treatment programs or at the university provided by the research study. The university-based psychosocial treatment consisted of 3 hours of group therapy, 1 hour of individual therapy, and 1 hour of case management for 6 weeks of intensive outpatient treatment followed by 20 weeks of outpatient treatment consisting of 1 hour of individual and 1 hour of case management per week. The psychosocial therapy provided by the community-based programs was similar in content, but typically included additional hours of group therapy.	Occupational: Percentage of participants employed, 6 months, naltrexone versus standard psychosocial TAU, RR: 1.27, 95% CI: (0.84, 1.93) Behavioral/social : Average number of charges (prior versus follow-up), 6 months, oral naltrexone plus standard psychosocial treatment versus standard psychosocial TAU, MD: -0.3, 95% CI: not calculable Average number of convictions (prior versus follow-up), 6 months, oral naltrexone plus standard psychosocial treatment versus standard psychosocial TAU, MD: 0, 95% CI: not calculable

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Coviello et al., 2011 Location: United States or Canada Study design: cohort comparison	Number enrolled: 490 Number completed: 230 Mean age: 44.2 Percentage female: 18 Race/ethnicity: White: 45%, Black: 51%, Hispanic: 4%, Asian: NR, other: NR Years of opioid use: heroin: 17.7, other opioids: 3.0 Inclusion criteria: All patients discharged from three methadone treatment programs in a 3-year period, regardless of whether they completed “successfully.” Exclusion criteria: NR	Intervention: methadone Setting: methadone clinic Route of administration: oral Duration (months): 3 Non-MAT comparator: 1. No methadone maintenance treatment in past 3 months.	Behavioral/social : Mean number of days of illegal activity in the past 30 days, 3 months, methadone—outreach case management (OCM) (treatment) versus methadone—passive referral (no treatment)—control, MD: -3.4, 95% CI: (-6.01, -0.79) Mean number of days of illegal activity in the past 30 days, 9 months, methadone—OCM (treatment) versus methadone—passive referral (no treatment)—control, RR: 0.94, 95% CI: (0.48, 1.83)
Reference: Crits-Christoph et al., 2015 Location: United States or Canada Study design: cohort comparison	Number enrolled: 2,882 Number completed: 2,882 Mean age: 34.7 (10.5) Percentage female: 25.2 Race/ethnicity: White: 57.1%, Black: 39.7%, Other: 3.2% Years of opioid use: NR Inclusion criteria: The analysis included men and women ages 18 and older, who reported opioid use as their primary, secondary, or tertiary substance use problem at admission and had intake data. The study group was limited to individuals who were under community supervision by the state correctional agency (i.e., on parole or probation). All patients were treated in outpatient centers that received funding from the Missouri Division of Behavioral Health during fiscal year 2013 (July 1, 2012–June 30, 2013) in a six-county area plus the city of St. Louis. Exclusion criteria: NR	Intervention: naltrexone Setting: NR Route of administration: implant Duration (months): 3 Intervention 2: naltrexone Setting: NR Route of administration: oral Duration (months): 2 Intervention 3: Suboxone Setting: NR Route of administration: NR Duration (months): 2.5 Non-MAT comparator: 1. Psychosocial only. Adjusted median durations of episode of care were: psychosocial only, 85 days	Behavioral/social: percentage arrested in past 30 days, extended-release naltrexone, oral naltrexone, Suboxone, versus psychosocial only; no difference among groups when controlling for baseline variables and propensity scores Occupational: percentage employed at discharge; extended-release naltrexone, oral naltrexone, Suboxone, versus psychosocial only; no difference among groups when controlling for baseline variables and propensity scores

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Dole et al., 1969 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 32 Number completed: NR Mean age: 30.0 (5.1) Percentage female: NR Race/ethnicity: White: 46.8%, Black: 31.2%, Asian: 21.9% Years of opioid use: 12.7 Inclusion criteria: Opioid dependence for 5 or more years, 5 or more previous convictions, not already committed to the custody of addiction services agency Exclusion criteria: NR	Intervention: methadone Setting: prison, then methadone clinic Route of administration: oral Duration (months): NR Non-MAT comparator: 1. Untreated controls	Behavioral/social: Percentage of participants convicted of crimes committed after release, 1 year, methadone versus placebo, RR: 0.27, 95% CI: (0.10, 0.71)
Reference: Farrell-MacDonald et al., 2014 Location: United States or Canada Study design: cohort comparison	Number enrolled: 137 Number completed: 137 Mean age: continued MMT: 33.0 (7.2); MMT-T: 34.5 (8.0); no MMT 31.3 (7.4) Percentage female: 100 Race/ethnicity: NR Years of opioid use: NR Inclusion criteria: Data for this retrospective study were drawn from administrative data on a sample of women federal offenders who were initiated on MMT (the only available OMT option during the study period) while incarcerated in a Correctional Service of Canada (CSC) facility between January 1, 2003, and December 31, 2008, and who were released into the community Exclusion criteria: NR	Intervention: methadone Setting: prison, then community Route of administration: oral Duration (months): NR Non-MAT comparator: 1. A comparison group of offenders admitted to CSC and community-released during the study period, who were assessed at correctional intake as having a moderate to severe substance abuse problem with opioid use confirmed as their primary problem drug, but who did not participate in MMT while incarcerated	Behavioral/social: Percentage participants reincarcerated, 27 weeks, continuing methadone versus no MAT, hazard ration (HR): 0.35, 95% CI: (0.13, 0.90) Percentage participants reincarcerated, 27 weeks, terminated methadone versus no MAT, HR: 1.20, 95% CI: (0.72, 2.02)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Fudala et al., 2003 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 326 Number completed: 243 Mean age: Suboxone: 38.1 (8.3), buprenorphine: 36.6 (8.9), placebo: 38.0 (9.3) Percentage female: 35.3 Race/ethnicity: White: 61%, Black: 28.5%, Hispanic: 7.1%, Asian: 2.2%, other: 1.2 (Native American)% Mean years of opioid use: 7 Inclusion criteria: Men and women who met the diagnostic criteria for opiate dependence according to the DSM-IV, who were seeking opiate-substitution pharmacotherapy, who were between the ages of 18 and 59 years, and who were able to give informed consent and comply with study procedures were eligible to participate. Exclusion criteria: Women who were pregnant or nursing were excluded. Other criteria for exclusion included any medical condition that made study participation medically hazardous; aspartate or alanine aminotransferase levels greater than three times the upper limit of normal; a current, primary, Axis I psychiatric diagnosis (according to the DSM-IV) other than opiate, caffeine, or nicotine dependence; and use of methadone, levomethadyl acetate, or naltrexone within the 14 days before enrollment.	Intervention: Suboxone Setting: doctor's office Route of administration: oral Duration (months): 1 Intervention 2: buprenorphine Setting: doctor's office Route of administration: oral Duration (months): 1 Non-MAT comparator: 1. Placebo: All the tablets were identical in appearance and taste. All the subjects received counseling regarding human immunodeficiency virus (HIV) infection and up to 1 hour of individualized counseling per week. Emergency counseling (e.g., after a relapse) and referrals (e.g., to community legal aid programs) could be provided, but no other counseling or services (e.g., regarding family or employment issues) were offered.	Physical : Number (percentage) of patients who reported experiencing pain during the trial, 4 weeks, Suboxone versus buprenorphine, RR: 1.22, 95% CI: (0.71, 2.08) Number (percentage) of patients who reported experiencing pain during the trial, 4 weeks, Suboxone versus placebo, RR: 1.2, 95% CI: (0.71, 2.03) Number (percentage) of patients who reported experiencing pain during the trial, 4 weeks, buprenorphine versus placebo, RR: 0.98, 95% CI: (0.56, 1.73) Percentage of participants reporting insomnia, 4 weeks, Suboxone versus buprenorphine, RR: 0.65, 95% CI: (0.36, 1.19) Percentage of participants reporting insomnia, 4 weeks, Suboxone versus placebo, RR: 0.88, 95% CI: (0.46, 1.67) Percentage of participants reporting insomnia, 4 weeks, buprenorphine versus placebo, RR: 1.35, 95% CI: (0.76, 2.38)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Gerra et al., 2007 Location: Europe Study design: randomized controlled trial	Number enrolled: 45 Number completed: NR Mean age: heroin users: 25 (5.3); healthy controls: 24.4 (3.6) Percentage female: 0 Race/ethnicity: NR Years of opioid use: 9.2 (3.9) Inclusion criteria: Heroin-dependent patients were randomly assigned to buprenorphine or methadone treatment after they contacted the outpatient center seeking treatment. All the subjects were available for methadone or buprenorphine treatment, without any preference for a specific substitution treatment. Fifteen healthy male volunteers, who were recruited from hospital staff, university students, and workers, were matched to the patients for age. Exclusion criteria: Exclusion criteria included severe chronic liver or renal diseases or other chronic physical disorders, recent weight loss or obesity, endocrinopathies, immunopathies and, in particular, HIV disease. The subjects treated with other prescribed drugs in association with psychosocial therapy were not included in the study.	Intervention: buprenorphine Setting: other (outpatient center) Route of administration: oral Duration (months): 3 Intervention 2: methadone Setting: other (outpatient center) Route of administration: oral Duration (months): 3 Non-MAT comparator: 1. Fifteen healthy male volunteers, who were recruited from hospital staff, university students, and workers, and who were matched to the patients for age (20–33 years: mean $\pm$ S.D. = 24.4 ± 3.6 years), served as controls. Subjects were confirmed to be healthy based on physical examination and routine biochemical tests.	Behaviorals/social : Aggressive responding—point subtraction aggression paradigm (PSAP) software program, 3 months, buprenorphine versus healthy controls (CONT), MD: 226.97, 95% CI: (187.54, 266.4) Aggressive responding—PSAP software program, 3 months, buprenorphine versus methadone, MD: -3.82, 95% CI: (-41.76, 34.12) Aggressive responding—PSAP software program, 3 months, methadone versus healthy controls, MD: 230.79, 95% CI: (188.79, 272.79) Escape responding—PSAP software program, 3 months, buprenorphine versus healthy controls, MD: 39.89, 95% CI: (-64.35, 144.13) Escape responding—PSAP software program, 3 months, buprenorphine versus methadone, MD: -42.9, 95% CI: (-132.96, 47.16) Escape responding—PSAP software program, 3 months, methadone versus healthy controls, MD: 82.79, 95% CI: (-1, 166.58) Point-maintained responses—PSAP software program, 3 months, buprenorphine versus healthy controls, MD: -579.66, 95% CI: (-786.67, -372.65) Point-maintained responses—PSAP software program, 3 months, buprenorphine versus methadone, MD: 1676.34, 95% CI: (1456.18, 1896.5) Point-maintained responses—PSAP software program, 3 months, methadone versus healthy controls, MD: -2256, 95% CI: (-2509.48, -2002.52)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Giacomuzzi et al., 2003 Location: Europe Study design: cohort comparison	Number enrolled: 67 Number completed: 53 Mean age: methadone: 26.2, buprenorphine: 30.2 Percentage female: 34 Race/ethnicity: NR Years of opioid use: methadone: 7.9, buprenorphine: 9.2 Inclusion criteria: The only requirements for inclusion in the methadone or buprenorphine program were a confirmed diagnosis of opioid dependence (DSM-IV 304.0) and informed consent. Exclusion criteria: Forced discharge criteria were limited only to drug trafficking in the clinical center and aggressive behavior.	Intervention: methadone Setting: university hospital Route of administration: oral Duration (months): 6 Intervention 2: buprenorphine Setting: university hospital Route of administration: oral Duration (months): 6 Non-MAT comparator: none	Occupational/employment: Job: Satisfaction scores of the Lancashire Quality of Life Profile, 6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), MD: 0.3, 95% CI: (−0.70, 1.30) Physical: Satisfaction scores of the Lancashire Quality of Life Profile, 6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), MD: 0.0, 95% CI: (−0.81, 0.81) Fatigue or tiredness: percentage of participants methadone—oral (treatment) versus buprenorphine—sublingual (control), RR: 0.53, 95% CI: (0.28, 1.02) Fatigue or tiredness: percentage of participants—6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), risk difference (RD): −0.26, 95% CI: (−0.52, −0.00) Insomnia: Percentage of participants—6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), RR: 0.83, 95% CI: (0.47, 1.46) Insomnia: percentage of participants—6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), RD: −0.09, 95% CI: (−0.36, 0.18) Neurological Aggression: percentage of participants—6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), RR: 0.54, 95% CI: (0.24, 1.19) Aggression: percentage of participants—6 months, methadone—oral (treatment) versus buprenorphine—sublingual (control), RD: −0.20, 95% CI: (−0.45, 0.05)
Reference: Giacomuzzi et al., 2005 Location: Europe Study design: randomized controlled trial	Number enrolled: 53 Number completed: 35 Mean Age: methadone: 29.9 (5.5); buprenorphine 33.4 (7.6) Percentage female: 40 Race/ethnicity: NR Years of opioid use: 11.3 Inclusion criteria: The only requirement for inclusion in the follow-up was participation in the first study period including a confirmed diagnosis of opioid dependence (DSM-IV 304.0) and informed consent. Exclusion criteria: NR	Intervention: buprenorphine Setting: NR Route of administration: oral Duration (months): 36 Intervention2: methadone Setting: NR Route of administration: NR Duration (months): 36 Non-MAT comparator: none	Physical: Fatigue: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RR: 0.41, 95% CI: (0.17, 0.99) Insomnia: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RR: 0.77, 95% CI: (0.31, 1.87) Fatigue: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RD: −0.44, 95% CI: (−0.79, −0.09) Insomnia: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RD: −0.11, 95% CI: (−0.50, 0.27) Neurological: Aggressiveness: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RR: 0.46, 95% CI: (0.10, 2.08) Aggressiveness: percentage participants, 3 years, methadone (treatment versus buprenorphine (control), RD: −0.18, 95% CI: (−0.51, 0.15)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Giacomuzzi et al., 2006 Location: Europe Study design: randomized controlled trial	Number enrolled: 240 Number completed: NR Mean age: methadone: 27.3 (6.4), buprenorphine: 26.3 (7.5), morphine: 27.8 (4.8), control at admission: 25.3 (7.1) Percentage female: 40.4 Race/ethnicity: NR Years of opioid use: 8.1 years (SD 5.6) Inclusion criteria: Current diagnosis of opioid dependence based on DSM-IV at admission or participation in a methadone, sublingual buprenorphine, or slow-release oral morphine maintenance program for 6 months; aged 17 years or older; live within commuting distance of the hospital; and mentally competent to give informed consent. Exclusion criteria: Having an acute medical condition, currently using antipsychotic medication, or in another clinical trial. Forced discharge criteria were limited to drug trafficking in the clinical center or aggressive behavior.	Intervention: methadone Setting: university psychiatry department Route of administration: oral Duration (months): 6 Intervention2: buprenorphine Setting: university psychiatry department Route of administration: oral Duration (months): 6 Non-MAT comparator: 120 opioid users seeking a maintenance treatment program served as a comparison.	Physical : Number of participants reporting fatigue or tiredness, 6 months, buprenorphine versus treatment seekers (control), RR: 0.47, 95% CI: (0.29, 0.77) Number of participants reporting fatigue or tiredness, 6 months, buprenorphine versus methadone, RR: 0.6, 95% CI: (0.34, 1.06) Number of participants reporting fatigue or tiredness, 6 months, methadone versus treatment seekers (control), RR: 0.78, 95% CI: (0.56, 1.09) Percentage of participants reporting inability to sleep, 6 months, buprenorphine versus treatment seekers (control), RR: 0.75, 95% CI: (0.52, 1.07) Percentage of participants reporting inability to sleep, 6 months, buprenorphine versus methadone, RR: 1.31, 95% CI: (0.77, 2.21) Percentage of participants reporting inability to sleep, 6 months, methadone versus treatment seekers (control), RR: 0.57, 95% CI: (0.37, 0.88) Percentage of participants reporting insomnia, 6 months, buprenorphine versus treatment seekers (control), RR: 0.71, 95% CI: (0.47, 1.08) Percentage of participants reporting insomnia, 6 months, buprenorphine versus methadone, RR: 1, 95% CI: (0.58, 1.71) Percentage of participants reporting insomnia, 6 months, methadone versus treatment seekers (control), RR: 0.71, 95% CI: (0.47, 1.08)
Reference: Gossop et al., 1999 Location: Europe Study design: cohort comparison	Number enrolled: 452 Number completed: 452 Mean age: 29 Percentage female: 26.5 Race/ethnicity: NR Years of opioid use: 8.8 Inclusion criteria: Criteria for agency participation were location (agencies were chosen throughout England), and capacity to recruit a sufficient number of cases within the time available. All consecutive patients treated at an agency during the time period were included. Exclusion criteria: NR	Intervention: methadone Setting: methadone clinic Route of administration: oral Duration (months): NR Intervention 2: methadone Setting: doctor's office Route of administration: oral Duration (months): NR Non-MAT comparator: none	Crime: crime (non-drug related) Mean score, 6 months, methadone from specialist drug clinic (treatment) versus methadone from GP (control), MD: -3.8, 95% CI: (-7.06, -0.54)

Reference: Kinlock et al., 2009 (also Gordon et al., 2008) Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 211 Number completed: 204 Mean age: 40.3 (7.1) Percentage female: 0 Race/ethnicity: White: 24.0%, Black: 69.6%, Hispanic: NR, Asian: NR, other: 6.4% Years of opioid use: NR Inclusion criteria: Eligibility criteria were (1) 3 to 6 months before anticipated release from prison; (2) meeting DSM-IV criteria for heroin dependence at time of incarceration and being physiologically dependent during the year prior to incarceration; (3) suitability for methadone maintenance as determined by medical evaluation; (4) willingness to enroll in a prison-based methadone maintenance treatment program; and, (5) residing in Baltimore following release. Individuals who did not meet the heroin-dependence criterion were eligible if they were enrolled in an opioid treatment program in the year before incarceration. Exclusion criteria: Any of the following conditions: (1) renal failure; (2) liver failure; (3) pending/unadjudicated charges, which could have resulted in transfer to another correctional facility and/or additional prison time; and (4) a pending parole hearing.	Intervention: methadone Setting: prison, then community Route of administration: oral Duration (months): 12 Non-MAT comparator: 1. Counseling only—counseling in prison, with passive referral to treatment upon release. All participants received an individual intake by the study counselor and were subsequently scheduled to receive, within treatment condition, 12 weekly sessions of group-based education and discussion on relapse and overdose prevention, cocaine and alcohol abuse, and other reentry issues. Immediately prior to release, all participants were scheduled to meet with the study's counselor to discuss plans for release, including housing, employment concerns, and treatment options. 2. Counseling and transfer—counseling in prison, with immediate access to methadone maintenance treatment upon release from prison, but no maintenance treatment in prison. All participants received an individual intake by the study counselor and were subsequently scheduled to receive, within treatment condition, 12 weekly sessions of group-based education and discussion on relapse and overdose prevention, cocaine and alcohol abuse, and other reentry issues. Immediately prior to release, all participants were scheduled to meet with the study's counselor to discuss plans for release, including housing, employment concerns, and treatment options.	Occupational : Number of days employed during past 30 (12-month post-release), 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release, MD: -1.8, 95% CI: (-5.35, 1.75) Number of days employed during past 30 (12-month post-release), 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: -3.5, 95% CI: (-7.01, 0.01) Number of days employed during past 30 (12-month post-release), 12 months, counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: -1.7, 95% CI: (-5.3, 1.9) Behavioral/social : Arrested (yes/no) during 12-month follow-up period, 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release, RR: 0.9, 95% CI: (0.67, 1.2) Arrested (yes/no) during 12-month follow-up period, 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling only—counseling in prison, with passive referral to treatment upon release, RR: 1.04, 95% CI: (0.75, 1.44) Arrested (yes/no) during 12-month follow-up period, 12 months, counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release versus counseling only—counseling in prison, with passive referral to treatment upon release, RR: 1.16, 95% CI: (0.85, 1.59) Frequency (number of days) committing crimes in past year, 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release, MD: 16.6, 95% CI: (-17.52, 50.72) Frequency (number of days) committing crimes in past year, 12 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: -24.9, 95% CI: (-65.43, 15.63) Frequency (number of days) committing crimes in past year, 12 months, Counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: -41.5, 95%
---	--	--	--

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
			CI: (–80.35, –2.65) Number of days in last 6 months committed crime, 6 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release, MD: –7.1, 95% CI: (–22.12, 7.92)
(Continued) Reference: Kinlock et al., 2009 (also Gordon et al., 2008)			Number of days in last 6 months committed crime, 6 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: –28, 95% CI: (–43.32, –12.68) Number of days in last 6 months committed crime, 6 months, counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: –20.9, 95% CI: (–36.32, –5.48) Number of days incarcerated in last 6 months, 6 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release, MD: –1.9, 95% CI: (–16.92, 13.12) Number of days incarcerated in last 6 months, 6 months, counseling + methadone—methadone maintenance and counseling in prison, continued in community versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: 0, 95% CI: (–15.32, 15.32) Number of days incarcerated in last 6 months, 6 months, counseling + transfer—counseling in prison, with transfer to methadone maintenance upon release versus counseling only—counseling in prison, with passive referral to treatment upon release, MD: 1.9, 95% CI: (–13.52, 17.32)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Lee et al., 2016 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 308 Number completed: 245 Mean age: Sample mean 44 years; intervention 44.4 (9.2); control 43.2 (9.4) Percentage female: 15 Race/ethnicity: White: intervention 20.4%; control 19.4%, Black: intervention 53.3%; control 47.7%, Hispanic: intervention 24.3%; control 29.0% Years of opioid use: NR Inclusion criteria: Eligibility criteria were current (within the previous 12 months) or lifetime (any previous) opioid dependence (as defined by DSM-IV); a stated goal of opiate-free treatment rather than opioid agonist or partial-agonist maintenance therapy; an opioid-free status as confirmed by negative urine toxicological screening for all opioids before randomization; residence in the community and receipt of an adjudicated sentence that included supervision (e.g., parole, probation, outpatient drug-court programs, or other court-mandated treatment) or, in the previous 12 months, release from jail or prison, a plea-bargain arrangement, or any community supervision as above; general good health as determined by history and physical examination; an age of 18 to 60 years; and the ability to provide written informed consent. Exclusion criteria: Exclusion criteria were other drug or alcohol dependence requiring a level of care that would interfere with trial participation; pregnancy or a plan to conceive during the 24-week treatment phase, lactation, or an inability to use adequate contraceptive methods; an untreated psychiatric disorder or medical condition that might make participation hazardous, including liver-enzyme levels more than three times the upper limit of the normal range and a body mass index of more than 40; allergy to naltrexone, polylactide-co-glycolide, carboxymethylcellulose, or other components of the diluent; a current diagnosis of chronic pain for which opioids were prescribed; or a drug overdose in the previous 3 years requiring inpatient hospitalization.	Intervention: naltrexone Setting: doctor's office Route of administration: injection Duration (months): 6 Non-MAT comparator: 1. Participants in the usual-treatment group received similar counseling that was focused on adverse events, the prevention of relapse and overdose, and support for community treatment involvement from the same trial personnel.	Behavioral/social: Mean number of days incarcerated in the last 78 weeks, 78 weeks, XR-NTX versus counseling + referral to community treatment, MD: -6.16, 95% CI: (-36.65, 24.33) Percentage participants reincarcerated, 78 weeks, XR-NTX versus counseling + referral to community treatment, RR: 0.79, 95% CI: (0.54, 1.16)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Ling et al., 2010 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 163 Number completed: 88 Mean age: buprenorphine: 35.8 (11.0), placebo: 39.3 (11.7) Percentage female: 31.3 Race/ethnicity: White: 74.8%, Black: 12.3%, Hispanic: 14.7%, other: 12.9% Years of opioid use: NR Inclusion criteria: Men or nonpregnant women, aged 18 to 65 years, were required to meet DSM-IV diagnosis of current opioid dependence. Exclusion criteria: Exclusion criteria were AIDS, met DSM-IV criteria for current dependence on psychoactive substances other than opioids or nicotine, currently using nonprescribed benzodiazepines, had received medication treatment for opioid dependence within the previous 90 days, or had a current diagnosis of chronic pain that required opioid treatment. Patients were also excluded if they had any of the following: aspartate aminotransferase levels at least three times higher than the upper limit of normal, alanine aminotransferase levels at least three times the upper limit of normal, total bilirubin levels of at least 1.5 times the upper limit of normal, or creatinine levels at least 1.5 times the upper limit of normal.	Intervention: buprenorphine Setting: VA Route of administration: implant Duration(months): 6 Non-MAT comparator: 1. At the end of the induction phase, patients were randomized (stratified by sex and site) at a 2:1 ratio to double blind treatment with four placebo implants. The 2:1 ratio was used to reduce patient exposure to placebo implants.	Physical: Percentage of participants reporting insomnia, 6 months, buprenorphine versus placebo implant, RR: 0.98, 95% CI: (0.53, 1.81)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Lobmaier et al., 2010 Location: Europe Study design: randomized controlled trial	Number enrolled: 46 Number completed: unclear Mean age: 35.1 (7.0) Percentage female: 6.5 Race/ethnicity: NR Years of opioid use: NR Inclusion criteria: Inclusion criteria were pre-incarceration heroin dependence and at least 2 months, sentence time remaining. Exclusion criteria: Individuals were excluded if they presented with untreated major depression or psychosis, severe hepatic impairment, or if they were already in agonist maintenance treatment or pregnant.	Intervention: naltrexone Setting: prison, then community Route of administration: implant Duration (months): 6 Intervention 2: methadone Setting: prison, then community Route of administration: oral Duration (months): 6 Non-MAT comparator: none	Behavioral/social: Mean days per month illegal activity, 6 months, naltrexone (implant) versus methadone, MD: 0.5, 95% CI: (-9.98, 10.98)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Magura et al., 2009 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 116 Number completed: 91 Mean age: buprenorphine: 38.4 (7.9) methadone: 40.7 (9.1) Percentage female: 0 Race/ethnicity: White: NR, Black: 25%, Hispanic: 63.8%, Asian: NR, other: NR Years of opioid use: NR Inclusion criteria: Inmates who were eligible for the Key Extended Entry Program (KEEP), 18–65 years of age, sentenced to at least 10 days but less than 90 days of jail time (this was more restrictive than KEEP, to allow time for postrelease follow-up), expected to reside in New York City after release Exclusion criteria: Receiving methadone treatment in the community at remand to Rikers (such inmates are offered continuity of methadone maintenance by KEEP at the doses they received in the community), took nonprescribed “street” methadone within the previous 3 days, currently receiving more than 20 mg per day of prescribed methadone, current psychotic symptoms (e.g., schizophrenia, schizoaffective disorder) requiring referral for psychiatric intervention or currently treated with antipsychotic medication, HIV infection with T-lymphocytes less than 200 mm of blood, and/or presence of a serious opportunistic infection requiring treatment, receiving the HIV medication atazanavir, unable to complete an English-language interview	Intervention: Suboxone Setting: prison, then community Route of administration: oral Duration (months): 3 Intervention 2: methadone Setting: prison, then community Route of administration: oral Duration (months): 3 Non-MAT comparator: none	Behavioral/social: Percentage of participants arrested after release, 3 months, buprenorphine/naloxone versus methadone, MD: -0.02, 95% CI: (-0.33, 0.29) Percentage of participants reincarcerated, 3 months, buprenorphine/naloxone versus methadone, RR: 0.8, 95% CI: (0.53, 1.2)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Mattick et al., 2003 Location: Australia, New Zealand Study design: randomized controlled trial	Number enrolled: 405 Number completed: 216 Mean age: 30 (8) Percentage female: 31 Race/ethnicity: other: English-speaking background: methadone 79%, buprenorphine: 79%; non-English-speaking background: methadone: 16% buprenorphine: 16%; indigenous: methadone: 5%, buprenorphine: 6% Years of opioid use: methadone: 7.6 (6.7), buprenorphine: 7.7 (7.0) Inclusion criteria: Patients were eligible for the study if they had a current diagnosis of opioid dependence using the criteria in the fourth edition of the DSM-IV); were aged 18 years or older; lived within commuting distance of the clinic; appeared mentally competent to give informed consent; and signed informed consent. Exclusion criteria: Patients were excluded from the study if they were pregnant or nursing, or were deemed likely to become pregnant in the study period; suffering an acute medical condition that could make participation in the study medically hazardous (e.g., active tuberculosis, unstable cardiovascular, or serious acute liver disease); currently using anticonvulsant medication, disulfiram, or antipsychotic medication; in opioid replacement treatment (methadone was the only available medication at the time of the study) in the preceding 30 days (and this status was verified from state department of health records); unable to attend the clinic daily for the study period; in a study of buprenorphine previously, or were currently in another clinical trial.	Intervention: buprenorphine Setting: methadone clinic Route of administration: oral Duration (months): 3.25 Intervention 2: methadone Setting: methadone clinic Route of administration: oral Duration (months): 3.25 Non-MAT comparator: None	Physical : Percentage reporting somnolence, 13 weeks, buprenorphine versus methadone, RR: 0.56, 95% CI: (0.26, 1.18) Number (percentage) of patients who reported experiencing pain during the trial, 13 weeks, buprenorphine versus methadone, RR: 0.93, 95% CI: (0.58, 1.51) Percentage of participants reporting insomnia, 13 weeks, buprenorphine versus methadone, RR: 1.3, 95% CI: (0.75, 2.26)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Metrebian, 2015 (also Gordon et al., 2008) Location: Europe Study design: randomized controlled trial	Number enrolled: 127 Number completed: NR Mean age: 37.2 (6.5) Percentage female: 27 Race/ethnicity: White: 96 Years of opioid use: 16.6 Inclusion criteria: Specific eligibility criteria were based on national guidance on injectable opiate treatment. The sample was thus locally residing, treatment-resistant, chronic, opiate-dependent patients receiving oral substitution treatment (methadone or buprenorphine treatment) for at least 6 months preceding recruitment to the trial and who, despite this, were still injecting street heroin on most days. Exclusion criteria: NR	Intervention: methadone Setting: methadone clinic Route of administration: injection Duration (months): 6 Intervention 2: methadone Setting: methadone clinic Route of administration: oral Duration (months): 6 Non-MAT comparator: 1. Supervised injectable heroin. This intervention is not used as MAT in the United States.	Physical : SF-36 physical function, 6 months, supervised injectable methadone versus optimized oral methadone, MD: 7.4, 95% CI: (−4.21, 19.01) Behavioral/social: SF-36—mental health, 6 months, supervised injectable methadone versus optimized oral methadone, MD: −2.16, 95% CI: (−8.94, 4.62) SF-36 social functioning, 6 months, supervised injectable methadone versus optimized oral methadone, MD: −5.2, 95% CI: (−16.57, 6.17) Criminal behavior in previous month—percentage of participants, 6 months, supervised injectable methadone versus optimized oral methadone, RR: 0.9, 95% CI: (0.4, 2.04) Percentage of participants committing crimes over follow-up period, 26 weeks, injectable methadone (supervised) versus oral methadone (optimized), RR: 1.05, 95% CI: (0.64, 1.72)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Neri et al., 2005 Location: Europe Study design: randomized controlled trial	Number enrolled: 62 Number completed: 57 Mean age: methadone: 27 (6), buprenorphine: 24 (5) Percentage female: 11.3 Race/ethnicity: NR Years of opioid use: methadone: 2, buprenorphine: 2 Inclusion criteria: Diagnosis according to the DSM-IV criteria Exclusion criteria: Subjects with severe psychiatric illness (dementia, psychosis, and cognitive impairment) who were unable to answer the questions were not enrolled in the investigation, as were individuals presenting with codependence of alcohol, amphetamines, cannabinoids, and benzodiazepines; however, sporadic use (less than a month with negative urine screen) was not considered an exclusion criterion. Subjects who missed medication for 3 consecutive days (for methadone) and once (for buprenorphine) were removed from the study.	Intervention: methadone Setting: other substance abuse treatment Route of administration: oral Duration (months): 12 Intervention 2: buprenorphine Setting: other substance abuse treatment Route of administration: oral Duration (months): 12 Non-MAT comparator: none	Physical: Number of participants reporting fatigue, 12 months, buprenorphine versus methadone, RR: 0.95, 95% CI: (0.23, 3.81) Number of participants reporting nausea, 12 months, buprenorphine versus methadone, RR: 1.01, 95% CI: (0.31, 3.33) Number of participants reporting sleepiness, 12 months, buprenorphine versus methadone, RR: 1.26, 95% CI: (0.28, 5.67) Percentage of participants reporting insomnia, 12 months, buprenorphine versus methadone, RR: 0.9, 95% CI: (0.33, 2.47)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Neumann et al., 2013 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 54 Number completed: 26 Mean age: 38.3 (9.7) Percentage female: 46.3 Race/ethnicity: White: 85.2% Years of opioid use: NR Inclusion criteria: Men and women ages 18 years or older with well-documented chronic nonmalignant pain related to the spine or a large joint (e.g., hip, knee, shoulder) and an addiction to prescription opioids. Exclusion criteria: Individuals were excluded from the study if (1) they were homeless or placed on parole; (2) were unable to give consent (e.g., because of neurological disorders, including dementia or cognitive dysfunction, psychosis) or lacked consent from the attending physician; (3) had a co-occurring psychiatric disorder (e.g., schizophrenia); (4) had an electrocardiogram showing prolonged QT and/or previous cardiac/pulmonary issues; (5) were taking a medication that is contraindicated with methadone or buprenorphine; (6) had a prior history of methadone or buprenorphine maintenance treatment; or (7) were pregnant.	Intervention: Suboxone Setting: doctor's office Route of administration: oral Duration (months): 6 Intervention 2: methadone Setting: doctor's office Route of administration: oral Duration (months): 6 Non-MAT comparator: none	Physical: Numerical rating scale pain score, 6 months, buprenorphine/naloxone versus methadone, MD: -0.96, 95% CI: (-3.54, 1.63)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Newman and Whitehill, 1979 Location: Asia Study design: randomized controlled trial	Number enrolled: 100 Number completed: 100 Mean age: 38 Percentage female: NR Race/ethnicity: NR Years of opioid use: methadone: 15; control: 12 Inclusion criteria: (1) Male, age 22–58 years, (2) documented history of heroin addiction for at least 4 years and at least one previous course of treatment, (3) evidence of current addiction to heroin as determined by 3 consecutive positive urine tests for morphine, (4) voluntary application for admission (referrals by the criminal justice system were excluded), (5) a resident with proven fixed address in Kowloon in a district near the treatment clinic, (6) absence of past or present major psychiatric or medical illness (for example, tuberculosis, peptic ulcer, psychosis). Exclusion criteria: NR	Intervention: methadone Setting: methadone clinic Route of administration: oral Duration (months): NR Non-MAT comparator: 1. Control group was weaned from methadone by a dose reduction of 1 mg/day, weaning (detoxification) being completed after 60 days. Thereafter, controls were maintained on a placebo solution that was indistinguishable by taste from the methadone solution.	Behavioral/social: Conviction rate per 100 man-months of study, 3 years, methadone versus placebo, incidence rate ratio: 0.45, 95% CI: not calculable

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Rapeli et al., 2009 (also Rapeli et al., 2011) Location: Europe Study design: case control	Number enrolled: 43 Number completed: unclear Mean age: methadone: 29.2 (6.8), buprenorphine: 27.7 (6.8), control: 28.7 (9.6) Percentage female: 44.2 Race/ethnicity: NR Years of opioid use: methadone: 15.0 (5.1), buprenorphine: 13.4 (5.2) Inclusion criteria: All participants included in the study were between 18 and 50 years old and native Finnish speakers. For opioid substitution patients, additional inclusion criteria were opioid dependence diagnosis, benzodiazepine dependence or abuse diagnosis, start of opioid substitute therapy during the last 2 months, and treatment using methadone, buprenorphine, or buprenorphine/naloxone. Exclusion criteria: Participants showing signs of current intoxication, ongoing binge on any substance of abuse, or any extra psychoactive drug dose within 24 hours were excluded. Participants with uncontrolled polysubstance abuse, acute alcohol abuse, or acute axis I psychiatric morbidity other than substance abuse related were excluded. Also excluded were participants with severe brain injury, chronic neurological disease, history of other than substance induced psychoses, epileptic seizures, HIV infection, pregnancy, or primary cognitive deficit. For these purposes, psychiatric interviews by clinical psychiatrist were conducted for all participants, and diagnostic criteria from the DSM-IV were applied.	Intervention: methadone Setting: other substance abuse treatment Route of administration: oral Duration (months): 6–9 Intervention 2: buprenorphine, Suboxone Setting: other substance abuse treatment Route of administration: oral Duration (months): 6–9 Non-MAT comparator: 1. Healthy nondrug-using adults recruited from adult education centers and by word of mouth	Cognitive : Go-NoGo errors: TAP, 12–17 months (T3), buprenorphine versus healthy controls, MD: 0.4, 95% CI: (–0.1, 0.9) Go-NoGo errors: TAP, 12–17 months (T3), buprenorphine versus methadone, MD: 0.1, 95% CI: (–0.59, 0.79) Go-NoGo errors: TAP, 12–17 months (T3), methadone versus healthy controls, MD: 0.3, 95% CI: (–0.26, 0.86) Go-NoGo errors: TAP, 6–9 months (T2), buprenorphine versus healthy controls, MD: 0, 95% CI: (–0.58, 0.58) Go-NoGo errors: TAP, 6–9 months (T2), buprenorphine versus methadone, MD: –0.5, 95% CI: (–1.12, 0.12) Go-NoGo errors: TAP, 6–9 months (T2), methadone versus healthy controls, MD: 0.5, 95% CI: (–0.13, 1.13) Go-NoGo reaction time (milliseconds [ms]): TAP, 12–17 months (T3), buprenorphine versus healthy controls, MD: 31, 95% CI: (–3.61, 65.61) Go-NoGo reaction time (ms): TAP, 12–17 months (T3), buprenorphine versus methadone, MD: –36, 95% CI: (–77, 5) Go-NoGo reaction time (ms): TAP, 12–17 months (T3), methadone versus healthy controls, MD: 67, 95% CI: (25.83, 108.17) Go-NoGo reaction time (ms): TAP, 6–9 months (T2), buprenorphine versus healthy controls, MD: 37, 95% CI: (–7.58, 81.58) Go-NoGo reaction time (ms): TAP, 6–9 months (T2), buprenorphine versus methadone, MD: –68, 95% CI: (–127.44, –8.56) Go-NoGo reaction time (ms): TAP, 6–9 months (T2), methadone versus healthy controls, MD: 105, 95% CI: (38.74, 171.26) Letter-number sequencing from the WMS-III, 12–17 months (T3), buprenorphine versus healthy controls, MD: –0.6, 95% CI: (–2.69, 1.49) Letter-number sequencing from the WMS-III, 12–17 months (T3), buprenorphine versus methadone, MD: 1.8, 95% CI: (0.03, 3.57) Letter-number sequencing from the WMS-III, 12–17 months (T3), methadone versus healthy controls, MD: –2.4, 95% CI: (–4.5, –0.3) Letter-number sequencing from the WMS-III, 6–9 months (T2), buprenorphine versus healthy controls, MD: –2.8, 95% CI: (–4.8, –0.8) Letter-number sequencing from the WMS-III, 6–9 months (T2), buprenorphine versus methadone, MD: 0.3, 95% CI: (–1.43, 2.03)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
(Continued) Reference: Rapeli et al., 2009 (also Rapeli et al., 2011)			Letter-number sequencing from the WMS-III, 6–9 months (T2), methadone versus healthy controls, MD: –3.1, 95% CI: (–5.08, –1.12) Letter-number sequencing from the WMS-III, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus healthy controls, MD: –2.4, 95% CI: (–4.27, –0.53) Letter-number sequencing from the WMS-III, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus methadone, MD: 0.6, 95% CI: (–1.03, 2.23) Letter-number sequencing from the WMS-III, 6–9 months, methadone versus healthy controls, MD: –3, 95% CI: (–4.86, –1.14) PASAT, 12–17 months (T3), buprenorphine versus healthy controls, MD: –14, 95% CI: (–21.17, –6.83) PASAT, 12–17 months (T3), buprenorphine versus methadone, MD: 0.9, 95% CI: (–7.18, 8.98) PASAT, 12–17 months (T3), methadone versus healthy controls, MD: –14.9, 95% CI: (–22.15, –7.65) PASAT, 6–9 months (T2), buprenorphine versus healthy controls, MD: –10.8, 95% CI: (–16.89, –4.71) PASAT, 6–9 months (T2), buprenorphine versus methadone, MD: 1.6, 95% CI: (–4.94, 8.14) PASAT, 6–9 months (T2), methadone versus healthy controls, MD: –12.4, 95% CI: (–19.49, –5.31) PASAT from the FORAMEN Rehab software package, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus healthy controls, MD: –11.9, 95% CI: (–18.02, –5.78) PASAT from the FORAMEN Rehab software package, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus methadone, MD: 2.5, 95% CI: (–3.82, 8.82) PASAT from the FORAMEN Rehab software package, 6–9 months, methadone versus healthy controls, MD: –14.4, 95% CI: (–20.82, –7.98) Phasic alertness/reaction time with warning signal (ms): TAP, 12–17 months (T3), buprenorphine versus healthy controls, MD: 4, 95% CI: (–11.76, 19.76) Phasic alertness/reaction time with warning signal (ms): TAP, 12–17 months (T3), buprenorphine versus methadone, MD: –25, 95% CI: (–50.91, 0.91) Phasic alertness/reaction time with warning signal (ms): TAP, 12–17 months (T3), methadone versus healthy controls, MD: 29, 95% CI: (2.76, 55.24) Phasic alertness/reaction time with warning signal (ms): TAP, 6–9 months (T2), buprenorphine versus healthy controls, MD: 5, 95% CI: (–11.19, 21.19) Phasic alertness/reaction time with warning signal (ms): TAP, 6–9 months (T2), buprenorphine versus methadone, MD: –26, 95% CI: (–44.88, –7.12)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
(Continued) Rapeli et al., 2009 (also Rapeli et al., 2011)			Phasic alertness/reaction time with warning signal (ms): TAP, 6–9 months (T2), methadone versus healthy controls, MD: 31, 95% CI: (12.67, 49.33) Tonic alertness/simple reaction time (ms): TAP, 12–17 months (T3), buprenorphine versus healthy controls, MD: 1, 95% CI: (–18.28, 20.28) Tonic alertness/simple reaction time (ms): TAP, 12–17 months (T3), buprenorphine versus methadone, MD: –25, 95% CI: (–48.57, –1.43) Tonic alertness/simple reaction time (ms): TAP, 12–17 months (T3), methadone versus healthy controls, MD: 26, 95% CI: (3.04, 48.96) Tonic alertness/simple reaction time (ms): TAP, 6–9 months (T2), buprenorphine versus healthy controls, MD: 3, 95% CI: (–11.99, 17.99) Tonic alertness/simple reaction time (ms): TAP, 6–9 months (T2), buprenorphine versus methadone, MD: –27, 95% CI: (–41.99, –12.01) Tonic alertness/simple reaction time (ms): (TAP, 6–9 months (T2), methadone versus healthy controls, MD: 30, 95% CI: (14.44, 45.56) List learning—Memory for Persons Data, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus healthy controls, MD: –0.3, 95% CI: (–0.67, 0.07) List learning—Memory for Persons Data, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus methadone, MD: –0.3, 95% CI: (–0.67, 0.07) List learning—Memory for Persons Data, 6–9 months, methadone versus healthy controls, MD: 0, 95% CI: (–0.15, 0.15) Logical memory, delayed—Logical Memory from the WMS-III, 12–17 months (T3), buprenorphine versus healthy controls, MD: –3.5, 95% CI: (–6.49, –0.51) Logical memory, delayed—Logical Memory from the WMS-III, 12–17 months (T3), buprenorphine versus methadone, MD: 0.6, 95% CI: (–2.78, 3.98) Logical memory, delayed—Logical Memory from the WMS-III, 12–17 months (T3), methadone versus healthy controls, MD: –4.1, 95% CI: (–7.2, –1) Logical memory, delayed—Logical Memory from the WMS-III, 6–9 months (T2), buprenorphine versus healthy controls, MD: –3.6, 95% CI: (–6.39, –0.81) Logical memory, delayed—Logical Memory from the WMS-III, 6–9 months (T2), buprenorphine versus methadone, MD: –1.7, 95% CI: (–4.78, 1.38) Logical memory, delayed—Logical Memory from the WMS-III, 6–9 months (T2), methadone versus healthy controls, MD: –1.9, 95% CI: (–4.55, 0.75)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
(Continued) Rapeli et al., 2009 (also Rapeli et al., 2011)			Logical memory, immediate—Logical Memory from the WMS-III, 12–17 months (T3), buprenorphine versus healthy controls, MD: -2.4, 95% CI: (-5.17, 0.37) Logical memory, immediate—Logical Memory from the WMS-III, 12–17 months (T3), buprenorphine versus methadone, MD: 1.2, 95% CI: (-2.03, 4.43) Logical memory, immediate—Logical Memory from the WMS-III, 12–17 months (T3), methadone versus healthy controls, MD: -3.6, 95% CI: (-6.32, -0.88) Logical memory, immediate—Logical Memory from the WMS-III, 6–9 months (T2), buprenorphine versus healthy controls, MD: -2.5, 95% CI: (-4.93, -0.07) Logical memory, immediate—Logical Memory from the WMS-III, 6–9 months (T2), buprenorphine versus methadone, MD: -1, 95% CI: (-3.61, 1.61) Logical memory, immediate—Logical Memory from the WMS-III, 6–9 months (T2), methadone versus healthy controls, MD: -1.5, 95% CI: (-4.06, 1.06) Raw score for Memory Complaint Questionnaire (MCQ)—Finnish version, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus healthy controls, MD: 4.1, 95% CI: (0.63, 7.57) Raw score for MCQ—Finnish version, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus methadone, MD: -1.1, 95% CI: (-4.91, 2.71) Raw score for MCQ—Finnish version, 6–9 months, methadone versus healthy controls, MD: 5.2, 95% CI: (3.3, 7.1) Story recall task—Logical Memory from the WMS III: immediate free recall, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus healthy controls, MD: -2.2, 95% CI: (-4.49, 0.09) Story recall task—Logic Memory from the WMS III: immediate free recall, 6–9 months, buprenorphine (buprenorphine/naloxone or buprenorphine only) versus methadone, MD: -0.1, 95% CI: (-2.47, 2.27) Story recall task—Logic Memory from the WMS III: immediate free recall, 6–9 months, methadone versus healthy controls, MD: -2.1, 95% CI: (-4.4, 0.2)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reijneveld and Plomp, 1993 Location: Europe Study design: cohort comparison	Number enrolled: 38 Number completed: 38 Mean age: current clients: 31.0 (1.5) and former clients 30.1 (1.2) Percentage female: 36.8 Race/ethnicity: NR Years of opioid use: current clients 14.6 and former clients 11.9 Inclusion criteria: Born in the Netherlands or a (former) Dutch colony (Surinam or the Dutch Antilles), registered inhabitants of the city of Amsterdam, and registered in the central methadone registry for the first time 5 years ago. Exclusion criteria: Clients who had left care were only included if they had received methadone in at least two instances 50 days apart to exclude clients only passing through the methadone maintenance system.	Intervention: methadone Setting: doctor's office Route of administration: oral Duration (months): varies Non-MAT comparator: 1. Individuals who enrolled in methadone maintenance but dropped out	Behavioral/social: Percentage of crime that reports some illegal activity in past week, patients continuing on methadone for 5 years versus dropouts (RD -0.04; CI -0.20, 0.13) Occupational: Percentage of employment reporting no time spent on job, past month, patients continuing on methadone for 5 years versus dropouts, RD -0.35; CI (-0.65, -0.05)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Rosenthal et al., 2013 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 287 Number completed: 163 Mean age: 35.7 Percentage female: 39 Race/ethnicity: White: 83.3%, Black: 13.0% Years of opioid use: NR Inclusion criteria: Men and nonpregnant women (aged 18–65 years) met the DSM-IV diagnosis of current opioid dependence as determined by the Mini International Neuropsychiatric Interview Exclusion criteria: Individuals were excluded if they had AIDS, a clinically low platelet count, substance dependence on other than opioids or nicotine, received methadone or buprenorphine for opioid dependence within 90 days, current diagnosis of chronic pain requiring opioid analgesics or currently using nonprescribed benzodiazepines. Subjects were also excluded with aspartate aminotransferase levels three times the upper limit of normal; alanine aminotransferase levels three times the upper limit of normal; total bilirubin one and a half times the upper limit of normal; and/or creatinine one and a half times the upper limit of normal.	Intervention: buprenorphine Setting: home Route of administration: implant Duration (months): 6 Intervention 2: Suboxone Setting: other substance abuse treatment Route of administration: oral Duration (months): 6 Non-MAT comparator: 1. Placebo implants. Manual-guided individual drug counseling sessions (10) were provided by experienced counselors twice weekly during weeks 1–12, and then weekly for the subsequent 12 weeks.	Physical: Percentage of participants, reporting insomnia, 6 months, buprenorphine—implant versus placebo implants (control), RR: 0.59, 95% CI: (0.23, 1.49) Percentage of participants reporting insomnia, 6 months, buprenorphine—implant versus buprenorphine/naloxone—sublingual, RR: 0.53, 95% CI: (0.25, 1.14) Percentage of participants reporting insomnia, 6 months, buprenorphine/naloxone—sublingual versus placebo implants (control), RR: 1.1, 95% CI: (0.49, 2.47)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Schwartz et al., 2009 (also Schwartz et al., 2007) Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 319 Number completed: 134 Mean age: 41.4 (6.0) Percentage female: 41 Race/ethnicity: White: 6.6%, Black: 93.4%, Hispanic: NR, Asian: NR, other: NR Years of opioid use: NR Inclusion criteria: NR Exclusion criteria: NR	Intervention: methadone Setting: clinic, unclear Route of administration: NR Duration (months): 10 Non-MAT comparator: 1. Waiting list	Behavioral/social : Days of illegal activities/last 30 days, 10 months, methadone—interim maintenance versus methadone—waiting list, MD: -5.2, 95% CI: (-7.37, -3.03) Days of illegal activities/last 30 days, 4 months, methadone—interim maintenance versus methadone—waiting list, MD: -5.2, 95% CI: (-7.14, -3.26) Percentage of participants arrested, 12 months, methadone—interim maintenance versus methadone—waiting list (control), MD: -0.06, 95% CI: (-0.34, 0.22) Percentage of participants arrested, 24 months, methadone—interim maintenance versus methadone—waiting list (control), MD: -0.15, 95% CI: (-0.6, 0.3) Percentage of participants arrested, 6 months, methadone—interim maintenance versus methadone—waiting list (control), MD : -0.14, 95% CI: (-0.35, 0.07) Percentage of participants with no arrest charge (0), 12 months, methadone—interim maintenance versus methadone—waiting list (control), RR : 0.99, 95% CI: (0.86, 1.14) Percentage of participants with no arrest charge (0), 24 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 1.12, 95% CI: (0.92, 1.36) Percentage of participants with no arrest charge (0), 6 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 1.06, 95% CI: (0.95, 1.18) Percentage of participants with nonsevere arrest charges (1–4), 12 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 0.99, 95% CI: (0.65, 1.49) Percentage of participants with nonsevere arrest charges (1–4), 24 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 0.84, 95% CI: (0.63, 1.12) Percentage of participants with non-severe arrest charges (1–4), 6 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 0.65, 95% CI: (0.39, 1.1) Percentage of participants with severe arrest charges (5–7), 12 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 1.4, 95% CI: (0.37, 5.34) Percentage of participants with severe arrest charges (5–7), 24 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 0.98, 95% CI: (0.37, 2.62) Percentage of participants with severe arrest charges (5–7), 6 months, methadone—interim maintenance versus methadone—waiting list (control), RR: 3.75, 95% CI: (0.44, 32.25)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Sees et al., 2000 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 179 Number completed: 134 Mean age: methadone detox 39.4 (7.91); methadone maintenance 39.4 (8.57) Percentage female: 41.3 Race/ethnicity: White: 51.4%, Black: 30.2%, Hispanic: 12.8%, other: 5.6% Years of opioid use: methadone detox 15.7 (9.26); MMT 16.6 (9.42) Inclusion criteria: DSM-III-R criteria for a diagnosis of opioid dependence and had an initial urine screening test positive for opioid other than methadone and negative for methadone. Women of childbearing age were required to be practicing birth control. Exclusion criteria: Medical or psychiatric conditions that contraindicated methadone treatment, enrolled in a substance abuse treatment program, had been in a methadone treatment program within the previous week or were in a follow-up phase of a previous methadone detoxification research protocol, could not be expected to remain in the study for 12 months, did not have signs of opioid withdrawal on three occasions, younger than 18 years old, or pregnant or breastfeeding.	Intervention: methadone detox only Setting: VA Route of administration: oral Duration (months): 14 Intervention 2: methadone Setting: VA Route of administration: oral Duration (months): 14 Non-MAT comparator: none	Physical : ASI medical score, 12 months, psychosocially enriched 180-day methadone-assisted detoxification versus methadone, MD: -0.06, 95% CI: (-0.16, 0.04) Occupational : ASI employment score, 12 months, psychosocially enriched 180-day methadone-assisted detoxification versus methadone, MD: -0.01, 95% CI: (-0.1, 0.08) Behavioral/social : ASI family function score, 12 months, psychosocially enriched 180-day methadone-assisted detoxification versus methadone, MD: -0.01, 95% CI: (-0.04, 0.02) ASI legal score, 12 months, psychosocially enriched 180-day methadone-assisted detoxification versus methadone, MD: 0.08, 95% CI: (0.02, 0.14) ASI psychiatric score, 12 months, psychosocially enriched 180-day methadone-assisted detoxification versus methadone, MD: -0.04, 95% CI: (-0.11, 0.03)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Reference: Senay et al., 1993 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 130 Number completed: 130 Mean age: 41.4 Percentage female: 33 Race/ethnicity: White: 70.8%, Black: 17.7%, Hispanic: 11.5% Years of opioid use: NR Inclusion criteria: Ages 21–60, in methadone treatment for past 1 year and no plans to attend detoxification in the next year, not on parole or probation at the time of entrance to the study, for the most recent 6-month period—all urines clean, employed or engaged in appropriate activity, no arrests, and compliance with treatment as evidenced by fulfilling all assignments in the treatment plan. Exclusion criteria: NR	Intervention: methadone Setting: home Route of administration: oral Duration (months): 6 Intervention 2: methadone Setting: methadone clinic Route of administration: oral Duration (months): 6 Non-MAT comparator: none	Physical : ASI medical score, 6 months, twice a month methadone take-home schedule versus control (twice or three times a week take-home schedule), MD: -0.01, 95% CI: (-0.04, 0.02) Occupational : ASI employment score, 6 months, twice a month methadone take-home schedule versus control (twice or three times a week take-home schedule), MD: 0, 95% CI: (-0.11, 0.11) Behavioral/social : ASI legal score, 6 months, twice a month methadone take-home schedule versus control (twice or three times a week take-home schedule), MD: 0, 95% CI: (-0.02, 0.02) ASI psychiatric score, 6 months, twice a month methadone take-home schedule versus control (twice or three times a week take-home schedule), MD: 0.02, 95% CI: (-0.01, 0.04)
Reference: Soyka et al., 2005 Location: Europe Study design: randomized controlled trial	Number enrolled: 62 Number completed: 46 Mean age: buprenorphine, 34.2; methadone: 32 Percentage female: 41.3 Race/ethnicity: NR Years of opioid use: buprenorphine: 11.5, methadone: 11 Inclusion criteria: All patients met the International Statistical Classification of Diseases, 10th revision, and DSM-IV criteria for opioid dependence. All patients were free of withdrawal symptoms when tested, had a driver's license (valid or withdrawn), or were experienced drivers. Exclusion criteria: Patients were excluded if they had a disabling physical disorder or organic brain disorder.	Intervention: buprenorphine Setting: unclear Route of administration, other: NR Duration (months): 2–2.5 (entire study) Intervention 2: methadone Setting: unclear Route of administration: NR Duration (months): follow up at 2–2.5 months (entire study 6 months) Non-MAT comparator: none	Cognitive : Attention test (Q1), correct answers, 8–10 weeks, buprenorphine versus methadone, MD: 7.7, 95% CI: (-29, 44.4) Decision and reaction test (DR2)—false decisions, 8–10 weeks, buprenorphine versus methadone, MD: 1.37, 95% CI: (-2.76, 0.02) DR2—time of decision, 8–10 weeks, buprenorphine versus methadone, MD: 13.88, 95% CI: (-34.45, 62.21) Peripheral vision test (PVT)—reaction time, total seconds, 8–10 weeks, buprenorphine versus methadone, MD: 0, 95% CI: (-0.23, 0.23) PVT—tracking performance (mean deviation), 8–10 weeks, buprenorphine versus methadone, MD: -1.1, 95% CI: (-2.2, 0) Visual perception, tachistoscope test (TT15), correct answers, 8–10 weeks, buprenorphine versus methadone, MD: -0.2, 95% CI: (-2.58, 2.18) Neurological : Reactive stress tolerance test (RST3), phase 1, omissions, 8–10 weeks, buprenorphine versus methadone, MD: 0.5, 95% CI: (-2, 3) RST3, phase 2, omissions, 8–10 weeks, buprenorphine versus methadone, MD: -2.4, 95% CI: (-14.28, 9.48) RST3, phase 3, omissions, 8–10 weeks, buprenorphine versus methadone, MD: 2.6, 95% CI: (-7.91, 13.11)

Reference: Soyka et al., 2008 Location: Europe Study design: randomized controlled trial	Number enrolled: 59 Number completed: 46 Mean age: NR Percentage female: NR Race/ethnicity: NR Years of opioid use: NR Inclusion criteria: (a) No confirmed subjective memory complaints or history of organic brain syndrome or seizures, (b) no measurable cognitive and memory impairment, (c) IQ of 85 or greater, and (d) neither neurological nor psychiatric diagnosis or history apart from the opioid dependence in the patient group. A group of healthy normal controls, matched for age, sex, and education level, was recruited, to detect possible impairments of cognitive ability in the patient group. Exclusion criteria: NR	Intervention: methadone Setting: outpatient clinic Route of administration: oral Duration (months): 6 Intervention 2: buprenorphine Setting: other, outpatient clinic Route of administration: oral Duration(months): 6 Non-MAT comparator: 1. A group of healthy normal controls, matched for age, sex, and education level, was recruited, to detect possible impairments of cognitive ability in the patient group.	Cognitive: Regensburg Word Fluency Test (RWT)—lexical, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -7.2, 95% CI: (-11.09, -3.31) RWT—lexical, 8–10 weeks (t2), buprenorphine versus methadone, MD: 0.9, 95% CI: (-3.57, 5.37) RWT—lexical, 8–10 weeks (t2), methadone versus healthy controls, MD: -8.1, 95% CI: (-13.13, -3.07) RWT—phonemic, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -3.1, 95% CI: (-6.66, 0.46) RWT—phonemic, 8–10 weeks (t2), buprenorphine versus methadone, MD: -0.1, 95% CI: (-3.21, 3.01) RWT—phonemic, 8–10 weeks (t2), methadone versus healthy controls, MD: -3, 95% CI: (-6.12, 0.12) RWT—shifting (lexical), 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -3.4, 95% CI: (-6.19, -0.61) RWT—shifting (lexical), 8–10 weeks (t2), buprenorphine versus methadone, MD: -1.1, 95% CI: (-4.45, 2.25) RWT—shifting (lexical), 8–10 weeks (t2), methadone versus healthy controls, MD: -2.3, 95% CI: (-5.27, 0.67) RWT—shifting (phonemic), 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -2.8, 95% CI: (-5.65, 0.05) RWT—shifting (phonemic), 8–10 weeks (t2), buprenorphine versus methadone, MD: -2.2, 95% CI: (-5.44, 1.04) RWT—shifting (phonemic), 8–10 weeks (t2), methadone versus healthy controls, MD: -0.6, 95% CI: (-3.57, 2.37) Trail-Making Test A (TMT-A), 8–10 weeks (t2), buprenorphine versus healthy controls, MD: 5.5, 95% CI: (0.32, 10.68) TMT-A, 8–10 weeks (t2), buprenorphine versus methadone, MD: -0.4, 95% CI: (-5.98, 5.18) TMT-A, 8–10 weeks (t2), methadone versus healthy controls, MD: 5.9, 95% CI: (1.49, 10.31) TMT-B, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: 26, 95% CI: (-43.35, 95.35) TMT-B, 8–10 weeks (t2), buprenorphine versus methadone, MD: 4.4, 95% CI: (-78.87, 87.67) TMT-B, 8–10 weeks (t2), methadone versus healthy controls, MD: 21.6, 95% CI: (-59.33, 102.53) VLMT—verbal digit span, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -0.2, 95% CI: (-2.34, 1.94) VLMT—verbal digit span, 8–10 weeks (t2), buprenorphine versus methadone, MD: 0.6, 95% CI: (-1.77, 2.97) VLMT—verbal digit span, 8–10 weeks (t2), methadone versus healthy controls, MD: -0.8, 95% CI: (-2.88, 1.28) VLMT—verbal learning, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -12.5, 95% CI: (-17.21, -7.79) VLMT—verbal learning, 8–10 weeks (t2), buprenorphine versus methadone, MD: -1.2, 95% CI: (-6.66, 4.26)
---	--	---	---

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
(Continued) Soyka et al., 2008			VLMT—verbal learning, 8–10 weeks (t2), methadone versus healthy controls, MD: -11.3, 95% CI: (-16.04, -6.56) VLMT—verbal memory, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: 1, 95% CI: (-0.3, 2.3) VLMT—verbal memory, 8–10 weeks (t2), buprenorphine versus methadone, MD: -0.5, 95% CI: (-1.88, 0.88) VLMT—verbal memory, 8–10 weeks (t2), methadone versus healthy controls, MD: 1.5, 95% CI: (0.48, 2.52) d2-Test of Attention, 8–10 weeks (t2), buprenorphine versus healthy controls, MD : -11.5, 95% CI: (-29.45, 6.45) d2-Test of Attention, 8–10 weeks (t2), buprenorphine versus healthy controls, MD: -42.4, 95% CI: (-86.46, 1.66) d2-Test of Attention, 8–10 weeks (t2), buprenorphine versus methadone, MD: -13.4, 95% CI: (-32.26, 5.46) d2-Test of Attention, 8–10 weeks (t2), buprenorphine versus methadone, MD: -29.2, 95% CI: (-75.11, 16.71) d2-Test of Attention, 8–10 weeks (t2), methadone versus healthy controls, MD: 1.9, 95% CI: (-17.81, 21.61) d2-Test of Attention, 8–10 weeks (t2), methadone versus healthy controls, MD: -13.2, 95% CI: (-53.45, 27.05)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Strain et al., 1993 Location: United States or Canada Study design: randomized controlled trial	Number enrolled: 95 Number completed: 95 Mean age: 34.7 (5.2) Percentage female: 32 Race/ethnicity: Black: 52% Years of opioid use: 7.4 Inclusion criteria: Ages between 18 and 50 years old, history of intravenous opioid dependence (including documentation of previous treatment for opioid dependence or legal involvement secondary to opioid use, a urine sample positive for opioids, and physical examination consistent with acute and chronic needle use), no chronic medical illnesses, absence of major mental illness, a negative pregnancy test for females, and at least 3 months since the patient's last treatment at the clinic. Exclusion criteria: NR	Intervention: methadone Setting: methadone clinic Route of administration: oral Duration (months): 9 Intervention 2: methadone Setting: methadone clinic Route of administration: oral Duration (months): 9 Non-MAT comparator: 1. Patients assigned to the 0 mg (placebo) condition received a 35-day methadone detoxification starting at 25 mg in the first week, and decreasing by 5 mg each week. Patients were assigned an individual counselor who set treatment goals and developed an individualized treatment plan. Patients received weekly group therapy focusing upon relapse prevention. On-site medical services were provided by a full-time internist and a part-time nurse practitioner.	Behavioral/social : Mean number of days' illegal activity in the past 30 days, 20 weeks, 20 mg methadone versus placebo, MD: 3, 95% CI: (-2.34, 8.34) Mean number of days' illegal activity in the past 30 days, 20 weeks, 50 mg methadone versus 20 mg methadone, MD: -5.5, 95% CI: (-9.6, -1.4) Mean number of days' illegal activity in the past 30 days, 20 weeks, 50 mg methadone versus placebo, MD: -2.5, 95% CI: (-7.63, 2.63) Number of crimes committed, 20 weeks, 20 mg methadone versus placebo, MD: -4, 95% CI: (not calculable) Number of crimes committed, 20 weeks, 50 mg methadone versus 20 mg methadone, MD: -5.23, 95% CI: (not calculable) Number of crimes committed, 20 weeks, 50 mg methadone versus placebo, MD: -9.23, 95% CI: (not calculable)

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Strang et al., 2000	Number enrolled: 37	Intervention: methadone	Physical :
Location: Europe	Number completed: 33	Setting: methadone clinic	Maudsley Addiction Profile (MAP): 10-item physical health symptom scale from the MAP measuring general health, cardiorespiratory, gastrointestinal, musculoskeletal, and neurological symptoms, 6 months; injectable methadone versus oral methadone, MD: -0.8, 95% CI: (-5.1, 3.5)
Study design: randomized controlled trial	Mean age: IV MMT, 31.9; oral MMT, 32.1	Route of administration: injection	Behavioral/social :
	Percentage female: IV MMT, 9.5; oral MMT, 6.3	Duration (months): 6	Number of days the individual reported committing shoplifting, robbery, burglary, and fraud (time period not reported), 6 months, injectable methadone versus oral methadone, MD: -3.6, 95% CI: (-10.02, 2.82)
	Race/ethnicity: other, nonwhite, IV MMT, 9.5%; oral MMT: 12.5%	Intervention 2: methadone	
	Years of opioid use: NR	Setting: methadone clinic	
	Inclusion criteria: Aged at least 23 years; illicit injecting for at least the last 3 years; at least one previous episode of opiate substitution treatment; adequate venous access on arms.	Route of administration: oral	
	Exclusion criteria: Current serious medical or psychiatric comorbidity or pregnant.	Duration (months): 6	
		Non-MAT comparator: none	

Study Details	Participants	Intervention Treatment	Functional Outcomes Results
Tiihonen et al., 2012 Location: Europe Study design: randomized controlled trial	Number enrolled: 100 Number completed: 40 Mean age: placebo: 29.3 (4.38), naltrexone: 28.0 (4.10) Percentage female: NR Race/ethnicity: NR Years of opioid use: placebo: 8.7 (2.83), naltrexone: 8.2 (3.75) Inclusion criteria: Primary DSM-IV diagnosis of concurrent amphetamine and opioid dependence, present for at least 1 year; age between 18 and 50 years; education level of high school graduate or above; negative urine toxicology and alcohol breath tests; no current use of psychotropic medications; at least one relative willing to participate in the treatment (e.g., to monitor the administration of medications, assist in follow-up, and provide outcome data); a stable address in St. Petersburg or in the nearest districts of Leningrad Region; a home telephone number at which the patient could be reached; willingness and ability to give informed consent and otherwise participate; and, for women of childbearing age, a negative pregnancy test and use of adequate contraception. Exclusion criteria: Clinically significant cognitive impairment, schizophrenia, a paranoid disorder, bipolar disorder, or a seizure disorder; advanced neurological, cardiovascular, renal, or hepatic disease; active tuberculosis; a current febrile illness; an AIDS-defining illness; a significant laboratory abnormality, such as severe anemia, unstable diabetes, or liver function test results greater than three times normal values; pregnancy; pending legal charges with potential impending incarceration; concurrent participation in another treatment study; and concurrent treatment in another substance abuse program.	Intervention: naltrexone Setting: other substance abuse treatment Route of administration: implant Duration (months): 2.5 Non-MAT comparator: 1. Placebo implant identical in appearance	Physical : Number of participants reporting fatigue, 10 weeks, naltrexone (sustained release implant) versus placebo, RR: 1, 95% CI: (0.06, 15.55) Percentage of participants reporting insomnia, 10 weeks, naltrexone (sustained release implant) versus placebo, RR: 1, 95% CI: (0.21, 4.72) Neurological : Percentage of participants reporting nervousness, irritability, 10 weeks, naltrexone (sustained release implant) versus placebo, RR: 1, 95% CI: (0.15, 6.82) Number of participants reporting nervousness, 10 weeks, naltrexone (sustained release implant) versus placebo, RR: 0, 95% CI: (not calculable)

Appendix E. Excluded Publications

Publications That Do Not Meet Inclusion Criteria (with Reasons for Exclusion)

- Abbott, P. J., B. Moore, H. Delaney, and S. Weller, “Retrospective Analyses of Additional Services for Methadone Maintenance Patients,” *Journal of Substance Abuse Treatment*, Vol. 17, No. 1–2, July–September 1999, pp. 129–137. Reason for exclusion: Only measures effects of a co-intervention.
- Abbott, P. J., S. B. Weller, H. D. Delaney, and B. A. Moore, “Community Reinforcement Approach in the Treatment of Opiate Addicts,” *American Journal of Drug and Alcohol Abuse*, Vol. 24, No. 1, February 1998, pp. 17–30. Reason for exclusion: Only measures effects of a co-intervention.
- Abidizadegan, Afsaneh, Alireza Moradi, and Robert Famam, “The Executive Functions in Methadone Maintenance Patients,” *Advances in Cognitive Science*, Vol. 10, No. 3, Fall 2008, 2008, pp. 75–82. Reason for exclusion: Wrong study design.
- Abrahms, J. L., “A Cognitive-Behavioural Versus Nondirective Group Treatment Program for Opioid Addicted Persons: An Adjunct to Methadone Maintenance,” *The International Journal of the Addictions*, Vol. 14, No. 4, 1979. Reason for exclusion: Only measures effects of a co-intervention.
- “Abuse-Deterrent Opioid Formulations,” *Medical Letter on Drugs and Therapeutics*, Vol. 57, No. 1476, August 31, 2015, pp. 119–121. Reason for exclusion: Wrong study design.
- Accurso, A. J., and D. A. Rastegar, “The Effect of a Payer-Mandated Decrease in Buprenorphine Dose on Aberrant Drug Tests and Treatment Retention Among Patients with Opioid Dependence,” *Journal of Substance Abuse Treatment*, Vol. 61, February 2016, pp. 74–79. Reason for exclusion: No functional outcomes.
- Achmad, Y. M., A. N. Istiqomah, S. Iskandar, R. Wisaksana, R. van Crevel, and T. Hidayat, “Integration of Methadone Maintenance Treatment and HIV Care for Injecting Drug Users: A Cohort Study in Bandung, Indonesia,” *Acta Medica Indonesiana*, Vol. 41, Supplement 1, July 2009, pp. 23–27. Reason for exclusion: No functional outcomes.
- Acosta, Michelle C., Lisa A. Marsch, Haiyi Xie, Honoria Guarino, and Yesenia Aponte-Melendez, “A Web-Based Behavior Therapy Program Influences the Association Between Cognitive Functioning and Retention and Abstinence in Clients Receiving Methadone Maintenance Treatment,” *Journal of Dual Diagnosis*, Vol. 8, No. 4, November 2012, pp. 283–293. Reason for exclusion: Only measures effects of a co-intervention.

- “Addiction,” *DATA: The Brown University Digest of Addiction Theory and Application*, Vol. 27, No. 1, 2008, p. 2p. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- “Addiction Research Roundup,” *DATA: The Brown University Digest of Addiction Theory and Application*, Vol. 27, No. 10, 2008, p. 2p. Reason for exclusion: Wrong study design.
- “Addiction: Research Roundup,” *DATA: The Brown University Digest of Addiction Theory and Application*, Vol. 29, No. 8, 2010, p. 2p. Reason for exclusion: Wrong study design.
- Adi, Y., A. Juarez-Garcia, D. Wang, S. Jowett, E. Frew, E. Day, S. Bayliss, T. Roberts, and A. Burls, “Oral Naltrexone as a Treatment for Relapse Prevention in Formerly Opioid-Dependent Drug Users: A Systematic Review and Economic Evaluation,” *Health Technology Assessment*, Vol. 11, No. 6, February 2007, pp. iii–iv, 1–85. Reason for exclusion: No functional outcomes.
- Ahmadi, J., “A Controlled Trial of Buprenorphine Treatment for Opium Dependence: The First Experience from Iran,” *Drug and Alcohol Dependence*, Vol. 66, No. 2, April 1, 2002, pp. 111–114. Reason for exclusion: No functional outcomes.
- Ahmadi, Jamshid, “A Randomized, Clinical Trial of Buprenorphine Maintenance Treatment for Iranian Patients with Opioid Dependency,” *Addictive Disorders and Their Treatment*, Vol. 1, No. 1, 2002, pp. 25–27. Reason for exclusion: Not English.
- Ahmadi, J., “Methadone Versus Buprenorphine Maintenance for the Treatment of Heroin-Dependent Outpatients,” *Journal of Substance Abuse Treatment*, Vol. 24, No. 3, April 2003, pp. 217–220. Reason for exclusion: No functional outcomes.
- Ahmadi, J., and K. Ahmadi, “Controlled Trial of Maintenance Treatment of Intravenous Buprenorphine Dependence,” *Irish Journal of Medical Science*, Vol. 172, No. 4, October–December 2003, pp. 171–173. Reason for exclusion: No functional outcomes.
- Ahmadi, J., K. Ahmadi, and J. Ohaeri, “Controlled, Randomized Trial in Maintenance Treatment of Intravenous Buprenorphine Dependence with Naltrexone, Methadone or Buprenorphine: A Novel Study,” *European Journal of Clinical Investigation*, Vol. 33, No. 9, September 2003, pp. 824–829. Reason for exclusion: No functional outcomes.
- Ahmadi, J., and M. Ahmadi, “Twelve-Month Maintenance Treatment of Heroin-Dependent Outpatients with Buprenorphine,” *Journal of Substance Use*, Vol. 8, No. 1, March 2003, pp. 39–41. Reason for exclusion: No functional outcomes.
- Ahmadi, J., M. Babaee-Beigi, M. Alishahi, I. Maany, and T. Hidari, “Twelve-Month Maintenance Treatment of Opium-Dependent Patients,” *Journal of Substance Abuse Treatment*, Vol. 26, No. 1, January 2004, pp. 363–366. Reason for exclusion: No functional outcomes.
- Ahmadi, J., and N. Bahrami, “Buprenorphine Treatment of Opium-Dependent Outpatients Seeking Treatment in Iran,” *Journal of Substance Abuse Treatment*, Vol. 23, No. 4, December 2002, pp. 415–417. Reason for exclusion: No functional outcomes.

- Ahmadi, Jamshid, Hassan Farrashbandi, Masoud Moosavinasab, Mohammadali Babaee, Ali Firoozabadi, Mohammadsadegh Mohagheghzadeh, Joel Porter, Saxby Pridmore, and Gholamreza Dehbozorgi, "Treatment of Heroin Dependence," *German Journal of Psychiatry*, Vol. 7, No. 2, 2004, pp. 1–5. Reason for exclusion: No functional outcomes.
- Ahmadi, J., I. Maany, and M. Ahmadi, "Treatment of Intravenous Buprenorphine Dependence: A Randomized Open Clinical Trial," *German Journal of Psychiatry*, Vol. 6, No. 1, 2003, pp. 23–29. Reason for exclusion: No functional outcomes.
- Alaee, A., M. Zarghami, S. Farnia, M. Khademloo, and T. Khoddad, "Comparison of Brain White Matter Hyperintensities in Methamphetamine and Methadone Dependent Patients and Healthy Controls," *Iranian Journal of Radiology*, Vol. 11, No. 2, May 2014, p. e14275. Reason for exclusion: No functional outcomes.
- Albayaty, M., M. Linden, H. Olsson, M. Johnsson, K. Strandgården, and F. Tiberg, "Pharmacokinetic Evaluation of Once-Weekly and Once-Monthly Buprenorphine Subcutaneous Injection Depots (CAM2038) Versus Intravenous and Sublingual Buprenorphine in Healthy Volunteers Under Naltrexone Blockade: An Open-Label Phase 1 Study," *Advances in Therapy*, 2017. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Ali, R., C. McGregor, P. Thomas, and L. Gowing, "A Randomised Controlled Trial of Antagonist-Precipitated Heroin Withdrawal Under Anaesthetic Prior to Naltrexone Maintenance: Outcomes at 6 and 12 Months," *Drug and Alcohol Dependence*, Vol. 63, Supplement 1, 2001. Reason for exclusion: Conference abstract.
- Altaf, A., S. A. Shah, and S. Vermund, "Importance of Research and Services Among People Who Inject Drugs in Pakistan," *Journal of Pakistan Medical Association*, Vol. 64, No. 12, December 2014, pp. 1413–1414. Reason for exclusion: Wrong study design.
- Altice, F. L., L. E. Sullivan, D. Smith-Rohrberg, S. Basu, S. Stancliff, and L. Eldred, "The Potential Role of Buprenorphine in the Treatment of Opioid Dependence in HIV-Infected Individuals and in HIV Infection Prevention," *Clinical Infectious Diseases*, Vol. 43, Supplement 4, December 15, 2006, pp. S178–S183. Reason for exclusion: Wrong study design.
- Amass, L., W. K. Bickel, S. T. Higgins, and J. R. Hughes, "Administering Twice the Daily Buprenorphine Dose Suppresses Opioid Withdrawal for 48 Hours in Opioid-Dependent Humans," *NIDA Research Monograph*, Vol. 141, 1994. Reason for exclusion: Conference abstract.
- Amass, L., W. K. Bickel, J. P. Crean, J. Blake, and S. T. Higgins, "Alternate-Day Buprenorphine Dosing Is Preferred to Daily Dosing by Opioid-Dependent Humans," *Psychopharmacology (Berl)*, Vol. 136, No. 3, April 1998, pp. 217–225. Reason for exclusion: Wrong study design.
- Amass, L., W. K. Bickel, S. T. Higgins, and G. J. Badger, "Alternate-Day Dosing During Buprenorphine Treatment of Opioid Dependence," *Life Sciences*, Vol. 54, No. 17, 1994, pp. 1215–1228. Reason for exclusion: No functional outcomes.

- Amass, L., J. B. Kamien, and S. K. Mikulich, "Thrice-Weekly Supervised Dosing with the Combination Buprenorphine-Naloxone Tablet Is Preferred to Daily Supervised Dosing by Opioid-Dependent Humans," *Drug and Alcohol Dependence*, Vol. 61, No. 2, 2001. Reason for exclusion: No functional outcomes.
- Amass, L., J. B. Kamien, and S. K. Mikulich, "Efficacy of Daily and Alternate-Day Dosing Regimens with the Combination Buprenorphine-Naloxone Tablet," *Drug and Alcohol Dependence*, Vol. 58, No. 1–2, February 1, 2000, pp. 143–152. Reason for exclusion: No functional outcomes.
- Amass, L., V. Pukeleviciene, E. Subata, A. R. Almeida, M. C. Pieri, P. D'Egidio, Z. Stankova, A. Costa, B. P. Smyth, S. Sakoman, Y. Wei, and J. Strang, "A Prospective, Randomized, Multicenter Acceptability and Safety Study of Direct Buprenorphine/Naloxone Induction in Heroin-Dependent Individuals," *Addiction*, Vol. 107, No. 1, January 2012, pp. 142–151. Reason for exclusion: No functional outcomes.
- "American Academy of Addiction Psychiatry 23rd Annual Meeting and Symposium Poster Presentations," *The American Journal on Addictions*, Vol. 22, No. 3, May–June 2013, pp. 305–321. Reason for exclusion: Conference abstract.
- "Analgesic Designed to Reduce Euphoric Effects of Abuse," *Journal of Gerontological Nursing*, Vol. 35, No. 12, 2009, p. 5. Reason for exclusion: Wrong study design.
- Anderson, B. A., M. L. Faulkner, J. J. Rilee, S. Yantis, and C. L. Marvel, "Attentional Bias for Nondrug Reward Is Magnified in Addiction," *Experimental and Clinical Psychopharmacology*, Vol. 21, No. 6, December 2013, pp. 499–506. Reason for exclusion: Wrong study design.
- Anderson, I. B., and T. E. Kearney, "Use of Methadone," *The Western Journal of Medicine*, Vol. 172, No. 1, January 2000, pp. 43–46. Reason for exclusion: Wrong study design.
- Andresz, V., N. Marcantoni, F. Binder, M. Velten, M. Alt, J. C. Weber, and D. Stephan, "Puffy Hand Syndrome Due to Drug Addiction: A Case-Control Study of the Pathogenesis," *Addiction*, Vol. 101, No. 9, September 2006, pp. 1347–1351. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Anglin, M. D., B. T. Conner, J. Annon, and D. Longshore, "Levo-Alpha-Acetylmethadol (LAAM) Versus Methadone Maintenance: 1-Year Treatment Retention, Outcomes and Status," *Addiction*, Vol. 102, No. 9, September 2007, pp. 1432–1442. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Anglin, M. D., B. T. Conner, J. J. Annon, and D. Longshore, "Longitudinal Effects of LAAM and Methadone Maintenance on Heroin Addict Behavior," *Journal of Behavioral Health Services and Research*, Vol. 36, No. 2, April 2009, pp. 267–282. Reason for exclusion: Wrong study design.

- Apelt, S. M., N. Scherbaum, and M. Soyka, "Induction and Switch to Buprenorphine-Naloxone in Opioid Dependence Treatment: Predictive Value of the First Four Weeks," *Heroin Addiction and Related Clinical Problems*, Vol. 16, No. 3, 2014. Reason for exclusion: Wrong study design.
- Appel, P. W., H. Joseph, A. Kott, W. Nottingham, E. Tasiny, and E. Habel, "Selected In-Treatment Outcomes of Long-Term Methadone Maintenance Treatment Patients in New York State," *Mount Sinai Journal of Medicine*, Vol. 68, No. 1, January 2001, pp. 55–61. Reason for exclusion: Wrong study design.
- Arkin, E., "Studies Confirm Effectiveness of Harm Reduction for People Who Inject Drugs," *HIV AIDS Policy Law Review*, Vol. 15, No. 3, October 2011, p. 29. Reason for exclusion: Wrong study design.
- "As Opioid Use Increases, MAT Must Expand as Well: The Vermont Role Model . . . Medication-Assisted Treatment," *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 46, 2014, pp. 1–3. Reason for exclusion: Wrong study design.
- Ashworth, O., "Methadone Maintenance Treatment as an Effective Harm Minimisation Intervention," *Mental Health Practice*, Vol. 8, No. 8, 2005, pp. 24–27. Reason for exclusion: Wrong study design.
- Athanasos, P., A. L. Farquharson, P. Compton, P. Psaltis, and J. Hay, "Electrocardiogram Characteristics of Methadone and Buprenorphine Maintained Subjects," *Journal of Addictive Diseases*, Vol. 27, No. 3, 2008, pp. 31–35. Reason for exclusion: No functional outcomes.
- Athanasos, P., G. Morrish, A. A. Somogyi, F. Bochner, and J. M. White, "Changes in Methadone Concentration, Opioid Effects, and Opioid Withdrawal During Induction onto Maintenance Treatment," *Addictive Disorders and Their Treatment*, Vol. 3, No. 3, 2004, pp. 122–128. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Athanasos, P., C. S. Smith, J. M. White, A. A. Somogyi, F. Bochner, and W. Ling, "Methadone Maintenance Patients Are Cross-Tolerant to the Antinociceptive Effects of Very High Plasma Morphine Concentrations," *Pain*, Vol. 120, No. 3, 2006. Reason for exclusion: Wrong study design.
- Avants, S. K., A. Margolin, and S. McKee, "A Path Analysis of Cognitive, Affective, and Behavioral Predictors of Treatment Response in a Methadone Maintenance Program," *Journal of Substance Abuse*, Vol. 11, No. 3, 2000, pp. 215–230. Reason for exclusion: Only measures effects of a co-intervention.
- Avants, S. K., A. Margolin, J. L. Sindelar, B. J. Rounsaville, R. Schottenfeld, S. Stine, N. L. Cooney, R. A. Rosenheck, S. H. Li, and T. R. Kosten, "Day Treatment Versus Enhanced Standard Methadone Services for Opioid-Dependent Patients: A Comparison of Clinical Efficacy and Cost," *American Journal of Psychiatry*, Vol. 156, No. 1, January 1999, pp. 27–33. Reason for exclusion: Only measures effects of a co-intervention.

- Awgu, Ezechukwu, Stephen Magura, and Andrew Rosenblum, "Heroin-Dependent Inmates' Experiences with Buprenorphine or Methadone Maintenance," *Journal of Psychoactive Drugs*, Vol. 42, No. 3, September 2010, pp. 329–346. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Ayres, R., J. Ingram, A. Rees, J. Neale, A. Beattie, and M. Telfer, "Enhancing Motivation Within a Rapid Opioid Substitution Treatment Feasibility RCT: A Nested Qualitative Study," *Substance Abuse Treatment, Prevention, and Policy*, Vol. 9, 2014, p. 44. Reason for exclusion: No functional outcomes.
- Azizi, A., A. Borjali, and M. Golzari, "The Effectiveness of Emotion Regulation Training and Cognitive Therapy on the Emotional and Addictional Problems of Substance Abusers," *Iranian Journal of Psychiatry*, Vol. 5, No. 2, Spring 2010, pp. 60–65. Reason for exclusion: Only measures effects of a co-intervention.
- Bach, P., S. Vollstadt-Klein, U. Frischknecht, M. Hoerst, F. Kiefer, K. Mann, G. Ende, and D. Hermann, "Diminished Brain Functional Magnetic Resonance Imaging Activation in Patients on Opiate Maintenance Despite Normal Spatial Working Memory Task Performance," *Clinical Neuropharmacology*, Vol. 35, No. 4, July–August 2012, pp. 153–160. Reason for exclusion: Wrong study design.
- Badrakalimuthu, Vellingiri Raja, Andrew Tarbuck, and Ajay Wagle, "Maintenance Treatment Programme for Opioid Dependence: Characteristics of 50+Age Group," *Drugs and Alcohol Today*, Vol. 12, No. 3, 2012, pp. 157–163. Reason for exclusion: Wrong study design.
- Baewert, A., W. Gombas, S. D. Schindler, A. Peternell-Moelzer, H. Eder, R. Jagsch, and G. Fischer, "Influence of Peak and Trough Levels of Opioid Maintenance Therapy on Driving Aptitude," *European Addiction Research*, Vol. 13, No. 3, 2007. Reason for exclusion: Wrong study design.
- Bailey, K. P., "Pharmacological Treatments for Substance Use Disorders," *Journal of Psychosocial Nursing and Ment Health Services*, Vol. 42, No. 8, August 2004, pp. 14–20. Reason for exclusion: Wrong study design.
- Baker, J. G., J. B. Rounds, and C. A. Carson, "Monitoring in Methadone Maintenance Treatment," *The International Journal of the Addictions*, Vol. 30, No. 9, July 1995, pp. 1177–1185. Reason for exclusion: Only measures effects of a co-intervention.
- Baker, J. R., A. M. Best, P. A. Pade, and E. F. McCance-Katz, "Effect of Buprenorphine and Antiretroviral Agents on the QT Interval in Opioid-Dependent Patients," *Annals of Pharmacotherapy*, Vol. 40, No. 3, March 2006, pp. 392–396. Reason for exclusion: Wrong study design.
- Baldacchino, A., D. J. Balfour, and K. Matthews, "Impulsivity and Opioid Drugs: Differential Effects of Heroin, Methadone and Prescribed Analgesic Medication," *Psychological Medicine*, Vol. 45, No. 6, April 2015, pp. 1167–1179. Reason for exclusion: Wrong study design.

- Balhara, Y. P., “Time to Include Buprenorphine-Naloxone Combination in the WHO Model List of Essential Medicines,” *Journal of Opioid Management*, Vol. 9, No. 4, July–August 2013, p. 237. Reason for exclusion: Wrong study design.
- Ballough, W., R. Crowe, and J. Elbert, “Buprenorphine Versus Methadone in Opioid Dependent Patients,” *Pharmacotherapy*, Vol. 33, No. 10, 2013. Reason for exclusion: Conference abstract.
- Balster, R. L., and G. E. Bigelow, “Guidelines and Methodological Reviews Concerning Drug Abuse Liability Assessment,” *Drug and Alcohol Dependence*, Vol. 70, 2003, pp. S13–40. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bandawar, M., A. Kandasamy, P. Chand, P. Murthy, and V. Benegal, “Adherence to Buprenorphine Maintenance Treatment in Opioid Dependence Syndrome: A Case Control Study,” *Indian Journal of Psychological Medicine*, Vol. 37, No. 3, July–September 2015, pp. 330–332. Reason for exclusion: No functional outcomes.
- Bandstra, E. S., “Maternal Opioid Treatment: Human Experimental Research (MOTHER) Study: Maternal, Fetal and Neonatal Outcomes from Secondary Analyses,” *Addiction*, Vol. 107, Supplement 1, November 2012, pp. 1–4. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Barnett, P. G., “Comparison of Costs and Utilization Among Buprenorphine and Methadone Patients,” *Addiction*, Vol. 104, No. 6, June 2009, pp. 982–992. Reason for exclusion: Wrong study design.
- Barnett, P. G., C. L. Masson, J. L. Sorensen, W. Wong, and S. Hall, “Linking Opioid-Dependent Hospital Patients to Drug Treatment: Health Care Use and Costs 6 Months After Randomization,” *Addiction*, Vol. 101, No. 12, December 2006, pp. 1797–1804. Reason for exclusion: Only measures effects of a co-intervention.
- Barratt, D. T., J. K. Collier, and A. A. Somogyi, “Association Between the DRD2 A1 Allele and Response to Methadone and Buprenorphine Maintenance Treatments,” *American Journal of Medical Genetics B: Neuropsychiatric Genetics*, Vol. 141b, No. 4, June 5, 2006, pp. 323–331. Reason for exclusion: No functional outcomes.
- Barry, D. T., C. J. Cutter, M. Beitel, C. Liong, and R. S. Schottenfeld, “Cognitive-Behavioral Therapy and Educational Counseling for Chronic Pain and Opioid Dependence,” *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Only measures effects of a co-intervention.
- Bart, G., “CSAT’s QT Interval Screening in Methadone Report: Outrageous Fortune or Sea of Troubles?” *Journal of Addictive Diseases*, Vol. 30, No. 4, October 2011, pp. 313–317. Reason for exclusion: Wrong study design.
- Baser, O., M. Chalk, D. A. Fiellin, and D. R. Gastfriend, “Cost and Utilization Outcomes of Opioid-Dependence Treatments,” *American Journal of Managed Care*, Vol. 17, Supplement 8, June 2011, pp. S235–248. Reason for exclusion: No functional outcomes.

- Battistella, S., N. Constantinou, C. J. Morgan, P. Davis, D. O’Ryan, and H. V. Curran, “Semantic Priming and Verbal Learning in Current Opiate Users, Ex-Users and Non-User Controls,” *Human Psychopharmacology*, Vol. 27, No. 5, September 2012, pp. 499–506. Reason for exclusion: Wrong study design.
- Bawor, M., B. B. Dennis, M. C. Samaan, C. Plater, A. Worster, M. Varenbut, J. Daiter, D. C. Marsh, D. Desai, M. Steiner, R. Anglin, M. Coote, G. Pare, L. Thabane, and Z. Samaan, “Methadone Induces Testosterone Suppression in Patients with Opioid Addiction,” *Scientific Reports*, Vol. 4, August 26, 2014, p. 6189. Reason for exclusion: No functional outcomes.
- Bearn, J., M. Gossop, and J. Strang, “Randomised Double-Blind Comparison of Lofexidine and Methadone in the In-Patient Treatment of Opiate Withdrawal,” *Drug and Alcohol Dependence*, Vol. 43, No. 1–2, December 2, 1996, pp. 87–91. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bearn, J., M. Gossop, J. Strang, and Y. Tang, “Lofexidine and Methadone in the Treatment of Opiate Addicts in Hospital: A Double Blind Randomized Controlled Study,” *Chinese Journal of Drug Abuse Prevention and Treatment*, No. 6, 1998. Reason for exclusion: Only measures effects of a co-intervention.
- Beck, M., A. Mirmohammadsadegh, B. Franz, J. Blanke, and U. R. Hengge, “Opioid Receptors on White Blood Cells: Effect of HIV Infection and Methadone Treatment,” *Pain*, Vol. 98, No. 1–2, July 2002, pp. 187–194. Reason for exclusion: Only measures effects of a co-intervention.
- Beck, T., C. Haasen, U. Verthein, S. Walcher, C. Schuler, M. Backmund, C. Ruckes, and J. Reimer, “Maintenance Treatment for Opioid Dependence with Slow-Release Oral Morphine: A Randomized Cross-Over, Non-Inferiority Study Versus Methadone,” *Addiction*, Vol. 109, No. 4, April 2014, pp. 617–626. Reason for exclusion: No functional outcomes.
- Becker, A. B., I. A. Liebson, and G. E. Bigelow, “Transdermal Buprenorphine: Abuse Potential Assessment in Non-Opioid Volunteers,” *Drug and Alcohol Dependence*, Vol. 63, Supplement 1, 2001. Reason for exclusion: Conference abstract.
- Beebe, K. L., S. Chavoustie, W. Ling, S. Sigmon, D. Leiderman, and G. Bailey, “Buprenorphine Implants for the Treatment of Opioid Dependence: Six and 12 Month Outcomes,” *Neuropsychopharmacology*, Vol. 38, 2012. Reason for exclusion: Conference abstract.
- Beitel, M., S. Peters, J. D. Savant, C. J. Cutter, J. J. Cecero, and D. T. Barry, “The Psychometric Properties of the Iowa Personality Disorder Screen in Methadone-Maintained Patients: An Initial Investigation,” *Journal of Personality Disorders*, Vol. 29, No. 1, February 2015, pp. 131–144. Reason for exclusion: Wrong study design.
- Bell, James, Gaye Byron, Amy Gibson, and Amanda Morris, “A Pilot Study of Buprenorphine-Naloxone Combination Tablet (Suboxone®) in Treatment of Opioid Dependence,” *Drug and Alcohol Review*, Vol. 23, No. 3, September 2004, pp. 311–317. Reason for exclusion: Wrong study design.

- Bell, J., J. Chan, and A. Kuk, "Investigating the Influence of Treatment Philosophy on Outcome of Methadone Maintenance," *Addiction*, Vol. 90, No. 6, June 1995, pp. 823–830. Reason for exclusion: Only measures effects of a co-intervention.
- Bell, J., M. Shanahan, C. Mutch, F. Rea, A. Ryan, R. Batey, A. Dunlop, and A. Winstock, "A Randomized Trial of Effectiveness and Cost-Effectiveness of Observed Versus Unobserved Administration of Buprenorphine-Naloxone for Heroin Dependence," *Addiction*, Vol. 102, No. 12, December 2007, pp. 1899–1907. Reason for exclusion: No functional outcomes.
- Bell, J., J. Shearer, A. Ryan, R. Graham, K. Korompay, S. Rizzo, D. Sindhusake, and A. A. Somogyi, "The Acceptability, Safety, and Tolerability of Methadone/Naloxone in a 50:1 Ratio," *Experimental and Clinical Psychopharmacology*, Vol. 17, No. 3, June 2009, pp. 146–153. Reason for exclusion: No functional outcomes.
- Bell, J., and D. Zador, "A Risk-Benefit Analysis of Methadone Maintenance Treatment," *Drug Safety*, Vol. 22, No. 3, March 2000, pp. 179–190. Reason for exclusion: Wrong study design.
- Bell, J. R., A. Ryan, C. Mutch, R. Batey, and F. Rea, "Optimising the Benefits of Unobserved Dose Administration for Stable Opioid Maintenance Patients: Follow-Up of a Randomised Trial," *Drug and Alcohol Dependence*, Vol. 96, No. 1–2, July 1, 2008, pp. 183–186. Reason for exclusion: Only measures effects of a co-intervention.
- Benko, L. B., "Drug-Free in Seattle. Harborview Hospital Takes on Heroin," *Modern Healthcare*, Vol. 30, No. 13, March 27, 2000, pp. 56–58. Reason for exclusion: Wrong study design.
- Bershad, A. K., J. H. Jaffe, E. Childs, and H. Wit, "Opioid Partial Agonist Buprenorphine Dampens Responses to Psychosocial Stress in Humans," *Psychoneuroendocrinology*, Vol. 52, No. 1, 2015. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bershad, A., J. Jaffe, and H. Wit, "Buprenorphine Dampens Responses to Psychosocial Stress in Healthy Adults," *Neuropsychopharmacology*, Vol. 39, 2014. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bickel, W. K., L. Amass, J. P. Crean, and G. J. Badger, "Buprenorphine Dosing Every 1, 2, or 3 Days in Opioid-Dependent Patients," *Psychopharmacology (Berl)*, Vol. 146, No. 2, September 1999, pp. 111–118. Reason for exclusion: Wrong study design.
- Bickel, W. K., S. T. Higgins, and M. L. Stitzer, "Choice of Blind Methadone Dose Increases by Methadone Maintenance Patients," *Drug and Alcohol Dependence*, Vol. 18, No. 2, October 1986, pp. 165–171. Reason for exclusion: Wrong study design.
- Bickel, W. K., L. A. Marsch, A. R. Buchhalter, and G. J. Badger, "Computerized Behavior Therapy for Opioid-Dependent Outpatients: A Randomized Controlled Trial," *Experimental and Clinical Psychopharmacology*, Vol. 16, No. 2, April 2008, pp. 132–143. Reason for exclusion: Only measures effects of a co-intervention.

- Bigelow, G., C. Lawrence, M. Stitzer, and D. Wells, "Behavioral Treatments During Outpatient Methadone Maintenance: A Controlled Evaluation," paper presented at the 84th Annual Convention of the American Psychological Association, Washington, D.C., September 3–7, 1976, p. 11. Reason for exclusion: Only measures effects of a co-intervention.
- Bigelow, G., M. Stitzer, and C. Lawrence, "Narcotics Addiction Treatment: Behavioral Methods Concurrent with Methadone Maintenance," *The International Journal of the Addictions*, Vol. 15, No. 3, 1980. Reason for exclusion: Only measures effects of a co-intervention.
- Binder, Philippe, Nassir Messaadi, Marie-Christine Perault-Pochat, Stéphanie Gagey, Yann Brabant, and Pierre Ingrand, "Preference for Brand-Name Buprenorphine Is Related to Severity of Addiction Among Outpatients in Opioid Maintenance Treatment," *Journal of Addictive Diseases*, Vol. 35, No. 2, 2016, pp. 101–108. Reason for exclusion: No functional outcomes.
- Binder, T., and B. Vavrinkova, "Prospective Randomised Comparative Study of the Effect of Buprenorphine, Methadone and Heroin on the Course of Pregnancy, Birthweight of Newborns, Early Postpartum Adaptation and Course of the Neonatal Abstinence Syndrome (NAS) in Women Followed Up in the Outpatient Department," *Neuro Endocrinology Letters*, Vol. 29, No. 1, February 2008, pp. 80–86. Reason for exclusion: Wrong population (i.e., women, adolescents).
- Bisaga, A., M. A. Sullivan, W. Y. Cheng, K. M. Carpenter, J. J. Mariani, F. R. Levin, W. N. Raby, and E. V. Nunes, "A Placebo Controlled Trial of Memantine as an Adjunct to Oral Naltrexone for Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 119, No. 1–2, December 1, 2011, pp. e23–29. Reason for exclusion: Only measures effects of a co-intervention.
- Bizzarri, J. V., V. Casetti, L. Sanna, A. G. I. Maremmanni, L. Rovai, S. Bacciardi, D. Piacentino, A. Conca, and I. Maremmanni, "The Newer Opioid Agonist Treatment with Lower Substitutive Opiate Doses Is Associated with Better Toxicology Outcome than the Older Harm Reduction Treatment," *Annals of General Psychiatry*, Vol. 15, No. 1, 2016. Reason for exclusion: No functional outcomes.
- Blanken, P., V. M. Hendriks, M. W. Koeter, J. M. van Ree, and W. van den Brink, "Craving and Illicit Heroin Use Among Patients in Heroin-Assisted Treatment," *Drug and Alcohol Dependence*, Vol. 120, No. 1–3, January 1, 2012, pp. 74–80. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bliesener, N., S. Albrecht, A. Schwager, K. Weckbecker, D. Lichtermann, and D. Klingmuller, "Plasma Testosterone and Sexual Function in Men Receiving Buprenorphine Maintenance for Opioid Dependence," *Journal of Clinical Endocrinology and Metabolism*, Vol. 90, No. 1, January 2005, pp. 203–206. Reason for exclusion: Wrong study design.

- Blokhina, E., E. Krupitsky, N. Bushara, T. Yaroslavl'tseva, A. Tiurina, V. Palatkin, E. Verbitskaya, E. Zvartau, and G. E. Woody, "Implantable and Oral Naltrexone for Preventing Relapse in Opiate Addicts: A Psychometric Evaluation," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- Blondell, R. D., L. Ashrafioun, C. M. Dambra, E. M. Foschio, A. L. Zielinski, and D. M. Salcedo, "A Clinical Trial Comparing Tapering Doses of Buprenorphine with Steady Doses for Chronic Pain and Co-Existent Opioid Addiction," *Journal of the American College of Cardiology*, Vol. 4, No. 3, 2010. Reason for exclusion: No functional outcomes.
- Blondell, R. D., L. M. Frydrych, B. C. Arber, H. L. Bashaw, A. Vexler, C. H. Purdy, R. M. Sawyer, and S. Okazaki, "A Randomized Trial of Extended Buprenorphine Detoxification for Opioid Dependency," *Journal of Addiction Medicine*, Vol. 2, No. 3, September 2008, pp. 139–146. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Blum, K., T. J. Chen, J. Bailey, A. Bowirrat, J. Femino, A. L. Chen, T. Simpatico, S. Morse, J. Giordano, U. Damle, M. Kerner, E. R. Braverman, F. Fornari, B. W. Downs, C. Rector, D. Barh, and M. Oscar-Berman, "Can the Chronic Administration of the Combination of Buprenorphine and Naloxone Block Dopaminergic Activity Causing Anti-Reward and Relapse Potential?" *Molecular Neurobiology*, Vol. 44, No. 3, December 2011, pp. 250–268. Reason for exclusion: Wrong study design.
- Blum, K., M. Oscar-Berman, J. Femino, R. L. Waite, L. Benya, J. Giordano, J. Borsten, W. B. Downs, E. R. Braverman, R. Loehmann, K. Dushaj, D. Han, T. Simpatico, M. Hauser, D. Barh, and T. McLaughlin, "Withdrawal from Buprenorphine/Naloxone and Maintenance with a Natural Dopaminergic Agonist: A Cautionary Note," *Journal of Addiction Research and Therapy*, Vol. 4, No. 2, April 23, 2013. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Boettiger, C. A., "Neurocognitive Effects of Naltrexone," *Neuropsychopharmacology*, Vol. 38, 2013. Reason for exclusion: Conference abstract.
- Boettiger, Charlotte A., Elizabeth A. Kelley, Jennifer M. Mitchell, Mark D'Esposito, and Howard L. Fields, "Now or Later? An fMRI Study of the Effects of Endogenous Opioid Blockade on a Decision-Making Network," *Pharmacology, Biochemistry, and Behavior*, Vol. 93, No. 3, September 2009, pp. 291–299. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Bogucka-Bonikowska, A., H. Baran-Furga, K. Chmielewska, B. Habrat, A. Scinska, A. Kukwa, E. Koros, W. Kostowski, E. Polanowska, and P. Bienkowski, "Taste Function in Methadone-Maintained Opioid-Dependent Men," *Drug and Alcohol Dependence*, Vol. 68, No. 1, September 1, 2002, pp. 113–117. Reason for exclusion: No functional outcomes.

- Booth, R. E., K. F. Corsi, and S. K. Mikulich-Gilbertson, "Factors Associated with Methadone Maintenance Treatment Retention Among Street-Recruited Injection Drug Users," *Drug and Alcohol Dependence*, Vol. 74, No. 2, May 10, 2004, pp. 177–185. Reason for exclusion: Only measures effects of a co-intervention.
- Bounes, V., A. Palmaro, M. Lapeyre-Mestre, and A. Roussin, "Long-Term Consequences of Acute Pain for Patients Under Methadone or Buprenorphine Maintenance Treatment," *Pain Physician*, Vol. 16, No. 6, 2013. Reason for exclusion: Wrong study design.
- Boyer, Jane L., "The Study of Cognitive Performance in Buprenorphine Treated Opioid Dependent Individuals: A Repeated Measurement Approach," *Dissertation Abstracts International: Section B: The Sciences and Engineering S2—Dissertation Abstracts International*, Vol. 70, No. 7-B, 2010, p. 4503. Reason for exclusion: Wrong study design.
- Bracken, B. K., G. H. Trksak, D. M. Penetar, W. L. Tartarini, M. A. Maywalt, C. M. Dorsey, and S. E. Lukas, "Response Inhibition and Psychomotor Speed During Methadone Maintenance: Impact of Treatment Duration, Dose, and Sleep Deprivation," *Drug and Alcohol Dependence*, Vol. 125, No. 1–2, September 1, 2012, pp. 132–139. Reason for exclusion: Wrong study design.
- Brady, T. M., S. Salvucci, L. S. Sverdlov, A. Male, H. Kyeyune, E. Sikali, S. DeSale, and P. Yu, "Methadone Dosage and Retention: An Examination of the 60 Mg/Day Threshold," *Journal of Addictive Diseases*, Vol. 24, No. 3, 2005, pp. 23–47. Reason for exclusion: Wrong study design.
- Brahen, L. S., T. Capone, R. C. Heller, S. L. Linden, H. J. Landy, and M. J. Lewis, "Controlled Clinical Study of Naltrexone Side Effects Comparing First-Day Doses and Maintenance Regimens," *The American Journal of Drug and Alcohol Abuse*, Vol. 5, No. 2, 1978, pp. 235–245. Reason for exclusion: Wrong setting (inpatient).
- Brahen, L. S., T. Capone, V. Wiechert, and D. Desiderio, "Naltrexone and Cyclazocine. A Controlled Treatment Study," *Archives of General Psychiatry*, Vol. 34, No. 10, October 1977, pp. 1181–1184. Reason for exclusion: No functional outcomes.
- Branstetter, S. A., E. H. Bower, J. Kamien, and L. Amass, "A History of Sexual, Emotional, or Physical Abuse Predicts Adjustment During Opioid Maintenance Treatment," *Journal of Substance Abuse Treatment*, Vol. 34, No. 2, March 2008, pp. 208–214. Reason for exclusion: Wrong study design.
- Bravo, M. J., N. Llorens, G. Barrio, M. T. Brugal, S. Santos, L. Sordo, and L. de la Fuente, "Methadone Maintenance Treatment: A Protective Factor for Cocaine Injection in a Street-Recruited Cohort of Heroin Users," *Drug and Alcohol Dependence*, Vol. 112, No. 1–2, November 1, 2010, pp. 62–68. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Breen, C. L., S. J. Harris, N. Lintzeris, R. P. Mattick, L. Hawken, J. Bell, A. J. Ritter, M. Lenne, and E. Mendoza, "Cessation of Methadone Maintenance Treatment Using Buprenorphine: Transfer from Methadone to Buprenorphine and Subsequent Buprenorphine Reductions," *Drug and Alcohol Dependence*, Vol. 71, No. 1, July 20, 2003, pp. 49–55. Reason for exclusion: No functional outcomes.
- Brewer, C., "Response to Degenhardt et al.: 'Depot Naltrexone Use for Opioid Dependence in Australia: Large-Scale Use of an Unregistered Medication in the Absence of Data on Safety and Efficacy,'" *Drug and Alcohol Review*, Vol. 27, No. 4, July 2008, pp. 447–448; author reply pp. 448–449. Reason for exclusion: Wrong study design.
- Brewer, C., "A 'Miracle Cure' Misunderstood," *Addiction*, Vol. 107, No. 8, August 2012, p. 1528. Reason for exclusion: Wrong study design.
- "Briefly Noted," *Alcoholism and Drug Abuse Weekly*, Vol. 19, No. 34, 2007, pp. 7–8. Reason for exclusion: Wrong study design.
- "BRIEFLY NOTED," *Alcoholism and Drug Abuse Weekly*, Vol. 24, No. 6, 2012, pp. 7–8.
Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Brink, W., "Heroin Assisted Treatment," *BMJ (Online)*, Vol. 339, No. 7734, 2009. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Brink, W., V. M. Hendriks, P. Blanken, and J. M. Ree, "Beneficial Effect of Heroin in Heroin Addicts," *European Neuropsychopharmacology*; 15th International Congress of the European College of Neuropsychopharmacology, October 5–9, Barcelona, Spain, Vol. 12, Supplement 3, 2002. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Brooks, Adam C., Sandra D. Comer, Maria A. Sullivan, Adam Bisaga, Kenneth M. Carpenter, Wilfrid M. Raby, Elmer Yu, Charles P. O'Brien, and Edward V. Nunes, "Long-Acting Injectable Versus Oral Naltrexone Maintenance Therapy with Psychosocial Intervention for Heroin Dependence: A Quasi-Experiment," *The Journal of Clinical Psychiatry*, Vol. 71, No. 10, October 2010, pp. 1371–1378. Reason for exclusion: Only measures effects of a co-intervention.
- Brooks, D. J., S. K. Vosburg, S. M. Evans, and F. R. Levin, "Assessment of Cognitive Functioning of Methadone-Maintenance Patients: Impact of Adult ADHD and Current Cocaine Dependence," *Journal of Addictive Diseases*, Vol. 25, No. 4, 2006, pp. 15–25. Reason for exclusion: Wrong study design.
- Brooner, R. K., M. S. Kidorf, V. L. King, J. Peirce, K. Neufeld, K. Stoller, and K. Kolodner, "Managing Psychiatric Comorbidity Within Versus Outside of Methadone Treatment Settings: A Randomized and Controlled Evaluation," *Addiction*, Vol. 108, No. 11, November 2013, pp. 1942–1951. Reason for exclusion: Only measures effects of a co-intervention.

- Brown, E. Sherwood, Carlos Tirado, Abu Minhajuddin, Maureen Hillhouse, Bryon Adinoff, Walter Ling, Geetha Doraimani, and Christie Thomas, "Association of Race and Ethnicity with Withdrawal Symptoms, Attrition, Opioid Use, and Side-Effects During Buprenorphine Therapy," *Journal of Ethnicity in Substance Abuse*, Vol. 9, No. 2, April 2010, pp. 106–114. Reason for exclusion: Wrong study design.
- Brown, R., M. Gassman, S. Hetzel, and L. Berger, "Community-Based Treatment for Opioid Dependent Offenders: A Pilot Study," *The American Journal on Addictions*, Vol. 22, No. 5, September–October 2013, pp. 500–502. Reason for exclusion: Only measures effects of a co-intervention.
- Bruce, R. D., "The Marketing of Methadone: How an Effective Medication Became Unpopular," *International Journal of Drug Policy*, Vol. 24, No. 6, November 2013, pp. e89–90. Reason for exclusion: Wrong study design.
- Brugal, M. T., A. Domingo-Salvany, R. Puig, G. Barrio, P. Garcia de Olalla, and L. de la Fuente, "Evaluating the Impact of Methadone Maintenance Programmes on Mortality Due to Overdose and Aids in a Cohort of Heroin Users in Spain," *Addiction*, Vol. 100, No. 7, July 2005, pp. 981–989. Reason for exclusion: Wrong study design.
- "Buprenorphine Implant Helps Opioid-Dependent Patients Stay Abstinent Longer," *Clinical Pharmacist*, Vol. 8, No. 8, 2016. Reason for exclusion: Wrong study design.
- "Buprenorphine + Naloxone: New Combination. Opiate Dependence: No Proof of Reduced Risk of Self-Administered Injection," *Prescrire International*, Vol. 16, No. 92, December 2007, pp. 232–235. Reason for exclusion: Wrong study design.
- "Buprenorphine Works, but Only If Patients Stay on It: Study," *Alcoholism and Drug Abuse Weekly*, Vol. 23, No. 44, 2011, pp. 4–5. Reason for exclusion: Wrong study design.
- Burns, L., D. Randall, W. D. Hall, M. Law, T. Butler, J. Bell, and L. Degenhardt, "Opioid Agonist Pharmacotherapy in New South Wales from 1985 to 2006: Patient Characteristics and Patterns and Predictors of Treatment Retention," *Addiction*, Vol. 104, No. 8, August 2009, pp. 1363–1372. Reason for exclusion: Wrong study design.
- Busch, S., K. Hawk, D. Fiellin, P. O'Connor, M. Chawarski, P. Owens, M. Pantalon, S. Bernstein, and G. D'Onofrio, "Health Service Use in a Randomized Clinical Trial Comparing Three Methods of Emergency Department Interventions for Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Byford, S., B. Barrett, N. Metrebian, T. Groshkova, M. Cary, V. Charles, N. Lintzeris, and J. Strang, "Cost-Effectiveness of Injectable Opioid Treatment v. Oral Methadone for Chronic Heroin Addiction," *The British Journal of Psychiatry*, Vol. 203, No. 5, November 2013, pp. 341–349. Reason for exclusion: Article reporting follow-up of included studies.

- Cagnacci, A., A. M. Paoletti, R. Soldani, F. Tuveri, and G. B. Melis, "Prolonged Opioid Blockade Does Not Influence Luteinizing Hormone Modifications of the Follicular and Luteal Menstrual Phases," *The Journal of Clinical Endocrinology and Metabolism*, Vol. 80, No. 3, March 1995, pp. 860–863. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Caldiero, R. M., T. V. Parran, Jr., C. L. Adelman, and B. Piche, "Inpatient Initiation of Buprenorphine Maintenance vs. Detoxification: Can Retention of Opioid-Dependent Patients in Outpatient Counseling Be Improved?" *The American Journal on Addictions*, Vol. 15, No. 1, January–February 2006, pp. 1–7. Reason for exclusion: Wrong study design.
- Callahan, E. J., R. A. Rawson, and B. McCleave, "The Treatment of Heroin Addiction: Naltrexone Alone and with Behavior Therapy," *The International Journal of the Addictions*, Vol. 15, No. 6, 1980. Reason for exclusion: Only measures effects of a co-intervention.
- Calsyn, D. A., E. A. Wells, A. J. Saxon, T. R. Jackson, A. F. Wrede, V. Stanton, and C. Fleming, "Contingency Management of Urinalysis Results and Intensity of Counseling Services Have an Interactive Impact on Methadone Maintenance Treatment Outcome," *Journal of Addictive Diseases*, Vol. 13, No. 3, 1994, pp. 47–63. Reason for exclusion: Only measures effects of a co-intervention.
- Cameron, I. M., C. I. Matheson, C. M. Bond, P. McNamee, T. Lawrie, A. Robinson, G. Robertson, and L. E. Eagles, "Pilot Randomised Controlled Trial of Community Pharmacy Administration of Buprenorphine Versus Methadone," *International Journal of Pharmacy Practice*, Vol. 14, No. 4, 2006. Reason for exclusion: No functional outcomes.
- Cami, J., D. Guerra, B. Ugena, J. Segura, and R. de la Torre, "Double-Blind Assessment of Buprenorphine Withdrawal in Opiate-Addicts," *NIDA Research Monograph*, Vol. 105, 1990, p. 345. Reason for exclusion: Conference abstract.
- Cami, J., X. Lamas, and M. Farre, "Acute Effects of Tramadol in Methadone-Maintained Volunteers," *Drugs*, Vol. 47, Supplement 1, 1994, pp. 39–43. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Caplehorn, J., "Comments on Mattick et al.: The Need For Independent Data Re-Analyses . . . Mattick RP, Ali R, White JM et al (2003) Buprenorphine Versus Methadone Maintenance Therapy: A Randomized Double-Blind Trial with 405 Opioid-Dependent Patients. *Addiction*, 98. 441–52," *Addiction*, Vol. 98, No. 11, 2003, pp. 1633–1636. Reason for exclusion: Wrong study design.
- Caplehorn, J. R., "A Comparison of Abstinence-Oriented and Indefinite Methadone Maintenance Treatment," *The International Journal of the Addictions*, Vol. 29, No. 11, September 1994, pp. 1361–1375. Reason for exclusion: Wrong study design.
- Caplehorn, J. R. M., M. S. Y. N. Dalton, F. Haldar, A. M. Petrenas, and J. G. Nisbet, "Methadone Maintenance and Addicts' Risk of Fatal Heroin Overdose," *Substance Use and Misuse*, Vol. 31, No. 2, 1996, pp. 177–196. Reason for exclusion: Wrong study design.

- Caplehorn, J. R. M., and M. W. Ross, "Methadone Maintenance and the Likelihood of Risky Needle-Sharing," *The International Journal of the Addictions*, Vol. 30, No. 6, 1995, pp. 685–698. Reason for exclusion: Wrong study design.
- Caputo, F., G. Addolorato, M. Domenicali, A. Mosti, M. Viaggi, F. Trevisani, G. Gasbarrini, M. Bernardi, and G. F. Stefanini, "Short-Term Methadone Administration Reduces Alcohol Consumption in Non-Alcoholic Heroin Addicts," *Alcohol and Alcoholism*, Vol. 37, No. 2, 2002. Reason for exclusion: No functional outcomes.
- Carrera, I., L. Sanchez, E. Sabater, C. Pereiro, G. Florez, M. Conde, C. Pino, M. Serrano, and M. A. Casado, "Study on Users' Perception of Agonist Opioid Treatment in the Galician Network of Drug Addiction," *Heroin Addiction and Related Clinical Problems*, Vol. 18, No. 3, 2016. Reason for exclusion: No functional outcomes.
- Carrieri, P. M., L. Michel, C. Lions, J. Cohen, M. Vray, M. Mora, F. Marcellin, B. Spire, A. Morel, and P. Roux, "Methadone Induction in Primary Care for Opioid Dependence: A Pragmatic Randomized Trial (ANRS Methaville)," *PLoS One*, Vol. 9, No. 11, 2014, p. e112328. Reason for exclusion: Wrong study design.
- Carroll, K. M., S. A. Ball, C. Nich, P. G. O'Connor, D. A. Egan, T. L. Frankforter, E. G. Triffleman, J. Shi, and B. J. Rounsaville, "Targeting Behavioral Therapies to Enhance Naltrexone Treatment of Opioid Dependence: Efficacy of Contingency Management and Significant Other Involvement," *Archives of General Psychiatry*, Vol. 58, No. 8, August 2001, pp. 755–761. Reason for exclusion: Only measures effects of a co-intervention.
- Carroll, K. M., R. Sinha, C. Nich, T. Babuscio, and B. J. Rounsaville, "Contingency Management to Enhance Naltrexone Treatment of Opioid Dependence: A Randomized Clinical Trial of Reinforcement Magnitude," *Experimental and Clinical Psychopharmacology*, Vol. 10, No. 1, February 2002, pp. 54–63. Reason for exclusion: Only measures effects of a co-intervention.
- Chakrabarti, A., G. E. Woody, M. L. Griffin, G. Subramaniam, and R. D. Weiss, "Predictors of Buprenorphine-Naloxone Dosing in a 12-Week Treatment Trial for Opioid-Dependent Youth: Secondary Analyses from a NIDA Clinical Trials Network Study," *Drug and Alcohol Dependence*, Vol. 107, No. 2–3, March 1, 2010, pp. 253–256. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Chan, Y. Y., Y. H. Chen, S. N. Yang, W. Y. Lo, and J. G. Lin, "Clinical Efficacy of Traditional Chinese Medicine, Suan Zao Ren Tang, for Sleep Disturbance During Methadone Maintenance: A Randomized, Double-Blind, Placebo-Controlled Trial," *Evidence Based Complementary and Alternative Medicine*, Vol. 710895, 2015. Reason for exclusion: Only measures effects of a co-intervention.
- Chan, Y. Y., S. N. Yang, J. C. Lin, J. L. Chang, J. G. Lin, and W. Y. Lo, "Inflammatory Response in Heroin Addicts Undergoing Methadone Maintenance Treatment," *Psychiatry Research*, Vol. 226, No. 1, March 30, 2015, pp. 230–234. Reason for exclusion: No functional outcomes.

- Chang, K. C., and C. Y. Lin, "Effects of Publicly Funded and Quality of Life on Attendance Rate Among Methadone Maintenance Treatment Patients in Taiwan: An 18-Month Follow-Up Study," *Harm Reduction Journal*, Vol. 12, October 16, 2015, p. 40. Reason for exclusion: Only measures effects of a co-intervention.
- Chang, L. R., Y. H. Lin, T. B. Kuo, Y. C. Ho, S. H. Chen, H. C. Wu Chang, C. M. Liu, and C. C. Yang, "Cardiac Autonomic Modulation During Methadone Therapy Among Heroin Users: A Pilot Study," *Progress in Neuro-Psychopharmacol Biological Psychiatry*, Vol. 37, No. 1, April 27, 2012, pp. 188–193. Reason for exclusion: Wrong study design.
- Chang, Y. H., S. L. Chen, S. Y. Lee, P. S. Chen, T. Y. Wang, I. H. Lee, K. C. Chen, Y. K. Yang, J. S. Hong, and R. B. Lu, "Low-Dose Add-On Memantine Treatment May Improve Cognitive Performance and Self-Reported Health Conditions in Opioid-Dependent Patients Undergoing Methadone-Maintenance-Therapy," *Scientific Reports*, Vol. 5, May 19, 2015, p. 9708. Reason for exclusion: Only measures effects of a co-intervention.
- Charles, M., C. E. Wellington, C. Mokrysz, T. P. Freeman, D. O’Ryan, and H. V. Curran, "Attentional Bias and Treatment Adherence in Substitute-Prescribed Opiate Users," *Addictive Behaviors*, Vol. 46, July 2015, pp. 100–105. Reason for exclusion: Wrong study design.
- Charney, D. S., D. E. Redmond, Jr., M. P. Galloway, H. D. Kleber, G. R. Heninger, M. Murberg, and R. H. Roth, "Naltrexone Precipitated Opiate Withdrawal in Methadone Addicted Human Subjects: Evidence for Noradrenergic Hyperactivity," *Life Sciences*, Vol. 35, No. 12, September 17, 1984, pp. 1263–1272. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Charuvastra, V. C., J. Panell, W. Ling, and J. Ouren, "Follow-Up Study of Subjects on Methadyl Acetate and Methadone," *Drug and Alcohol Dependence*, Vol. 6, No. 5, November 1980, pp. 273–284. Reason for exclusion: Wrong study design.
- Chatham, L. R., K. Knight, G. W. Joe, and D. D. Simpson, "Suicidality in a Sample of Methadone Maintenance Clients," *American Journal of Drug and Alcohol Abuse*, Vol. 21, No. 3, August 1995, pp. 345–361. Reason for exclusion: No functional outcomes.
- Chawarski, M. C., M. Mazlan, and R. S. Schottenfeld, "Behavioral Drug and HIV Risk Reduction Counseling (BDRC) with Abstinence-Contingent Take-Home Buprenorphine: A Pilot Randomized Clinical Trial," *Drug and Alcohol Dependence*, Vol. 94, No. 1–3, April 1, 2008, pp. 281–284. Reason for exclusion: Only measures effects of a co-intervention.
- Chawarski, M. C., D. E. Moody, J. Pakes, P. G. O’Connor, and R. S. Schottenfeld, "Buprenorphine Tablet Versus Liquid: A Clinical Trial Comparing Plasma Levels, Efficacy, and Symptoms," *The Journal of Substance Abuse Treatment*, Vol. 29, No. 4, December 2005, pp. 307–312. Reason for exclusion: No functional outcomes.
- Chawarski, M. C., R. S. Schottenfeld, P. G. O’Connor, and J. Pakes, "Plasma Concentrations of Buprenorphine 24 to 72 Hours After Dosing," *Drug and Alcohol Dependence*, Vol. 55, No. 1–2, June 1, 1999, pp. 157–163. Reason for exclusion: No functional outcomes.

- Chen, S. L., S. Y. Lee, P. L. Tao, Y. H. Chang, S. H. Chen, C. H. Chu, P. S. Chen, I. H. Lee, T. L. Yeh, Y. K. Yang, J. S. Hong, and R. B. Lu, "Dextromethorphan Attenuated Inflammation and Combined Opioid Use in Humans Undergoing Methadone Maintenance Treatment," *Journal of Neuroimmune Pharmacology*, Vol. 7, No. 4, December 2012, pp. 1025–1033. Reason for exclusion: Only measures effects of a co-intervention.
- Chen, V. C., M. H. Wu, T. Y. Lin, Y. F. Ho, H. Y. Wang, and M. Gossop, "Comparison of Socio-Demographic Characteristics, Substance, and Depression Among Male Heroin Users Attending Therapeutic Community and Methadone Maintenance Treatment Program in Nantou, Taiwan," *Substance Abuse Treatment, Prevention, and Policy*, Vol. 10, October 27, 2015, p. 41. Reason for exclusion: Wrong study design.
- Chen, W., Y. Hong, X. Zou, M. M. McLaughlin, Y. Xia, and L. Ling, "Effectiveness of Prize-Based Contingency Management in a Methadone Maintenance Program in China," *Drug and Alcohol Dependence*, Vol. 133, No. 1, November 1, 2013, pp. 270–274. Reason for exclusion: Only measures effects of a co-intervention.
- Chen, Y. Z., W. L. Huang, J. C. Shan, Y. H. Lin, H. C. Chang, and L. R. Chang, "Self-Reported Psychopathology and Health-Related Quality of Life in Heroin Users Treated with Methadone," *Neuropsychiatric Disease and Treatment*, Vol. 9, 2013, pp. 41–48. Reason for exclusion: Wrong study design.
- Cho, Elizabeth, Helen Dermatis, and Marc Galanter, "Editorial: Buprenorphine for Heroin Addicts: The Issue of Illicit Opioid Abuse During Maintenance," *Substance Abuse*, Vol. 27, No. 1–2, 2006, pp. 1–3. Reason for exclusion: Wrong study design.
- Chopra, M. P., R. D. Landes, K. M. Gatchalian, L. C. Jackson, A. R. Buchhalter, M. L. Stitzer, L. A. Marsch, and W. K. Bickel, "Buprenorphine Medication Versus Voucher Contingencies in Promoting Abstinence from Opioids and Cocaine," *Experimental and Clinical Psychopharmacology*, Vol. 17, No. 4, August 2009, pp. 226–236. Reason for exclusion: Only measures effects of a co-intervention.
- Chutuape, M. A., K. Silverman, and M. L. Stitzer, "Use of Methadone Take-Home Contingencies with Persistent Opiate and Cocaine Abusers," *Journal of Substance Abuse Treatment*, Vol. 16, No. 1, January 1999, pp. 23–30. Reason for exclusion: Only measures effects of a co-intervention.
- Chutuape, M. A., K. Silverman, and M. L. Stitzer, "Effects of Urine Testing Frequency on Outcome in a Methadone Take-Home Contingency Program," *Drug and Alcohol Dependence*, Vol. 62, No. 1, March 1, 2001, pp. 69–76. Reason for exclusion: Only measures effects of a co-intervention.
- Cicero, T. J., and J. A. Inciardi, "Diversion and Abuse of Methadone Prescribed for Pain Management," *JAMA*, Vol. 293, No. 3, January 19, 2005, pp. 297–298. Reason for exclusion: Wrong study design.

- Cicero, T. J., H. L. Surratt, and J. Inciardi, "Use and Misuse of Buprenorphine in the Management of Opioid Addiction," *Journal of Opioid Management*, Vol. 3, No. 6, November–December 2007, pp. 302–308. Reason for exclusion: Wrong study design.
- Cioe, P. A., B. J. Anderson, and M. D. Stein, "Change in Symptoms of Erectile Dysfunction in Depressed Men Initiating Buprenorphine Therapy," *Journal of Substance Abuse Treatment*, Vol. 45, No. 5, November–December 2013, pp. 451–456. Reason for exclusion: Only measures effects of a co-intervention.
- Clair, J., L. Martin, A. J. Bond, D. O’Ryan, P. Davis, and H. V. Curran, "An Experimental Study of Aggressive and Neutral Interpretative Bias in Opiate-Dependent and Opiate-Abstinent Men," *Journal of Psychopharmacology*, Vol. 23, No. 4, June 2009, pp. 428–435. Reason for exclusion: Wrong study design.
- Clark, H. W., C. L. Masson, K. L. Delucchi, S. M. Hall, and K. L. Sees, "Violent Traumatic Events and Drug Abuse Severity," *Journal of Substance Abuse Treatment*, Vol. 20, No. 2, March 2001, pp. 121–127. Reason for exclusion: Wrong study design.
- Clark, N. C., P. Dietze, M. G. Lenne, and J. R. Redman, "Effect of Opioid Substitution Therapy on Alcohol Metabolism," *Journal of Substance Abuse Treatment*, Vol. 30, No. 3, April 2006, pp. 191–196. Reason for exclusion: No functional outcomes.
- Clarke, T. K., R. C. Crist, A. Ang, L. M. Ambrose-Lanci, F. W. Lohoff, A. J. Saxon, W. Ling, M. P. Hillhouse, R. D. Bruce, G. Woody, and W. H. Berrettini, "Genetic Variation in OPRD1 and the Response to Treatment for Opioid Dependence with Buprenorphine in European-American Females," *Pharmacogenomics Journal*, Vol. 14, No. 3, 2014. Reason for exclusion: No functional outcomes.
- "Clinical Evaluation of Naltrexone Treatment of Opiate-Dependent Individuals. Report of the National Research Council Committee on Clinical Evaluation of Narcotic Antagonists," *Archives of General Psychiatry*, Vol. 35, No. 3, March 1978, pp. 335–340. Reason for exclusion: No functional outcomes.
- Cobb, M. N., J. Desai, L. S. Brown, Jr., P. N. Zannikos, and P. M. Rainey, "The Effect of Fluconazole on the Clinical Pharmacokinetics of Methadone," *Clinical Pharmacology and Therapeutics*, Vol. 63, No. 6, June 1998, pp. 655–662. Reason for exclusion: Only measures effects of a co-intervention.
- Cohen, J., "Law Enforcement and Drug Treatment: A Culture Clash," *Science*, Vol. 329, No. 5988, July 9, 2010, p. 169. Reason for exclusion: Wrong study design.
- Cohen, J., "No Opiate Substitutes for the Masses of IDUs," *Science*, Vol. 329, No. 5988, July 9, 2010, pp. 165–167. Reason for exclusion: Wrong study design.
- Cohen, L. J., E. Gertmenian-King, L. Kunik, C. Weaver, E. D. London, and I. Galynker, "Personality Measures in Former Heroin Users Receiving Methadone or in Protracted Abstinence from Opiates," *Acta Psychiatrica Scandinavica*, Vol. 112, No. 2, August 2005, pp. 149–158. Reason for exclusion: Wrong study design.

- Cohen, M., D. Korts, R. Hanbury, V. Sturiano, G. Jackson, and B. Stimmel, "The Effect of Alcoholism in Methadone-Maintained Persons on Productive Activity: A Randomized Control Trial," *Alcoholism: Clinical and Experimental Research*, Vol. 6, No. 3, Summer 1982, pp. 358–361. Reason for exclusion: Wrong study design.
- Colameco, S., and M. Pohl, "Buprenorphine in the Workers' Compensation Setting," *Journal of Opioid Management*, Vol. 10, No. 4, July–August 2014, pp. 277–283. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Coller, J. K., D. T. Barratt, K. Dahlen, M. H. Loennechen, and A. A. Somogyi, "ABCB1 Genetic Variability and Methadone Dosage Requirements in Opioid-Dependent Individuals," *Clinical Pharmacology and Therapeutics*, Vol. 80, No. 6, December 2006, pp. 682–690. Reason for exclusion: No functional outcomes.
- Collins, D., "A Shift in Focus," *Canadian Nurse*, Vol. 104, No. 6, June 2008, p. 6. Reason for exclusion: Wrong study design.
- Colom Farran, J., M. Casas, J. Perez de Los Cobos, M. Del Rio, C. Roncero, X. Castells, S. Valero, F. J. Eiroa-Orosa, F. Batlle, and J. Trujols, "Feasibility of Double-Blind Clinical Trials with Oral Diacetylmorphine: A Randomized Controlled Phase II Study in an Inpatient Setting," *European Addiction Research*, Vol. 18, No. 6, 2012, pp. 279–287. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Colquhoun, R., D. Y. Tan, and S. Hull, "A Comparison of Oral and Implant Naltrexone Outcomes at 12 Months," *Journal of Opioid Management*, Vol. 1, No. 5, 2005, 2005, pp. 249–256. Reason for exclusion: No functional outcomes.
- Comer, S. D., and E. D. Collins, "Self-Administration of Intravenous Buprenorphine and the Buprenorphine/Naloxone Combination by Recently Detoxified Heroin Abusers," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 303, No. 2, 2002. Reason for exclusion: Wrong setting (i.e., inpatient).
- Comer, S. D., E. D. Collins, H. D. Kleber, E. S. Nuwayser, J. H. Kerrigan, and M. W. Fischman, "Depot Naltrexone: Long-Lasting Antagonism of the Effects of Heroin in Humans," *Psychopharmacology*, Vol. 159, No. 4, 2002. Reason for exclusion: Wrong setting (i.e., inpatient).
- Comer, S. D., M. A. Sullivan, and E. A. Walker, "Comparison of Intravenous Buprenorphine and Methadone Self-Administration by Recently Detoxified Heroin-Dependent Individuals," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 315, No. 3, 2005. Reason for exclusion: Wrong setting (i.e., inpatient).
- Comer, S. D., E. A. Walker, and E. D. Collins, "Buprenorphine/Naloxone Reduces the Reinforcing and Subjective Effects of Heroin in Heroin-Dependent Volunteers," *Psychopharmacology*, Vol. 181, No. 4, 2005. Reason for exclusion: Only measures effects of a co-intervention.

- Comer, S. D., E. D. Collins, and M. W. Fischman, "Buprenorphine Sublingual Tablets: Effects on IV Heroin Self-Administration by Humans," *Psychopharmacology (Berl)*, Vol. 154, No. 1, February 2001, pp. 28–37. Reason for exclusion: No functional outcomes.
- Comer, Sandra D., Eric D. Collins, and Marian W. Fischman, "Intravenous Buprenorphine Self-Administration by Detoxified Heroin Abusers," *The Journal of Pharmacology and Experimental Therapeutics*, Vol. 301, No. 1, April 2002, pp. 266–276. Reason for exclusion: Wrong study design.
- Comer, S. D., V. E. Metz, Z. D. Cooper, W. J. Kowalczyk, J. D. Jones, M. A. Sullivan, J. M. Manubay, S. K. Vosburg, M. E. Smith, D. Peyser, and P. A. Saccone, "Comparison of a Drug Versus Money and Drug Versus Drug Self-Administration Choice Procedure with Oxycodone and Morphine in Opioid Addicts," *Behavioural Pharmacology*, Vol. 24, No. 5–6, September 2013, pp. 504–516. Reason for exclusion: Only measures effects of a co-intervention.
- Comer, S. D., M. A. Sullivan, E. Yu, J. L. Rothenberg, H. D. Kleber, K. Kampman, C. Dackis, and C. P. O'Brien, "Injectable, Sustained-Release Naltrexone for the Treatment of Opioid Dependence: A Randomized, Placebo-Controlled Trial," *Archives of General Psychiatry*, Vol. 63, No. 2, February 2006, pp. 210–218. Reason for exclusion: No functional outcomes.
- Comiskey, C. M., and G. Cox, "Analysis of the Impact of Treatment Setting on Outcomes from Methadone Treatment," *Journal of Substance Abuse Treatment*, Vol. 39, No. 3, October 2010, pp. 195–201. Reason for exclusion: Only measures effects of a co-intervention.
- "Community-Based Opioid Overdose Prevention Programs Providing Naloxone—United States, 2010," *Morbidity and Mortality Weekly Report*, Vol. 61, No. 6, February 17, 2012, pp. 101–105. Reason for exclusion: Wrong study design.
- Compton, P., V. C. Charuvastra, K. Kintaudi, and W. Ling, "Pain Responses in Methadone-Maintained Opioid Abusers," *Journal of Pain and Symptom Management*, Vol. 20, No. 4, 2000. Reason for exclusion: Wrong study design.
- Compton, P., V. C. Charuvastra, and W. Ling, "Pain Intolerance in Opioid-Maintained Former Opiate Addicts: Effect of Long-Acting Maintenance Agent," *Drug and Alcohol Dependence*, Vol. 63, No. 2, July 1, 2001, pp. 139–146. Reason for exclusion: Wrong study design.
- Compton, P., D. H. Geschwind, and M. Alarcon, "Association Between Human Mu-Opioid Receptor Gene Polymorphism, Pain Tolerance, and Opioid Addiction," *American Journal of Medical Genetics B: Neuropsychiatric Genetics*, Vol. 121b, No. 1, August 15, 2003, pp. 76–82. Reason for exclusion: Wrong study design.
- Compton, P., W. Ling, D. Moody, and N. Chiang, "Pharmacokinetics, Bioavailability and Opioid Effects of Liquid Versus Tablet Buprenorphine," *Drug and Alcohol Dependence*, Vol. 82, No. 1, March 15, 2006, pp. 25–31. Reason for exclusion: Wrong setting (inpatient).

- Compton, P. A., D. R. Wesson, V. C. Charuvastra, and W. Ling, "Buprenorphine as a Pharmacotherapy for Opiate Addiction: What Dose Provides a Therapeutic Response?" *The American Journal on Addictions*, Vol. 5, No. 3, 1996. Reason for exclusion: No functional outcomes.
- "Connecticut Addiction Counselors Train CPS Workforce on MAT . . . Medication-Assisted Treatment," *Alcoholism and Drug Abuse Weekly*, Vol. 27, No. 41, 2015, pp. 1–3. Reason for exclusion: Wrong study design.
- Constantinou, N., C. J. Morgan, S. Battistella, D. O’Ryan, P. Davis, and H. V. Curran, "Attentional Bias, Inhibitory Control and Acute Stress in Current and Former Opiate Addicts," *Drug and Alcohol Dependence*, Vol. 109, No. 1–3, June 1, 2010, pp. 220–225. Reason for exclusion: Wrong study design.
- "Controversy Boiling Over as SAMHSA Considers Raising Bupe Cap," *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 28, 2014, pp. 4–6. Reason for exclusion: Wrong study design.
- Cook, S., D. Indig, J. Gray, and D. McGrath, "Opiate Overdose and Health Treatment Options for Opiate Users in New South Wales, 1999–2002," *NSW Public Health Bulletin*, Vol. 15, No. 7–8, July–August 2004, pp. 125–131. Reason for exclusion: Wrong study design.
- Cornish, J., C. D. Nordeck, D. D. Langleben, D. Coviello, K. Lynch, T. Boney, and C. P. O’Brien, "Six-Month Depot Naltrexone Treatment Reduces Relapse in Parolees Formerly Addicted to Opioids," *Neuropsychopharmacology*, Vol. 38, 2012. Reason for exclusion: Conference abstract.
- Cornish, J. W., B. H. Herman, R. N. Ehrman, S. J. Robbins, A. R. Childress, V. Bead, C. A. Esmonde, K. Martz, S. Poole, F. S. Caruso, and C. P. O’Brien, "A Randomized, Double-Blind, Placebo-Controlled Safety Study of High-Dose Dextromethorphan in Methadone-Maintained Male Inpatients," *Drug and Alcohol Dependence*, Vol. 67, No. 2, July 1, 2002, pp. 177–183. Reason for exclusion: Only measures effects of a co-intervention.
- Correia, C. J., S. L. Walsh, G. E. Bigelow, and E. C. Strain, "Effects Associated with Double-Blind Omission of Buprenorphine/Naloxone over a 98-H Period," *Psychopharmacology (Berl)*, Vol. 189, No. 3, December 2006, pp. 297–306. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Cosentino, B. W., "New Hope for Heroin Addicts," *Nursing Spectrum—New York and New Jersey Edition*, Vol. 14A, No. 2, 2002, p. 4p. Reason for exclusion: Wrong study design.
- Cottrill, C. B., and S. C. Matson, "Medication-Assisted Treatment of Opioid Use Disorder in Adolescents and Young Adults," *Adolescent Medicine: State of the Art Reviews*, Vol. 25, No. 2, August 2014, pp. 251–265. Reason for exclusion: Wrong population (pregnant women, adolescents).
- Counselman, Francis L., "Malpractice Counsel: Child Abuse, Methadone Overdose. Methadone Overdose: Intentional or Not," *Emergency Medicine (00136654)*, Vol. 46, No. 11, November 2014, pp. 506–507. Reason for exclusion: Wrong study design.

- Courtney, K. E., D. G. Ghahremani, and L. A. Ray, "The Effects of Pharmacological Opioid Blockade on Neural Measures of Drug Cue-Reactivity in Humans," *Neuropsychopharmacology*, Vol. 41, No. 12, November 2016, pp. 2872–2881. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Courtwright, D. T., "Preventing and Treating Narcotic Addiction—Century of Federal Drug Control," *New England Journal of Medicine*, Vol. 373, No. 22, November 26, 2015, pp. 2095–2097. Reason for exclusion: Wrong study design.
- Cousins, S. J., L. Denering, D. Crevecœur-MacPhail, J. Viernes, W. Sugita, J. Barger, T. Kim, S. Weimann, and R. A. Rawson, "A Demonstration Project Implementing Extended-Release Naltrexone in Los Angeles County," *Substance Abuse*, Vol. 37, No. 1, 2016, pp. 54–62. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Coviello, D. M., D. A. Zanis, S. A. Wesnoski, and S. W. Domis, "An Integrated Drug Counseling and Employment Intervention for Methadone Clients," *Journal of Psychoactive Drugs*, Vol. 41, No. 2, June 2009, pp. 189–197. Reason for exclusion: Only measures effects of a co-intervention.
- Cowan, A., "Buprenorphine: New Pharmacological Aspects," *International Journal of Clinical Practice Supplement*, No. 133, February 2003, pp. 3–8; discussion 23–24. Reason for exclusion: Wrong study design.
- Coyle, M. G., A. L. Salisbury, B. M. Lester, H. E. Jones, H. Lin, K. Graf-Rohrmeister, and G. Fischer, "Neonatal Neurobehavior Effects Following Buprenorphine Versus Methadone Exposure," *Addiction*, Vol. 107, Supplement 1, November 2012, pp. 63–73. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Crandall, C., R. D. Crosby, and G. A. Carlson, "Does Pregnancy Affect Outcome of Methadone Maintenance Treatment?" *Journal of Substance Abuse Treatment*, Vol. 26, No. 4, June 2004, pp. 295–303. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Crettol, S., J. Besson, M. Croquette-Krokar, R. Hämmig, I. Gothuey, M. Monnat, J. J. Déglon, M. Preisig, and C. B. Eap, "Association of Dopamine and Opioid Receptor Genetic Polymorphisms with Response to Methadone Maintenance Treatment," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, Vol. 32, No. 7, 2008. Reason for exclusion: No functional outcomes.
- Crist, R. C., T. K. Clarke, A. Ang, L. M. Ambrose-Lanci, F. W. Lohoff, A. J. Saxon, W. Ling, M. P. Hillhouse, R. D. Bruce, G. Woody, and W. H. Berrettini, "An Intronic Variant in OPRD1 Predicts Treatment Outcome for Opioid Dependence in African-Americans," *Neuropsychopharmacology*, Vol. 38, No. 10, September 2013, pp. 2003–2010. Reason for exclusion: No functional outcomes.

- Cropsey, K. L., P. S. Lane, G. J. Hale, D. O. Jackson, C. B. Clark, K. S. Ingersoll, M. A. Islam, and M. L. Stitzer, "Results of a Pilot Randomized Controlled Trial of Buprenorphine for Opioid Dependent Women in the Criminal Justice System," *Drug and Alcohol Dependence*, Vol. 119, No. 3, December 15, 2011, pp. 172–178. Reason for exclusion: Wrong population (pregnant women, adolescents).
- Cruts, G., M. Buster, J. Vicente, I. Deerenberg, and M. Van Laar, "Estimating the Total Mortality Among Problem Drug Users," *Substance Use and Misuse*, Vol. 43, No. 5, 2008, pp. 733–747. Reason for exclusion: No functional outcomes.
- Cullen, W., S. O'Brien, A. O'Carroll, F. D. O'Kelly, and G. Bury, "Chronic Illness and Multimorbidity Among Problem Drug Users: A Comparative Cross Sectional Pilot Study in Primary Care," *BMC Family Practice*, Vol. 10, April 21, 2009, p. 25. Reason for exclusion: Wrong study design.
- Cunningham, C. O., A. Giovanniello, H. V. Kunins, R. J. Roose, A. D. Fox, and N. L. Sohler, "Buprenorphine Treatment Outcomes Among Opioid-Dependent Cocaine Users and Non-Users," *The American Journal on Addictions/American Academy of Psychiatrists in Alcoholism and Addictions*, Vol. 22, No. 4, 2013. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Curran, H. V., J. Bolton, S. Wanigaratne, and C. Smyth, "Additional Methadone Increases Craving for Heroin: A Double-Blind, Placebo-Controlled Study of Chronic Opiate Users Receiving Methadone Substitution Treatment," *Addiction*, Vol. 94, No. 5, May 1999, pp. 665–674. Reason for exclusion: Wrong study design.
- Cutter, C. J., B. A. Moore, D. Barry, L. E. Fiellin, R. S. Schottenfeld, D. Fiellin, and P. O'Connor, "Cognitive Behavioral Therapy Improves Treatment Outcome for Prescription Opioid Users in Primary Care Based Buprenorphine/Naloxone Treatment," *Drug and Alcohol Dependence*, Vol. 146, No. 1, 2015, p. e255. Reason for exclusion: Only measures effects of a co-intervention.
- Dakwar, E., and H. D. Kleber, "Naltrexone-Facilitated Buprenorphine Discontinuation: A Feasibility Trial," *Journal of Substance Abuse Treatment*, Vol. 53, 2015. Reason for exclusion: Wrong study design.
- Dale, Robert T., and Farley R. Dale, "The Use of Methadone in a Representative Group of Heroin Addicts," *The International Journal of the Addictions*, Vol. 8, No. 2, 1973, pp. 293–308. Reason for exclusion: Wrong study design.
- Dansereau, D. F., G. W. Joe, and D. D. Simpson, "Attentional Difficulties and the Effectiveness of a Visual Representation Strategy for Counseling Drug-Addicted Clients," *The International Journal of the Addictions*, Vol. 30, No. 4, March 1995, pp. 371–386. Reason for exclusion: Only measures effects of a co-intervention.

- Darke, S., S. McDonald, S. Kaye, and M. Torok, "Comparative Patterns of Cognitive Performance Amongst Opioid Maintenance Patients, Abstinent Opioid Users and Non-Opioid Users," *Drug and Alcohol Dependence*, Vol. 126, No. 3, December 1, 2012, pp. 309–315. Reason for exclusion: Wrong study design.
- Darke, S., J. Ross, M. Lynskey, and M. Teesson, "Attempted Suicide Among Entrants to Three Treatment Modalities for Heroin Dependence in the Australian Treatment Outcome Study (ATOS): Prevalence and Risk Factors," *Drug and Alcohol Dependence*, Vol. 73, No. 1, January 7, 2004, pp. 1–10. Reason for exclusion: Wrong study design.
- Darke, S., J. Sims, S. McDonald, and W. Wickes, "Cognitive Impairment Among Methadone Maintenance Patients," *Addiction*, Vol. 95, No. 5, May 2000, pp. 687–695. Reason for exclusion: Wrong study design.
- Dasgupta, N., E. J. Bailey, T. Cicero, J. Inciardi, M. Parrino, A. Rosenblum, and R. C. Dart, "Post-Marketing Surveillance of Methadone and Buprenorphine in the United States," *Pain Medicine*, Vol. 11, No. 7, 2010, pp. 1078–1091. Reason for exclusion: No functional outcomes.
- Dashe, J. S., J. S. Sheffield, and G. D. Wendel, Jr., "Improving the Management of Opioid-Dependent Pregnancies," *American Journal of Obstetrics and Gynecology*, Vol. 190, No. 6, June 2004, p. 1806; author reply 1806–1807. Reason for exclusion: Wrong population (pregnant women, adolescents).
- D'Aunno, T., N. Folz-Murphy, and X. Lin, "Changes in Methadone Treatment Practices: Results from a Panel Study, 1988–1995," *The American Journal of Drug and Alcohol Abuse*, Vol. 25, No. 4, November 1999, pp. 681–699. Reason for exclusion: Wrong study design.
- Davis, P. E., H. Liddiard, and T. M. McMillan, "Neuropsychological Deficits and Opiate Abuse," *Drug and Alcohol Dependence*, Vol. 67, No. 1, June 1, 2002, pp. 105–108. Reason for exclusion: Wrong study design.
- Davoli, M., A. M. Bargagli, C. A. Perucci, P. Schifano, V. Belleudi, M. Hickman, G. Salamina, R. Diecidue, F. Vigna-Taglianti, and F. Faggiano, "Risk of Fatal Overdose During and After Specialist Drug Treatment: The Vedette Study, a National Multi-Site Prospective Cohort Study," *Addiction*, Vol. 102, No. 12, December 2007, pp. 1954–1959. Reason for exclusion: No functional outcomes.
- Davoli, M., and C. A. Perucci, "Appropriateness of Methadone Maintenance Treatment for Opiate Addiction: Treatment Goals and Effectiveness," *Sozial-Und Praventivmedizin*, Vol. 48, Supplement 1, 2003, pp. S21–22; discussion S25–27. Reason for exclusion: Wrong study design.
- Dawe, S., and P. Harnett, "Reducing Potential for Child Abuse Among Methadone-Maintained Parents: Results from a Randomized Controlled Trial," *Journal of Substance Abuse Treatment*, Vol. 32, No. 4, June 2007, pp. 381–390. Reason for exclusion: Only measures effects of a co-intervention.

- Day, E., "Maximising the Treatment Outcomes of Opioid Substitution Treatment," *Evidence-Based Mental Health*, Vol. 13, No. 1, February 2010, pp. 5–7. Reason for exclusion: Wrong study design.
- Day, E., J. Ison, and J. Strang, "Inpatient Versus Other Settings for Detoxification for Opioid Dependence," *The Cochrane Database of Systematic Reviews*, No. 2, 2005, p. Cd004580. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- De Jong, C. A., H. G. Roozen, L. G. van Rossum, P. F. Krabbe, and A. J. Kerkhof, "High Abstinence Rates in Heroin Addicts by a New Comprehensive Treatment Approach," *American Journal of Addiction*, Vol. 16, No. 2, March–April 2007, pp. 124–130. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- De Leon, G., G. L. Staines, T. E. Perlis, S. Sacks, K. McKendrick, R. Hilton, and R. Brady, "Therapeutic Community Methods in Methadone Maintenance (Passages): An Open Clinical Trial," *Drug and Alcohol Dependence*, Vol. 37, No. 1, January 1995, pp. 45–57. Reason for exclusion: Only measures effects of a co-intervention.
- de los Cobos, José Pérez, Silvia Martin, Alvaro Etcheberrigaray, Joan Trujols, Fanny Batlle, Antoni Tejero, Josep M. Queraltó, and Miquel Casas, "A Controlled Trial of Daily Versus Thrice-Weekly Buprenorphine Administration for the Treatment of Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 59, No. 3, June 2000, pp. 223–233. Reason for exclusion: No functional outcomes.
- De, S., R. Jain, R. Ray, A. Dhawan, and S. T. Varghese, "Assessment of Differential Doses of Buprenorphine for Long Term Pharmacotherapy Among Opiate Dependent Subjects," *Indian Journal of Physiology and Pharmacology*, Vol. 52, No. 1, January–March 2008, pp. 53–63. Reason for exclusion: Wrong setting (inpatient).
- de Vos, J. W., J. G. Ufkes, C. D. Kaplan, M. Tursch, J. K. Krause, H. van Wilgenburg, B. G. Woodcock, and A. H. Staib, "L-Methadone and D,L-Methadone in Methadone Maintenance Treatment: A Comparison of Therapeutic Effectiveness and Plasma Concentrations," *European Addiction Research*, Vol. 4, No. 3, September 1998, pp. 134–141. Reason for exclusion: Only measures effects of a co-intervention.
- Dean, A. J., J. Bell, M. J. Christie, and R. P. Mattick, "Depressive Symptoms During Buprenorphine vs. Methadone Maintenance: Findings from a Randomised, Controlled Trial in Opioid Dependence," *European Psychiatry*, Vol. 19, No. 8, December 2004, pp. 510–513. Reason for exclusion: No functional outcomes.
- Dean, A. J., J. B. Saunders, R. T. Jones, R. M. Young, J. P. Connor, and B. R. Lawford, "Does Naltrexone Treatment Lead to Depression? Findings from a Randomized Controlled Trial in Subjects with Opioid Dependence," *Journal of Psychiatry Neuroscience*, Vol. 31, No. 1, January 2006, pp. 38–45. Reason for exclusion: No functional outcomes.

- Deck, D., W. Wiitala, B. McFarland, K. Campbell, J. Mullooly, A. Krupski, and D. McCarty, "Medicaid Coverage, Methadone Maintenance, and Felony Arrests: Outcomes of Opiate Treatment in Two States," *Journal of Addictive Diseases*, Vol. 28, No. 2, 2009, pp. 89–102. Reason for exclusion: Wrong study design.
- DeFulio, A., J. J. Everly, J. M. Leoutsakos, A. Umbricht, M. Fingerhood, G. E. Bigelow, and K. Silverman, "Employment-Based Reinforcement of Adherence to an FDA Approved Extended Release Formulation of Naltrexone in Opioid-Dependent Adults: A Randomized Controlled Trial," *Drug and Alcohol Dependence*, Vol. 120, No. 1–3, January 1, 2012, pp. 48–54. Reason for exclusion: Only measures effects of a co-intervention.
- Dehghani-Arani, F., R. Rostami, and H. Nadali, "Neurofeedback Training for Opiate Addiction: Improvement of Mental Health and Craving," *Applied Psychophysiology and Biofeedback*, Vol. 38, No. 2, June 2013, pp. 133–141. Reason for exclusion: Only measures effects of a co-intervention.
- Demaret, Isabelle, Etienne Quertemont, Géraldine Litran, Cécile Magoga, Clémence Deblire, Nathalie Dubois, Corinne Charlier, André Lemaitre, and Marc Ansseau, "Loss of Treatment Benefit When Heroin-Assisted Treatment Is Stopped After 12 Months," *Journal of Substance Abuse Treatment*, Vol. 69, October 2016, pp. 72–75. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Demarie, D., G. Marletta, M. Imazio, C. Cappa, S. Ferro, R. Compostino, E. De Vivo, R. Trincherio, and E. Bignamini, "Cardiovascular-Associated Disease in an Addicted Population: An Observation Study," *Journal of Cardiovascular Medicine (Hagerstown)*, Vol. 12, No. 1, January 2011, pp. 51–54. Reason for exclusion: No functional outcomes.
- Des Jarlais, D. C., T. M. Hammett, L. Wei, L. K. Van, M. Donghua, and D. Ngu, "Opiate Agonist Maintenance Treatment for Injecting Drug User Peer Educators," *Addiction*, Vol. 99, No. 10, October 2004, pp. 1355–1356. Reason for exclusion: Wrong study design.
- Desmond, D. P., and J. F. Maddux, "Compulsory Supervision and Methadone Maintenance," *Journal of Substance Abuse Treatment*, Vol. 13, No. 1, January–February 1996, pp. 79–83. Reason for exclusion: Only measures effects of a co-intervention.
- Desrosiers, A., E. Blokhina, E. Krupitsky, and M. C. Chawarski, "Gender Differences in Psychiatric Symptoms Among Opiate-Dependent Patients in Russia," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- deWit, H., "A New Role for Clonidine in Addictions: Catching Relapses Before They Happen," *American Journal of Psychiatry*, Vol. 172, No. 8, August 1, 2015, pp. 700–701. Reason for exclusion: Wrong study design.
- Dhingra, L., C. Masson, D. C. Perlman, R. M. Seewald, J. Katz, C. McKnight, P. Homel, E. Wald, A. E. Jordan, C. Young, and R. K. Portenoy, "Epidemiology of Pain Among Outpatients in Methadone Maintenance Treatment Programs," *Drug and Alcohol Dependence*, Vol. 128, No. 1–2, February 1, 2013, pp. 161–165. Reason for exclusion: Only measures effects of a co-intervention.

- Di Paola, A., T. Lincoln, D. J. Skiest, M. Desabrais, F. L. Altice, and S. A. Springer, “Design and Methods of a Double Blind Randomized Placebo-Controlled Trial of Extended-Release Naltrexone for HIV-Infected, Opioid Dependent Prisoners and Jail Detainees Who Are Transitioning to the Community,” *Contemporary Clinical Trials*, Vol. 39, No. 2, November 2014, pp. 256–268. Reason for exclusion: Wrong study design.
- Diaz-Flores, J. F., R. I. Sanudo, E. M. Rodriguez, and C. D. Romero, “Serum Concentrations of Macro and Trace Elements in Heroin Addicts of the Canary Islands,” *Journal of Trace Elements in Medicine and Biology*, Vol. 17, No. 4, 2004, pp. 235–242. Reason for exclusion: No functional outcomes.
- Dick, Andrew W., Rosalie L. Pacula, Adam J. Gordon, Mark Sorbero, Rachei M. Burns, Douglas Leslie, and Bradley D. Stein, “Access to Care. Growth in Buprenorphine Waivers for Physicians Increased Potential Access to Opioid Agonist Treatment, 2002–11,” *Health Affairs*, Vol. 34, No. 6, 2015, pp. 1028–1034. Reason for exclusion: Wrong study design.
- Dijkgraaf, M. G., B. P. van der Zanden, C. A. de Borgie, P. Blanken, J. M. van Ree, and W. van den Brink, “Cost Utility Analysis of Co-Prescribed Heroin Compared with Methadone Maintenance Treatment in Heroin Addicts in Two Randomised Trials,” *BMJ*, Vol. 330, No. 7503, June 4, 2005, p. 1297. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Dolan, K. A., J. Shearer, M. MacDonald, R. P. Mattick, W. Hall, and A. D. Wodak, “A Randomised Controlled Trial of Methadone Maintenance Treatment Versus Wait List Control in an Australian Prison System,” *Drug and Alcohol Dependence*, Vol. 72, No. 1, October 24, 2003, pp. 59–65. Reason for exclusion: Wrong setting (i.e., inpatient).
- Dolan, K. A., J. Shearer, B. White, J. Zhou, J. Kaldor, and A. D. Wodak, “Four-Year Follow-Up of Imprisoned Male Heroin Users and Methadone Treatment: Mortality, Re-Incarceration and Hepatitis C Infection,” *Addiction*, Vol. 100, No. 6, June 2005, pp. 820–828. Reason for exclusion: Wrong study design.
- Dolgin, E., “Technology: Barriers to Misuse,” *Nature*, Vol. 522, No. 7557, June 25, 2015, pp. S60–61. Reason for exclusion: Wrong study design.
- Dong, Z. L., S. F. Wang, and L. Zhou, “A Clinical Study on the Efficacy of Methadone Treatment Supplemented by Transdermal Application of Fentanyl in Detoxifying Patients with Severe Heroin Abuse,” *Chinese Journal of Drug Abuse Prevention and Treatment*, Vol. 9, No. 5, 2003. Reason for exclusion: Only measures effects of a co-intervention.
- Donny, E. C., M. L. Stitzer, G. E. Bigelow, and S. L. Walsh, “Methadone Maintenance Inpatient Opiate Abusers Suppresses Withdrawal, but Fails to Completely Block Heroin’s Effects,” *Drug and Alcohol Dependence*, Vol. 36, Supplement 1, 2001. Reason for exclusion: Conference abstract.

- Donny, E. C., S. L. Walsh, G. E. Bigelow, T. Eissenberg, and M. L. Stitzer, "High-Dose Methadone Produces Superior Opioid Blockade and Comparable Withdrawal Suppression to Lower Doses in Opioid-Dependent Humans," *Psychopharmacology*, Vol. 161, No. 2, 2002. Reason for exclusion: Wrong study design.
- D'Onofrio, G., P. O'Connor, M. Pantalon, M. Chawarski, S. Busch, P. Owens, S. Bernstein, and D. Fiellin, "A Randomized Clinical Trial of Emergency Department Initiated Treatment for Opioid Dependence: Two and Six Month Outcomes," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- D'Onofrio, G., P. G. O'Connor, M. V. Pantalon, M. C. Chawarski, S. H. Busch, P. H. Owens, S. L. Bernstein, and D. A. Fiellin, "Emergency Department-Initiated Buprenorphine/Naloxone Treatment for Opioid Dependence: A Randomized Clinical Trial," *JAMA*, Vol. 313, No. 16, April 28, 2015, pp. 1636–1644. Reason for exclusion: No functional outcomes.
- Doran, C. M., "Buprenorphine, Buprenorphine/Naloxone and Methadone Maintenance: A Cost-Effectiveness Analysis," *Expert Review of Pharmacoeconomics and Outcomes Research*, Vol. 5, No. 5, October 2005, pp. 583–591. Reason for exclusion: No functional outcomes.
- Doran, C. M., M. Shanahan, R. P. Mattick, R. Ali, J. White, and J. Bell, "Buprenorphine Versus Methadone Maintenance: A Cost-Effectiveness Analysis," *Drug and Alcohol Dependence*, Vol. 71, No. 3, September 10, 2003, pp. 295–302. Reason for exclusion: No functional outcomes.
- "Douglas' Case Hoped to Bring Methadone Treatment Behind Bars," *Alcoholism and Drug Abuse Weekly*, Vol. 24, No. 21, 2012, pp. 1–3. Reason for exclusion: Wrong study design.
- Downey, K. K., T. C. Helmus, and C. R. Schuster, "Treatment of Heroin-Dependent Poly-Drug Abusers with Contingency Management and Buprenorphine Maintenance," *Experimental and Clinical Psychopharmacology*, Vol. 8, No. 2, 2000. Reason for exclusion: Only measures effects of a co-intervention.
- Downey, K. K., J. A. Hopper, M. Henderson, E. Corby, T. Helmus, and C. R. Schuster, "Treatment of Heroin Dependent Poly-Drug Abusers with Contingent Voucher and Buprenorphine," *NIDA Research Monograph*, 1999. Reason for exclusion: Only measures effects of a co-intervention.
- Drummond, Colin, Christos Kouimtsidis, Martina Reynolds, Ian Russell, Christine Godfrey, Monica McCusker, Simon Coulton, Steve Parrott, Paul Davis, Nick Tarrier, Douglas Turkington, Louise Sell, John Merrill, Hugh Williams, Mohammed Abou-Saleh, Hamid Ghodse, and Sally Porter, "The Effectiveness and Cost Effectiveness of Cognitive Behaviour Therapy for Opiate Misusers in Methadone Maintenance Treatment: A Multicentre, Randomised, Controlled Trial UKCBTMM Study: United Kingdom Cognitive Behaviour Therapy Study in Methadone Maintenance Treatment," *Drugs: Education, Prevention, and Policy*, Vol. 12, Supplement 1, December 2005, pp. 69–76. Reason for exclusion: Only measures effects of a co-intervention.

- Ducharme, L. J., H. K. Knudsen, and P. M. Roman, "Evidence-Based Treatment for Opiate-Dependent Clients: Availability, Variation, and Organizational Correlates," *The American Journal of Drug and Alcohol Abuse*, Vol. 32, No. 4, 2006, pp. 569–576. Reason for exclusion: No functional outcomes.
- Duke, A. N., C. J. Correia, S. L. Walsh, G. E. Bigelow, and E. C. Strain, "Acute Effects of Intramuscular and Sublingual Buprenorphine and Buprenorphine/Naloxone in Non-Dependent Opioid Abusers," *Psychopharmacology*, Vol. 211, No. 3, 2010. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Dunn, K., A. DeFulio, J. J. Everly, W. D. Donlin, W. M. Aklin, P. A. Nuzzo, J. M. S. Leoutsakos, A. Umbricht, M. Fingerhood, G. E. Bigelow, and K. Silverman, "Employment-Based Reinforcement of Adherence to Oral Naltrexone in Unemployed Injection Drug Users: 12-Month Outcomes," *Psychology of Addictive Behaviors*, Vol. 18, 2014. Reason for exclusion: Only measures effects of a co-intervention.
- Dunn, K. E., A. Defulio, J. J. Everly, W. D. Donlin, W. M. Aklin, P. A. Nuzzo, J. M. Leoutsakos, A. Umbricht, M. Fingerhood, G. E. Bigelow, and K. Silverman, "Employment-Based Reinforcement of Adherence to Oral Naltrexone Treatment in Unemployed Injection Drug Users," *Experimental and Clinical Psychopharmacology*, Vol. 21, No. 1, February 2013, pp. 74–83. Reason for exclusion: Only measures effects of a co-intervention.
- Dvir, Y., and P. Smallwood, "Serotonin Syndrome: A Complex but Easily Avoidable Condition," *General Hospital Psychiatry*, Vol. 30, No. 3, May–June 2008, pp. 284–287. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Dyer, K. R., J. M. White, D. J. Foster, F. Bochner, A. Menelaou, and A. A. Somogyi, "The Relationship Between Mood State and Plasma Methadone Concentration in Maintenance Patients," *Journal of Clinical Psychopharmacology*, Vol. 21, No. 1, February 2001, pp. 78–84. Reason for exclusion: Wrong study design.
- Dyer, O., "Doctors Should Do More for Drug Addicts, Committee Says," *BMJ*, Vol. 324, No. 7349, June 1, 2002, p. 1295. Reason for exclusion: Wrong study design.
- Dzialdowski, A., and M. London, "A Cognitive Behavioural Intervention in the Context of Methadone Tapering Treatment for Opiate Addiction—Two Single Cases," *Clinical Psychology and Psychotherapy*, Vol. 6, No. 4, 1999. Reason for exclusion: Only measures effects of a co-intervention.
- Edelman, E. J., and L. E. Fiellin, "Opioid Substitution Therapy Is Associated with Decreased HIV Transmission Among People Who Inject Drugs," *Evidence-Based Medicine*, Vol. 18, No. 5, October 2013, pp. 177–178. Reason for exclusion: Wrong study design.
- Eder, H., G. Fischer, W. Gombas, R. Jagsch, G. Stühlinger, and S. Kasper, "Comparison of Buprenorphine and Methadone Maintenance in Opiate Addicts," *European Addiction Research*, Vol. 4, Supplement 1, 1998. Reason for exclusion: No functional outcomes.

- Eder, H., R. Jagsch, D. Kraigher, A. Primorac, N. Ebner, and G. Fischer, "Comparative Study of the Effectiveness of Slow-Release Morphine and Methadone for Opioid Maintenance Therapy," *Addiction*, Vol. 100, No. 8, August 2005, pp. 1101–1109. Reason for exclusion: Wrong study design.
- Edwards, G., "Seeing America—Diary of a Drug-Focused Study Tour Made in 1967," *Addiction*, Vol. 105, No. 6, June 2010, pp. 984–990. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Eghbali, H., M. Zare, A. Bakhtiari, N. Monirpoor, and A. Ganjali, "The Effectiveness of Matrix Interventions in Improving Methadone Treatment," *International Journal of High Risk Behaviors and Addiction*, Vol. 1, No. 4, Winter 2013, pp. 159–165. Reason for exclusion: Only measures effects of a co-intervention.
- Ehret, G. B., C. Voide, M. Gex-Fabry, J. Chabert, D. Shah, B. Broers, V. Piguet, T. Musset, J. M. Gaspoz, A. Perrier, P. Dayer, and J. A. Desmeules, "Drug-Induced Long QT Syndrome in Injection Drug Users Receiving Methadone: High Frequency in Hospitalized Patients and Risk Factors," *Archives of Internal Medicine*, Vol. 166, No. 12, June 26, 2006, pp. 1280–1287. Reason for exclusion: No functional outcomes.
- Eissenberg, T., R. E. Johnson, G. E. Bigelow, S. L. Walsh, I. A. Liebson, E. C. Strain, and M. L. Stitzer, "Controlled Opioid Withdrawal Evaluation During 72 H Dose Omission in Buprenorphine-Maintained Patients," *Drug and Alcohol Dependence*, Vol. 45, No. 1–2, April 14, 1997, pp. 81–91. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Eissenberg, T., M. L. Stitzer, G. E. Bigelow, A. R. Buchhalter, and S. L. Walsh, "Relative Potency of Levo-Alpha-Acetylmethadol and Methadone in Humans Under Acute Dosing Conditions," *The Journal of Pharmacology and Experimental Therapeutics*, Vol. 289, No. 2, 1999. Reason for exclusion: Wrong setting (inpatient).
- Elkader, A., and B. Sproule, "Buprenorphine: Clinical Pharmacokinetics in the Treatment of Opioid Dependence," *Clinical Pharmacokinetics*, Vol. 44, No. 7, 2005, pp. 661–680. Reason for exclusion: Wrong study design.
- Elkader, A. K., B. Brands, P. Selby, and B. A. Sproule, "Methadone-Nicotine Interactions in Methadone Maintenance Treatment Patients," *Journal of Clinical Psychopharmacology*, Vol. 29, No. 3, June 2009, pp. 231–238. Reason for exclusion: No functional outcomes.
- Elmoghazy, E., B. D. Johnson, and F. A. Alling, "A Pilot Study of a Neuro-Stimulator Device vs. Methadone in Alleviating Opiate Withdrawal Symptoms," *NIDA Research Monographs*, Vol. 95, 1989, pp. 388–389. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- "Emergency Department Visits Involving Nonmedical Use of Selected Prescription Drugs—United States, 2004–2008," *Morbidity and Mortality Weekly Report*, Vol. 59, No. 23, June 18, 2010, pp. 705–709. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Epstein, D. H., K. L. Preston, and J. Schmittner, "114 Cocaine and Heroin Abusers, 226 Palmpilots: Initial Experiences with Ecological Momentary Assessment at a Methadone Clinic," Proceedings of the 69th Annual Scientific Meeting of the College on Problems of Drug Dependence; Quebec City, Canada, June 16–21, 2007. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Epstein, D. H., J. Schmittner, A. Umbricht, J. R. Schroeder, E. T. Moolchan, and K. L. Preston, "Promoting Abstinence from Cocaine and Heroin with a Methadone Dose Increase and a Novel Contingency," *Drug and Alcohol Dependence*, Vol. 101, No. 1–2, April 1, 2009, pp. 92–100. Reason for exclusion: Only measures effects of a co-intervention.
- Ersche, K. D., P. C. Fletcher, J. P. Roiser, T. D. Fryer, M. London, T. W. Robbins, and B. J. Sahakian, "Differences in Orbitofrontal Activation During Decision-Making Between Methadone-Maintained Opiate Users, Heroin Users and Healthy Volunteers," *Psychopharmacology (Berl)*, Vol. 188, No. 3, October 2006, pp. 364–373. Reason for exclusion: Wrong study design.
- Ersche, K. D., J. P. Roiser, L. Clark, M. London, T. W. Robbins, and B. J. Sahakian, "Punishment Induces Risky Decision-Making in Methadone-Maintained Opiate Users but Not in Heroin Users or Healthy Volunteers," *Neuropsychopharmacology*, Vol. 30, No. 11, November 2005, pp. 2115–2124. Reason for exclusion: Wrong study design.
- Escher, M., Y. Daali, J. Chabert, G. Hopfgartner, P. Dayer, and J. Desmeules, "Pharmacokinetic and Pharmacodynamic Properties of Buprenorphine After a Single Intravenous Administration in Healthy Volunteers: A Randomized, Double-Blind, Placebo-Controlled, Crossover Study," *Clinical Therapeutics*, Vol. 29, No. 8, August 2007, pp. 1620–1631. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Evans, E., A. Kelleghan, L. Li, J. Min, D. Huang, D. Urada, Y. I. Hser, and B. Nosyk, "Gender Differences in Mortality Among Treated Opioid Dependent Patients," *Drug Alcohol Depend*, Vol. 155, October 1, 2015, pp. 228–235. Reason for exclusion: No functional outcomes.
- Evans, M., P. J. Robson, M. A. Chadd, C. M. Evans, and D. M. Fry, "Administration of Meptazinol to Opiate-Dependent Patients," *Postgraduate Medical Journal*, Vol. 59, Supplement 1, 1983, pp. 78–84. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Everly, J. J., A. DeFulio, M. N. Koffarnus, J. M. Leoutsakos, W. D. Donlin, W. M. Aklin, A. Umbricht, M. Fingerhood, G. E. Bigelow, and K. Silverman, "Employment-Based Reinforcement of Adherence to Depot Naltrexone in Unemployed Opioid-Dependent Adults: A Randomized Controlled Trial," *Addiction*, Vol. 106, No. 7, July 2011, pp. 1309–1318. Reason for exclusion: Only measures effects of a co-intervention.

- Everly, Jeffrey J., Anthony DeFulio, Mikhail N. Koffarnus, Jeannie-Marie S. Leoutsakos, Wendy D. Donlin, Will M. Aklin, Annie Umbricht, Michael Fingerhood, George E. Bigelow, and Kenneth Silverman, "Employment-Based Reinforcement of Adherence to Depot Naltrexone in Unemployed Opioid-Dependent Adults: A Randomized Controlled Trial," *Addiction*, Vol. 106, No. 7, July 2011, pp. 1309–1318. Reason for exclusion: Only measures effects of a co-intervention.
- Fadardi, J. S., and S. S. Ziaee, "A Comparative Study of Drug-Related Attentional Bias: Evidence from Iran," *Experimental and Clinical Psychopharmacology*, Vol. 18, No. 6, December 2010, pp. 539–545. Reason for exclusion: Wrong study design.
- Falcato, L. M., T. Beck, J. Reimer, and U. Verthein, "Self-Reported Cravings for Heroin and Cocaine During Maintenance Treatment with Slow-Release Oral Morphine Compared to Methadone: A Randomized Crossover Clinical Trial," *Sucht*, Vol. 60, 2014. Reason for exclusion: Wrong study design.
- Falcato, L., T. Beck, J. Reimer, and U. Verthein, "Self-Reported Cravings for Heroin and Cocaine During Maintenance Treatment with Slow-Release Oral Morphine Compared with Methadone: A Randomized, Crossover Clinical Trial," *Journal of Clinical Psychopharmacology*, Vol. 35, No. 2, April 2015, pp. 150–157. Reason for exclusion: No functional outcomes.
- Fals-Stewart, W., T. J. O'Farrell, and G. R. Birchler, "Behavioral Couples Therapy for Male Methadone Maintenance Patients: Effects on Drug-Using Behavior and Relationship Adjustment," *Behavior Therapy*, Vol. 32, No. 2, 2001. Reason for exclusion: Only measures effects of a co-intervention.
- Fals-Stewart, W., and T. J. O'Farrell, "Behavioral Family Counseling and Naltrexone for Male Opioid-Dependent Patients," *Journal of Consulting and Clinical Psychology*, Vol. 71, No. 3, June 2003, pp. 432–442. Reason for exclusion: Only measures effects of a co-intervention.
- Fan, C. Y., H. K. Tan, I. C. Chien, and S. Y. Chou, "Prevalence of Psychiatric Disorders Among Heroin Users Who Received Methadone Maintenance Therapy in Taiwan," *The American Journal on Addictions*, Vol. 23, No. 3, May–June 2014, pp. 249–256. Reason for exclusion: Wrong study design.
- Fareed, Ayman, Jennifer Casarella, Richard Amar, Sreedevi Vayalapalli, and Karen Drexler, "Methadone Maintenance Dosing Guideline for Opioid Dependence, a Literature Review," *Journal of Addictive Diseases*, Vol. 29, No. 1, January 2010, pp. 1–14. Reason for exclusion: No functional outcomes.
- Fareed, A., J. Casarella, M. Roberts, M. Sleboda, R. Amar, S. Vayalapalli, and K. Drexler, "High Dose Versus Moderate Dose Methadone Maintenance: Is There a Better Outcome?" *Journal of Addictive Diseases*, Vol. 28, No. 4, October 2009, pp. 399–405. Reason for exclusion: No functional outcomes.

- Fareed, A., S. Vayalapalli, S. Stout, J. Casarella, K. Drexler, and S. P. Bailey, "Effect of Methadone Maintenance Treatment on Heroin Craving, a Literature Review," *Journal of Addictive Diseases*, Vol. 30, No. 1, January 2011, pp. 27–38. Reason for exclusion: No functional outcomes.
- Farrell, M., and W. Hall, "The Swiss Heroin Trials: Testing Alternative Approaches," *BMJ*, Vol. 316, No. 7132, February 28, 1998, p. 639. Reason for exclusion: Wrong study design.
- Farrell, M., J. Ward, R. Mattick, W. Hall, G. V. Stimson, D. des Jarlais, M. Gossop, and J. Strang, "Methadone Maintenance Treatment in Opiate Dependence: A Review," *BMJ*, Vol. 309, No. 6960, October 15, 1994, pp. 997–1001. Reason for exclusion: Wrong study design.
- Farren, C. K., and S. O'Malley, "A Pilot Double Blind Placebo Controlled Trial of Sertraline with Naltrexone in the Treatment of Opiate Dependence," *The American Journal on Addictions*, Vol. 11, No. 3, Summer 2002, pp. 228–234. Reason for exclusion: Only measures effects of a co-intervention.
- Fatseas, M., C. Denis, Z. Massida, M. Verger, P. Franques-Rénéric, and M. Auriacombe, "Cue-Induced Reactivity, Cortisol Response and Substance Use Outcome in Treated Heroin Dependent Individuals," *Biological Psychiatry*, Vol. 70, No. 8, 2011. Reason for exclusion: No functional outcomes.
- Fellows-Smith, J., "Opioid-Dependent Error Processing," *Journal of Opioid Management*, Vol. 7, No. 6, November–December 2011, pp. 443–449. Reason for exclusion: No functional outcomes.
- Ferner, R. E., and A. M. Daniels, "Office-Based Treatment of Opioid-Dependent Patients," *New England Journal of Medicine*, Vol. 348, No. 1, January 2, 2003, pp. 81–82; author reply 81–82. Reason for exclusion: Wrong study design.
- Ferrara, M. M., "The Disparate Treatment of Medications and Opiate Pain Medications Under The Law: Permitting the Proliferation of Opiates and Limiting Access to Treatment," *Seton Hall Law Review*, Vol. 42, No. 2, 2012, pp. 741–767. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Ferri, M., A. J. Finlayson, L. Wang, and P. R. Martin, "Predictive Factors for Relapse in Patients on Buprenorphine Maintenance," *The American Journal on Addictions*, Vol. 23, No. 1, January–February 2014, pp. 62–67. Reason for exclusion: Wrong study design.
- Fiellin, D., C. J. Cutter, B. A. Moore, D. Barry, P. O'Connor, and R. S. Schottenfeld, "Primary Care Buprenorphine Detoxification vs. Maintenance for Prescription Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- Fiellin, D. A., "Methadone Syrup Injection in Australia: A Sentinel Finding?" *Addiction*, Vol. 98, No. 4, April 2003, pp. 385–386. Reason for exclusion: Wrong study design.

- Fiellin, D. A., D. T. Barry, L. E. Sullivan, C. J. Cutter, B. A. Moore, P. G. O'Connor, and R. S. Schottenfeld, "A Randomized Trial of Cognitive Behavioral Therapy in Primary Care-Based Buprenorphine," *The American Journal of Medicine*, Vol. 126, No. 1, January 2013, pp. 74.e11–74.e77. Reason for exclusion: Only measures effects of a co-intervention.
- Fiellin, D. A., and P. G. O'Connor, "Clinical Practice. Office-Based Treatment of Opioid-Dependent Patients," *New England Journal of Medicine*, Vol. 347, No. 11, September 12, 2002, pp. 817–823. Reason for exclusion: Wrong study design.
- Fiellin, D. A., P. G. O'Connor, M. Chawarski, J. P. Pakes, M. V. Pantalon, and R. S. Schottenfeld, "Methadone Maintenance in Primary Care: A Randomized Controlled Trial," *JAMA*, Vol. 286, No. 14, October 10, 2001, pp. 1724–1731. Reason for exclusion: Only measures effects of a co-intervention.
- Fiellin, D. A., M. V. Pantalon, M. C. Chawarski, B. A. Moore, L. E. Sullivan, P. G. O'Connor, and R. S. Schottenfeld, "Counseling Plus Buprenorphine-Naloxone Maintenance Therapy for Opioid Dependence," *New England Journal of Medicine*, Vol. 355, No. 4, July 27, 2006, pp. 365–374. Reason for exclusion: Only measures effects of a co-intervention.
- Fiellin, D. A., R. A. Rosenheck, and T. R. Kosten, "Office-Based Treatment for Opioid Dependence: Reaching New Patient Populations," *American Journal of Psychiatry*, Vol. 158, No. 8, August 2001, pp. 1200–1204. Reason for exclusion: Wrong study design.
- Fiellin, D. A., R. S. Schottenfeld, C. J. Cutter, B. A. Moore, D. T. Barry, and P. G. O'Connor, "Primary Care-Based Buprenorphine Taper vs Maintenance Therapy for Prescription Opioid Dependence: A Randomized Clinical Trial," *JAMA Internal Medicine*, Vol. 174, No. 12, December 2014, pp. 1947–1954. Reason for exclusion: No functional outcomes.
- Fingelkurts, A. A., A. A. Fingelkurts, R. Kivisaari, T. Autti, S. Borisov, V. Puuskari, O. Jokela, and S. Kahkonen, "Reorganization of The Composition of Brain Oscillations and Their Temporal Characteristics in Opioid Dependent Patients," *Progress in Neuro-Psychopharmacology and Biologica Psychiatry*, Vol. 30, No. 8, December 30, 2006, pp. 1453–1465. Reason for exclusion: No functional outcomes.
- Fingerhood, M. I., M. R. Thompson, and D. R. Jasinski, "A Comparison of Clonidine and Buprenorphine in the Outpatient Treatment of Opiate Withdrawal," *Substance Abuse*, Vol. 22, No. 3, 2001. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Fischer, B., A. T. Chin, I. Kuo, M. Kirst, and D. Vlahov, "Canadian Illicit Opiate Users' Views on Methadone and Other Opiate Prescription Treatment: An Exploratory Qualitative Study," *Substance Use and Misuse*, Vol. 37, No. 4, 2002, pp. 495–522. Reason for exclusion: Wrong study design.
- Fischer, G., W. Gombas, H. Eder, R. Jagsch, A. Peternell, G. Stuhlinger, L. Pezawas, H. N. Aschauer, and S. Kasper, "Buprenorphine Versus Methadone Maintenance for the Treatment of Opioid Dependence," *Addiction*, Vol. 94, No. 9, September 1999, pp. 1337–1347. Reason for exclusion: No functional outcomes.

- Fischer, G., R. Ortner, K. Rohrmeister, R. Jagsch, A. Baewert, M. Langer, and H. Aschauer, "Methadone Versus Buprenorphine in Pregnant Addicts: A Double-Blind, Double-Dummy Comparison Study," *Addiction*, Vol. 101, No. 2, February 2006, pp. 275–281. Reason for exclusion: Wrong population (pregnant women, adolescents).
- Fischer, G., B. Winklbaaur-Hausknost, R. Jagsch, K. Graf-Rohrmeister, A. Unger, A. Baewert, M. Langer, and K. Thau, "Lessons Learned from Acomparison of Evidence-Based Research in Pregnant Opioid-Dependent Women," *Alcohol and Alcoholism*, Vol. 49, Supplement 1, 2014, p. i31. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Fishman, M. J., L. T. Wu, and G. E. Woody, "Buprenorphine for Prescription Opioid Addiction in a Patient with Depression and Alcohol Dependence," *American Journal of Psychiatry*, Vol. 168, No. 7, July 2011, pp. 675–679. Reason for exclusion: Wrong study design.
- Fitzpatrick, M., "Drugs—Facing Facts," *British Journal of General Practice*, Vol. 57, No. 537, April 2007, p. 335. Reason for exclusion: Wrong study design.
- Flannery, W., and M. Farrell, "Harm Reduction the Key to Managing Problem Drug Users," *Practitioner*, Vol. 251, No. 1694, May 2007, pp. 99, 101–106. Reason for exclusion: Wrong study design.
- Flynn, P. M., G. W. Joe, K. M. Broome, D. D. Simpson, and B. S. Brown, "Recovery from Opioid Addiction in DATOS," *Journal of Substance Abuse Treatment*, Vol. 25, No. 3, October 2003, pp. 177–186. Reason for exclusion: Wrong study design.
- Foltin, R. W., and M. W. Fischman, "Interaction of Buprenorphine with Cocaine-Morphine Combinations," *Experimental and Clinical Psychopharmacology*, Vol. 3, No. 3, 1995. Reason for exclusion: No functional outcomes.
- Franckowiak, Bonnie A., and Doris F. Glick, "The Effect of Self-Efficacy on Treatment," *Journal of Addictions Nursing (Lippincott Williams and Wilkins)*, Vol. 26, No. 2, 2015, pp. 62–70. Reason for exclusion: Only measures effects of a co-intervention.
- Frankland, L., B. P. Bradley, and K. Mogg, "Time Course of Attentional Bias to Drug Cues in Opioid Dependence," *Psychology of Addictive Behaviors*, Vol. 30, No. 5, August 2016, pp. 601–606. Reason for exclusion: Wrong study design.
- Freedman, R. R., and G. Czertko, "A Comparison of Thrice Weekly LAAM and Daily Methadone in Employed Heroin Addicts," *Drug and Alcohol Dependence*, Vol. 8, No. 3, November 1981, pp. 215–222. Reason for exclusion: No functional outcomes.
- Friedmann, P. D., J. D. Lee, E. V. Nunes, T. W. Kinlock, and C. P. O'Brien, "Patient Selection for Extended-Release Naltrexone Among Criminal Justice-Involved Persons with Opioid Use Disorder," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Friedmann, P. D., J. C. Hendrickson, D. R. Gerstein, Z. Zhang, and M. D. Stein, "Do Mechanisms That Link Addiction Treatment Patients to Primary Care Influence Subsequent Utilization of Emergency and Hospital Care?" *Medical Care*, Vol. 44, No. 1, January 2006, pp. 8–15. Reason for exclusion: Only measures effects of a co-intervention.

- “From the Field . . . AATOD’s 5-Year Plan Includes Support for Buprenorphine, Risk Planning,” *Alcoholism and Drug Abuse Weekly*, Vol. 19, No. 15, 2007, p. 5. Reason for exclusion: Wrong study design.
- Fudala, P. J., J. H. Jaffe, E. M. Dax, and R. E. Johnson, “Use of Buprenorphine in the Treatment of Opioid Addiction. II. Physiologic and Behavioral Effects of Daily and Alternate-Day Administration and Abrupt Withdrawal,” *Clinical Pharmacology and Therapeutics*, Vol. 47, No. 4, 1990. Reason for exclusion: Wrong setting (i.e., inpatient).
- Fudala, P. J., E. Yu, W. Macfadden, C. Boardman, and C. N. Chiang, “Effects of Buprenorphine and Naloxone in Morphine-Stabilized Opioid Addicts,” *Drug and Alcohol Dependence*, Vol. 50, No. 1, March 1, 1998, pp. 1–8. Reason for exclusion: Wrong setting (i.e., inpatient).
- Fugelstad, A., M. Stenbacka, A. Leifman, M. Nylander, and I. Thiblin, “Methadone Maintenance Treatment: The Balance Between Life-Saving Treatment and Fatal Poisonings,” *Addiction*, Vol. 102, No. 3, March 2007, pp. 406–412. Reason for exclusion: Wrong study design.
- Gaalema, D. E., T. L. Scott, S. H. Heil, M. G. Coyle, K. Kaltenbach, G. J. Badger, A. M. Arria, S. M. Stine, P. R. Martin, and H. E. Jones, “Differences in the Profile of Neonatal Abstinence Syndrome Signs in Methadone-Versus Buprenorphine-Exposed Neonates,” *Addiction*, Vol. 107, Supplement 1, November 2012, pp. 53–62. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Gabbay, Mark Benjamin, Sinead Clarke, Emma Willert, and Aneez Esmail, “Shared Care Methadone Clinics: A Survey of Patient Satisfaction Behaviour Change and Staff Views,” *Addiction Research and Theory*, Vol. 7, No. 2, April 1999, pp. 129–147. Reason for exclusion: Only measures effects of a co-intervention.
- Gaertner, J., R. Voltz, and C. Ostgathe, “Methadone: A Closer Look at the Controversy,” *Journal of Pain and Symptom Management*, Vol. 36, No. 2, August 2008, pp. e4–e7. Reason for exclusion: Wrong study design.
- Galanter, M., H. Dermatis, L. Glickman, R. Maslansky, S. M. Brealyn, and C. Rahman-Dujarric, “Network Therapy and Buprenorphine Maintenance for the Treatment of Heroin Addiction,” *Proceedings of the 66th Annual Scientific Meeting of College on Problems of Drug Dependence; June 12–17, 2004; San Juan, Puerto Rico, 2004*. Reason for exclusion: Conference abstract.
- Galanter, M., H. Dermatis, L. Glickman, R. Maslansky, M. B. Sellers, E. Neumann, and C. Rahman-Dujarric, “Network Therapy: Decreased Secondary Opioid Use During Buprenorphine Maintenance,” *Journal of Substance Abuse Treatment*, Vol. 26, No. 4, June 2004, pp. 313–318. Reason for exclusion: Only measures effects of a co-intervention.
- Galanter, M., H. Dermatis, R. Resnick, R. Maslansky, and E. Neumann, “Short-Term Buprenorphine Maintenance: Treatment Outcome,” *Journal of Addictive Diseases*, Vol. 22, No. 3, 2003, pp. 39–49. Reason for exclusion: Wrong study design.

- Galynker, I. I., D. Eisenberg, J. A. Matochik, E. Gertmenian-King, L. Cohen, A. S. Kimes, C. Contoreggi, V. Kurian, M. Ernst, R. N. Rosenthal, J. Prosser, and E. D. London, "Cerebral Metabolism and Mood in Remitted Opiate Dependence," *Drug and Alcohol Dependence*, Vol. 90, No. 2–3, October 8, 2007, pp. 166–174. Reason for exclusion: No functional outcomes.
- Galynker, II, S. Watras-Ganz, C. Miner, R. N. Rosenthal, D. C. Des Jarlais, B. L. Richman, and E. London, "Cerebral Metabolism in Opiate-Dependent Subjects: Effects of Methadone Maintenance," *Mount Sinai Journal of Medicine*, Vol. 67, No. 5–6, October–November 2000, pp. 381–387. Reason for exclusion: No functional outcomes.
- Gastfriend, D., B. Silverman, A. Memisoglu, E. Krupitsky, and W. Ling, "Continuity of Clinical Efficacy with Injectable Extended-Release Naltrexone (XR-NTX)," *Journal of Addiction Medicine*, Vol. 7, No. 4, 2013. Reason for exclusion: Conference abstract.
- Gastfriend, D. R., "Intramuscular Extended-Release Naltrexone: Current Evidence," *Annals of the New York Academy of Sciences*, Vol. 1216, January 2011, pp. 144–166. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gaughwin, M., P. Solomon, and R. Ali, "Correlates of Retention on the South Australian Methadone Program 1981–91," *Australian and New Zealand Journal Public Health*, Vol. 22, No. 7, December 1998, pp. 771–776. Reason for exclusion: Wrong study design.
- Geng, Liuna, and Bojian Qian, "Implicit and Explicit Cognition of Chinese Heroin Abusers," *Social Behavior and Personality*, Vol. 39, No. 4, 2011, pp. 433–444. Reason for exclusion: Wrong study design.
- George, S., and H. Ekhtiari, "Naltrexone in the Treatment of Opioid Dependence," *British Journal of Hospital Medicine (London)*, Vol. 71, No. 10, October 2010, pp. 568–570. Reason for exclusion: Wrong study design.
- Gerra, G., F. Borella, A. Zaimovic, G. Moi, M. Bussandri, C. Bubici, and S. Bertacca, "Buprenorphine Versus Methadone for Opioid Dependence: Predictor Variables for Treatment Outcome," *Drug and Alcohol Dependence*, Vol. 75, No. 1, 2004. Reason for exclusion: No functional outcomes.
- Gerra, G., M. Ferri, E. Polidori, G. Santoro, A. Zaimovic, and E. Sternieri, "Long-Term Methadone Maintenance Effectiveness: Psychosocial and Pharmacological Variables," *Journal of Substance Abuse Treatment*, Vol. 25, No. 1, July 2003, pp. 1–8. Reason for exclusion: No functional outcomes.
- Gerra, G., G. Fertonani, A. Zaimovic, I. Rota-Graziosi, P. Avanzini, R. Caccavari, R. Delsignore, and A. Lucchini, "Hostility in Heroin Abusers Subtypes: Fluoxetine and Naltrexone Treatment," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, Vol. 19, No. 8, 1995. Reason for exclusion: No functional outcomes.

- Gerra, G., M. Manfredini, L. Somaini, I. Maremmanni, C. Leonardi, and C. Donnini, "Sexual Dysfunction in Men Receiving Methadone Maintenance Treatment: Clinical History and Psychobiological Correlates," *European Addiction Research*, Vol. 22, No. 3, 2016, pp. 163–175. Reason for exclusion: No functional outcomes.
- Gerra, G., E. Saenz, A. Busse, I. Maremmanni, R. Ciccocioppo, A. Zaimovic, M. L. Gerra, M. Amore, M. Manfredini, C. Donnini, and L. Somaini, "Supervised Daily Consumption, Contingent Take-Home Incentive and Non-Contingent Take-Home in Methadone Maintenance," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, Vol. 35, No. 2, 2011. Reason for exclusion: Only measures effects of a co-intervention.
- Gerra, G., A. Zaimovic, M. A. Raggi, F. Giusti, R. Delsignore, S. Bertacca, and F. Brambilla, "Aggressive Responding of Male Heroin Addicts Under Methadone Treatment: Psychometric and Neuroendocrine Correlates," *Drug and Alcohol Dependence*, Vol. 65, No. 1, December 1, 2001, pp. 85–95. Reason for exclusion: Wrong study design.
- Ghitza, U. E., D. H. Epstein, and K. L. Preston, "Self-Report of Illicit Benzodiazepine Use on the Addiction Severity Index Predicts Treatment Outcome," *Drug and Alcohol Dependence*, Vol. 97, No. 1–2, September 1, 2008, pp. 150–157. Reason for exclusion: Wrong study design.
- Ghitza, U. E., D. H. Epstein, J. Schmittner, M. Vahabzadeh, J. L. Lin, and K. L. Preston, "Randomized Trial of Prize-Based Reinforcement Density for Simultaneous Abstinence from Cocaine and Heroin," *Journal of Consulting and Clinical Psychology*, Vol. 75, No. 5, October 2007, pp. 765–774. Reason for exclusion: Only measures effects of a co-intervention.
- Giacomuzzi, S., W. Haaser, L. Pilsz, and Y. Riemer, "Driving Impairment on Buprenorphine and Slow-Release Oral Morphine in Drug-Dependent Patients," *Forensic Science International*, Vol. 152, No. 2–3, September 10, 2005, pp. 323–324. Reason for exclusion: Wrong study design.
- Giacomuzzi, S. M., M. Ertl, A. Vigl, Y. Riemer, V. Gunther, M. Kopp, W. Pilsz, and W. Haaser, "Driving Capacity of Patients Treated with Methadone and Slow-Release Oral Morphine," *Addiction*, Vol. 100, No. 7, July 2005, p. 1027. Reason for exclusion: Wrong study design.
- Gibson, A., L. Degenhardt, R. P. Mattick, R. Ali, J. White, and S. O'Brien, "Exposure to Opioid Maintenance Treatment Reduces Long-Term Mortality," *Addiction*, Vol. 103, No. 3, March 2008, pp. 462–468. Reason for exclusion: No functional outcomes.
- Gimbel, J., E. L. Spierings, N. Katz, Q. Xiang, E. Tzanis, and A. Finn, "Efficacy and Tolerability of Buccal Buprenorphine in Opioid-Experienced Patients with Moderate to Severe Chronic Low Back Pain: Results of a Phase 3, Enriched Enrollment, Randomized Withdrawal Study," *Pain*, Vol. 157, No. 11, November 2016, pp. 2517–2526. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Glaizal, M., V. Gazin, I. Aymard, C. Messina-Gourlot, N. Richard, M. Mallaret, P. Saviuc, and L. de Haro, "Suicidal Poisonings with Methadone in France: Results of a Two-Year National Survey by the Toxicovigilance Network," *Clinical Toxicology (Philadelphia)*, Vol. 50, No. 9, November 2012, pp. 841–846. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Goldstein, Avram, "Blind Dosage Comparisons and Other Studies in a Large Methadone Program," *Journal of Psychedelic Drugs*, Vol. 4, No. 2, Winter 1971, pp. 177–181. Reason for exclusion: No functional outcomes.
- Goldstein, A., "Blind Comparison of Once-Daily and Twice-Daily Dosage Schedules in a Methadone Program," *Clinical Pharmacology and Therapeutics*, Vol. 13, No. 1, January–February 1972, pp. 59–63. Reason for exclusion: No functional outcomes.
- Goldstein, A., and B. A. Judson, "Proceedings: Efficacy and Side Effects of Three Widely Different Methadone Doses," *Proceedings. National Conference on Methadone Treatment*, Vol. 1, 1973. Reason for exclusion: No functional outcomes.
- Goldstein, M. F., S. Deren, S. Y. Kang, D. C. Des Jarlais, and S. Magura, "Evaluation of an Alternative Program for MMTP Drop-Outs: Impact on Treatment Re-Entry," *Drug and Alcohol Dependence*, Vol. 66, No. 2, April 1, 2002, pp. 181–187. Reason for exclusion: Only measures effects of a co-intervention.
- Goli, V., L. R. Webster, M. J. Lamson, J. M. Cleveland, K. W. Sommerville, and E. Carter, "Effects of Concurrent Intravenous Morphine Sulfate and Naltrexone Hydrochloride on End-Tidal Carbon Dioxide," *Harm Reduction Journal*, Vol. 9, 2012. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gölz, J., and G. Partecke, "Catamnestic Development of Opiate Addicts After Naltrexone Induced Detoxification Under Anaesthesia, Naltrexone Supported Relapse Prevention, and Psychosocial Outpatient Aftercare," *Suchttherapie*, Vol. 1, No. 3, 2000. Reason for exclusion: Not English.
- Gombas, W., G. Fischer, H. Eder, R. Jagsch, S. Schindler, L. Pezawas, and S. Kasper, "Comparison of Buprenorphine and Methadone in Opiate Addicts," *Pharmacopsychiatry*, Vol. 30, 1997. Reason for exclusion: No functional outcomes.
- Gonzalez, G., G. DiGirolamo, M. E. Kolodziej, D. Smelson, and M. Romero-Gonzalez, "Memantine Improves Buprenorphine Treatment for Opioid-Dependent Young Adults," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- Gonzalez, G., G. DiGirolamo, M. Romero-Gonzalez, D. Smelson, D. Ziedonis, and M. Kolodziej, "Memantine Improves Buprenorphine/Naloxone Treatment for Opioid Dependent Young Adults," *Drug and Alcohol Dependence*, Vol. 156, November 1, 2015, pp. 243–253. Reason for exclusion: Only measures effects of a co-intervention.
- Gonzalez, G., A. Oliveto, and T. R. Kosten, "Combating Opiate Dependence: A Comparison Among the Available Pharmacological Options," *Expert Opinion on Pharmacotherapy*, Vol. 5, No. 4, April 2004, pp. 713–725. Reason for exclusion: Wrong study design.

- Goodarzi, A., N. Vouseoghi, M. Sedaghati, A. Mokri, and M. R. Zarrindast, "Dopamine Receptors in Human Peripheral Blood Lymphocytes: Changes in Mrna Expression in Opioid Addiction," *European Journal of Pharmacology*, Vol. 615, No. 1–3, August 1, 2009, pp. 218–222. Reason for exclusion: No functional outcomes.
- Goonoo, N., A. Bhaw-Luximon, R. Ujoodha, A. Jhugroo, G. K. Hulse, and D. Jhurry, "Naltrexone: A Review of Existing Sustained Drug Delivery Systems and Emerging Nano-Based Systems," *Journal of Controlled Release*, Vol. 183, June 10, 2014, pp. 154–166. Reason for exclusion: Wrong study design.
- Gordon, Michael S., Timothy W. Kinlock, Kathryn A. Couvillion, Robert P. Schwartz, and Kevin O'Grady, "A Randomized Clinical Trial of Methadone Maintenance for Prisoners: Prediction of Treatment Entry and Completion in Prison," *Journal of Offender Rehabilitation*, Vol. 51, No. 4, May 2012, pp. 222–238. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Gordon, Michael S., Timothy W. Kinlock, Robert P. Schwartz, Kathryn A. Couvillion, and Kevin E. O'Grady, "The Severity, Frequency, and Variety of Crime in Heroin-Dependent Prisoners Enrolled in a Buprenorphine Clinical Trial," *The Prison Journal*, Vol. 93, No. 4, December 2013, pp. 390–410. Reason for exclusion: Wrong study design.
- Gordon, M. S., T. W. Kinlock, R. P. Schwartz, T. T. Fitzgerald, K. E. O'Grady, and F. J. Vocci, "A Randomized Controlled Trial of Prison-Initiated Buprenorphine: Prison Outcomes and Community Treatment Entry," *Drug and Alcohol Dependence*, Vol. 142, September 1, 2014, pp. 33–40. Reason for exclusion: No functional outcomes.
- Gordon, M. S., T. W. Kinlock, R. P. Schwartz, and K. E. O'Grady, "A Randomized Clinical Trial of Methadone Maintenance for Prisoners: Findings at 6 Months Post-Release," *Addiction*, Vol. 103, No. 8, August 2008, pp. 1333–1342. Reason for exclusion: Article reporting follow up of included studies.
- Gordon, M. S., T. W. Kinlock, R. P. Schwartz, K. E. O'Grady, T. T. Fitzgerald, and F. J. Vocci, "A Randomized Clinical Trial of Buprenorphine for Prisoners: Findings at 12-Months Post-Release," *Drug and Alcohol Dependence*, Vol. 172, 1, 2017, pp. 34–42. Reason for exclusion: Only measures effects of a co-intervention.
- Gossop, M., "The Dutch Heroin Trial: Issues and Implications," *Sucht*, Vol. 48, No. 4, 2002, pp. 304–306. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gossop, M., "The National Treatment Outcomes Research Study (NTORS) and Its Influence on Addiction Treatment Policy in the United Kingdom," *Addiction*, Vol. 110, Supplement 2, July 2015, pp. 50–53. Reason for exclusion: Wrong study design.
- Gossop, M., and B. Bradley, "Insomnia Among Addicts During Supervised Withdrawal from Opiates: A Comparison of Oral Methadone and Electrostimulation," *Drug and Alcohol Dependence*, Vol. 13, No. 2, 1984. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Gossop, M., P. Griffiths, B. Bradley, and J. Strang, "Opiate Withdrawal Symptoms in Response to 10-Day and 21-Day Methadone Withdrawal Programmes," *British Journal of Psychiatry*, Vol. 154, March 1989, pp. 360–363. Reason for exclusion: Wrong setting (i.e., inpatient).
- Gossop, M., J. Marsden, D. Stewart, and S. Treacy, "Outcomes After Methadone Maintenance and Methadone Reduction Treatments: Two-Year Follow-Up Results from the National Treatment Outcome Research Study," *Drug and Alcohol Dependence*, Vol. 62, No. 3, 2001. Reason for exclusion: Wrong study design.
- Gossop, M., J. Strang, and B. Bradley, "The Bethlem Study on NET: Putting the Record Straight," *British Journal of Addiction*, Vol. 84, No. 7, July 1989, pp. 818–819. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gotthardt, F., C. Huber, C. Thierfelder, L. Grize, M. Kraenzlin, C. Scheidegger, and C. Meier, "Bone Mineral Density and Its Determinants in Men with Opioid Dependence," *Journal of Bone and Mineral Metabolism*, January 8, 2016. Reason for exclusion: No functional outcomes.
- Gourevitch, M. N., D. Hartel, P. Tenore, K. Freeman, I. Marion, J. Hecht, and J. Lowinson, "Three Oral Formulations of Methadone. A Clinical and Pharmacodynamic Comparison," *Journal of Substance Abuse Treatment*, Vol. 17, No. 3, October 1999, pp. 237–241. Reason for exclusion: Wrong study design.
- Gouveia, M., R. Sousa, J. Costa, and M. Borges, "Economic Evaluation of Suboxone for Substitution Treatment of Opioid Drug Dependence in Portugal," *Heroin Addiction and Related Clinical Problems*, Vol. 17, No. 1, 2015. Reason for exclusion: Wrong study design.
- Gowing, L., M. Farrell, R. Ali, and J. M. White, "Alpha(2)-Adrenergic Agonists for the Management of Opioid Withdrawal," *The Cochrane Database of Systematic Reviews*, No. 5, May 3, 2016, p. Cd002024. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gowing, L. R., "The Role of Opioid Substitution Treatment in Reducing HIV Transmission," *BMJ*, Vol. 345, October 3, 2012, p. e6425. Reason for exclusion: Only measures effects of a co-intervention.
- Gowing, L. R., R. L. Ali, and J. M. White, "Which Substitution Pharmacotherapy Is Most Effective in Treating Opioid Dependence?" *Medical Journal of Australia*, Vol. 176, No. 10, 20, 2002, pp. 493–494. Reason for exclusion: Wrong study design.
- Gowing, L. R., M. Hickman, and L. Degenhardt, "Mitigating the Risk of HIV Infection with Opioid Substitution Treatment," *Bulletin of the World Health Organization*, Vol. 91, No. 2, February 1, 2013, pp. 148–149. Reason for exclusion: Wrong study design.
- Gowing, L. R., and O. Lopatko, "Office Based Treatment with Buprenorphine Plus Naloxone, or Buprenorphine Alone Reduces Opiate Use and Craving," *Evidence Based Mental Health*, Vol. 7, No. 2, 2004, p. 54. Reason for exclusion: Wrong study design.

- Gradin, V. B., A. Baldacchino, D. Balfour, K. Matthews, and J. D. Steele, "Abnormal Brain Activity During a Reward and Loss Task in Opiate-Dependent Patients Receiving Methadone Maintenance Therapy," *Neuropsychopharmacology*, Vol. 39, No. 4, March 2014, pp. 885–894. Reason for exclusion: No functional outcomes.
- Graham, A. W., "Buprenorphine Alone or in Combination with Naloxone Reduced the Use of and Craving for Opiates in Opiate Addiction," *ACP Journal Club*, Vol. 140, No. 3, May–June 2004, p. 65. Reason for exclusion: Wrong study design.
- Greenwald, M. K., "Opioid Abstinence Reinforcement Delays Heroin Lapse During Buprenorphine Dose Tapering," *Journal of Applied Behavior Analysis*, Vol. 41, No. 4, 2008. Reason for exclusion: Only measures effects of a co-intervention.
- Greenwald, M., C. E. Johanson, J. Bueller, Y. Chang, D. E. Moody, M. Kilbourn, R. Koeppe, and J. K. Zubieta, "Buprenorphine Duration of Action: Mu-Opioid Receptor Availability and Pharmacokinetic and Behavioral Indices," *Biological Psychiatry*, Vol. 61, No. 1, January 1, 2007, pp. 101–110. Reason for exclusion: Wrong study design.
- Greenwald, M. K., C. R. Schuster, and C. E. Johanson, "Opiate Craving During Naloxone-Precipitated Withdrawal in Methadone-Maintained Volunteers," *NIDA Research Monograph*, Vol. 162, 1996. Reason for exclusion: Wrong study design.
- Greenwald, M. K., E. A. Woodcock, D. Khatib, and J. A. Stanley, "Methadone Maintenance Dose Modulates Anterior Cingulate Glutamate Levels in Heroin-Dependent Individuals: A Preliminary in Vivo H MRS Study," *Psychiatry Research—Neuroimaging*, Vol. 233, No. 2, 2015. Reason for exclusion: Wrong study design.
- Greenwald, M. K., "Early Impact of Methadone Induction for Heroin Dependence: Differential Effects of Two Dose Sequences in a Randomized Controlled Study," *Experimental and Clinical Psychopharmacology*, Vol. 14, No. 1, February 2006, pp. 52–67. Reason for exclusion: No functional outcomes.
- Greenwald, M. K., "Effects of Experimental Unemployment, Employment and Punishment Analogs on Opioid Seeking and Consumption in Heroin-Dependent Volunteers," *Drug and Alcohol Dependence*, Vol. 111, No. 1–2, September 1, 2010, pp. 64–73. Reason for exclusion: Only measures effects of a co-intervention.
- Greenwald, M. K., C. E. Johanson, and C. R. Schuster, "Opioid Reinforcement in Heroin-Dependent Volunteers During Outpatient Buprenorphine Maintenance," *Drug and Alcohol Dependence*, Vol. 56, No. 3, October 1, 1999, pp. 191–203. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Greenwald, M. K., K. J. Schuh, J. A. Hopper, C. R. Schuster, and C. E. Johanson, "Effects of Buprenorphine Sublingual Tablet Maintenance on Opioid Drug-Seeking Behavior by Humans," *Psychopharmacology (Berl)*, Vol. 160, No. 4, April 2002, pp. 344–352. Reason for exclusion: Only measures effects of a co-intervention.

- Greenwald, M. K., E. A. Woodcock, D. Khatib, and J. A. Stanley, "Methadone Maintenance Dose Modulates Anterior Cingulate Glutamate Levels in Heroin-Dependent Individuals: A Preliminary in Vivo H MRS Study," *Psychiatry Research*, Vol. 233, No. 2, August 30, 2015, pp. 218–224. Reason for exclusion: Wrong study design.
- Grey, A., K. Rix-Trott, A. Horne, G. Gamble, M. Bolland, and I. R. Reid, "Decreased Bone Density in Men on Methadone Maintenance Therapy," *Addiction*, Vol. 106, No. 2, February 2011, pp. 349–354. Reason for exclusion: No functional outcomes.
- Griffin, M. L., H. E. Bennett, G. M. Fitzmaurice, K. P. Hill, S. E. Provost, and R. D. Weiss, "Health-Related Quality of Life Among Prescription Opioid-Dependent Patients: Results from a Multi-Site Study," *American Journal on Addictions*, Vol. 24, No. 4, 2015. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Griffin, M. L., D. R. Dodd, J. S. Potter, L. S. Rice, W. Dickinson, S. Sparenborg, and R. D. Weiss, "Baseline Characteristics and Treatment Outcomes in Prescription Opioid Dependent Patients with and Without Co-Occurring Psychiatric Disorder," *The American Journal of Drug and Alcohol Abuse*, Vol. 40, No. 2, March 2014, pp. 157–162. Reason for exclusion: Only measures effects of a co-intervention.
- Gronbladh, L., and L. Gunne, "Methadone-Assisted Rehabilitation of Swedish Heroin Addicts," *Drug and Alcohol Dependence*, Vol. 24, No. 1, August 1989, pp. 31–37. Reason for exclusion: Only measures effects of a co-intervention.
- Gronbladh, L., L. S. Ohlund, and L. M. Gunne, "Mortality in Heroin Addiction: Impact of Methadone Treatment," *Acta Psychiatrica Scandinavica*, Vol. 82, No. 3, 1990. Reason for exclusion: No functional outcomes.
- Gross, A., E. A. Jacobs, N. M. Petry, G. J. Badger, and W. K. Bickel, "Limits to Buprenorphine Dosing: A Comparison Between Quintuple and Sextuple the Maintenance Dose Every 5 Days," *Drug and Alcohol Dependence*, Vol. 64, No. 1, September 1, 2001, pp. 111–116. Reason for exclusion: No functional outcomes.
- Gross, A., L. A. Marsch, G. J. Badger, and W. K. Bickel, "A Comparison Between Low-Magnitude Voucher and Buprenorphine Medication Contingencies in Promoting Abstinence from Opioids and Cocaine," *Experimental and Clinical Psychopharmacology*, Vol. 14, No. 2, May 2006, pp. 148–156. Reason for exclusion: Only measures effects of a co-intervention.
- Gruber, K., M. A. Chutuape, and M. L. Stitzer, "Reinforcement-Based Intensive Outpatient Treatment for Inner City Opiate Abusers: A Short-Term Evaluation," *Drug and Alcohol Dependence*, Vol. 57, No. 3, January 1, 2000, pp. 211–223. Reason for exclusion: Only measures effects of a co-intervention.
- Gruber, V. A., P. M. Rainey, D. E. Moody, G. D. Morse, Q. Ma, S. Prathikanti, P. A. Pade, A. A. Alvanzo, and E. F. McCance-Katz, "Interactions Between Buprenorphine and the Protease Inhibitors Darunavir-Ritonavir and Fosamprenavir-Ritonavir," *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, Vol. 54, No. 3, 2012. Reason for exclusion: Only measures effects of a co-intervention.

- Gruber, V. A., K. L. Delucchi, A. Kielstein, and S. L. Batki, "A Randomized Trial of 6-Month Methadone Maintenance with Standard or Minimal Counseling Versus 21-Day Methadone Detoxification," *Drug and Alcohol Dependence*, Vol. 94, No. 1–3, April 1, 2008, pp. 199–206. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gryczynski, Jan, Jerome H. Jaffe, Robert P. Schwartz, Kristi A. Dušek, Nishan Gugsu, Cristin L. Monroe, Kevin E. O'Grady, Yngvild K. Olsen, and Shannon Gwin Mitchell, "Patient Perspectives on Choosing Buprenorphine over Methadone in an Urban, Equal-Access System," *The American Journal on Addictions*, Vol. 22, No. 3, May–June 2013, pp. 285–291. Reason for exclusion: Wrong study design.
- Gryczynski, J., S. G. Mitchell, J. H. Jaffe, S. M. Kelly, C. P. Myers, K. E. O'Grady, Y. K. Olsen, and R. P. Schwartz, "Retention in Methadone and Buprenorphine Treatment Among African Americans," *Journal of Substance Abuse Treatment*, Vol. 45, No. 3, September 2013, pp. 287–292. Reason for exclusion: No functional outcomes.
- Gu, J., J. T. Lau, H. Xu, Y. Zhong, Y. Hao, Y. Zhao, L. Fan, and W. Ling, "A Randomized Controlled Trial to Evaluate the Relative Efficacy of the Addition of a Psycho-Social Intervention to Standard-of-Care Services in Reducing Attrition and Improving Attendance Among First-Time Users of Methadone Maintenance Treatment in China," *AIDS and Behavior*, Vol. 17, No. 6, July 2013, pp. 2002–2010. Reason for exclusion: Only measures effects of a co-intervention.
- Gu, J., H. Xu, J. T. Lau, L. Chen, Z. Wang, C. Hao, and Y. Hao, "Situation-Specific Factors Predicting Nonadherence to Methadone Maintenance Treatment: A Cross-Sectional Study Using the Case-Crossover Design in Guangzhou, China," *AIDS Care*, Vol. 26, Supplement 1, 2014, pp. S107–112. Reason for exclusion: Wrong study design.
- Gunderson, E. W., and M. Sumner, "Efficacy of Buprenorphine/Naloxone Rapidly Dissolving Sublingual Tablets (BNX-RDT) After Switching from BNX Sublingual Film," *Journal of Addiction Medicine*, Vol. 10, No. 2, 2016. Reason for exclusion: No functional outcomes.
- Gunderson, E. W., P. Hjelmstrom, and M. Sumner, "Effects of a Higher-Bioavailability Buprenorphine/Naloxone Sublingual Tablet Versus Buprenorphine/Naloxone Film for the Treatment of Opioid Dependence During Induction and Stabilization: A Multicenter, Randomized Trial," *Clinical Therapeutics*, Vol. 37, No. 10, October 1, 2015, pp. 2244–2255. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Gunderson, E. W., X. Q. Wang, D. A. Fiellin, B. Bryan, and F. R. Levin, "Unobserved Versus Observed Office Buprenorphine/Naloxone Induction: A Pilot Randomized Clinical Trial," *Addictive Behaviors*, Vol. 35, No. 5, May 2010, pp. 537–540. Reason for exclusion: Only measures effects of a co-intervention.
- Gunne, L. M., and L. Gronblad, "The Efficacy of Methadone Maintenance Treatment," *Lakartidningen*, Vol. 77, No. 4, 1980. Reason for exclusion: Not English.

- Gunne, L. M., and L. Gronbladh, "The Swedish Methadone Maintenance Program: A Controlled Study," *Drug and Alcohol Dependence*, Vol. 7, No. 3, June 1981, pp. 249–256. Reason for exclusion: No functional outcomes.
- Guo, S., Z. Jiang, and Y. Wu, "Efficacy of Naltrexone Hydrochloride for Preventing Relapse Among Opiate-Dependent Patients After Detoxification," *Hong Kong Journal of Psychiatry*, Vol. 11, No. 4, 2001. Reason for exclusion: No functional outcomes.
- Gupta, S., J. E. Iudicello, C. Shi, S. Letendre, A. Knight, J. Li, P. K. Riggs, D. R. Franklin, Jr., N. Duarte, H. Jin, J. Hampton Atkinson, X. Yu, Z. Wu, I. Grant, and R. K. Heaton, "Absence of Neurocognitive Impairment in a Large Chinese Sample of HCV-Infected Injection Drug Users Receiving Methadone Treatment," *Drug and Alcohol Dependence*, Vol. 137, April 1, 2014, pp. 29–35. Reason for exclusion: Wrong study design.
- Gutwinski, S., L. K. Bald, A. Heinz, C. A. Muller, A. K. Schmidt, C. Wiers, F. BERPpohl, and J. Gallinat, "In Reply," *Deutsches Ärzteblatt International*, Vol. 110, No. 41, October 2013, pp. 688–689. Reason for exclusion: Wrong study design.
- Haasen, C., F. J. Eiroa-Orosa, U. Verthein, M. Soyka, C. Dilg, I. Schafer, and J. Reimer, "Effects of Heroin-Assisted Treatment on Alcohol Consumption: Findings of the German Randomized Controlled Trial," *Alcohol*, Vol. 43, No. 4, June 2009, pp. 259–264. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Haasen, C., and W. van den Brink, "Innovations in Agonist Maintenance Treatment of Opioid-Dependent Patients," *Current Opinion in Psychiatry*, Vol. 19, No. 6, November 2006, pp. 631–636. Reason for exclusion: Wrong study design.
- Haasen, C., U. Verthein, P. Degkwitz, J. Berger, M. Krausz, and D. Naber, "Heroin-Assisted Treatment for Opioid Dependence: Randomised Controlled Trial," *British Journal of Psychiatry*, Vol. 191, July 2007, pp. 55–62. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Haasen, C., U. Verthein, P. Degkwitz, S. Kuhn, C. Hartwig, and J. Reimer, "A Multicentre, Randomised Controlled Study of Diamorphine Maintenance Therapy for Opiate-Dependent Patients: Target Group-Specific Results. [German]," *Sucht*, Vol. 53, No. 5, 2007. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Haig, T., "Randomized Controlled Trial Proves Effectiveness of Methadone Maintenance Treatment in Prison," *Canadian HIV/AIDS Policy and Law Review*, Vol. 8, No. 3, December 2003, p. 48. Reason for exclusion: Wrong study design.
- Haigney, M. C., "First, Do No Harm: QT Interval Screening in Methadone Maintenance Treatment," *Journal of Addictive Diseases*, Vol. 30, No. 4, October 2011, pp. 309–312. Reason for exclusion: Wrong study design.
- Hales, T. G., "Arresting the Development of Morphine Tolerance and Dependence," *British Journal Anaesthesia*, Vol. 107, No. 5, November 2011, pp. 653–655. Reason for exclusion: Wrong study design.

- Hall, S. M., P. Loeb, M. LeVois, and J. Cooper, "Increasing Employment in Ex-Heroin Addicts: II Methadone Maintenance Sample," *Behavior Therapy*, Vol. 12, No. 4, 1981. Reason for exclusion: Only measures effects of a co-intervention.
- Hall, S. M., P. C. Loeb, and M. Kushner, "Methadone Dose Decreases and Anxiety Reduction," *Addictive Behaviors*, Vol. 9, No. 1, 1984, pp. 11–19. Reason for exclusion: Only measures effects of a co-intervention.
- Hall, W., and A. Carter, "Is Deep Brain Stimulation for Addiction an Acceptable Crime Control Measure?" *Addiction*, Vol. 107, No. 7, July 2012, p. 1360; author reply 1361. Reason for exclusion: Wrong study design.
- Hall, Wayne, and Adrian Carter, "Is Deep Brain Stimulation for Addiction an Acceptable Crime Control Measure? . . . Stephen J. H., Halpern C. H., Barrios C. J., Balmuri U., Pisapia J. M., Wolf J. A. et al. Deep Brain Stimulation Compared with Methadone Maintenance for the Treatment of Heroin Dependence: A Threshold and Cost-Effectiveness Analysis. *Addiction* 2012; 107: 624–34," *Addiction*, Vol. 107, No. 7, 2012, p. 1360. Reason for exclusion: Wrong study design.
- Hall, Wayne, Jeff Ward, and Richard P. Mattick, eds. *The Effectiveness of Methadone Maintenance Treatment I: Heroin Use and Crime*, Amsterdam, Netherlands: Harwood Academic Publishers, 1998.
- Halliday, L., "Published Research," *Paediatric Nursing*, Vol. 17, No. 5, 2005, pp. 10–10. Reason for exclusion: Wrong study design.
- Hameedi, F. A., S. W. Woods, M. I. Rosen, H. R. Pearsall, and T. R. Kosten, "Dose Dependent Effects of Yohimbine on Methadone Maintained Patients," *The American Journal of Drug and Alcohol Abuse*, Vol. 23, No. 2, May 1997, pp. 327–333. Reason for exclusion: Only measures effects of a co-intervention.
- Hamilton, S. P., E. V. Nunes, M. Janal, and L. Weber, "The Effect of Sertraline on Methadone Plasma Levels in Methadone-Maintenance Patients," *American Journal of Addiction*, Vol. 9, No. 1, Winter 2000, pp. 63–69. Reason for exclusion: Only measures effects of a co-intervention.
- Hammig, R., W. Kohler, K. Bonorden-Kleij, B. Weber, K. Lebentrau, T. Berthel, L. Babic-Hohnjec, C. Vollmert, D. Hopner, N. Gholami, U. Verthein, C. Haasen, J. Reimer, and C. Ruckes, "Safety and Tolerability of Slow-Release Oral Morphine Versus Methadone in the Treatment of Opioid Dependence," *Journal of Substance Abuse Treatment*, Vol. 47, No. 4, October 2014, pp. 275–281. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Hammond, Christopher J., and Kevin M. Gray, "Pharmacotherapy for Substance Use Disorders in Youths," *Journal of Child and Adolescent Substance Abuse*, Vol. 25, No. 4, 2016, pp. 292–316. Reason for exclusion: Wrong population (pregnant women, adolescents).

- Hansen, L. E., G. L. Stone, C. A. Matson, D. J. Tybor, M. E. Pevear, and E. L. Smith, "Total Joint Arthroplasty in Patients Taking Methadone or Buprenorphine/Naloxone Preoperatively for Prior Heroin Addiction: A Prospective Matched Cohort Study," *The Journal of Arthroplasty*, Vol. 31, No. 8, August 2016, pp. 1698–1701. Reason for exclusion: No functional outcomes.
- Hao, S. Q., M. Zhao, R. W. Zhang, J. C. Zhang, J. Zhang, and X. S. Feng, "The Effectiveness Comparison of Jitai Tablets Versus Methadone in Community-Based Drug Treatment: A 1-Year Follow-Up Study," *Addictive Behaviors*, Vol. 38, No. 10, 2013. Reason for exclusion: No functional outcomes.
- Harlow, S. D., M. Cohen, S. E. Ohmit, P. Schuman, S. Cu-Uvin, X. Lin, R. Greenblatt, A. Gurtman, A. Khalsa, H. Minkoff, M. A. Young, and R. S. Klein, "Substance Use and Psychotherapeutic Medications: A Likely Contributor to Menstrual Disorders in Women Who Are Seropositive for Human Immunodeficiency Virus," *American Journal of Obstetrics and Gynecology*, Vol. 188, No. 4, April 2003, pp. 881–886. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Harris, A. H., E. Gospodarevskaya, and A. J. Ritter, "A Randomised Trial of the Cost Effectiveness of Buprenorphine as an Alternative to Methadone Maintenance Treatment for Heroin Dependence in a Primary Care Setting," *Pharmacoeconomics*, Vol. 23, No. 1, 2005, pp. 77–91. Reason for exclusion: No functional outcomes.
- Harris, D. S., R. T. Jones, S. Welm, R. A. Upton, E. Lin, and J. Mendelson, "Buprenorphine and Naloxone Co-Administration in Opiate-Dependent Patients Stabilized on Sublingual Buprenorphine," *Drug and Alcohol Dependence*, Vol. 61, No. 1, 2000. Reason for exclusion: Wrong study design.
- Hartnoll, R. L., M. C. Mitcheson, A. Battersby, G. Brown, M. Ellis, P. Fleming, and N. Hedley, "Evaluation of Heroin Maintenance in Controlled Trial," *Archives of General Psychiatry*, Vol. 37, No. 8, August 1980, pp. 877–884. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Hasan, S. I., R. Abd Rashid, M. Ahmad Zahari, C. G. Ng, A. H. Sulaiman, F. Ibrahim, N. Robson, R. Abdul Kadir, M. Mustafa, M. H. Habil, and W. A. Irnee Wan Hussin, "A Study on the Effectiveness of Electroacupuncture as Adjunctive Treatment Among Methadone Maintenance Therapy Clients in University of Malaya Medical Center, Kuala Lumpur, Malaysia," *International Journal of Neuropsychopharmacology*, Vol. 15, Supplement 1, June 2012, p. 70. Reason for exclusion: Only measures effects of a co-intervention.
- Hasanovic, M., I. Pajevic, A. Kuldija, and A. Delic, "Medically Assisted Treatment for Opiate Addiction—Suboxone Method as Prevention of Social Exclusion of Youth—Tuzla Model," *Psychiatria Danubina*, Vol. 24, Supplement 3, October 2012, pp. S398–404. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).

- Havassy, B., and W. A. Hargreaves, "Allowing Methadone Clients Control over Dosage: A 48-Week Controlled Trial," *Addictive Behaviors*, Vol. 6, No. 4, 1981, pp. 283–288. Reason for exclusion: Only measures effects of a co-intervention.
- Havassy, B., W. A. Hargreaves, and L. De Barros, "Self-Regulation of Dose in Methadone Maintenance with Contingent Privileges," *Addictive Behaviors*, Vol. 4, No. 1, 1979, pp. 31–38. Reason for exclusion: Only measures effects of a co-intervention.
- Havassy, B. E., and J. M. Tschann, "Chronic Heroin Use During Methadone Treatment: A Test of the Efficacy of High Maintenance Doses," *Addictive Behaviors*, Vol. 9, No. 1, 1984, pp. 57–65. Reason for exclusion: Only measures effects of a co-intervention.
- Hay, J. L., J. M. White, F. Bochner, A. A. Somogyi, T. J. Semple, and B. Rounsefell, "Hyperalgesia in Opioid-Managed Chronic Pain and Opioid-Dependent Patients," *The Journal of Pain*, Vol. 10, No. 3, March 2009, pp. 316–322. Reason for exclusion: Wrong study design.
- He, J., and S. M. Apelt, "Association Between Buprenorphine/Naloxone Sublingual Tablet and All-Cause Mortality in Patients Receiving Medication Assisted Treatment for Opioid Addiction in UK," *Pharmacoepidemiology and Drug Safety*, Vol. 25, Supplement 3, 2016, p. 581. Reason for exclusion: Conference abstract.
- Healey, A., M. Knapp, J. Marsden, M. Gossop, and D. Stewart, "Criminal Outcomes and Costs of Treatment Services for Injecting and Non-Injecting Heroin Users: Evidence from a National Prospective Cohort Survey," *Journal of Health Services Research and Policy*, Vol. 8, No. 3, July 2003, pp. 134–141. Reason for exclusion: Wrong study design.
- Heimer, R., H. Catania, R. G. Newman, J. Zambrano, A. Brunet, and A. M. Ortiz, "Methadone Maintenance in Prison: Evaluation of a Pilot Program in Puerto Rico," *Drug and Alcohol Dependence*, Vol. 83, No. 2, 2006, pp. 122–129. Reason for exclusion: Wrong setting (i.e., inpatient).
- Henney, J. E., "From the Food and Drug Administration," *JAMA*, Vol. 284, No. 17, November 1, 2000, p. 2178. Reason for exclusion: Wrong study design.
- Henriksen, K., H. Waal, and P. Krajci, "Will Non-Compliant 'Hard-to-Treat' Opioid-Dependent Patients Profit from Low Threshold Methadone Treatment? A Prospective 15-Month Evaluation of Patients on Low Dosage Methadone Treatment at Oslo University Hospital," *Heroin Addiction and Related Clinical Problems*, Vol. 18, No. 3, 2016. Reason for exclusion: Wrong study design.
- Herning, R. I., W. E. Better, K. Tate, A. Umbricht, K. L. Preston, and J. L. Cadet, "Methadone Treatment Induces Attenuation of Cerebrovascular Deficits Associated with the Prolonged Abuse of Cocaine and Heroin," *Neuropsychopharmacology*, Vol. 28, No. 3, March 2003, pp. 562–568. Reason for exclusion: No functional outcomes.

- Hess, Melanie, Lukas Boesch, Ria Leisinger, and Rudolf Stohler, "Transdermal Buprenorphine to Switch Patients from Higher Dose Methadone to Buprenorphine Without Severe Withdrawal Symptoms," *The American Journal on Addictions*, Vol. 20, No. 5, September–October 2011, pp. 480–481. Reason for exclusion: No functional outcomes.
- Hesse, Morten, and Mads U. Pedersen, "Easy-Access Services in Low-Threshold Opiate Agonist Maintenance," *International Journal of Mental Health and Addiction*, Vol. 6, No. 3, July 2008, pp. 316–324. Reason for exclusion: Only measures effects of a co-intervention.
- Highfield, D. A., R. P. Schwartz, J. H. Jaffe, and K. E. O'Grady, "Intravenous and Intranasal Heroin-Dependent Treatment-Seekers: Characteristics and Treatment Outcome," *Addiction*, Vol. 102, No. 11, November 2007, pp. 1816–1823. Reason for exclusion: Wrong study design.
- Hill, E., D. Han, P. Dumouchel, N. Dehak, T. Quatieri, C. Moehs, M. Oscar-Berman, J. Giordano, T. Simpatico, D. Barh, and K. Blum, "Long Term Suboxone Emotional Reactivity as Measured by Automatic Detection in Speech," *PLoS One*, Vol. 8, No. 7, 2013, p. e69043. Reason for exclusion: Wrong study design.
- Hiller, M. L., G. A. Rowan-Szal, N. G. Bartholomew, and D. D. Simpson, "Effectiveness of a Specialized Women's Intervention in a Residential Treatment Program," *Substance Use and Misuse*, Vol. 31, No. 6, May 1996, pp. 771–783. Reason for exclusion: Only measures effects of a co-intervention.
- Hines, S., S. Theodorou, A. Williamson, D. Fong, and K. Curry, "Management of Acute Pain in Methadone Maintenance Therapy In-Patients," *Drug and Alcohol Review*, Vol. 27, No. 5, September 2008, pp. 519–523. Reason for exclusion: Wrong setting (inpatient).
- Hobbins, D. F., "Case Report: Methadone Maintenance Dose of 1200 Mg/Day in an Ultra-Rapid Metabolizer," *Journal of Addictions Nursing (Taylor & Francis Ltd)*, Vol. 21, No. 1, 2010, pp. 22–36. Reason for exclusion: Wrong study design.
- Hoefler, M., and I. Petrakis, "Prescription Drug Monitoring Programs and Buprenorphine Maintenance: Clinical Considerations," *The American Journal on Addictions*, Vol. 23, No. 6, November–December 2014, pp. 616–617. Reason for exclusion: Wrong study design.
- Hoffmann, O. M., T. Ljungberg, and F. Frisell, "Drug-Free Period from Non-Opioids Before Start of the Treatment with Methadone or Buprenorphine for Heroin-Dependence Improves Efficacy," *Drug and Alcohol Dependence*, Vol. 156, No. 1, 2015, pp. e95–e96. Reason for exclusion: Wrong study design.
- Hoflich, A. S., M. Langer, R. Jagsch, A. Bawert, B. Winklbaier, G. Fischer, and A. Unger, "Peripartum Pain Management in Opioid Dependent Women," *European Journal of Pain*, Vol. 16, No. 4, April 2012, pp. 574–584. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).

- Holland, R., V. Maskrey, L. Swift, C. Notley, A. Robinson, J. Nagar, T. Gale, and C. Kouimtsidis, "Treatment Retention, Drug Use and Social Functioning Outcomes in Those Receiving 3 Months Versus 1 Month of Supervised Opioid Maintenance Treatment. Results from the Super C Randomized Controlled Trial," *Addiction*, Vol. 109, No. 4, April 2014, pp. 596–607. Reason for exclusion: Only measures effects of a co-intervention.
- Holland, R., C. Matheson, G. Anthony, K. Roberts, S. Priyadarshi, A. Macrae, E. Whitelaw, S. Appavoo, and C. Bond, "A Pilot Randomised Controlled Trial of Brief Versus Twice Weekly Versus Standard Supervised Consumption in Patients on Opiate Maintenance Treatment," *Drug and Alcohol Review*, Vol. 31, No. 4, June 2012, pp. 483–491. Reason for exclusion: Only measures effects of a co-intervention.
- Hollander, P., A. K. Gupta, R. Plodkowski, F. Greenway, H. Bays, C. Burns, P. Klassen, and K. Fujioka, "Effects of Naltrexone Sustained-Release/Bupropion Sustained-Release Combination Therapy on Body Weight and Glycemic Parameters in Overweight and Obese Patients with Type 2 Diabetes," *Diabetes Care*, Vol. 36, No. 12, 2013. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Holtyn, A. F., M. N. Koffarnus, A. DeFulio, S. O. Sigurdsson, E. C. Strain, R. P. Schwartz, J. M. Leoutsakos, and K. Silverman, "The Therapeutic Workplace to Promote Treatment Engagement and Drug Abstinence in Out-of-Treatment Injection Drug Users: A Randomized Controlled Trial," *Preventive Medicine*, Vol. 68, November 2014, pp. 62–70. Reason for exclusion: Only measures effects of a co-intervention.
- Hopfer, C. J., Khuri, T. J. Crowley, and S. Hooks, "Adolescent Heroin Use: A Review of the Descriptive and Treatment Literature (Structured Abstract)," *Journal of Substance Abuse Treatment*, Vol. 23, No. 3, 2002. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Housova, J., H. Wilczek, M. M. Haluzik, J. Kremen, J. Krizova, and M. Haluzik, "Adipocyte-Derived Hormones in Heroin Addicts: The Influence of Methadone Maintenance Treatment," *Physiological Research*, Vol. 54, No. 1, 2005, pp. 73–78. Reason for exclusion: No functional outcomes.
- Hser, Y. I., A. J. Saxon, D. Huang, A. Hasson, C. Thomas, M. Hillhouse, P. Jacobs, C. Teruya, P. McLaughlin, K. Wiest, A. Cohen, and W. Ling, "Treatment Retention Among Patients Randomized to Buprenorphine/Naloxone Compared to Methadone in a Multi-Site Trial," *Addiction*, Vol. 109, No. 1, January 2014, pp. 79–87. Reason for exclusion: No functional outcomes.
- Huang, W. L., L. R. Chang, Y. Z. Chen, H. C. Chang, M. H. Hsieh, C. H. Lin, and Y. H. Lin, "The Tridimensional Personality of Male Heroin Users Treated with Methadone in Taiwan," *Comprehensive Psychiatry*, Vol. 55, No. 5, July 2014, pp. 1220–1226. Reason for exclusion: No functional outcomes.

- Huang, Y. F., H. S. Kuo, C. Y. Lew-Ting, F. Tian, C. H. Yang, T. I. Tsai, S. J. Gange, and K. E. Nelson, "Mortality Among a Cohort of Drug Users After Their Release from Prison: An Evaluation of the Effectiveness of a Harm Reduction Program in Taiwan," *Addiction*, Vol. 106, No. 8, August 2011, pp. 1437–1445. Reason for exclusion: Only measures effects of a co-intervention.
- Hubbard, R. L., S. G. Craddock, and J. Anderson, "Overview of 5-Year Followup Outcomes in the Drug Abuse Treatment Outcome Studies (DATOS)," *Journal of Substance Abuse Treatment*, Vol. 25, No. 3, October 2003, pp. 125–134. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Hulse, G. K., N. Morris, D. Arnold-Reed, and R. J. Tait, "Improving Clinical Outcomes in Treating Heroin Dependence: Randomized, Controlled Trial of Oral or Implant Naltrexone," *Archives of General Psychiatry*, Vol. 66, No. 10, October 2009, pp. 1108–1115. Reason for exclusion: No functional outcomes.
- Hulse, G. K., H. T. Ngo, and R. J. Tait, "Risk Factors for Craving and Relapse in Heroin Users Treated with Oral or Implant Naltrexone," *Biological Psychiatry*, Vol. 68, No. 3, August 1, 2010, pp. 296–302. Reason for exclusion: No functional outcomes.
- Hulse, G. K., R. J. Tait, S. D. Comer, M. A. Sullivan, I. G. Jacobs, and D. Arnold-Reed, "Reducing Hospital Presentations for Opioid Overdose in Patients Treated with Sustained Release Naltrexone Implants," *Drug and Alcohol Dependence*, Vol. 79, No. 3, September 1, 2005, pp. 351–357. Reason for exclusion: No functional outcomes.
- Hyman, S. M., K. I. Hong, T. M. Chaplin, Z. Dabre, A. D. Comegys, A. Kimmerling, and R. Sinha, "A Stress-Coping Profile of Opioid Dependent Individuals Entering Naltrexone Treatment: A Comparison with Healthy Controls," *Psychology of Addictive Behaviors*, Vol. 23, No. 4, December 2009, pp. 613–619. Reason for exclusion: No functional outcomes.
- Ignjatova, L., and M. Raleva, "Gender Difference in the Treatment Outcome of Patients Served in the Mixed-Gender Program," *Bratislavské Lekárske Listy*, Vol. 110, No. 5, 2009, pp. 285–289. Reason for exclusion: Wrong study design.
- Iguchi, M., "Methadone Treatment and Needle Use Prevention," *Focus*, Vol. 17, No. 3, February 2002, pp. 5–6. Reason for exclusion: Wrong study design.
- Iguchi, M. Y., M. L. Stitzer, G. E. Bigelow, and I. A. Liebson, "Contingency Management in Methadone Maintenance: Effects of Reinforcing and Aversive Consequences on Illicit Polydrug Use," *Drug and Alcohol Dependence*, Vol. 22, No. 1–2, October 1988, pp. 1–7. Reason for exclusion: Only measures effects of a co-intervention.
- Ilgén, M., A. Jain, H. M. Kim, and J. A. Trafton, "The Effect of Stress on Craving for Methadone Depends on the Timing of Last Methadone Dose," *Behaviour Research and Therapy*, Vol. 46, No. 10, October 2008, pp. 1170–1175. Reason for exclusion: Wrong study design.
- "Improving Treatment of Opioid Addiction and Preventing Abuse," Plainsboro, N.J., 2016, pp. 8–9.

- “In Case You Haven’t Heard,” *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 46, 2014, p. 8.
Reason for exclusion: Wrong study design.
- Ista, E., E. Wildschut, and D. Tibboel, “Creating or Preventing Opioid Addiction, Finding the Right Dose,” *Pediatric Critical Care Medicine*, Vol. 12, No. 5, September 2011, pp. 590–592. Reason for exclusion: Wrong study design.
- Jackson, G., D. Korts, R. Hanbury, V. Sturiano, L. Wolpert, M. Cohen, and B. Stimmel, “Alcohol Consumption in Persons on Methadone Maintenance Therapy,” *The American Journal of Drug and Alcohol Abuse*, Vol. 9, No. 1, 1982, pp. 69–76. Reason for exclusion: Wrong study design.
- Jaffe, J. H., S. G. Mitchell, J. Gryczynski, R. P. Schwartz, K. E. O’Grady, and Y. K. Olsen, “IOP vs. OP with Buprenorphine: 6-Month Treatment Outcomes,” *American Journal on Addictions*, Vol. 22, No. 3, 2013. Reason for exclusion: Only measures effects of a co-intervention.
- Jaffe, J. H., C. R. Schuster, B. B. Smith, and P. H. Blachley, “Comparison of Acetylmethadol and Methadone in the Treatment of Long-Term Heroin Users. A Pilot Study,” *JAMA*, Vol. 211, No. 11, March 16, 1970, pp. 1834–1836. Reason for exclusion: No functional outcomes.
- Jaffe, Jerome H., and Edward C. Senay, “Methadone and l-Methadyl Acetate: Use in Management of Narcotics Addicts,” *JAMA: Journal of the American Medical Association*, Vol. 216, No. 8, May 1971, pp. 1303–1305. Reason for exclusion: Not English.
- Jaffe, J. H., E. C. Senay, C. R. Schuster, P. F. Renault, B. Smith, and S. Dimenza, “Methadyl Acetate vs Methadone: A Double-Blind Study in Heroin Users. 1972 [Proceedings],” *NIDA Research Monograph*, No. 8, July 1976, pp. 76–78. Reason for exclusion: Conference abstract.
- Jaffe, J. H., E. C. Senay, C. R. Schuster, P. R. Renault, B. Smith, and S. DiMenza, “Methadyl Acetate vs Methadone. A Double-Blind Study in Heroin Users,” *JAMA*, Vol. 222, No. 4, October 1972, pp. 437–442. Reason for exclusion: No functional outcomes.
- Jagsch, R., W. Gombas, S. D. Schindler, H. Eder, D. E. Moody, and G. Fischer, “Opioid Plasma Concentrations in Methadone-and Buprenorphine-Maintained Patients,” *Addiction Biology*, Vol. 10, No. 4, December 2005, pp. 365–371. Reason for exclusion: No functional outcomes.
- Jaremko, K. M., R. C. Sterling, and E. J. Van Bockstaele, “Psychological and Physiological Stress Negatively Impacts Early Engagement and Retention of Opioid-Dependent Individuals on Methadone Maintenance,” *Journal of Substance Abuse Treatment*, Vol. 48, No. 1, January 2015, pp. 117–127. Reason for exclusion: Wrong study design.
- Jasinski, D. R., J. J. Boren, J. E. Henningfield, R. E. Johnson, W. R. Lange, and S. E. Lukas, “Progress Report from the NIDA Addiction Research Center, Baltimore, Maryland,” *NIDA Research Monograph*, Vol. 49, March 1984, pp. 69–76. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Jasinski, D. R., P. J. Fudala, and R. E. Johnson, "Sublingual Versus Subcutaneous Buprenorphine in Opiate Abusers," *Clinical Pharmacology Therapy*, Vol. 45, No. 5, May 1989, pp. 513–519. Reason for exclusion: Wrong setting (i.e., inpatient).
- Jasinski, D. R., J. S. Pevnick, S. C. Clark, and J. D. Griffith, "Therapeutic Usefulness of Propoxyphene Napsylate in Narcotic Addiction," *Archives of General Psychiatry*, Vol. 34, No. 2, February 1977, pp. 227–233. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Jasinski, D. R., J. S. Pevnick, and J. D. Griffith, "Human Pharmacology and Abuse Potential of the Analgesic Buprenorphine: A Potential Agent for Treating Narcotic Addiction," *Archives of General Psychiatry*, Vol. 35, No. 4, April 1978, pp. 501–516. Reason for exclusion: Wrong setting (i.e., inpatient).
- Jia, H., M. K. Greenwald, B. D. Vince, P. J. Fudala, C. Heidbreder, and A. F. Nasser, "A Multiple-Dose Study of Blockade of Opioid Subjective Effects by Subcutaneous Injections of Depot Buprenorphine in Subjects with Opioid Use Disorder," *Clinical Pharmacology and Therapeutics*, Vol. 97, Supplement 1, February 2015, p. S75. Reason for exclusion: Conference abstract.
- Jiang, G. H., Y. W. Qiu, X. L. Zhang, L. J. Han, X. F. Lv, L. M. Li, C. L. Lin, F. Z. Zhuo, S. Y. Hu, and J. Z. Tian, "Amplitude Low-Frequency Oscillation Abnormalities in the Heroin Users: A Resting State fMRI Study," *Neuroimage*, Vol. 57, No. 1, July 1, 2011, pp. 149–154. Reason for exclusion: No functional outcomes.
- Jiang, H., J. Du, F. Wu, Z. Wang, S. Fan, Z. Li, Y. I. Hser, and M. Zhao, "Efficacy of Contingency Management in Improving Retention and Compliance to Methadone Maintenance Treatment: A Random Controlled Study," *Shanghai Archives of Psychiatry*, Vol. 24, No. 1, February 2012, pp. 11–19. Reason for exclusion: Only measures effects of a co-intervention.
- Jin, H., J. H. Atkinson, N. A. Duarte, X. Yu, C. Shi, P. K. Riggs, J. Li, S. Gupta, T. Wolfson, A. F. Knight, D. Franklin, S. Letendre, Z. Wu, I. Grant, and R. K. Heaton, "Risks and Predictors of Current Suicidality in HIV-Infected Heroin Users in Treatment in Yunnan, China: A Controlled Study," *Journal of Acquired Immune Deficiency Syndrome*, Vol. 62, No. 3, March 1, 2013, pp. 311–316. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Johnson, F. K., S. Ciric, S. Boudriau, J. Kisicki, and J. Stauffer, "Effects of Alcohol on the Pharmacokinetics of Morphine Sulfate and Naltrexone Hydrochloride Extended Release Capsules," *Journal of Clinical Pharmacology*, Vol. 52, No. 5, May 2012, pp. 747–756. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Johnson, R. E., M. A. Chutuape, E. C. Strain, S. L. Walsh, M. L. Stitzer, and G. E. Bigelow, "A Comparison of Levomethadyl Acetate, Buprenorphine, and Methadone for Opioid Dependence," *New England Journal of Medicine*, Vol. 343, No. 18, November 2, 2000, pp. 1290–1297. Reason for exclusion: No functional outcomes.

- Johnson, R. E., T. Eissenberg, M. L. Stitzer, E. C. Strain, I. A. Liebson, and G. E. Bigelow, "Buprenorphine Treatment of Opioid Dependence: Clinical Trial of Daily Versus Alternate-Day Dosing," *Drug and Alcohol Dependence*, Vol. 40, No. 1, November 1995, pp. 27–35. Reason for exclusion: No functional outcomes.
- Johnson, R. E., T. Eissenberg, M. L. Stitzer, E. C. Strain, I. A. Liebson, and G. E. Bigelow, "A Placebo Controlled Clinical Trial of Buprenorphine as a Treatment for Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 40, No. 1, November 1995, pp. 17–25. Reason for exclusion: No functional outcomes.
- Johnson, R. E., P. J. Fudala, and J. H. Jaffe, "Outpatient Comparison of Buprenorphine and Methadone Maintenance. I. Effects on Opiate Use and Self-Reported Adverse Effects and Withdrawal Symptomatology," *NIDA Research Monograph*, Vol. 105, 1990, pp. 585–586. Reason for exclusion: Conference abstract.
- Johnson, R. E., J. H. Jaffe, and P. J. Fudala, "A Controlled Trial of Buprenorphine Treatment for Opioid Dependence," *JAMA*, Vol. 267, No. 20, May 27, 1992, pp. 2750–2755. Reason for exclusion: No functional outcomes.
- Johnson, Steven Ross, "Why a Proven Tool to Combat Opioid Abuse Is Underutilized by Healthcare Professionals," *Modern Healthcare*, Vol. 46, No. 14, 2016, p. 0008. Reason for exclusion: Wrong study design.
- Jones, H. E., H. Fitzgerald, and R. E. Johnson, "Males and Females Differ in Response to Opioid Agonist Medications," *The American Journal on Addictions*, Vol. 14, No. 3, May–June 2005, pp. 223–233. Reason for exclusion: No functional outcomes.
- Jones, H. E., N. A. Haug, M. L. Stitzer, and D. S. Svikis, "Improving Treatment Outcomes for Pregnant Drug-Dependent Women Using Low-Magnitude Voucher Incentives," *Addictive Behaviors*, Vol. 25, No. 2, March–April 2000, pp. 263–267. Reason for exclusion: Only measures effects of a co-intervention.
- Jones, H. E., L. M. Jansson, K. E. O'Grady, and K. Kaltenbach, "The Relationship Between Maternal Methadone Dose at Delivery and Neonatal Outcome: Methodological and Design Considerations," *Neurotoxicology and Teratology*, Vol. 39, September–October 2013, pp. 110–115. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Jones, H. E., E. C. Strain, G. E. Bigelow, S. L. Walsh, M. L. Stitzer, T. Eissenberg, and R. E. Johnson, "Induction with Levomethadyl Acetate: Safety and Efficacy," *Archives of General Psychiatry*, Vol. 55, No. 8, August 1998, pp. 729–736. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Jones, J., and S. D. Comer, "The Reinforcing and Subjective Effects of Intravenous and Intranasal Buprenorphine in Heroin Users," *Neuropsychopharmacology*, Vol. 38, 2012. Reason for exclusion: Wrong setting (i.e., inpatient).

- Jones, J. B., F. Johnson, G. Wagner, J. Stauffer, and E. Sellers, "Relative Abuse Potential of ALO-01, Extended-Release Morphine Sulfate Plus Sequestered Naltrexone," *Proceedings of the 70th Annual Scientific Meeting of the College on Problems of Drug Dependence; 2008 June 14–19; San Juan, Puerto Rico, USA*, 2008. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Jones, J., M. A. Sullivan, S. K. Vosburg, Z. Cooper, and S. D. Comer, "Comparison of the Abuse Liability of Intranasal Buprenorphine Versus Buprenorphine/Naloxone in Buprenorphine-Dependent Heroin Abusers," *Neuropsychopharmacology*, Vol. 35, 2010. Reason for exclusion: Wrong study design.
- Jones, J. D., G. Madera, and S. D. Comer, "The Reinforcing and Subjective Effects of Intravenous and Intranasal Buprenorphine in Heroin Users," *Pharmacology, Biochemistry, and Behavior*, Vol. 122, July 2014, pp. 299–306. Reason for exclusion: Wrong study design.
- Judson, B. A., T. M. Carney, and A. Goldstein, "Naltrexone Treatment of Heroin Addiction: Efficacy and Safety in a Double-Blind Dosage Comparison," *Drug and Alcohol Dependence*, Vol. 7, No. 4, July 1981, pp. 325–346. Reason for exclusion: Wrong study design.
- Judson, B. A., W. H. Horns, and A. Goldstein, "Side Effects of Levomethadone and Racemic Methadone in a Maintenance Program," *Clinical Pharmacology and Therapeutics*, Vol. 20, No. 4, October 1976, pp. 445–449. Reason for exclusion: No functional outcomes.
- Jürgens, Ralf, and Joanne Csete, "In the Name of Treatment: Ending Abuses in Compulsory Drug Detention Centers," *Addiction*, Vol. 107, No. 4, 2012, pp. 689–691. Reason for exclusion: Wrong study design.
- Kagerer, S., B. Hock, B. Wiese, R. Lehnert, and M. Soyka, "Attention, Concentration and Memory Performance in Opiate Addicts Under Methadone and Under Buprenorphine," *Suchtmedizin in Forschung Und Praxis*, Vol. 6, No. 1, 2004. Reason for exclusion: Not English.
- Kahan, M., A. Srivastava, and K. Shen, "Why We Object to NAOMI. Heroin Maintenance in Canada . . . North American Opiate Medication Initiative," *Canadian Family Physician*, Vol. 52, 2006, pp. 705–706. Reason for exclusion: Wrong study design.
- Kakko, J., L. Gronbladh, K. D. Svanborg, J. von Wachenfeldt, C. Ruck, B. Rawlings, L. H. Nilsson, and M. Heilig, "A Stepped Care Strategy Using Buprenorphine and Methadone Versus Conventional Methadone Maintenance in Heroin Dependence: A Randomized Controlled Trial," *The American Journal of Psychiatry*, Vol. 164, No. 5, May 2007, pp. 797–803. Reason for exclusion: Only measures effects of a co-intervention.
- Kakko, J., K. D. Svanborg, M. J. Kreek, and M. Heilig, "1-Year Retention and Social Function After Buprenorphine-Assisted Relapse Prevention Treatment for Heroin Dependence on Sweden: A Randomised, Placebo-Controlled Trial," *Lancet*, Vol. 361, No. 9358, February 22, 2003, pp. 662–668. Reason for exclusion: Other.

- Kamien, J. B., S. A. Branstetter, and L. Amass, “Buprenorphine-Naloxone Versus Methadone Maintenance Therapy: A Randomised Double-Blind Trial with Opioid-Dependent Patients,” *Heroin Addiction and Related Clinical Problems*, Vol. 10, No. 4, 2008. Reason for exclusion: No functional outcomes.
- Kampman, K., and M. Jarvis, “American Society of Addiction Medicine (ASAM) National Practice Guideline for the Use of Medications in the Treatment of Addiction Involving Opioid Use,” *Journal of Addiction Medicine*, Vol. 9, No. 5, September–October 2015, pp. 358–367. Reason for exclusion: Wrong study design.
- Kaplan, A., “No More Constant Craving? Opioid Dependence and XR Naltrexone,” *Psychiatric Times*, Vol. 27, No. 8, 2010, p. 8. Reason for exclusion: Wrong study design.
- Karch, S. B., “Is It Time to Reformulate Racemic Methadone?” *Journal of Addiction Medicine*, Vol. 5, No. 3, September 2011, pp. 229–231. Reason for exclusion: Wrong study design.
- Karoly, Hollis C., Sophie L. YorkWilliams, and Kent E. Hutchison, “Clinical Neuroscience of Addiction: Similarities and Differences Between Alcohol and Other Drugs,” *Alcoholism: Clinical and Experimental Research*, Vol. 39, No. 11, 2015, pp. 2073–2084. Reason for exclusion: Wrong study design.
- Karow, A., J. Reimer, I. Schafer, M. Krausz, C. Haasen, and U. Verthein, “Quality of Life Under Maintenance Treatment with Heroin Versus Methadone in Patients with Opioid Dependence,” *Drug and Alcohol Dependence*, Vol. 112, No. 3, December 1, 2010, pp. 209–215. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Karow, A., U. Verthein, R. Pukrop, J. Reimer, C. Haasen, M. Krausz, and I. Schafer, “Quality of Life Profiles and Changes in the Course of Maintenance Treatment Among 1,015 Patients with Severe Opioid Dependence,” *Substance Use and Misuse*, Vol. 46, No. 6, 2011, pp. 705–715. Reason for exclusion: No functional outcomes.
- Karp-Gelernter, E., C. Savage, and O. L. McCabe, “Evaluation of Clinic Attendance Schedules for LAAM and Methadone: A Controlled Study,” *The International Journal of the Addictions*, Vol. 17, No. 5, July 1982, pp. 805–813. Reason for exclusion: No functional outcomes.
- Karuntzos, G. T., J. M. Caddell, and M. L. Dennis, “Gender Differences in Vocational Needs and Outcomes for Methadone Treatment Clients,” *Journal of Psychoactive Drugs*, Vol. 26, No. 2, April–June 1994, pp. 173–180. Reason for exclusion: No functional outcomes.
- Kastelic, A., G. Dubajic, and E. Strbad, “Slow-Release Oral Morphine for Maintenance Treatment of Opioid Addicts Intolerant to Methadone or with Inadequate Withdrawal Suppression,” *Addiction*, Vol. 103, No. 11, November 2008, pp. 1837–1846. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Katz, J. G., “Heroin Maintenance Treatment: Its Effectiveness and the Legislative Changes Necessary to Implement It in the U.S.,” *Journal Contemporary Health and Law Policy*, Vol. 26, No. 2, Spring 2010, pp. 300–342. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kay, D. C., W. B. Pickworth, G. L. Neidert, D. Falcone, P. M. Fishman, and E. Othmer, “Opioid Effects on Computer-Derived Sleep and EEG Parameters in Nondependent Human Addicts,” *Sleep*, Vol. 2, No. 2, 1979, pp. 175–191. Reason for exclusion: Wrong study design.
- Keen, J., G. Rowse, N. Mathers, M. Campbell, and N. Seivewright, “Can Methadone Maintenance for Heroin-Dependent Patients Retained in General Practice Reduce Criminal Conviction Rates and Time Spent in Prison?” *British Journal of General Practice*, Vol. 50, No. 450, January 2000, pp. 48–49. Reason for exclusion: Wrong study design.
- Keen, M. R., A. Owen, and R. C. Hill, “Comparison of the Effects of Intramuscular Meptazinol, Methadone and Placebo in Opiate Addicts (Editors’ Abstract),” *Postgraduate Medical Journal*, Vol. 61, Supplement 2, 1985, pp. 27–28. Reason for exclusion: Wrong study design.
- Kelly, S. M., K. E. O’Grady, J. H. Jaffe, D. Gandhi, and R. P. Schwartz, “Improvements in Outcomes in Methadone Patients on Probation/Parole Regardless of Counseling Early in Treatment,” *Journal of Addiction Medicine*, Vol. 7, No. 2, March–April 2013, pp. 133–138. Reason for exclusion: Only measures effects of a co-intervention.
- Kelly, S. M., R. P. Schwartz, K. E. O’Grady, D. Gandhi, and J. H. Jaffe, “Impact of Methadone with Versus Without Drug Abuse Counseling on HIV Risk: 4- and 12-Month Findings from a Clinical Trial,” *Journal of Addiction Medicine*, Vol. 6, No. 2, June 2012, pp. 145–152. Reason for exclusion: Only measures effects of a co-intervention.
- Kelty, E., H. Ngo, and G. Hulse, “Assessing the Usefulness of Health Data Linkage in Obtaining Adverse Event Data in a Randomised Controlled Trial of Oral and Implant Naltrexone in the Treatment of Heroin Dependence,” *Clinical Trials*, Vol. 10, No. 1, February 2013, pp. 170–180. Reason for exclusion: No functional outcomes.
- Kennedy, A. P., K. A. Phillips, D. H. Epstein, D. A. Reamer, J. Schmittner, and K. L. Preston, “A Randomized Investigation of Methadone Doses at or over 100 Mg/Day, Combined with Contingency Management,” *Drug and Alcohol Dependence*, Vol. 130, No. 1–3, June 1, 2013, pp. 77–84. Reason for exclusion: No functional outcomes.
- Kerr, T., A. Wodak, R. Elliott, J. S. Montaner, and E. Wood, “Opioid Substitution and HIV/AIDS Treatment and Prevention,” *Lancet*, Vol. 364, No. 9449, November 27–December 3, 2004, pp. 1918–1919. Reason for exclusion: Only measures effects of a co-intervention.
- Kettler, L., M. Nikic, D. Newcombe, and R. Ali, “Mood Disturbance and Withdrawal Severity in Substitution Treatment for Opioid Dependence: Their Association and Impact on Continued Illicit Drug Use,” *Addictive Disorders and Their Treatment*, Vol. 9, No. 1, 2010, pp. 1–7. Reason for exclusion: Wrong study design.

- Keyser-Marcus, L., D. Miles, L. Jansson, H. Jones, and D. Svikis, "Perinatal Opiate Dependence: Methadone and Birth Outcomes," *Drug and Alcohol Dependence*, Vol. 66 Supplement 1, 2002. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Khanna, N., S. Arnold, S. Sadaphal, A. Joshi, D. Stewart, and D. Gandhi, "Nicotine Dependence and Depression Among Women Smokers on Methadone Maintenance," *European Journal of General Practice*, Vol. 16, No. 4, December 2010, pp. 222–228. Reason for exclusion: No functional outcomes.
- Khemiri, A., E. Kharitonova, V. Zah, J. Ruby, and M. Toumi, "Analysis of Buprenorphine/Naloxone Dosing Impact on Treatment Duration, Resource Use and Costs in the Treatment of Opioid-Dependent Adults: A Retrospective Study of U.S. Public and Private Health Care Claims," *Postgraduate Medicine*, Vol. 126, No. 5, September 2014, pp. 113–120. Reason for exclusion: No functional outcomes.
- Khodabandeh, F., S. Kahani, S. Shadnia, and M. Abdollahi, "Comparison of the Efficacy of Methadone Maintenance Therapy vs. Narcotics Anonymous in the Treatment of Opioid Addiction: A 2-Year Survey," *International Journal of Pharmacology*, Vol. 8, No. 5, 2012, pp. 445–449. Reason for exclusion: No functional outcomes.
- Khodadadi, Arash, Mohammad Mahdi Keramati, Amir Dezfouli, Houman Safaie, and Hamed Ekhtiari, "Analysis of Decision-Making Processes in Drug-Abusers Before and After Maintenance Treatment with Methadone," *Advances in Cognitive Science*, Vol. 12, No. 1, Spring 2010, pp. 26–42. Reason for exclusion: Wrong study design.
- Kidorf, M., R. K. Brooner, N. Gandotra, D. Antoine, V. L. King, J. Peirce, and S. Ghazarian, "Reinforcing Integrated Psychiatric Service Attendance in an Opioid-Agonist Program: A Randomized and Controlled Trial," *Drug and Alcohol Dependence*, Vol. 133, No. 1, November 1, 2013, pp. 30–36. Reason for exclusion: Only measures effects of a co-intervention.
- Kidorf, M., V. L. King, J. Peirce, N. Gandotra, S. Ghazarian, and R. K. Brooner, "Substance Use and Response to Psychiatric Treatment in Methadone-Treated Outpatients with Comorbid Psychiatric Disorder," *Journal of Substance Abuse Treatment*, Vol. 51, April 2015, pp. 64–69. Reason for exclusion: Only measures effects of a co-intervention.
- Kidorf, M., and M. L. Stitzer, "Contingent Use of Take-Homes and Split-Dosing to Reduce Illicit Drug Use of Methadone Patients," *Behavior Therapy*, Vol. 27, No. 1, 1996. Reason for exclusion: Only measures effects of a co-intervention.
- Kidorf, M., M. L. Stitzer, R. K. Brooner, and J. Goldberg, "Contingent Methadone Take-Home Doses Reinforce Adjunct Therapy Attendance of Methadone Maintenance Patients," *Drug and Alcohol Dependence*, Vol. 36, No. 3, December 1994, pp. 221–226. Reason for exclusion: Only measures effects of a co-intervention.

- Kienbaum, P., T. Heuter, M. C. Michel, N. Scherbaum, M. Gastpar, and J. Peters, “Chronic Mu-Opioid Receptor Stimulation in Humans Decreases Muscle Sympathetic Nerve Activity,” *Circulation*, Vol. 103, No. 6, Feb 13, 2001, pp. 850–855. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kienbaum, P., T. Heuter, N. Scherbaum, M. Gastpar, and J. Peters, “Chronic Mu-Opioid Receptor Stimulation Alters Cardiovascular Regulation in Humans: Differential Effects on Muscle Sympathetic and Heart Rate Responses to Arterial Hypotension,” *Journal of Cardiovascular Pharmacology*, Vol. 40, No. 3, September 2002, pp. 363–369. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kiepek, Niki, Bobbi Groom, Debbie Toppozini, Kara Kakekagumick, Jill Muileboom, and Len Kelly, “Evaluation of an Inpatient Medical Withdrawal Program in Rural Ontario: A 1-Year Prospective Study,” *Canadian Journal of Rural Medicine*, Vol. 20, No. 3, Summer 2015, pp. 92–97. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kim, S. J., L. A. Marsch, H. Guarino, M. C. Acosta, and Y. Aponte-Melendez, “Predictors of Outcome from Computer-Based Treatment for Substance Use Disorders: Results from a Randomized Clinical Trial,” *Drug and Alcohol Dependence*, Vol. 157, December 1, 2015, pp. 174–178. Reason for exclusion: Only measures effects of a co-intervention.
- Kimber, J., S. Larney, M. Hickman, D. Randall, and L. Degenhardt, “Mortality Risk of Opioid Substitution Therapy with Methadone Versus Buprenorphine: A Retrospective Cohort Study,” *Lancet Psychiatry*, Vol. 2, No. 10, October 2015, pp. 901–908. Reason for exclusion: No functional outcomes.
- King, J. B., A. M. Sainski-Nguyen, and B. K. Bellows, “Office-Based Buprenorphine Versus Clinic-Based Methadone: A Cost-Effectiveness Analysis,” *Journal of Pain and Palliative Care Pharmacotherapy*, Vol. 30, No. 1, 2016. Reason for exclusion: Wrong study design.
- King, V. L., M. S. Kidorf, K. B. Stoller, R. Schwartz, K. Kolodner, and R. K. Brooner, “A 12-Month Controlled Trial of Methadone Medical Maintenance Integrated into an Adaptive Treatment Model,” *Journal Substance Abuse Treatment*, Vol. 31, No. 4, December 2006, pp. 385–393. Reason for exclusion: Only measures effects of a co-intervention.
- King, V. L., K. B. Stoller, M. Hayes, A. Umbricht, M. Currens, M. S. Kidorf, J. A. Carter, R. Schwartz, and R. K. Brooner, “A Multicenter Randomized Evaluation of Methadone Medical Maintenance,” *Drug and Alcohol Dependence*, Vol. 65, No. 2, January 1, 2002, pp. 137–148. Reason for exclusion: Only measures effects of a co-intervention.
- Kinlock, T. W., M. S. Gordon, R. P. Schwartz, K. O’Grady, T. T. Fitzgerald, and M. Wilson, “A Randomized Clinical Trial of Methadone Maintenance for Prisoners: Results at 1-Month Post-Release,” *Drug and Alcohol Dependence*, Vol. 91, No. 2–3, December 1, 2007, pp. 220–227. Reason for exclusion: No functional outcomes.

- Kirtadze, I., D. Otiashvili, K. E. O'Grady, and H. E. Jones, "Behavioral Treatment + Naltrexone Reduces Drug Use and Legal Problems in the Republic of Georgia," *American Journal of Drug and Alcohol Abuse*, Vol. 38, No. 2, March 2012, pp. 171–175. Reason for exclusion: Only measures effects of a co-intervention.
- Kleykamp, B. A., R. G. Vandrey, G. E. Bigelow, E. C. Strain, and M. Z. Mintzer, "Effects of Methadone Plus Alcohol on Cognitive Performance in Methadone-Maintained Volunteers," *American Journal of Drug and Alcohol Abuse*, Vol. 41, No. 3 // K24 DA023186 (NIDA) *National Institute on Drug Abuse* // R01 DA17688 (NIDA) *National Institute on Drug Abuse* // T32 DA07209 (NIDA) *National Institute on Drug Abuse*, 2015. Reason for exclusion: Wrong setting (i.e., inpatient).
- Knopf, Alison, "Buprenorphine Weekly, Monthly Injection Studies Underway," *Alcoholism and Drug Abuse Weekly*, Vol. 28, No. 42, 2016, pp. 7–8. Reason for exclusion: Wrong study design.
- Kolarzyk, E., D. Targosz, D. Pach, and L. Misiolek, "Nervous Regulation of Breathing in Opiate Dependent Patient. Part II. Respiratory System Efficiency and Breathing Regulation of Persons Classified to the Methadone Maintenance Treatment," *Przegląd Lekarski*, Vol. 57, No. 10, 2000, pp. 536–538. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Kornor, H., and H. Nordvik, "Five-Factor Model Personality Traits in Opioid Dependence," *BMC Psychiatry*, Vol. 7, August 6, 2007, p. 37. Reason for exclusion: No functional outcomes.
- Kornor, H., and H. Waal, "From Opioid Maintenance to Abstinence: A Literature Review (Structured Abstract)," *Drug and Alcohol Review*, Vol. 24, No. 3, 2005. Reason for exclusion: Wrong study design.
- Kornreich, C., M. L. Foisy, P. Philippot, B. Dan, J. Tecco, X. Noel, U. Hess, I. Pelc, and P. Verbanck, "Impaired Emotional Facial Expression Recognition in Alcoholics, Opiate Dependence Subjects, Methadone Maintained Subjects and Mixed Alcohol-Opiate Antecedents Subjects Compared with Normal Controls," *Psychiatry Research*, Vol. 119, No. 3, August 1, 2003, pp. 251–260. Reason for exclusion: Wrong study design.
- Kosten, T., J. Falcioni, A. Oliveto, and A. Feingold, "Depression Predicts Higher Rates of Heroin Use on Desipramine with Buprenorphine than with Methadone," *The American Journal on Addictions*, Vol. 13, No. 2, March–April 2004, pp. 191–201. Reason for exclusion: Only measures effects of a co-intervention.
- Kosten, T. R., S. T. Higgins, S. Sigmon, and G. Badger, "Gender Effects of Desipramine on Opiate Use in Buprenorphine- vs. Methadone-Maintained Cocaine Abusers," *NIDA Research Monograph*, 2000. Reason for exclusion: Only measures effects of a co-intervention.

- Kosten, T., A. Oliveto, A. Feingold, J. Poling, K. Sevarino, E. McCance-Katz, S. Stine, G. Gonzalez, and K. Gonsai, "Desipramine and Contingency Management for Cocaine and Opiate Dependence in Buprenorphine Maintained Patients," *Drug and Alcohol Dependence*, Vol. 70, No. 3, June 5, 2003, pp. 315–325. Reason for exclusion: Only measures effects of a co-intervention.
- Kosten, T., J. Poling, and A. Oliveto, "Effects of Reducing Contingency Management Values on Heroin and Cocaine Use for Buprenorphine- and Desipramine-Treated Patients," *Addiction*, Vol. 98, No. 5, May 2003, pp. 665–671. Reason for exclusion: Only measures effects of a co-intervention.
- Kosten, T. R., "Buprenorphine for Benzodiazepine-Abusing Heroin Addicts," *American Journal of Psychiatry*, Vol. 151, No. 1, January 1994, p. 151. Reason for exclusion: Wrong study design.
- Kosten, T. R., J. H. Krystal, D. S. Charney, L. H. Price, C. H. Morgan, and H. D. Kleber, "Opioid Antagonist Challenges in Buprenorphine Maintained Patients," *Drug and Alcohol Dependence*, Vol. 25, No. 1, February 1990, pp. 73–78. Reason for exclusion: Wrong setting (i.e., inpatient).
- Kosten, T. R., C. Morgan, and H. D. Kleber, "Phase II Clinical Trials of Buprenorphine: Detoxification and Induction onto Naltrexone," *NIDA Research Monograph*, Vol. 121, 1992, pp. 101–119. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kosten, T. R., and B. S. Rayford, "Effects of Ethnicity on Low-Dose Opiate Stabilization," *Journal of Substance Abuse Treatment*, Vol. 12, No. 2, March–April 1995, pp. 111–116. Reason for exclusion: No functional outcomes.
- Kosten, T. R., R. Schottenfeld, D. Ziedonis, and J. Falcioni, "Buprenorphine Versus Methadone Maintenance for Opioid Dependence," *The Journal of Nervous and Mental Diseases*, Vol. 181, No. 6, June 1993, pp. 358–364. Reason for exclusion: No functional outcomes.
- Kouri, E. M., S. E. Lukas, and J. H. Mendelson, "P300 Assessment of Opiate and Cocaine Users: Effects of Detoxification and Buprenorphine Treatment," *Biological Psychiatry*, Vol. 40, No. 7, October 1, 1996, pp. 617–628. Reason for exclusion: Wrong setting (i.e., inpatient).
- Kovacs, S. J., J. Ke, A. Barve, Y. Cheng, J. Zhang, R. Maietta, G. Sunkara, and D. S. Stein, "Effect of Therapeutic Doses of Alisporivir (ALV) on Methadone Pharmacokinetics in Healthy Subjects and Patients on Stable Methadone Maintenance Therapy (MMT)," *Journal of Hepatology*, Vol. 58, 2013. Reason for exclusion: Only measures effects of a co-intervention.

- Kovacs, S. J., J. Ke, A. Barve, L. R. Webster, Y. Cheng, J. Zhang, R. Maietta, G. Sunkara, and D. S. Stein, "Effect of Alisporivir (ALV) on the Pharmacokinetics (PK), Safety, and Tolerability of Methadone in Healthy Subjects and Opioid-Dependent Patients on Stable Methadone Maintenance Therapy (MMT)," *Journal of Hepatology*, Vol. 60, No. 1, Supplement 1, 2014. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kowalczyk, W. J., M. A. Furnari, K. A. Phillips, M. L. Jobes, U. Ghitza, D. Epstein, and K. L. Preston, "Reducing the Cost of Free Time: Treatment Success in a Randomized Trial of Clonidine as Adjunct to Buprenorphine Maintenance Is Associated with More Leisure Activities in the Clonidine Condition," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Kowalczyk, W. J., K. A. Phillips, M. L. Jobes, A. P. Kennedy, U. E. Ghitza, D. A. Agage, J. P. Schmittner, D. H. Epstein, and K. L. Preston, "Clonidine Maintenance Prolongs Opioid Abstinence and Decouples Stress from Craving in Daily Life: A Randomized Controlled Trial with Ecological Momentary Assessment," *The American Journal of Psychiatry*, Vol. 172, No. 8, August 1, 2015, pp. 760–767. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kraft, W. K., K. Dysart, J. S. Greenspan, E. Gibson, K. Kaltenbach, and M. E. Ehrlich, "Revised Dose Schema of Sublingual Buprenorphine in the Treatment of the Neonatal Opioid Abstinence Syndrome," *Addiction*, Vol. 106, No. 3, March 2011, pp. 574–580. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Krantz, Mori J., "Heterogeneous Impact of Methadone on the QTc Interval: What Are the Practical Implications?" *Journal of Addictive Diseases*, Vol. 27, No. 4, 2008, pp. 5–9. Reason for exclusion: Wrong study design
- Krantz, M. J., and P. S. Mehler, "QTc Prolongation: Methadone's Efficacy-Safety Paradox," *Lancet*, Vol. 368, No. 9535, August 12, 2006, pp. 556–557. Reason for exclusion: Wrong study design.
- Krebs, E., T. Kerr, J. Montaner, E. Wood, and B. Nosyk, "Dynamics in the Costs of Criminality Among Opioid Dependent Individuals," *Drug and Alcohol Dependence*, Vol. 144, November 1, 2014, pp. 193–200. Reason for exclusion: Wrong study design.
- Kreek, M. J., R. Jones, H. D. Kleber, T. Kosten, and C. O'Brien, "NIDA's Commitment to Tackling Drug Abuse," *Nature*, Vol. 430, No. 7003, August 26, 2004, p. 965. Reason for exclusion: Wrong study design.
- Krishnan, S., A. Salter, T. Sullivan, M. Gentgall, J. White, and P. Rolan, "Comparison of Pain Models to Detect Opioid-Induced Hyperalgesia," *Journal of Pain Research*, Vol. 5, 2012, pp. 99–106. Reason for exclusion: Wrong study design.

- Kristensen, O., O. Espegren, R. Asland, E. Jakobsen, O. Lie, and S. Seiler, "A Randomised Clinical Trial of Methadone vs. Buprenorphine to Opioid Dependents. [Norwegian]," *Tidsskrift for den Norske laegeforening*, Vol. 125, No. 2, 2005. Reason for exclusion: Not English.
- Krook, A. L., O. Brors, J. Dahlberg, K. Grouff, P. Magnus, E. Roysamb, and H. Waal, "A Placebo-Controlled Study of High Dose Buprenorphine in Opiate Dependents Waiting for Medication-Assisted Rehabilitation in Oslo, Norway," *Addiction*, Vol. 97, No. 5, May 2002, pp. 533–542. Reason for exclusion: No functional outcomes.
- Krupitsky, E. M., E. A. Blokhina, E. V. Verbitskaya, O. V. Mamontova, E. E. Zvartau, and J. Tiihonen, "Naltrexone Implant Treatment of Polydrug Dependence," *Zhurnal Nevrologii i Psikiatrii imeni S.S. Korsakova*, Vol. 2014, No. 5, 2014. Reason for exclusion: Not English.
- Krupitsky, E., A. Kibitov, E. Zvartau, E. Blokhina, E. Verbitskaya, Palatkin, E. Nielsen, T. Kosten, and G. E. Woody, "Pharmacogenetics of Treatment of Opioid Dependence with Oral Naltrexone and Long-Acting Sustained-Release Naltrexone Implant," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- Krupitsky, E., E. V. Nunes, W. Ling, D. R. Gastfriend, A. Memisoglu, and B. L. Silverman, "Injectable Extended-Release Naltrexone (XR-NTX) for Opioid Dependence: Long-Term Safety and Effectiveness," *Addiction*, Vol. 108, No. 9, September 2013, pp. 1628–1637. Reason for exclusion: No functional outcomes.
- Krupitsky, E., E. V. Nunes, W. Ling, A. Illeperuma, D. R. Gastfriend, and B. L. Silverman, "Injectable Extended-Release Naltrexone for Opioid Dependence: A Double-Blind, Placebo-Controlled, Multicentre Randomised Trial," *Lancet*, Vol. 377, No. 9776, April 30, 2011, pp. 1506–1513. Reason for exclusion: No functional outcomes
- Krupitsky, E., G. E. Woody, E. Zvartau, D. Mazalov, M. Tsoi, V. Egorova, A. Burakov, T. Didenko, T. Romanova, E. Verbitskaya, A. Bespalov, N. Neznanov, T. Slavina, A. Grinenko, and C. O'Brien, "A Placebo-Controlled Trial of Naltrexone and Fluoxetine for Opioid Addiction: Analysis of Medication Responders," *Proceedings of the 69th Annual Scientific Meeting of the College on Problems of Drug Dependence; 2007 June 16–21; Quebec City, Canada*, 2007. Reason for exclusion: Only measures effects of a co-intervention.
- Krupitsky, E., G. E. Woody, E. Zvartau, and C. P. O'Brien, "Addiction Treatment in Russia," *Lancet*, Vol. 376, No. 9747, October 2, 2010, p. 1145. Reason for exclusion: Wrong study design.
- Krupitsky, E., E. Zvartau, E. Blokhina, D. Nielsen, T. Kosten, and G. Woody, "Pharmacogenetics of Treatment of Opioid Dependence with Oral Naltrexone and Long Acting Sustained Release Naltrexone Implant," *European Neuropsychopharmacology*, Vol. 23, 2013. Reason for exclusion: Conference abstract.

- Krupitsky, E., E. Zvartau, E. Blokhina, E. Verbitskaya, G. Sulimov, and G. Woody, “Anhedonia, Depression, Anxiety, and Craving for Opiates in Opiate Addicts Stabilized on Oral Naltrexone and Long Acting Naltrexone Implant,” *European Neuropsychopharmacology*, Vol. 25, 2015. Reason for exclusion: Conference abstract.
- Krupitsky, E., E. Zvartau, E. Blokhina, E. Verbitskaya, M. Tsoy, V. Wahlgren, A. Burakov, D. Masalov, T. N. Romanova, V. Palatkin, A. Tyurina, T. Yaroslavtseva, R. Sinha, and T. R. Kosten, “Naltrexone with or Without Guanfacine for Preventing Relapse to Opiate Addiction in St.-Petersburg, Russia,” *Drug and Alcohol Dependence*, Vol. 132, No. 3, October 1, 2013, pp. 674–680. Reason for exclusion: Only measures effects of a co-intervention.
- Krupitsky, E., E. Zvartau, E. Blokhina, E. Verbitskaya, V. Wahlgren, M. Tsoy-Podosenin, N. Bushara, A. Burakov, D. Masalov, T. Romanova, A. Tyurina, V. Palatkin, T. Slavina, A. Pecoraro, and G. E. Woody, “Randomized Trial of Long-Acting Sustained-Release Naltrexone Implant vs Oral Naltrexone or Placebo for Preventing Relapse to Opioid Dependence,” *Archives of General Psychiatry*, Vol. 69, No. 9, September 2012, pp. 973–981. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Krupitsky, Evgeny, Edwin Zvartau, Elena Blokhina, Elena Verbitskaya, Valentina Wahlgren, Marina Tsoy-Podosenin, Natalia Bushara, Andrey Burakov, Dmitry Masalov, Tatyana Romanova, Arina Tyurina, Vladimir Palatkin, Tatyana Yaroslavtseva, Anna Pecoraro, and George Woody, “Anhedonia, Depression, Anxiety, and Craving in Opiate Dependent Patients Stabilized on Oral Naltrexone or an Extended Release Naltrexone Implant,” *The American Journal of Drug and Alcohol Abuse*, Vol. 42, No. 5, September 2016, pp. 614–620. Reason for exclusion: No functional outcomes.
- Krupitsky, E., E. Zvartau, D. Masalov, M. Tsoi, A. Burakov, T. Didenko, T. Romanova, V. Egorova, E. Ivanova, A. Beshpalov, N. Neznanov, A. Martynikhin, A. Grinenko, and G. Woody, “Double-Blind Placebo-Controlled Randomized Clinical Trial of Naltrexone for Heroin Addiction and HIV Risk Reduction in Russia,” *Drug and Alcohol Dependence*, Vol. 66, Supplement 1, 2002. Reason for exclusion: Conference abstract.
- Krupitsky, E. M., E. E. Zvartau, N. G. Neznanov, A. M. Burakov, D. V. Masalov, S. M. Tsoi, T. Y. Didenko, T. N. Romanova, E. B. Ivanova, Egorova Vy, Y. Beshpalov, A. V. Martynikhin, A. Y. Grinenko, and G. Woody, “A Double-Blind Placebo Controlled Clinical Trial of Naltrexone for Heroin Addiction in Russia: Sample Characteristics and Short Term Follow-Up,” *Drug and Alcohol Dependence*, Vol. 63 Supplement 1, 2001. Reason for exclusion: Conference abstract.
- Krupitsky, E., E. Zvartau, E. Verbitskaya, V. Egorova, D. Masalov, and M. Tsoi, “Long-Acting vs. Oral Naltrexone for Preventing Heroin Addiction Relapse,” *Proceedings of the 71th Annual Scientific Meeting of the College on Problems of Drug Dependence; 2009 June 20–25; Reno/Sparks, Nevada, USA*, 2009. Reason for exclusion: Conference abstract.

- Krupitsky, E., E. E. Zvartau, and G. Woody, "Long Acting Naltrexone Implants for Heroin Dependence," *European Neuropsychopharmacology*, Vol. 19, 2009. Reason for exclusion: No functional outcomes.
- Krupitsky, E., E. Zvartau, and G. Woody, "Use of Naltrexone to Treat Opioid Addiction in a Country in Which Methadone and Buprenorphine Are Not Available," *Current Psychiatry Reports*, Vol. 12, No. 5, October 2010, pp. 448–453. Reason for exclusion: Wrong study design.
- Krupitsky, E. M., and E. A. Blokhina, "Long-Acting Depot Formulations of Naltrexone for Heroin Dependence: A Review," *Current Opinion in Psychiatry*, Vol. 23, No. 3, 2010, pp. 210–214. Reason for exclusion: Wrong study design.
- Krupitsky, E. M., E. E. Zvartau, D. V. Masalov, M. V. Tsoi, A. M. Burakov, V. Y. Egorova, T. Y. Didenko, T. N. Romanova, E. B. Ivanova, A. Y. Bespalov, E. V. Verbitskaya, N. G. Neznanov, A. Y. Grinenko, C. P. O'Brien, and G. E. Woody, "Naltrexone for Heroin Dependence Treatment in St. Petersburg, Russia," *Journal of Substance Abuse Treatment*, Vol. 26, No. 4, June 2004, pp. 285–294. Reason for exclusion: No functional outcomes.
- Krupitsky, E. M., E. E. Zvartau, D. V. Masalov, M. V. Tsoy, A. M. Burakov, V. Y. Egorova, T. Y. Didenko, T. N. Romanova, E. B. Ivanova, A. Y. Bespalov, E. V. Verbitskaya, N. G. Neznanov, A. Y. Grinenko, C. P. O'Brien, and G. E. Woody, "Naltrexone with or Without Fluoxetine for Preventing Relapse to Heroin Addiction in St. Petersburg, Russia," *Journal of Substance Abuse Treatment*, Vol. 31, No. 4, December 2006, pp. 319–328. Reason for exclusion: Only measures effects of a co-intervention.
- Krystal, J. H., S. W. Woods, T. R. Kosten, M. I. Rosen, J. P. Seibyl, C. C. van Dyck, L. H. Price, I. G. Zubal, P. B. Hoffer, and D. S. Charney, "Opiate Dependence and Withdrawal: Preliminary Assessment Using Single Photon Emission Computerized Tomography (SPECT)," *The American Journal of Drug and Alcohol Abuse*, Vol. 21, No. 1, February 1995, pp. 47–63. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kuehn, B. M., "Patients Warned About Risks of Drugs Used for Analgesia, Fevers, Addiction," *JAMA*, Vol. 301, No. 22, June 10, 2009, pp. 2315–2316. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kuehn, B. M., "Buprenorphine May Boost HIV Treatment," *JAMA*, Vol. 304, No. 3, July 21, 2010, pp. 261–263. Reason for exclusion: Only measures effects of a co-intervention.
- Kuhlman, J. J., Jr., B. Levine, R. E. Johnson, P. J. Fudala, and E. J. Cone, "Relationship of Plasma Buprenorphine and Norbuprenorphine to Withdrawal Symptoms During Dose Induction, Maintenance and Withdrawal from Sublingual Buprenorphine," *Addiction*, Vol. 93, No. 4, April 1998, pp. 549–559. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Kuhn, S., M. Schu, I. Vogt, M. Schmid, R. Simmedinger, G. Schlanstedt, G. Farnbacher, U. Verthein, and C. Haasen, "Psychosocial Care in the German Model Project on Heroin-Maintenance Therapy for Opiate Dependence. [German]," *Sucht*, Vol. 53, No. 5, 2007. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Kumari, S., P. Manalai, S. Leong, A. Wooditch, M. Malik, and W. B. Lawson, "Factors Associated with Non-Adherence to Buprenorphine-Naloxone Among Opioid Dependent African-Americans: A Retrospective Chart Review," *The American Journal on Addictions*, Vol. 25, No. 2, March 2016, pp. 110–117. Reason for exclusion: Wrong study design.
- Kunoe, N., P. Lobmaier, J. K. Vederhus, B. Hjerkin, S. Hegstad, M. Gossop, O. Kristensen, and H. Waal, "Naltrexone Implants After In-Patient Treatment for Opioid Dependence: Randomised Controlled Trial," *British Journal of Psychiatry*, Vol. 194, No. 6, June 2009, pp. 541–546. Reason for exclusion: No functional outcomes.
- Kunoe, N., A. Opheim, K. K. Solli, Z. Gaulen, K. Sharma-Haase, Z.-E. Latif, and L. Tanum, "Design of a Randomized Controlled Trial of Extended-Release Naltrexone Versus Daily Buprenorphine-Naloxone for Opioid Dependence in Norway (NTX-SBX)," *BMC Pharmacology and Toxicology*, Vol. 17, No. 1, 2016. Reason for exclusion: Wrong study design.
- Kwiatkowski, C. F., and R. E. Booth, "Methadone Maintenance as HIV Risk Reduction with Street-Recruited Injecting Drug Users," *Journal of Acquired Immune Deficiency Syndrome*, Vol. 26, No. 5, April 15, 2001, pp. 483–489. Reason for exclusion: Only measures effects of a co-intervention.
- Lacroix, I., A. Berrebi, L. Schmitt, D. Garripuy, M. Lapeyre-Mestre, J. L. Montastruc, and C. Damase-Michel, "Buprenorphine High Dosage in Pregnancy: First Data of a Prospective Study," *Drug and Alcohol Dependence*, Vol. 66, Supplement 1, 2002. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Lamas, X., M. Farre, and J. Cami, "Acute Effects of Pentazocine, Naloxone and Morphine in Opioid-Dependent Volunteers," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 268, No. 3, March 1994, pp. 1485–1492. Reason for exclusion: Wrong study design.
- Lamas, X., M. Farre, T. Teran, B. Ugena, and J. Cami, "Comparative Effects of Pentazocine, Naloxone and Morphine in Opioid-Dependent Human Subjects," *NIDA Research Monograph Series*, Vol. 132, 1993. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Landabaso, M. A., I. Iraurgi, J. M. Jimenez-Lerma, J. Sanz, B. Fernandez de Corres, K. Araluce, R. Calle, and M. Gutierrez-Fraile, "A Randomized Trial of Adding Fluoxetine to a Naltrexone Treatment Programme for Heroin Addicts," *Addiction*, Vol. 93, No. 5, May 1998, pp. 739–744. Reason for exclusion: Only measures effects of a co-intervention.

- Landau, C. J., W. D. Carr, A. J. Razzetti, N. E. Sessler, C. Munera, and S. R. Ripa, "Buprenorphine Transdermal Delivery System in Adults with Persistent Noncancer-Related Pain Syndromes Who Require Opioid Therapy: A Multicenter, 5-Week Run-In and Randomized, Double-Blind Maintenance-of-Analgesia Study," *Clinical Therapeutics*, Vol. 29, No. 10, October 2007, pp. 2179–2193. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Lange, W. R., P. J. Fudala, E. M. Dax, and R. E. Johnson, "Safety and Side-Effects of Buprenorphine in the Clinical Management of Heroin Addiction," *Drug and Alcohol Dependence*, Vol. 26, No. 1, 1990. Reason for exclusion: Wrong study design.
- Langendam, M. W., G. H. van Brussel, R. A. Coutinho, and E. J. van Ameijden, "Methadone Maintenance and Cessation of Injecting Drug Use: Results from the Amsterdam Cohort Study," *Addiction*, Vol. 95, No. 4, April 2000, pp. 591–600. Reason for exclusion: Wrong study design.
- Langleben, D. D., E. L. Busch, C. P. O'Brien, and I. Elman, "Depot Naltrexone Decreases Rewarding Properties of Sugar in Patients with Opioid Dependence," *Psychopharmacology (Berl)*, Vol. 220, No. 3, April 2012, pp. 559–564. Reason for exclusion: Wrong study design.
- Langleben, D. D., K. Ruparel, I. Elman, S. Busch-Winokur, R. Pratiwadi, J. Loughhead, C. P. O'Brien, and A. R. Childress, "Acute Effect of Methadone Maintenance Dose on Brain fMRI Response to Heroin-Related Cues," *American Journal of Psychiatry*, Vol. 165, No. 3, 2008, pp. 390–394. Reason for exclusion: Wrong study design.
- Lapeyre-Mestre, M., "Supervised Prescription of Methadone Plus Heroin Is More Effective than Methadone Alone," *Evidence-Based Healthcare*, Vol. 8, No. 2, April 2004, pp. 110–111. Reason for exclusion: Only measures effects of a co-intervention.
- Lapeyre-Mestre, M., M. E. Llau, M. Gony, A. M. Navel, J. Bez, M. Grau, and J. L. Montastruc, "Opiate Maintenance with Buprenorphine in Ambulatory Care: A 24-Week Follow-Up Study of New Users," *Drug and Alcohol Dependence*, Vol. 72, No. 3, December 11, 2003, pp. 297–303. Reason for exclusion: No functional outcomes.
- Law, F. D., J. S. Myles, and D. J. Nutt, "BD rct of Suboxone (buprenorphine/naloxone) vs. Methadone/Lofexidine for Stabilization and Withdrawal of Low-Dose Outpatient Opiate Addicts," *Drug and Alcohol Dependence*, Vol. 66, Supplement 1, 2002. Reason for exclusion: Conference abstract.
- Lawental, M., "Effectiveness of Rapid Intake into Methadone Treatment: A Natural Experiment in Israel," *European Addiction Research*, Vol. 21, No. 4, 2015, pp. 211–216. Reason for exclusion: Wrong study design.
- Lawford, B. R., R. M. Young, E. P. Noble, J. Sargent, J. Rowell, S. Shadforth, X. Zhang, and T. Ritchie, "The D(2) Dopamine Receptor A(1) Allele and Opioid Dependence: Association with Heroin Use and Response to Methadone Treatment," *American Journal of Medical Genetics*, Vol. 96, No. 5, October 9, 2000, pp. 592–598. Reason for exclusion: Wrong study design.

- Lee, C. E., "Tackling Subutex Abuse in Singapore," *Singapore Medical Journal*, Vol. 47, No. 11, November 2006, pp. 919–921. Reason for exclusion: Wrong study design.
- Lee, J. D., P. D. Friedmann, T. W. Kinlock, E. V. Nunes, M. S. Gordon, and C. P. O'Brien, "Extended-Release Naltrexone for Opioid Relapse Prevention Among Opioid-Dependent, Criminal Justice-Involved Adults," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Lee, Joshua D., Ryan McDonald, Ellie Grossman, Jennifer McNeely, Eugene Laska, John Rotrosen, and Marc N. Gourevitch, "Opioid Treatment at Release from Jail Using Extended-Release Naltrexone: A Pilot Proof-of-Concept Randomized Effectiveness Trial," *Addiction*, Vol. 110, No. 6, June 2015, pp. 1008–1014. Reason for exclusion: No functional outcomes.
- Lee, S. Y., S. L. Chen, Y. H. Chang, P. S. Chen, S. Y. Huang, N. S. Tzeng, L. J. Wang, I. H. Lee, T. Y. Wang, K. C. Chen, Y. K. Yang, J. S. Hong, and R. B. Lu, "Low-Dose Memantine Attenuated Methadone Dose in Opioid-Dependent Patients: A 12-Week Double-Blind Randomized Controlled Trial," *Scientific Reports*, Vol. 5, May 19, 2015, p. 10140. Reason for exclusion: Only measures effects of a co-intervention.
- Lee, S. Y., S. L. Chen, Y. H. Chang, C. H. Chu, S. H. Chen, P. S. Chen, S. Y. Huang, N. S. Tzeng, L. J. Wang, I. H. Lee, T. Y. Wang, K. C. Chen, Y. K. Yang, J. S. Hong, and R. B. Lu, "A Placebo-Controlled Trial of Dextromethorphan as an Adjunct in Opioid-Dependent Patients Undergoing Methadone Maintenance Treatment," *International Journal of Neuropsychopharmacology*, Vol. 18, No. 7, February 25, 2015, p. pyv008. Reason for exclusion: Only measures effects of a co-intervention.
- Leece, P., C. Cavacuiti, E. M. Macdonald, T. Gomes, M. Kahan, A. Srivastava, L. Steele, J. Luo, M. M. Mamdani, and D. N. Juurlink, "Predictors of Opioid-Related Death During Methadone Therapy," *Journal of Substance Abuse Treatment*, Vol. 57, October 2015, pp. 30–35. Reason for exclusion: Wrong study design.
- Lehotay, D. C., S. George, M. L. Etter, K. Graybiel, J. C. Eichhorst, B. Fern, W. Wildenboer, P. Selby, and B. Kapur, "Free and Bound Enantiomers of Methadone and Its Metabolite, EDDP in Methadone Maintenance Treatment: Relationship to Dosage," *Clinical Biochemistry*, Vol. 38, No. 12, December 2005, pp. 1088–1094. Reason for exclusion: Wrong study design.
- Lenne, M. G., P. Dietze, G. R. Rumbold, J. R. Redman, and T. J. Triggs, "The Effects of the Opioid Pharmacotherapies Methadone, LAAM and Buprenorphine, Alone and in Combination with Alcohol, on Simulated Driving," *Drug and Alcohol Dependence*, Vol. 72, No. 3, December 11, 2003, pp. 271–278. Reason for exclusion: Wrong study design.
- Lerner, A., M. Sigal, A. Bacalu, R. Shiff, I. Burganski, and M. Gelkopf, "A Naltrexone Double Blind Placebo Controlled Study in Israel," *The Israel Journal of Psychiatry and Related Sciences*, Vol. 29, No. 1, 1992, pp. 36–43. Reason for exclusion: No functional outcomes.

- Lerner, A. G., M. Gelkopf, I. Oyffe, and M. Sigal, "Home-Based Inpatient Treatment vs. Outpatient Day Clinic Treatment: A Preliminary Report in Opiate-Dependent Patients," *The Journal of Nervous and Mental Diseases*, Vol. 183, No. 11, November 1995, p. 715. Reason for exclusion: Conference abstract.
- Lerner, Arturo G., Marc Gelkopf, Mircea Sigal, and Igor Oyffe, "Indicators of Good Prognosis in Naltrexone Treatment: A Five-Year Prospective Study," *Addiction Research S2—Addiction Research and Theory*, Vol. 4, No. 4, February 1997, pp. 385–391. Reason for exclusion: Wrong study design.
- Leslie, D. L., W. Milchak, D. R. Gastfriend, P. L. Herschman, E. O. Bixler, D. L. Velott, and R. E. Meyer, "Effects of Injectable Extended-Release Naltrexone (XR-NTX) for Opioid Dependence on Residential Rehabilitation Outcomes and Early Follow-Up," *The American Journal on Addictions*, Vol. 24, No. 3, April 2015, pp. 265–270. Reason for exclusion: No functional outcomes.
- Levrán, O., E. Peles, S. Hamon, M. Randesi, M. Adelson, and M. J. Kreek, "CYP2B6 SNPs Are Associated with Methadone Dose Required for Effective Treatment of Opioid Addiction," *Addiction Biology*, Vol. 18, No. 4, July 2013, pp. 709–716. Reason for exclusion: Wrong study design.
- Li, L., Z. Wu, L. J. Liang, C. Lin, L. Zhang, S. Guo, K. Rou, and J. Li, "An Intervention Targeting Service Providers and Clients for Methadone Maintenance Treatment in China: A Cluster-Randomized Trial," *Addiction*, Vol. 108, No. 2, February 2013, pp. 356–366. Reason for exclusion: Only measures effects of a co-intervention.
- Li, Q., W. Li, H. Wang, Y. Wang, Y. Zhang, J. Zhu, Y. Zheng, D. Zhang, L. Wang, Y. Li, X. Yan, H. Chang, M. Fan, Z. Li, J. Tian, M. S. Gold, W. Wang, and Y. Liu, "Predicting Subsequent Relapse by Drug-Related Cue-Induced Brain Activation in Heroin Addiction: An Event-Related Functional Magnetic Resonance Imaging Study," *Addiction Biology*, Vol. 20, No. 5, 61131003 (NSFC) *National Natural Science Foundation of China* // 81071142 (NSFC) *National Natural Science Foundation of China* // 81071143 (NSFC) *National Natural Science Foundation of China* // 81201081 (NSFC) *National Natural Science Foundation of China* // 81271549 (NSFC) *National Natural Science Foundation of China* // 81371532 (NSFC) *National Natural Science Foundation of China*, 2015. Reason for exclusion: No functional outcomes.
- Liang, C. S., P. S. Ho, C. H. Yen, Y. W. Yeh, S. C. Kuo, C. C. Huang, C. Y. Chen, M. C. Shih, K. H. Ma, and S. Y. Huang, "Reduced Striatal Dopamine Transporter Density Associated with Working Memory Deficits in Opioid-Dependent Male Subjects: A SPECT Study," *Addiction Biology*, Vol. 21, No. 1, January 2016, pp. 196–204. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Liao, D. L., C. Y. Huang, S. Hu, S. C. Fang, C. S. Wu, W. T. Chen, T. S. Lee, P. C. Chen, and C. S. Li, "Cognitive Control in Opioid Dependence and Methadone Maintenance Treatment," *PLoS One*, Vol. 9, No. 4, 2014, p. e94589. Reason for exclusion: Wrong study design.

- Lidz, V., D. M. Sorrentino, L. Robison, and S. Bunce, "Learning from Disappointing Outcomes: An Evaluation of Prevocational Interventions for Methadone Maintenance Patients," *Substance Use and Misuse*, Vol. 39, No. 13–14, 2004, pp. 2287–2308. Reason for exclusion: Only measures effects of a co-intervention.
- Liebschutz, J. M., D. Crooks, D. S. Herman, B. J. Anderson, L. Meshesha, S. Dossabhoy, and M. D. Stein, "Initiating Buprenorphine Maintenance for Opiate-Dependent Hospitalized Patients: A Randomized Controlled Trial," *Journal of General Internal Medicine*, Vol. 28, 2013. Reason for exclusion: Conference abstract.
- Lin, C. Y., K. C. Chang, J. D. Wang, and L. J. Lee, "Quality of Life and Its Determinants for Heroin Addicts Receiving a Methadone Maintenance Program: Comparison with Matched Referents from the General Population," *Journal of the Formosan Medical Association*, Vol. 115, No. 9, September 2016, pp. 714–727. Reason for exclusion: Wrong study design.
- Lin, S. H., K. C. Chen, S. Y. Lee, C. Y. Hsiao, I. H. Lee, T. L. Yeh, P. S. Chen, R. B. Lu, and Y. K. Yang, "The Economic Cost of Heroin Dependency and Quality of Life Among Heroin Users in Taiwan," *Psychiatry Research*, Vol. 209, No. 3, October 30, 2013, pp. 512–517. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Lin, S. H., L. T. Lee, H. C. Tsai, K. C. Chen, W. T. Chen, I. H. Lee, R. B. Lu, P. S. Chen, and Y. K. Yang, "Association Between Blood Level of Plasma Oxytocin and Novelty Seeking Among Methadone-Maintained Heroin Users," *Neuropsychobiology*, Vol. 71, No. 2, April 9, 2015, pp. 65–69. Reason for exclusion: Only measures effects of a co-intervention.
- Lin, W. C., K. H. Chou, C. C. Chen, C. C. Huang, H. L. Chen, C. H. Lu, S. H. Li, Y. L. Wang, Y. F. Cheng, and C. P. Lin, "White Matter Abnormalities Correlating with Memory and Depression in Heroin Users Under Methadone Maintenance Treatment," *PLoS One*, Vol. 7, No. 4, 2012, p. e33809. Reason for exclusion: Wrong study design.
- Lin, W. C., K. H. Chou, H. L. Chen, C. C. Huang, C. H. Lu, S. H. Li, Y. L. Wang, Y. F. Cheng, C. P. Lin, and C. C. Chen, "Structural Deficits in the Emotion Circuit and Cerebellum Are Associated with Depression, Anxiety and Cognitive Dysfunction in Methadone Maintenance Patients: A Voxel-Based Morphometric Study," *Psychiatry Research*, Vol. 201, No. 2, February 28, 2012, pp. 89–97. Reason for exclusion: No functional outcomes.
- Lindgren, Britt-Marie, Margita Eklund, Ylva Melin, and Ulla Hällgren Graneheim, "From Resistance to Existence—Experiences of Medication-Assisted Treatment as Disclosed by People with Opioid Dependence," *Issues in Mental Health Nursing*, Vol. 36, No. 12, 2015, pp. 963–970. Reason for exclusion: Wrong study design.
- Ling, W., M. Blakis, E. D. Holmes, C. J. Klett, and W. E. Carter, "Restabilization with Methadone After Methadyl Acetate Maintenance," *Archives of General Psychiatry*, Vol. 37, No. 2, February 1980, pp. 194–196. Reason for exclusion: Wrong study design.

- Ling, W., C. Charuvastra, J. F. Collins, S. Batki, L. S. Brown, Jr., P. Kintaudi, D. R. Wesson, L. McNicholas, D. J. Tusel, U. Malkerneker, J. A. Renner, Jr., E. Santos, P. Casadonte, C. Fye, S. Stine, R. I. Wang, and D. Segal, "Buprenorphine Maintenance Treatment of Opiate Dependence: A Multicenter, Randomized Clinical Trial," *Addiction*, Vol. 93, No. 4, April 1998, pp. 475–486. Reason for exclusion: No functional outcomes.
- Ling, W., C. Charuvastra, S. C. Kaim, and C. J. Klett, "Methadyl Acetate and Methadone as Maintenance Treatments for Heroin Addicts. A Veterans Administration Cooperative Study," *Archives of General Psychiatry*, Vol. 33, No. 6, June 1976, pp. 709–720. Reason for exclusion: No functional outcomes.
- Ling, W., M. Hillhouse, A. Ang, J. Jenkins, and J. Fahey, "Comparison of Behavioral Treatment Conditions in Buprenorphine Maintenance," *Addiction*, Vol. 108, No. 10, October 2013, pp. 1788–1798. Reason for exclusion: Only measures effects of a co-intervention.
- Ling, W., M. Hillhouse, C. Domier, G. Doraimani, J. Hunter, C. Thomas, J. Jenkins, A. Hasson, J. Annon, A. Saxon, J. Selzer, J. Boverman, and R. Bilangi, "Buprenorphine Tapering Schedule and Illicit Opioid Use," *Addiction*, Vol. 104, No. 2, February 2009, pp. 256–265. Reason for exclusion: No functional outcomes.
- Ling, Walter, Maureen P. Hillhouse, Andrew J. Saxon, Larissa J. Mooney, Christie M. Thomas, Alfonso Ang, Abigail G. Matthews, Albert Hasson, Jeffrey Annon, Steve Sparenborg, David S. Liu, Jennifer McCormack, Sarah Church, William Swafford, Karen Drexler, Carolyn Schuman, Stephen Ross, Katharina Wiest, P. Todd Korthuis, William Lawson, Gregory S. Brigham, Patricia C. Knox, Michael Dawes, and John Rotrosen, "Buprenorphine + Naloxone Plus Naltrexone for the Treatment of Cocaine Dependence: The Cocaine Use Reduction with Buprenorphine (CURB) Study," *Addiction*, Vol. 111, No. 8, August 2016, 2016, pp. 1416–1427. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Ling, W., D. R. Wesson, C. Charuvastra, and C. J. Klett, "A Controlled Trial Comparing Buprenorphine and Methadone Maintenance in Opioid Dependence," *Archives of General Psychiatry*, Vol. 53, No. 5, May 1996, pp. 401–407. Reason for exclusion: No functional outcomes.
- Lintzeris, N., S. Y. Leung, A. J. Dunlop, B. Larance, N. White, G. R. Rivas, R. M. Holland, L. Degenhardt, P. Muhleisen, M. Hurley, and R. Ali, "A Randomised Controlled Trial of Sublingual Buprenorphine-Naloxone Film Versus Tablets in the Management of Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 131, No. 1–2, July 1, 2013, pp. 119–126. Reason for exclusion: No functional outcomes.
- Lintzeris, N., T. B. Mitchell, A. Bond, L. Nestor, and J. Strang, "Interactions on Mixing Diazepam with Methadone or Buprenorphine in Maintenance Patients," *Journal of Clinical Psychopharmacology*, Vol. 26, No. 3, June 2006, pp. 274–283. Reason for exclusion: Only measures effects of a co-intervention.

- Lintzeris, N., T. B. Mitchell, A. J. Bond, L. Nestor, and J. Strang, "Pharmacodynamics of Diazepam Co-Administered with Methadone or Buprenorphine Under High Dose Conditions in Opioid Dependent Patients," *Drug and Alcohol Dependence*, Vol. 91, No. 2-3, December 1, 2007, pp. 187–194. Reason for exclusion: Only measures effects of a co-intervention.
- Liu, H., L. Li, Y. Hao, D. Cao, L. Xu, R. Rohrbaugh, Z. Xue, W. Hao, B. Shan, and Z. Liu, "Disrupted White Matter Integrity in Heroin Dependence: A Controlled Study Utilizing Diffusion Tensor Imaging," *The American Journal of Drug and Alcohol Abuse*, Vol. 34, No. 5, 2008. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Lobmaier, P., M. Gossop, H. Waal, and J. Bramness, "The Pharmacological Treatment of Opioid Addiction—A Clinical Perspective," *European Journal of Clinical Pharmacology*, Vol. 66, No. 6, June 2010, pp. 537–545. Reason for exclusion: Wrong study design.
- Lobmaier, Philipp Paul, Nikolaj Kunøe, and Helge Waal, "Treatment Research in Prison: Problems and Solutions in a Randomized Trial," *Addiction Research and Theory*, Vol. 18, No. 1, 2010, pp. 1–13. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Lobmann, R., "Diamorphine Substitution Therapy and Criminal Activity. [German]," *Sucht*, Vol. 53, No. 5, 2007. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Lobmann, R., and U. Verthein, "Explaining the Effectiveness of Heroin-Assisted Treatment on Crime Reductions," *Law and Human Behavior*, Vol. 33, No. 1, February 2009, pp. 83–95. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Loeber, S., A. Kniest, A. Diehl, K. Mann, and B. Croissant, "Neuropsychological Functioning of Opiate-Dependent Patients: A Nonrandomized Comparison of Patients Preferring Either Buprenorphine or Methadone Maintenance Treatment," *The American Journal of Drug and Alcohol Abuse*, Vol. 34, No. 5, 2008. Reason for exclusion: Wrong study design.
- Loimer, N., R. Schmid, J. Grünberger, and L. Linzmayer, "Naloxone Induces Miosis in Normal Subjects," *Psychopharmacology*, Vol. 101, No. 2, 1990. Reason for exclusion: Wrong study design.
- Longshore, D., J. Annon, M. D. Anglin, and R. A. Rawson, "Levo-Alpha-Acetylmethadol (LAAM) Versus Methadone: Treatment Retention and Opiate Use," *Addiction*, Vol. 100, No. 8, August 2005, pp. 1131–1139. Reason for exclusion: No functional outcomes.
- Loreck, David, Nicole J. Brandt, and Bethany DiPaula, "Managing Opioid Abuse in Older Adults," *Journal of Gerontological Nursing*, Vol. 42, No. 4, 2016, pp. 10–15. Reason for exclusion: Wrong study design.
- Lorenzetti, V., D. I. Lubman, D. Velakoulis, and M. Yucel, "Pituitary Gland Volume Among Heroin Users Stabilised on Substitution Pharmacotherapy," *Drug and Alcohol Dependence*, Vol. 110, No. 1–2, July 1, 2010, pp. 164–166. Reason for exclusion: No functional outcomes.

- Lott, D. C., E. C. Strain, R. K. Brooner, G. E. Bigelow, and R. E. Johnson, "HIV Risk Behaviors During Pharmacologic Treatment for Opioid Dependence: A Comparison of Levomethadyl Acetate [Corrected] Buprenorphine, and Methadone," *Journal of Substance Abuse Treatment*, Vol. 31, No. 2, September 2006, pp. 187–194. Reason for exclusion: No functional outcomes.
- Lott, Davic C., Eric C. Strain, Robert K. Brooner, George E. Bigelow, and Rolley E. Johnson, "HIV Risk Behaviors During Pharmacologic Treatment for Opioid Dependence: A Comparison of Levomethadyl Acetate Hydrochloride, Buprenorphine, and Methadone: 'Erratum,'" *Journal of Substance Abuse Treatment*, Vol. 31, No. 3, October 2006, p. 317. Reason for exclusion: Other.
- Loue, S., and B. Ioan, "Legal and Ethical Issues in Heroin Diagnosis, Treatment, and Research," *The Journal of Legal Medicine*, Vol. 28, No. 2, April–June 2007, pp. 193–221. Reason for exclusion: Wrong study design.
- Lu, L., D. Zhao, Y. P. Bao, and J. Shi, "Methadone Maintenance Treatment of Heroin Abuse in China," *The American Journal of Drug and Alcohol Abuse*, Vol. 34, No. 2, 2008, pp. 127–131. Reason for exclusion: Wrong study design.
- Lu, W. J., N. Thong, and D. A. Flockhart, "Reduced Methadone Clearance During Aromatase Inhibition," *Journal of Clinical Psychopharmacology*, Vol. 32, No. 4, August 2012, pp. 511–517. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Lubman, D. I., N. B. Allen, L. A. Peters, and J. F. Deakin, "Electrophysiological Evidence of the Motivational Salience of Drug Cues in Opiate Addiction," *Psychological Medicine*, Vol. 37, No. 8, August 2007, pp. 1203–1209. Reason for exclusion: Wrong study design.
- Lubman, D. I., L. A. Peters, K. Mogg, B. P. Bradley, and J. F. Deakin, "Attentional Bias for Drug Cues in Opiate Dependence," *Psychological Medicine*, Vol. 30, No. 1, January 2000, pp. 169–175. Reason for exclusion: Wrong study design.
- Lucas, G. M., A. Chaudhry, J. Hsu, T. Woodson, B. Lau, Y. Olsen, J. C. Keruly, D. A. Fiellin, R. Finkelstein, P. Barditch-Crovo, K. Cook, and R. D. Moore, "Clinic-Based Treatment of Opioid-Dependent HIV-Infected Patients Versus Referral to an Opioid Treatment Program: A Randomized Trial," *Annals of Internal Medicine*, Vol. 152, No. 11, June 1, 2010, pp. 704–711. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Lucas, G. M., A. Young, D. Donnell, P. Richardson, A. Aramrattana, Y. Shao, Y. Ruan, W. Liu, L. Fu, J. Ma, D. D. Celentano, D. Metzger, J. B. Jackson, and D. Burns, "Hepatotoxicity in a 52-Week Randomized Trial of Short-Term Versus Long-Term Treatment with Buprenorphine/Naloxone in HIV-Negative Injection Opioid Users in China and Thailand," *Drug and Alcohol Dependence*, Vol. 142, September 1, 2014, pp. 139–145. Reason for exclusion: No functional outcomes.

- Lugoboni, F., S. Carli, G. Bissoli, G. Musi, E. Florio, G. Civitelli, M. Brizio, C. Smacchia, C. Biasin, G. Cifelli, G. Manzato, S. Rosa, and E. Manzato, "Evaluation of the Quality of Life in 171 Patients Undergoing Methadone Maintenance Treatment and in 46 Monodependent Benzodiazepine Patients," *Heroin Addiction and Related Clinical Problems*, Vol. 16, No. 4, 2014, pp. 5–14. Reason for exclusion: No functional outcomes.
- Lundgren, L. M., R. F. Schilling, and S. D. Peloquin, "Evidence-Based Drug Treatment Practice and the Child Welfare System: The Example of Methadone," *Social Work*, Vol. 50, No. 1, January 2005, pp. 53–63. Reason for exclusion: Wrong study design.
- Lyvers, M., and M. Yakimoff, "Neuropsychological Correlates of Opioid Dependence and Withdrawal," *Addictive Behaviors*, Vol. 28, No. 3, April 2003, pp. 605–611. Reason for exclusion: Wrong study design.
- Ma, N., Y. Liu, N. Li, C. X. Wang, H. Zhang, X. F. Jiang, H. S. Xu, X. M. Fu, X. Hu, and D. R. Zhang, "Addiction Related Alteration in Resting-State Brain Connectivity," *Neuroimage*, Vol. 49, No. 1, January 1, 2010, pp. 738–744. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Maarefvand, M., H. R. Ghiasvand, and H. Ekhtiari, "Drug Craving Terminology Among Opiate Dependents; A Mixed Method Study," *Iranian Journal of Psychiatry*, Vol. 8, No. 2, June 2013, pp. 97–103. Reason for exclusion: Wrong study design.
- Macswain, Mary-Ann, Shanna Farrell-MacDonald, Madelon Cheverie, and Benedikt Fischer, "Assessing the Impact of Methadone Maintenance Treatment (MT) on Post-Release Recidivism Among Male Federal Correctional Inmates in Canada," *Criminal Justice and Behavior*, Vol. 41, No. 3, March 2014, pp. 380–394. Reason for exclusion: No functional outcomes.
- Maddux, J. F., D. P. Desmond, and M. Esquivel, "Rapid Admission and Retention on Methadone," *The American Journal of Drug and Alcohol Abuse*, Vol. 21, No. 4, November 1995, pp. 533–547. Reason for exclusion: No functional outcomes.
- Maddux, James F., David P. Desmond, and Kenneth N. Vogtsberger, "Patient-Regulated Methadone Dose and Optional Counseling in Methadone Maintenance," *The American Journal on Addictions*, Vol. 4, No. 1, Winter 1995, pp. 18–32. Reason for exclusion: Only measures effects of a co-intervention.
- Malhotra, A., D. Basu, S. K. Mattoo, N. Gupta, and R. Malhotra, "Acceptability of Naltrexone by Patients with Opioid Dependence: An Indian Perspective," *Hong Kong Journal of Psychiatry S2–East Asian Archives of Psychiatry*, Vol. 13, No. 4, December 2003, pp. 6–13. Reason for exclusion: Wrong study design.
- Mancino, M., G. Curran, X. Han, E. Allee, K. Humphreys, and B. M. Booth, "Predictors of Attrition from a National Sample of Methadone Maintenance Patients," *The American Journal of Drug and Alcohol Abuse*, Vol. 36, No. 3, May 2010, pp. 155–160. Reason for exclusion: Wrong study design.

- Mann, D. J., I. D. Montoya, C. Contoreggi, P. A. Ellison, W. R. Lange, and K. L. Preston, "Inpatient Medically Supervised Opioid Withdrawal with Buprenorphine Alone or in Combination with Naltrexone," *NIDA Research Monograph*, Vol. 153, 1994. Reason for exclusion: Only measures effects of a co-intervention.
- Mannelli, Paolo, Ashwin A. Patkar, Kathleen Peindl, Heather W. Murray, Li-Tzy Wu, and Robert Hubbard, "Effectiveness of Low-Dose Naltrexone in the Post-Detoxification Treatment of Opioid Dependence," *Journal of Clinical Psychopharmacology*, Vol. 27, No. 5, October 2007, pp. 468–474. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mannelli, Paolo, Li-Tzy Wu, Kathleen S. Peindl, Marvin S. Swartz, and George E. Woody, "Extended Release Naltrexone Injection Is Performed in the Majority of Opioid Dependent Patients Receiving Outpatient Induction: A Very Low Dose Naltrexone and Buprenorphine Open Label Trial," *Drug and Alcohol Dependence*, Vol. 138, May 1, 2014, pp. 83–88. Reason for exclusion: Wrong study design.
- "Manufacturers Respond to Opioid Abuse Through Risk Management Programs," *Alcoholism and Drug Abuse Weekly*, Vol. 19, No. 35, 2007, pp. 1–4. Reason for exclusion: Wrong study design.
- March, J. C., E. Oviedo-Joekes, E. Perea-Milla, and F. Carrasco, "Controlled Trial of Prescribed Heroin in the Treatment of Opioid Addiction," *Journal of Substance Abuse Treatment*, Vol. 31, No. 2, September 2006, pp. 203–211. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Marchand, K., E. Oviedo-Joekes, D. Guh, D. C. Marsh, S. Brissette, and M. T. Schechter, "Sex Work Involvement Among Women with Long-Term Opioid Injection Drug Dependence Who Enter Opioid Agonist Treatment," *Harm Reduction Journal*, Vol. 9, 2012. Reason for exclusion: No functional outcomes.
- Marchand, K. I., E. Oviedo-Joekes, D. Guh, S. Brissette, D. C. Marsh, and M. T. Schechter, "Client Satisfaction Among Participants in a Randomized Trial Comparing Oral Methadone and Injectable Diacetylmorphine for Long-Term Opioid-Dependency," *BMC Health Services Research*, Vol. 11, 2011, p. 174. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Marcovitz, David E., R. Kathryn McHugh, Julie Volpe, Victoria Votaw, and Hilary S. Connery, "Predictors of Early Dropout in Outpatient Buprenorphine/Naloxone Treatment," *American Journal on Addictions*, Vol. 25, No. 6, September 1, 2016, pp. 472–477. Reason for exclusion: Wrong study design.
- Marcucci, C., J. Fudin, P. Thomas, N. B. Sandson, and C. Welsh, "A New Pattern of Buprenorphine Misuse May Complicate Perioperative Pain Control," *Anesthesia and Analgesia*, Vol. 108, No. 6, June 2009, pp. 1996–1997. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Maremmani, I., M. Pacini, D. Popovic, A. Romano, A. G. Maremmani, G. Perugi, J. Deltito, K. Akiskal, and H. Akiskal, "Affective Temperaments in Heroin Addiction," *Journal of Affective Disorders*, Vol. 117, No. 3, October 2009, pp. 186–192. Reason for exclusion: No functional outcomes.
- Maremmani, I., P. P. Pani, A. Tagliamonte, and G. L. Gessa, "Buprenorphine Vs Methadone: An Italian Multicenter Study [Italian]," *Giornale Italiano di Farmacia Clinica*, Vol. 13, No. 4, 1999. Reason for exclusion: Not English.
- Maremmani, I., P. P. Pani, A. Mellini, M. Pacini, G. Marini, M. Lovrecic, G. Perugi, and M. Shinderman, "Alcohol and Cocaine Use and Abuse Among Opioid Addicts Engaged in a Methadone Maintenance Treatment Program," *Journal of Addictive Diseases*, Vol. 26, No. 1, 2007, pp. 61–70. Reason for exclusion: Wrong study design.
- Margolin, A., M. Beitel, Z. Schuman-Olivier, and S. K. Avants, "A Controlled Study of a Spirituality-Focused Intervention for Increasing Motivation for HIV Prevention Among Drug Users," *AIDS Education and Prevention: Official Publication of the International Society for AIDS Education*, Vol. 18, No. 4, 2006. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Marie-Claire, C., S. Crettol, N. Cagnard, V. Bloch, S. Mouly, J. L. Laplanche, F. Bellivier, J. P. Lepine, C. Eap, and F. Vorspan, "Variability of Response to Methadone: Genome-Wide DNA Methylation Analysis in Two Independent Cohorts," *Epigenomics*, Vol. 8, No. 2, 2016, pp. 181–195. Reason for exclusion: Wrong study design.
- Marini, J., M. Sullivan, A. Bisaga, K. Carpenter, K. A. Murray, F. R. Levin, and E. V. Nunes, "Naltrexone-Induced Protracted Opioid Withdrawal," *Proceedings of the 71th Annual Scientific Meeting of the College on Problems of Drug Dependence*, Reno/Sparks, Nev., 2009. Reason for exclusion: Conference abstract.
- Markwick, N., R. McNeil, S. Anderson, W. Small, and T. Kerr, "Communicating Risk in the Context of Methadone Formulation Changes: A Qualitative Study of Overdose Warning Posters in Vancouver, Canada," *International Journal of Drug Policy*, Vol. 27, January 2016, pp. 178–181. Reason for exclusion: Only measures effects of a co-intervention.
- Marsch, L. A., "Leveraging Technology in the Behavioral Treatment of Substance Use Disorders," *Alcoholism: Clinical and Experimental Research*, Vol. 36, 2012. Reason for exclusion: Conference abstract.
- Marsch, L. A., W. K. Bickel, G. J. Badger, and E. A. Jacobs, "Buprenorphine Treatment for Opioid Dependence: The Relative Efficacy of Daily, Twice and Thrice Weekly Dosing," *Drug and Alcohol Dependence*, Vol. 77, No. 2, February 14, 2005, pp. 195–204. Reason for exclusion: No functional outcomes.

- Marsch, L. A., W. K. Bickel, G. J. Badger, M. E. Stothart, K. J. Quesnel, C. Stanger, and J. Brooklyn, "Comparison of Pharmacological Treatments for Opioid-Dependent Adolescents: A Randomized Controlled Trial," *Archives of General Psychiatry*, Vol. 62, No. 10, October 2005, pp. 1157–1164. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Marsch, L. A., M. A. Stephens, T. Mudric, E. C. Strain, G. E. Bigelow, and R. E. Johnson, "Predictors of Outcome in LAAM, Buprenorphine, and Methadone Treatment for Opioid Dependence," *Experimental and Clinical Psychopharmacology*, Vol. 13, No. 4, November 2005, pp. 293–302. Reason for exclusion: No functional outcomes.
- Martin, K. R., "Combining Medications May Be Effective Treatment for 'Speedball' Abuse," *Nida Notes*, Vol. 17, No. 3, 2002, p. 2p. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Martin, L., J. Clair, P. Davis, D. O'Ryan, R. Hoshi, and H. V. Curran, "Enhanced Recognition of Facial Expressions of Disgust in Opiate Users Receiving Maintenance Treatment," *Addiction*, Vol. 101, No. 11, November 2006, pp. 1598–1605. Reason for exclusion: Wrong study design.
- Martin, W. R., D. R. Jasinski, C. A. Haertzen, D. C. Kay, B. E. Jones, P. A. Mansky, and R. W. Carpenter, "Methadone—A Reevaluation," *Archives of General Psychiatry*, Vol. 28, No. 2, February 1973, pp. 286–295. Reason for exclusion: Wrong setting (i.e., inpatient).
- Martin, W. R., D. R. Jasinski, and P. A. Mansky, "Naltrexone, an Antagonist for the Treatment of Heroin Dependence. Effects in Man," *Archives of General Psychiatry*, Vol. 28, No. 6, June 1973, pp. 784–791. Reason for exclusion: Wrong setting (i.e., inpatient).
- Martin-Soelch, C., A. F. Chevalley, G. Kunig, J. Missimer, S. Magyar, A. Mino, W. Schultz, and K. L. Leenders, "Changes in Reward-Induced Brain Activation in Opiate Addicts," *European Journal of Neuroscience*, Vol. 14, No. 8, October 2001, pp. 1360–1368. Reason for exclusion: No functional outcomes.
- Maruyama, Anna, Scott Macdonald, Elizabeth Borycki, and Jinhui Zhao, "Hypertension, Chronic Obstructive Pulmonary Disease, Diabetes and Depression Among Older Methadone Maintenance Patients in British Columbia," *Drug and Alcohol Review*, Vol. 32, No. 4, July 2013, pp. 412–418. Reason for exclusion: No functional outcomes.
- Marvel, C. L., M. L. Faulkner, E. C. Strain, M. Z. Mintzer, and J. E. Desmond, "An fMRI Investigation of Cerebellar Function During Verbal Working Memory in Methadone Maintenance Patients," *Cerebellum*, Vol. 11, No. 1, March 2012, pp. 300–310. Reason for exclusion: No functional outcomes.
- Masson, C. L., P. G. Barnett, K. L. Sees, K. L. Delucchi, A. Rosen, W. Wong, and S. M. Hall, "Cost and Cost-Effectiveness of Standard Methadone Maintenance Treatment Compared to Enriched 180-Day Methadone Detoxification," *Addiction*, Vol. 99, No. 6, June 2004, pp. 718–726. Reason for exclusion: No functional outcomes.

- Masson, Carmen L., Kevin L. Delucchi, Courtney McKnight, Jennifer Hettema, Mandana Khalili, Albert Min, Ashly E. Jordan, Nicole Pepper, Jessica Hall, Nicholas S. Hengl, Christopher Young, Michael S. Shopshire, Jennifer K. Manuel, Lara Coffin, Hali Hammer, Bradley Shapiro, Randy M. Seewald, Henry C. Bodenheimer, Jr., James L. Sorensen, Don C. Des Jarlais, and David C. Perlman, "A Randomized Trial of a Hepatitis Care Coordination Model in Methadone Maintenance Treatment," *American Journal of Public Health*, Vol. 103, No. 10, October 2013, pp. e81–e88. Reason for exclusion: Only measures effects of a co-intervention.
- Matheson, C., M. Jaffray, C. Bond, A. Leea, A. Johnstone, L. Skea, and B. Davidson, "Improving Outcomes and Quality of Life for People on Methadone Maintenance Therapy (MMT): The Enhanced Pharmacy Services (EPS) Randomised Controlled Trial," *International Journal of Pharmacy Practice*, Vol. 18, Supplement 2, September 2010, pp. 75–76. Reason for exclusion: Only measures effects of a co-intervention.
- Matsuyama, S. S., V. C. Charuvastra, J. Ouren, J. Schwartz, and L. Jarvik, "Immunoglobulin Levels in Heroin Addicts After Treatment with Methadone and Methadyl Acetate," *Drug and Alcohol Dependence*, Vol. 6, No. 5, 1980. Reason for exclusion: No functional outcomes.
- Mattick, R. P., R. Ali, G. Bammer, J. Bell, C. Breen, E. Digiusto, C. Doran, J. Gates, N. Glasgow, E. Gospodarevskaya, A. Harris, L. Hawken, N. Henderson, J. Kimber, N. Lintzeris, P. Marshall, S. O'Brien, A. Quigley, A. Ritter, J. B. Saunders, A. Shakeshaft, M. Shanahan, J. Shearer, and J. White, "Maintenance Pharmacotherapy for Opioid Dependence and SF-36 Health Status: A Comparison with General Population Norms and Other Chronic Disorders," *Addictive Disorders and Their Treatment*, Vol. 5, No. 4, 2006. Reason for exclusion: Wrong study design.
- Maxwell, S., and M. Shinderman, "Optimizing Response to Methadone Maintenance Treatment: Use of Higher-Dose Methadone," *Journal of Psychoactive Drugs*, Vol. 31, No. 2, April–June 1999, pp. 95–102. Reason for exclusion: No functional outcomes.
- Maxwell, S., and M. S. Shinderman, "Optimizing Long-Term Response to Methadone Maintenance Treatment: A 152-Week Follow-Up Using Higher-Dose Methadone," *Journal of Addictive Diseases*, Vol. 21, No. 3, 2002, pp. 1–12. Reason for exclusion: No functional outcomes.
- Mazhari, S., Z. Keshvari, A. Sabahi, and S. Mottaghian, "Assessment of Cognitive Functions in Methadone Maintenance Patients," *Addiction Health*, Vol. 7, No. 3–4, Summer–Autumn 2015, pp. 109–116. Reason for exclusion: Wrong study design.
- McCaffrey, B. R., "Methadone Treatment: Our Vision for the Future," *Journal of Addictive Diseases*, Vol. 20, No. 1, 2001, pp. 93–101. Reason for exclusion: Wrong study design.
- McCance-Katz, E. F., D. E. Moody, S. Prathikanti, G. Friedland, and P. M. Rainey, "Rifampin, but Not Rifabutin, May Produce Opiate Withdrawal in Buprenorphine-Maintained Patients," *Drug and Alcohol Dependence*, Vol. 118, No. 2–3, November 1, 2011, pp. 326–334. Reason for exclusion: Only measures effects of a co-intervention.

- McCarthy, J. J., and O. T. Borders, "Limit Setting on Drug Abuse in Methadone Maintenance Patients," *American Journal of Psychiatry*, Vol. 142, No. 12, December 1985, pp. 1419–1423. Reason for exclusion: Only measures effects of a co-intervention.
- McCaul, M. E., G. E. Bigelow, M. L. Stitzer, and I. Liebson, "Short-Term Effects of Oral Methadone in Methadone Maintenance Subjects," *Clinical Pharmacology and Therapeutics*, Vol. 31, No. 6, 1982. Reason for exclusion: Wrong setting (i.e., inpatient).
- McDermott, K. A., M. L. Griffin, H. S. Connery, E. Y. Hilario, D. A. Fiellin, G. M. Fitzmaurice, and R. D. Weiss, "Initial Response as a Predictor of 12-Week Buprenorphine-Naloxone Treatment Response in a Prescription Opioid-Dependent Population," *Journal of Clinical Psychiatry*, Vol. 76, No. 2, February 2015, pp. 189–194. Reason for exclusion: No functional outcomes.
- McDonald, S., S. Darke, S. Kaye, and M. Torok, "Deficits in Social Perception in Opioid Maintenance Patients, Abstinent Opioid Users and Non-Opioid Users," *Addiction*, Vol. 108, No. 3, March 2013, pp. 566–574. Reason for exclusion: Wrong study design.
- McKeganey, Neil, Christopher Russell, and Lucinda Cockayne, "Medically Assisted Recovery from Opiate Dependence Within the Context of the UK Drug Strategy: Methadone and Suboxone (Buprenorphine–Naloxone) Patients Compared," *Journal of Substance Abuse Treatment*, Vol. 44, No. 1, January 2013, pp. 97–102. Reason for exclusion: No functional outcomes.
- McKenzie, M., N. Zaller, S. L. Dickman, T. C. Green, A. Parihk, P. D. Friedmann, and J. D. Rich, "A Randomized Trial of Methadone Initiation Prior to Release from Incarceration," *Substance Abuse*, Vol. 33, No. 1, 2012, pp. 19–29. Reason for exclusion: Only measures effects of a co-intervention.
- McLellan, A. T., I. O. Arndt, D. S. Metzger, G. E. Woody, and C. P. O'Brien, "The Effects of Psychosocial Services in Substance Abuse Treatment," *JAMA*, Vol. 269, No. 15, April 21, 1993, pp. 1953–1959. Reason for exclusion: Only measures effects of a co-intervention.
- McRae, A. L., S. C. Sonne, K. T. Brady, V. Durkalski, and Y. Palesch, "A Randomized, Placebo-Controlled Trial of Buspirone for the Treatment of Anxiety in Opioid-Dependent Individuals," *The American Journal on Addictions*, Vol. 13, No. 1, January–February 2004, pp. 53–63. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Meadows, M., "Prescription Drug Abuse: FDA and SAMHSA Join Forces," *FDA Consumer*, Vol. 37, No. 2, March–April 2003, p. 36. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mehic-Basara, N., L. Oruc, L. Kapur-Pojkskic, and J. Ramic, "Association of Dopamine Receptor Gene Polymorphism and Psychological Personality Traits in Liability for Opioid Addiction," *Bosnian Journal of Basic Medical Sciences*, Vol. 13, No. 3, August 2013, pp. 158–162. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Mello, N. K., and J. H. Mendelson, "Buprenorphine Suppresses Heroin Use by Heroin Addicts," *Science*, Vol. 207, No. 4431, February 8, 1980, pp. 657–659. Reason for exclusion: Wrong setting (i.e., inpatient).
- Mello, N. K., J. H. Mendelson, and J. C. Kuehnle, "Buprenorphine Effects on Human Heroin Self-Administration: An Operant Analysis," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 223, No. 1, October 1982, pp. 30–39. Reason for exclusion: Wrong setting (i.e., inpatient).
- Mello, N. K., J. H. Mendelson, J. C. Kuehnle, and M. S. Sellers, "Operant Analysis of Human Heroin Self-Administration and the Effects of Naltrexone," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 216, No. 1, January 1981, pp. 45–54. Reason for exclusion: Wrong setting (i.e., inpatient).
- Mendelson, J., R. T. Jones, S. Welm, M. Baggott, I. Fernandez, A. K. Melby, and R. P. Nath, "Buprenorphine and Naloxone Combinations: The Effects of Three Dose Ratios in Morphine-Stabilized, Opiate-Dependent Volunteers," *Psychopharmacology*, Vol. 141, No. 1, 1999. Reason for exclusion: No functional outcomes.
- Mendelson, J., R. T. Jones, S. Welm, J. Brown, and S. L. Batki, "Buprenorphine and Naloxone Interactions in Methadone Maintenance Patients," *Biological Psychiatry*, Vol. 41, No. 11, 1997. Reason for exclusion: Wrong study design.
- Mendelson, J., R. T. Jones, I. Fernandez, S. Welm, A. K. Melby, and M. J. Baggott, "Buprenorphine and Naloxone Interactions in Opiate-Dependent Volunteers," *Clinical Pharmacology and Therapeutics*, Vol. 60, No. 1, July 1996, pp. 105–114. Reason for exclusion: No functional outcomes.
- Mendelson, J., R. A. Upton, E. T. Everhart, I. P. Jacob, and R. T. Jones, "Bioavailability of Sublingual Buprenorphine," *Journal of Clinical Pharmacology*, Vol. 37, No. 1, 1997. Reason for exclusion: Wrong study design.
- Merrill, Joseph O., and Mark H. Duncan, "Addiction Disorders," *Medical Clinics of North America*, Vol. 98, No. 5, 2014, pp. 1097–1122. Reason for exclusion: Wrong study design.
- Messinis, L., E. Lyros, V. Andrian, P. Katsakiori, G. Panagis, V. Georgiou, and P. Papathanasopoulos, "Neuropsychological Functioning in Buprenorphine Maintained Patients Versus Abstinent Heroin Abusers on Naltrexone Hydrochloride Therapy," *Human Psychopharmacology*, Vol. 24, No. 7, October 2009, pp. 524–531. Reason for exclusion: Wrong study design.
- "Methadone Overdoses: Take-Homes, Benzodiazepines, Induction Phase Debated," *Alcoholism & Drug Abuse Weekly*, Vol. 19, No. 44, 2007, pp. 1–3. Reason for exclusion: Wrong study design.
- "Methadone Replacement Therapy: Facilitating Immediate Access," *Prescribe International*, Vol. 16, No. 89, June 2007, p. 122. Reason for exclusion: Wrong study design.

- Metrebian, N., “Supervised Co-Prescription of Heroin to Treatment-Resistant Heroin Addicts Is More Effective than Treatment with Methadone Alone,” *Evidence Based Mental Health*, Vol. 7, No. 1, 2004, p. 23. Reason for exclusion: Only measures effects of a co-intervention.
- Metrebian, Nicky, Gerry V. Stimson, William Shanahan, Ron Alcorn, Louise Sell, Andrew Nunn, Rhian Gabe, Martin Knapp, Sarah Byford, David Epstein, Simon Thompson, Mani Medhikhani, Dan Harley, Mark Medjesi-Jones, Gill Campbell, Tom Carnwath, and Sue Ruben, “Pilot UK-Injectable Methadone Trial (Pilot UK-INJECT),” *Drugs: Education, Prevention and Policy*, Vol. 12, Supplement 1, December 2005, pp. 111–114. Reason for exclusion: Wrong study design.
- Metzger, D. S., J. Cornish, G. E. Woody, A. T. McLellan, P. Druley, and C. P. O’Brien, “Naltrexone in Federal Probationers,” *NIDA Research Monograph*, Vol. 95, 1989, pp. 465–466. Reason for exclusion: Conference abstract.
- Metzger, David S., Deborah Donnell, David D. Celentano, J. Brooks Jackson, Yiming Shao, Apinun Aramrattana, Liu Wei, Liping Fu, Jun Ma, Gregory M. Lucas, Marek Chawarski, Yuhua Ruan, Paul Richardson, Katherine Shin, Ray Y. Chen, Jeremy Sugarman, Bonnie J. Dye, Scott M. Rose, Geetha Beauchamp, and David N. Burns, “Expanding Substance Use Treatment Options for HIV Prevention with Buprenorphine–Naloxone: HIV Prevention Trials Network 058,” *JAIDS Journal of Acquired Immune Deficiency Syndromes*, Vol. 68, No. 5, April 15, 2015, pp. 554–561. Reason for exclusion: Only measures effects of a co-intervention.
- Meyer, A. C., M. E. Miller, and S. C. Sigmon, “Lifetime History of Heroin Use Is Associated with Greater Drug Severity Among Prescription Opioid Abusers,” *Addictive Behaviors*, Vol. 42, March 2015, pp. 189–193. Reason for exclusion: Wrong study design.
- Milby, J. B., C. Garrett, C. English, O. Fritschi, and C. Clarke, “Take-Home Methadone: Contingency Effects on Drug-Seeking and Productivity of Narcotic Addicts,” *Addictive Behaviors*, Vol. 3, No. 3–4, 1978, pp. 215–220. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Miller, P., M. Jarvis, and L. Strain, “News and Notes,” *Addiction*, Vol. 104, No. 12, 2009, pp. 2121–2124. Reason for exclusion: Wrong study design.
- Miller, Peter, Stephen McKenzie, Jane Walker, Nicholas Lintzeris, and John Strang, “Investigating the Effect on Public Behaviour of Patients of a Medically Supervised Injectable Maintenance Clinic,” *Drugs and Alcohol Today*, Vol. 11, No. 4, 2011, pp. 204–209. Reason for exclusion: No functional outcomes.
- Milos, G., L. M. Gallo, B. Sasic, D. Uebelhart, G. Goerres, H. J. Haeuselmann, and D. Eich, “Bone Mineral Density in Young Women on Methadone Substitution,” *Calcified Tissue International*, Vol. 89, No. 3, September 2011, pp. 228–233. Reason for exclusion: No functional outcomes.

- Mintz, J., C. P. O'Brien, K. O'Hare, and J. Goldschmidt, "Double-Blind Detoxification of Methadone Maintenance Patients," *The International Journal of the Addictions*, Vol. 10, No. 5, 1975, pp. 815–824. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mintzer, M. Z., M. L. Copersino, and M. L. Stitzer, "Opioid Abuse and Cognitive Performance," *Drug and Alcohol Dependence*, Vol. 78, No. 2, May 9, 2005, pp. 225–230. Reason for exclusion: Wrong study design.
- Mintzer, M. Z., C. J. Correia, and E. C. Strain, "A Dose-Effect Study of Repeated Administration of Buprenorphine/Naloxone on Performance in Opioid-Dependent Volunteers," *Drug and Alcohol Dependence*, Vol. 74, No. 2, May 10, 2004, pp. 205–209. Reason for exclusion: Wrong setting (i.e., inpatient).
- Mintzer, M. Z., and M. L. Stitzer, "Cognitive Impairment in Methadone Maintenance Patients," *Drug and Alcohol Dependence*, Vol. 67, No. 1, June 1, 2002, pp. 41–51. Reason for exclusion: Wrong study design.
- Miotto, K., M. Hillhouse, R. Donovan, J. Cunningham-Rathner, C. Charuvastra, M. Torrington, A. E. Esagoff, and W. Ling, "Comparison of Buprenorphine Treatment for Opioid Dependence in 3 Settings," *Journal of Addiction Medicine*, Vol. 6, No. 1, March 2012, pp. 68–76. Reason for exclusion: Only measures effects of a co-intervention.
- Mitchell, M. C., A. Memisoglu, and B. L. Silverman, "Hepatic Safety of Injectable Extended-Release Naltrexone in Patients with Chronic Hepatitis C and HIV Infection," *Journal of Studies on Alcohol and Drugs*, Vol. 73, No. 6, November 2012, pp. 991–997. Reason for exclusion: No functional outcomes.
- Mitchell, S. G., S. M. Kelly, J. Gryczynski, C. P. Myers, J. H. Jaffe, K. E. O'Grady, Y. K. Olsen, and R. P. Schwartz, "African American Patients Seeking Treatment in the Public Sector: Characteristics of Buprenorphine vs. Methadone Patients," *Drug and Alcohol Dependence*, Vol. 122, No. 1–2, 2012. Reason for exclusion: No functional outcomes.
- Mitchell, S. G., J. Gryczynski, R. P. Schwartz, K. E. O'Grady, Y. K. Olsen, and J. H. Jaffe, "A Randomized Trial of Intensive Outpatient (IOP) vs. Standard Outpatient (OP) Buprenorphine Treatment for African Americans," *Drug and Alcohol Dependence*, Vol. 128, No. 3, March 1, 2013, pp. 222–229. Reason for exclusion: Only measures effects of a co-intervention.
- Mitchell, S. G., S. M. Kelly, B. S. Brown, K. E. O'Grady, and R. P. Schwartz, "HIV Sex-Risk Behaviors Among In- Versus Out-of-Treatment Heroin-Addicted Adults," *The American Journal of Drug and Alcohol Abuse*, Vol. 38, No. 4, July 2012, pp. 328–333. Reason for exclusion: Only measures effects of a co-intervention.
- Mitchell, T. B., K. R. Dyer, D. Newcombe, A. Salter, A. A. Somogyi, F. Bochner, and J. M. White, "Subjective and Physiological Responses Among Racemic-Methadone Maintenance Patients in Relation to Relative (S)- vs. (R)-Methadone Exposure," *British Journal of Clinical Pharmacology*, Vol. 58, No. 6, December 2004, pp. 609–617. Reason for exclusion: Wrong study design.

- Mitchell, T. B., J. M. White, A. A. Somogyi, and F. Bochner, "Slow-Release Oral Morphine Versus Methadone: A Crossover Comparison of Patient Outcomes and Acceptability as Maintenance Pharmacotherapies for Opioid Dependence," *Addiction*, Vol. 99, No. 8, August 2004, pp. 940–945. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Modesto-Lowe, V., "Divided Doses for Methadone Maintenance," *American Journal of Psychiatry*, Vol. 165, No. 10, October 2008, pp. 1358; author reply 1358–1359. Reason for exclusion: Wrong study design.
- Mogali, S., N. A. Khan, E. S. Drill, M. Pavlicova, M. A. Sullivan, E. Nunes, and A. Bisaga, "Baseline Characteristics of Patients Predicting Suitability for Rapid Naltrexone Induction," *The American Journal on Addictions*, December 30, 2014. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mogali, S., N. A. Khan, E. S. Drill, M. Pavlicova, M. A. Sullivan, E. Nunes, and A. Bisaga, "Baseline Characteristics of Patients Predicting Suitability for Rapid Naltrexone Induction," *The American Journal on Addictions*, Vol. 24, No. 3, April 2015, pp. 258–264. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mohlman, M. K., B. Tanzman, K. Finison, M. Pinette, and C. Jones, "Impact of Medication-Assisted Treatment for Opioid Addiction on Medicaid Expenditures and Health Services Utilization Rates in Vermont," *Journal of Substance Abuse Treatment*, Vol. 67, August 2016, pp. 9–14. Reason for exclusion: No functional outcomes.
- Moitra, E., B. J. Anderson, and M. D. Stein, "Perceived Stress and Substance Use in Methadone-Maintained Smokers," *Drug and Alcohol Dependence*, Vol. 133, No. 2, December 1, 2013, pp. 785–788. Reason for exclusion: Only measures effects of a co-intervention.
- Mok, M. S., M. Lippmann, and S. N. Steen, "Multidose/Observational, Comparative Clinical Analgetic Evaluation of Buprenorphine," *Journal of Clinical Pharmacology*, Vol. 21, No. 7, July 1981, pp. 323–329. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Mokri, Azarakhsh, Marek C. Chawarski, Hamidreza Taherinakhost, and Richard S. Schottenfeld, "Medical Treatments for Opioid Use Disorder in Iran: A Randomized, Double-Blind Placebo-Controlled Comparison of Buprenorphine/Naloxone and Naltrexone Maintenance Treatment," *Addiction*, Vol. 111, No. 5, May 2016, pp. 874–882. Reason for exclusion: No functional outcomes.
- Monico, L. B., J. Gryczynski, S. G. Mitchell, R. P. Schwartz, K. E. O'Grady, and J. H. Jaffe, "Buprenorphine Treatment and 12-Step Meeting Attendance: Conflicts, Compatibilities, and Patient Outcomes," *Journal of Substance Abuse Treatment*, Vol. 57, 2015. Reason for exclusion: Only measures effects of a co-intervention.

- Monico, L. B., S. G. Mitchell, J. Gryczynski, R. P. Schwartz, K. E. O'Grady, Y. K. Olsen, and J. H. Jaffe, "Prior Experience with Non-Prescribed Buprenorphine: Role in Treatment Entry and Retention," *Journal of Substance Abuse Treatment*, Vol. 57, October 2015, pp. 57–62. Reason for exclusion: No functional outcomes.
- Montesano, F., D. Zaccone, E. Battaglia, F. Genco, and V. Mellace, "Therapeutic Switch to Buprenorphine/Naloxone from Buprenorphine Alone: Clinical Experience in an Italian Addiction Centre," *Clinical Drug Investigation*, Vol. 30 Supplement 1, 2010, pp. 13–19. Reason for exclusion: Wrong study design.
- Montoya, I. D., D. A. Gorelick, K. L. Preston, J. R. Schroeder, A. Umbricht, L. J. Cheskin, W. R. Lange, C. Contoreggi, R. E. Johnson, and P. J. Fudala, "Randomized Trial of Buprenorphine for Treatment of Concurrent Opiate and Cocaine Dependence," *Clinical Pharmacology and Therapeutics*, Vol. 75, No. 1, January 2004, pp. 34–48. Reason for exclusion: No functional outcomes.
- Mooney, L., M. P. Hillhouse, C. Thomas, A. Ang, A. Hasson, J. Annon, S. Reed, and W. Ling, "Psychiatric Symptoms and Treatment Outcomes in Cocaine-Dependent Adults Treated with Buprenorphine and Long-Acting Naltrexone," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Moore, B. A., D. T. Barry, L. E. Sullivan, P. G. O'Connor, C. J. Cutter, R. S. Schottenfeld, and D. A. Fiellin, "Counseling and Directly Observed Medication for Primary Care Buprenorphine Maintenance: A Pilot Study," *Journal of Addiction Medicine*, Vol. 6, No. 3, 2012. Reason for exclusion: Only measures effects of a co-intervention.
- Moore, B. A., T. Fazzino, D. T. Barry, D. A. Fiellin, C. J. Cutter, R. S. Schottenfeld, and S. A. Ball, "The Recovery Line: A Pilot Trial of Automated, Telephone-Based Treatment for Continued Drug Use in Methadone Maintenance," *Journal of Substance Abuse Treatment*, Vol. 45, No. 1, July 2013, pp. 63–69. Reason for exclusion: Only measures effects of a co-intervention.
- Moore, B. A., D. A. Fiellin, C. J. Cutter, F. D. Buono, D. T. Barry, L. E. Fiellin, P. G. O'Connor, and R. S. Schottenfeld, "Cognitive Behavioral Therapy Improves Treatment Outcomes for Prescription Opioid Users in Primary Care Buprenorphine Treatment," *Journal of Substance Abuse Treatment*, Vol. 71, No. 1, 2016, pp. 54–57. Reason for exclusion: Only measures effects of a co-intervention.
- Moore, B. C., C. J. Easton, and T. J. McMahon, "Drug Abuse and Intimate Partner Violence: A Comparative Study of Opioid-Dependent Fathers," *American Journal of Orthopsychiatry*, Vol. 81, No. 2, April 2011, pp. 218–227. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Moore, T. J., A. Ritter, and J. P. Caulkins, "The Costs and Consequences of Three Policy Options for Reducing Heroin Dependency," *Drug and Alcohol Review*, Vol. 26, No. 4, July 2007, pp. 369–378. Reason for exclusion: No functional outcomes.

- “More Convenient Preparations of Buprenorphine Pass Test,” *Nida Notes*, 2012, p. 1. Reason for exclusion: Wrong study design.
- Morgan, O., C. Griffiths, and M. Hickman, “Association Between Availability of Heroin and Methadone and Fatal Poisoning in England and Wales 1993–2004,” *International Journal of Epidemiology*, Vol. 35, No. 6, December 2006, pp. 1579–1585. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Morral, A. R., M. A. Belding, and M. Y. Iguchi, “Identifying Methadone Maintenance Clients at Risk for Poor Treatment Response: Pretreatment and Early Progress Indicators,” *Drug and Alcohol Dependence*, Vol. 55, No. 1–2, June 1, 1999, pp. 25–33. Reason for exclusion: Only measures effects of a co-intervention.
- Moszynski, P., “Methadone Is Trialled in Afghanistan to Try to Counter ‘Exponential Surge’ in Heroin Use,” *BMJ*, Vol. 342, April 7, 2011, p. d2281. Reason for exclusion: Wrong study design.
- Msekela, S., E. Krupitsky, E. Zvartau, E. Blokhina, A. Kibitov, and G. Woody, “Stabilization of Remission from Opioid Dependence with Long-Acting Naltrexone Implant,” *European Neuropsychopharmacology*, Vol. 24, 2014. Reason for exclusion: Conference abstract.
- Mulder, R., “Is It Time for Treatment Trials in Addiction?” *The New Zealand Medical Journal*, Vol. 117, No. 1190, March 12, 2004, p. U790. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Muller, B. W., M. Specka, N. Steinchen, D. Zerbin, E. Lodemann, T. Finkbeiner, and N. Scherbaum, “Auditory Target Processing in Methadone Substituted Opiate Addicts: The Effect of Nicotine in Controls,” *BMC Psychiatry*, Vol. 7, November 6, 2007, p. 63. Reason for exclusion: Wrong study design.
- Murray, D. E., T. C. Durazzo, T. P. Schmidt, C. Abe, J. Guydish, and D. J. Meyerhoff, “Frontal Metabolite Concentration Deficits in Opiate Dependence Relate to Substance Use, Cognition, and Self-Regulation,” *Journal of Addiction Research and Therapy*, Vol. 7, No. 4, August 2016. Reason for exclusion: No functional outcomes.
- Musshoff, F., K. Lachenmeier, D. Lichtermann, and B. Madea, “Cocaine and Opiate Concentrations in Hair from Subjects in a Heroin Maintenance Program in Comparison to a Methadone Substituted Group,” *International Journal of Legal Medicine*, Vol. 123, No. 5, September 2009, pp. 363–369. Reason for exclusion: No functional outcomes.
- Mustafa, M., S. I. Hasan, R. Abd Rashid, A. Mohd Ali, M. K. Mohd Ishak, M. A. Said, and M. H. Habil, “Effectiveness of Electroacupuncture as Adjunctive Therapy in Relieving Opiate Withdrawal Symptoms Among Methadone Maintenance Treatment (MMT) Patient: A Single Blinded, Randomized Study,” *European Psychiatry*, Vol. 28, No. 1, 2013, pp. 2001–2004. Reason for exclusion: Only measures effects of a co-intervention.

- Myers, C. E., J. Rego, P. Haber, K. Morley, K. D. Beck, L. Hogarth, and A. A. Moustafa, "Learning and Generalization from Reward and Punishment in Opioid Addiction," *Behavioural Brain Research*, Vol. 317, January 15, 2017, pp. 122–131. Reason for exclusion: Wrong study design.
- Nahvi, S., K. Richter, X. Li, L. Modali, and J. Arnsten, "Cigarette Smoking and Interest in Quitting in Methadone Maintenance Patients," *Addictive Behaviors*, Vol. 31, No. 11, November 2006, pp. 2127–2134. Reason for exclusion: Only measures effects of a co-intervention.
- Nasser, A. F., M. K. Greenwald, B. Vince, P. J. Fudala, P. Twumasi-Ankrah, Y. Liu, J. P. Jones, 3rd, and C. Heidbreder, "Sustained-Release Buprenorphine (RBP-6000) Blocks the Effects of Opioid Challenge with Hydromorphone in Subjects with Opioid Use Disorder," *Journal of Clinical Psychopharmacology*, Vol. 36, No. 1, February 2016, pp. 18–26. Reason for exclusion: Wrong study design.
- Nasser, A. F., C. Heidbreder, Y. Liu, and P. J. Fudala, "Pharmacokinetics of Sublingual Buprenorphine and Naloxone in Subjects with Mild to Severe Hepatic Impairment (Child-Pugh Classes A, B, and C), in Hepatitis C Virus-Seropositive Subjects, and in Healthy Volunteers," *Clinical Pharmacokinetics*, Vol. 54, No. 8, August 2015, pp. 837–849. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Nava, F., E. Caldiroli, S. Premi, and A. Lucchini, "Relationship Between Plasma Cortisol Levels, Withdrawal Symptoms and Craving in Abstinent and Treated Heroin Addicts," *Journal of Addictive Diseases*, Vol. 25, No. 2, 2006, pp. 9–16. Reason for exclusion: No functional outcomes.
- Nava, F., E. Manzato, C. Leonardi, and A. Lucchini, "Opioid Maintenance Therapy Suppresses Alcohol Intake in Heroin Addicts with Alcohol Dependence: Preliminary Results of an Open Randomized Study," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, Vol. 32, No. 8, December 12, 2008, pp. 1867–1872. Reason for exclusion: No functional outcomes.
- Nava-Mesa, Mo, Mr Lamprea, and A Munera, "Divergent Short- and Long-Term Effects of Acute Stress in Object Recognition Memory Are Mediated by Endogenous Opioid System Activation," *Neurobiology of Learning and Memory*, Vol. 106, 2013. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Navaratnam, V., A. Jamaludin, N. Raman, M. Mohamed, and S. M. Mansor, "Determination of Naltrexone Dosage for Narcotic Agonist Blockade in Detoxified Asian Addicts," *Drug and Alcohol Dependence*, Vol. 34, No. 3, February 1994, pp. 231–236. Reason for exclusion: No functional outcomes.
- Ndefo, U. A., and G. I. Erowele, "Focus on Methylnaltrexone: A Selective Mu-Opioid Receptor Antagonist for the Treatment of Opioid-Induced Constipation," *Journal of Pharmacy Technology*, Vol. 27, No. 1, 2011, pp. 15–18. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Neufeld, K. J., M. S. Kidorf, K. Kolodner, V. L. King, M. Clark, and R. K. Brooner, “A Behavioral Treatment for Opioid-Dependent Patients with Antisocial Personality,” *Journal of Substance Abuse Treatment*, Vol. 34, No. 1, January 2008, pp. 101–111. Reason for exclusion: Only measures effects of a co-intervention.
- Newcombe, D. A., F. Bochner, J. M. White, and A. A. Somogyi, “Evaluation of Levo-Alpha-Acetylmethadol (LAAM) as an Alternative Treatment for Methadone Maintenance Patients Who Regularly Experience Withdrawal: A Pharmacokinetic and Pharmacodynamic Analysis,” *Drug and Alcohol Dependence*, Vol. 76, No. 1, October 5, 2004, pp. 63–72. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Newman, R. G., “Comparing Buprenorphine and Methadone Maintenance,” *Journal of Nervous and Mental Disease*, Vol. 182, No. 4, April 1994, pp. 245–246. Reason for exclusion: Wrong study design.
- Newman, R. G., “Opiate Agonist Treatment for Addiction,” *Lancet*, Vol. 372, No. 9654, December 6, 2008, pp. 1951–1952; author reply 1952. Reason for exclusion: Wrong study design.
- Newman, R. G., “Risk Reduction with Buprenorphine-Naloxone and Methadone: Patient’s Choice,” *Journal of Acquired Immune Deficiency Syndrome*, Vol. 67, No. 5, December 15, 2014, p. e142. Reason for exclusion: Wrong study design.
- “News and Innovations,” *Journal of Pain and Palliative Care Pharmacotherapy*, Vol. 25, No. 1, 2011, pp. 90–109. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Nielsen, D. A., S. Hamon, V. Yuferov, C. Jackson, A. Ho, J. Ott, and M. J. Kreek, “Ethnic Diversity of DNA Methylation in the OPRM1 Promoter Region in Lymphocytes of Heroin Addicts,” *Human Genetics*, Vol. 127, No. 6, June 2010, pp. 639–649. Reason for exclusion: No functional outcomes.
- Nielsen, D. A., A. Ho, A. Bahl, P. Varma, S. Kellogg, L. Borg, and M. J. Kreek, “Former Heroin Addicts with or Without a History of Cocaine Dependence Are More Impulsive than Controls,” *Drug and Alcohol Dependence*, Vol. 124, No. 1–2, July 1, 2012, pp. 113–120. Reason for exclusion: Wrong study design.
- Nielsen, D. A., V. Yuferov, S. Hamon, C. Jackson, A. Ho, J. Ott, and M. J. Kreek, “Increased OPRM1 DNA Methylation in Lymphocytes of Methadone-Maintained Former Heroin Addicts,” *Neuropsychopharmacology*, Vol. 34, No. 4, March 2009, pp. 867–873. Reason for exclusion: No functional outcomes.
- Nielsen, S., M. Hillhouse, L. Mooney, A. Ang, and W. Ling, “Buprenorphine Pharmacotherapy and Behavioral Treatment: Comparison of Outcomes Among Prescription Opioid Users, Heroin Users and Combination Users,” *Journal of Substance Abuse Treatment*, Vol. 48, No. 1, January 2015, pp. 70–76. Reason for exclusion: Only measures effects of a co-intervention.

- Nielsen, S., M. Hillhouse, C. Thomas, A. Hasson, and W. Ling, "A Comparison of Buprenorphine Taper Outcomes Between Prescription Opioid and Heroin Users," *Journal of Addiction Medicine*, Vol. 7, No. 1, January–February 2013, pp. 33–38. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Nielsen, S., M. Hillhouse, R. D. Weiss, L. Mooney, J. Sharpe Potter, J. Lee, M. N. Gourevitch, and W. Ling, "The Relationship Between Primary Prescription Opioid and Buprenorphine-Naloxone Induction Outcomes in a Prescription Opioid Dependent Sample," *The American Journal on Addictions*, Vol. 23, No. 4, July–August 2014, pp. 343–348. Reason for exclusion: No functional outcomes.
- Nigam, A. K., R. Ray, and B. M. Tripathi, "Buprenorphine in Opiate Withdrawal: A Comparison with Clonidine," *Journal of Substance Abuse Treatment*, Vol. 10, No. 4, July–August 1993, pp. 391–394. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Nikolaou, P. D., I. I. Papoutsis, C. P. Maravelias, C. A. Spiliopoulou, C. M. Pistos, A. C. Calokerinos, and J. Atta-Politou, "Development and Validation of an EI-GC-MS Method for the Determination of Methadone and Its Major Metabolites (EDDP and EMDP) in Human Breast Milk," *Journal of Analytical Toxicology*, Vol. 32, No. 7, September 2008, pp. 478–484. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Nolan, L. J., and L. M. Scagnelli, "Preference for Sweet Foods and Higher Body Mass Index in Patients Being Treated in Long-Term Methadone Maintenance," *Substance Use and Misuse*, Vol. 42, No. 10, 2007, pp. 1555–1566. Reason for exclusion: No functional outcomes.
- Noori, R., H. Narenjiha, H. Aghabakhshi, G. Habibi, and B. Khoshkrood Mansoori, "Methadone Maintenance Therapy Outcomes in Iran," *Substance Use and Misuse*, Vol. 47, No. 7, June 2012, pp. 767–773. Reason for exclusion: Wrong study design.
- Nordmann, S., C. Lions, A. Vilotitch, L. Michel, M. Mora, B. Spire, G. Maradan, A. Morel, P. Roux, and Carrieri M. Patrizia, "A Prospective, Longitudinal Study of Sleep Disturbance and Comorbidity in Opiate Dependence (the ANRS Methaville Study)," *Psychopharmacology*, Vol. 233, No. 7, 2016. Reason for exclusion: Only measures effects of a co-intervention.
- Nordt, C., and R. Stohler, "Combined Effects of Law Enforcement and Substitution Treatment on Heroin Mortality," *Drug and Alcohol Review*, Vol. 29, No. 5, September 2010, pp. 540–545. Reason for exclusion: Wrong study design.
- Nosyk, B., J. W. Bray, E. Wittenberg, B. Aden, A. A. Eggman, R. D. Weiss, J. Potter, A. Ang, Y. I. Hser, W. Ling, and B. R. Schackman, "Short Term Health-Related Quality of Life Improvement During Opioid Agonist Treatment," *Drug and Alcohol Dependence*, Vol. 157, December 1, 2015, pp. 121–128. Reason for exclusion: No functional outcomes.

- Nosyk, B., J. Geller, D. P. Guh, E. Oviedo-Joeckes, S. Brissette, D. C. Marsh, M. T. Schechter, and A. H. Anis, "The Effect of Motivational Status on Treatment Outcome in the North American Opiate Medication Initiative (NAOMI) Study," *Drug and Alcohol Dependence*, Vol. 111, No. 1–2, September 1, 2010, pp. 161–165. Reason for exclusion: Only measures effects of a co-intervention.
- Nosyk, B., D. P. Guh, H. Sun, E. Oviedo-Joeckes, S. Brissette, D. C. Marsh, M. T. Schechter, and A. H. Anis, "Health Related Quality of Life Trajectories of Patients in Opioid Substitution Treatment," *Drug and Alcohol Dependence*, Vol. 118, No. 2–3, 2011. Reason for exclusion: Wrong study design.
- Notley, C., R. Holland, C. Kouimtsidis, V. Maskrey, A. Robinson, L. Swift, I. Harvey, F. Schifano, A. Lowe, and K. Rosenbloom, "Pharmacist and Patient Views of Supervised Consumption—Findings from a Randomised Controlled Trial and Qualitative Study of Supervised Consumption in Patients Managed with Opiate Maintenance Treatment," *International Journal of Pharmacy Practice*, Vol. 22, Supplement 1, April 2014, p. 19. Reason for exclusion: Conference abstract.
- Notley, C., R. Holland, V. Maskrey, J. Nagar, and C. Kouimtsidis, "Regaining Control: The Patient Experience of Supervised Compared with Unsupervised Consumption in Opiate Substitution Treatment," *Drug and Alcohol Review*, Vol. 33, No. 1, January 2014, pp. 64–70. Reason for exclusion: Only measures effects of a co-intervention.
- Nunes, E. V., E. Krupitsky, W. Ling, J. Zummo, A. Memisoglu, B. L. Silverman, and D. R. Gastfriend, "Treating Opioid Dependence with Injectable Extended-Release Naltrexone (XR-NTX): Who Will Respond?" *Journal of Addiction Medicine*, Vol. 9, No. 3, May–June 2015, pp. 238–243. Reason for exclusion: No functional outcomes.
- Nunes, E. V., J. L. Rothenberg, M. A. Sullivan, K. M. Carpenter, and H. D. Kleber, "Behavioral Therapy to Augment Oral Naltrexone for Opioid Dependence: A Ceiling on Effectiveness?" *The American Journal of Drug and Alcohol Abuse*, Vol. 32, No. 4, 2006, pp. 503–517. Reason for exclusion: Only measures effects of a co-intervention.
- Nurco, D. N., B. J. Primm, M. Lerner, P. Stephenson, L. S. Brown, and D. C. Ajuluchukwu, "Changes in Locus-of-Control Attitudes About Drug Misuse in a Self-Help Group in a Methadone Maintenance Clinic," *The International Journal of the Addictions*, Vol. 30, No. 6, May 1995, pp. 765–778. Reason for exclusion: Only measures effects of a co-intervention.
- Nurhidayat, Aw, "Efficacy of Behavioural Drug and Risk Counseling for Patients in Methadone Maintenance Treatment in Jakarta, Indonesia: A Randomised Clinical Trial," *European Neuropsychopharmacology*, Vol. 25, 2015. Reason for exclusion: Only measures effects of a co-intervention.
- Nutt, J. G., and D. R. Jasinski, "Methadone-Naloxone Mixtures for Use in Methadone Maintenance Programs I: An Evaluation in Man of Their Pharmacological Feasibility 11. Demonstration of Acute Physical Dependence," *Clinical Pharmacology and Therapeutics*, Vol. 15, No. 2, 1974, pp. 156–166. Reason for exclusion: No functional outcomes.

- Nyamathi, Adeline, Cynthia R. Albarrán, Catherine Branson, Mary Marfisee, Farinaz Khalilifard, and Barbara Leake, "Correlates of Depressive Symptoms Among Alcohol-Using Methadone Maintained Adults," *The American Journal on Addictions*, Vol. 21, No. Supplement 1, November 2012, pp. S49–S55. Reason for exclusion: Wrong study design.
- Nyamathi, A. M., K. Nandy, B. Greengold, M. Marfisee, F. Khalilifard, A. Cohen, and B. Leake, "Effectiveness of Intervention on Improvement of Drug Use Among Methadone Maintained Adults," *Journal of Addictive Diseases*, Vol. 30, No. 1, January 2011, pp. 6–16. Reason for exclusion: Only measures effects of a co-intervention.
- O'Brien, C. P., P. D. Friedmann, E. Nunes, J. D. Lee, and T. W. Kinlock, "Depot Naltrexone as Relapse Prevention for Opioid-Dependent Parolees," *Drug and Alcohol Dependence*, Vol. 146, 2015. Reason for exclusion: Conference abstract.
- Ochoa, E., A. Madoz-Gúrpide, and E. Salvador, "Gender Differences in the Treatment of the Opiate Dependence with Naltrexone," *Actas Españolas de Psiquiatría*, Vol. 36, No. 4, July–August 2008, pp. 197–204. Reason for exclusion: Wrong study design.
- O'Connor, P. G., A. H. Oliveto, J. M. Shi, E. G. Triffleman, K. M. Carroll, T. R. Kosten, B. J. Rounsaville, J. A. Pakes, and R. S. Schottenfeld, "A Randomized Trial of Buprenorphine Maintenance for Heroin Dependence in a Primary Care Clinic for Substance Users Versus a Methadone Clinic," *The American Journal of Medicine*, Vol. 105, No. 2, August 1998, pp. 100–105. Reason for exclusion: Only measures effects of a co-intervention.
- Ohlin, L., M. Fridell, and A. Nyhlen, "Buprenorphine Maintenance Program with Contracted Work/Education and Low Tolerance for Non-Prescribed Drug Use: A Cohort Study of Outcome for Women and Men After Seven Years," *BMC Psychiatry*, Vol. 15, March 24, 2015, p. 56. Reason for exclusion: Wrong study design.
- Oliveira, J., P. Lopes, P. Gamito, H. Trigo, P. Sargento, B. Rosa, and R. Coutinho, "Neuropsychological Status of Heroin Users Undergoing Methadone Maintenance in Harm Reduction Program and Therapeutic Community," *International Journal of Mental Health and Addiction*, Vol. 14, No. 2, April 2016, pp. 141–148. Reason for exclusion: Wrong study design.
- "ONDCP Responds to Opioid Trends: Naloxone and MAT . . . Office of National Drug Control Policy," *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 7, 2014, pp. 3–4. Reason for exclusion: Wrong study design.
- "Optimum Methadone Compliance Testing (Provisional Abstract)," *Database of Abstracts of Reviews of Effects*, No. 2, 2006. Reason for exclusion: Only measures effects of a co-intervention.
- Orman, J. S., and G. M. Keating, "Buprenorphine/Naloxone: A Review of Its Use in the Treatment of Opioid Dependence," *Drugs*, Vol. 69, No. 5, 2009, pp. 577–607. Reason for exclusion: Wrong study design.

- Orman, Jennifer S., and Gillian M. Keating, "Spotlight on Buprenorphine/Naloxone in the Treatment of Opioid Dependence," *CNS Drugs*, Vol. 23, No. 10, October 1, 2009, pp. 899–902. Reason for exclusion: Wrong study design.
- Oslin, D. W., "Evidence-Based Treatment of Geriatric Substance Abuse," *Psychiatric Clinics of North America*, Vol. 28, No. 4, December 2005, pp. 897–911, ix. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Oslin, D. W., C. P. O'Brien, W. Berrettini, and A. Gordon, "Prospective Study of Asn40Asp as a Moderator of Naltrexone Treatment of Alcohol Dependence," *Alcoholism: Clinical and Experimental Research*, Vol. 38 Supplement 1, June 2014, p. 327A. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Otiashvili, D., G. Piralishvili, Z. Sikharulidze, G. Kamkamidze, S. Poole, and G. E. Woody, "Methadone and Buprenorphine-Naloxone Are Effective in Reducing Illicit Buprenorphine and Other Opioid Use, and Reducing HIV Risk Behavior—Outcomes of a Randomized Trial," *Drug and Alcohol Dependence*, Vol. 133, No. 2, Dec 1, 2013, pp. 376–382. Reason for exclusion: No functional outcomes.
- "OTP Banks on Effectiveness of Integrated Primary Care . . . Opioid Treatment Program," *Alcoholism and Drug Abuse Weekly*, Vol. 24, No. 21, 2012, pp. 1–6. Reason for exclusion: Wrong study design.
- Oviedo-Joekes, E., S. Brissette, D. C. Marsh, P. Lauzon, D. Guh, A. Anis, and M. T. Schechter, "Diacetylmorphine Versus Methadone for the Treatment of Opioid Addiction," *New England Journal of Medicine*, Vol. 361, No. 8, August 20, 2009, pp. 777–786. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Oviedo-Joekes, E., D. Guh, S. Brissette, K. Marchand, D. Marsh, J. Chettiar, B. Nosyk, M. Krausz, A. Anis, and M. T. Schechter, "Effectiveness of Diacetylmorphine Versus Methadone for the Treatment of Opioid Dependence in Women," *Drug and Alcohol Dependence*, Vol. 111, No. 1–2, September 1, 2010, pp. 50–57. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Oviedo-Joekes, E., D. Guh, K. Marchand, D. C. Marsh, K. Lock, S. Brissette, A. H. Anis, and M. T. Schechter, "Differential Long-Term Outcomes for Voluntary and Involuntary Transition from Injection to Oral Opioid Maintenance Treatment," *Substance Abuse Treatment, Prevention, and Policy*, Vol. 9, 2014, p. 23. Reason for exclusion: No functional outcomes.
- Oviedo-Joekes, E., J. C. March, M. Romero, and E. Perea-Milla, "The Andalusian Trial on Heroin-Assisted Treatment: A 2 Year Follow-Up," *Drug and Alcohol Review*, Vol. 29, No. 1, January 2010, pp. 75–80. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Oviedo-Joekes, E., K. Marchand, D. Guh, D. C. Marsh, S. Brissette, M. Krausz, A. Anis, and M. T. Schechter, "History of Reported Sexual or Physical Abuse Among Long-Term Heroin Users and Their Response to Substitution Treatment," *Addictive Behaviors*, Vol. 36, No. 1–2, January–February 2011, pp. 55–60. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Oviedo-Joekes, E., K. Marchand, K. Lock, J. Chettiar, D. C. Marsh, S. Brissette, A. H. Anis, and M. T. Schechter, "A Chance to Stop and Breathe: Participants' Experiences in the North American Opiate Medication Initiative Clinical Trial," *Addiction Science and Clinical Practice*, Vol. 9, 2014. Reason for exclusion: Wrong study design.
- Oviedo-Joekes, E., L. Sordo, D. Guh, D. C. Marsh, K. Lock, S. Brissette, A. H. Anis, and M. T. Schechter, "Predictors of Non-Use of Illicit Heroin in Opioid Injection Maintenance Treatment of Long-Term Heroin Dependence," *Addictive Behaviors*, Vol. 41, February 2015, pp. 81–86. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Oyemade, A., "Buprenorphine/Naloxone Treatment in the Correctional Setting: Use or Misuse," *Innovations in Clinical Neuroscience*, Vol. 10, November–December 2013, pp. 13–14. Reason for exclusion: Wrong study design.
- Paderni, C., G. Campisi, A. Schumacher, T. Gottsche, L. I. Giannola, V. De Caro, and A. Wolff, "Controlled Delivery of Naltrexone by an Intraoral Device: In Vivo Study on Human Subjects," *International Journal of Pharmaceutics*, Vol. 452, No. 1–2, August 16, 2013, pp. 128–134. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Palatkin, V., E. Krupitsky, E. Zvartau, E. Blokhina, N. Bushara, A. Tyurina, T. Yaroslavl'tseva, and E. Verbitskaya, "Stress-Protective and Anticraving Effects of Guanfacine in Opiate Dependent Patients Stabilized on Oral Naltrexone," *European Neuropsychopharmacology*, Vol. 24, 2014. Reason for exclusion: Conference abstract.
- Panell, J., V. C. Charuvastra, and J. Ouren, "Methadyl Acetate Versus Methadone: The Experience of One Hospital," *The Medical Journal of Australia*, Vol. 2, No. 5, July 30, 1977, pp. 150–152. Reason for exclusion: No functional outcomes.
- Pani, P. P., I. Maremmanni, R. Pirastu, A. Tagliamonte, and G. L. Gessa, "Buprenorphine: A Controlled Clinical Trial in the Treatment of Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 60, No. 1, July 1, 2000, pp. 39–50. Reason for exclusion: No functional outcomes.
- Parran, T. V., C. A. Adelman, B. Merkin, M. E. Pagano, R. Defranco, R. A. Ionescu, and A. G. Mace, "Long-Term Outcomes of Office-Based Buprenorphine/Naloxone Maintenance Therapy," *Drug and Alcohol Dependence*, Vol. 106, No. 1, January 1, 2010, pp. 56–60. Reason for exclusion: Wrong study design.

- Parwatikar, S. D., and R. R. Knowles, "Methadone-Naloxone in Combination for the Treatment of Heroin Addicts," *Clinical Pharmacology and Therapeutics*, Vol. 14, No. 6, 1973. Reason for exclusion: No functional outcomes.
- Payne, L., "Sinking Success Rates of Treatment of Heroin Addicts: Are We Missing the Boat?" *Journal of Addictions Nursing*, Vol. 13, No. 1, 2001, pp. 3–8. Reason for exclusion: Wrong study design.
- Pecoraro, A., M. Ma, and G. E. Woody, "The Science and Practice of Medication-Assisted Treatments for Opioid Dependence," *Substance Use and Misuse*, Vol. 47, No. 8–9, June–July 2012, pp. 1026–1040. Reason for exclusion: Wrong study design.
- Pedrero-Perez, E. J., and L. G. MethaQo, "Methadone Dosage and Its Relationship to Quality of Life, Satisfaction, Psychopathology, Cognitive Performance and Additional Consumption of Non-Prescribed Drugs," *Adicciones*, June 14, 2016, p. 831. Reason for exclusion: Only measures effects of a co-intervention.
- Peirce, J. M., N. M. Petry, M. L. Stitzer, J. Blaine, S. Kellogg, F. Satterfield, M. Schwartz, J. Krasnansky, E. Pencer, L. Silva-Vazquez, K. C. Kirby, C. Royer-Malvestuto, J. M. Roll, A. Cohen, M. L. Copersino, K. Kolodner, and R. Li, "Effects of Lower-Cost Incentives on Stimulant Abstinence in Methadone Maintenance Treatment: A National Drug Abuse Treatment Clinical Trials Network Study," *Archives of General Psychiatry*, Vol. 63, No. 2, February 2006, pp. 201–208. Reason for exclusion: Only measures effects of a co-intervention.
- Pektas, O., A. O. Kalyoncu, H. Mirsal, A. Pektas, T. T. Satir, and M. Beyazyrek, "Different Forms (Oral or Implant) of Naltrexone Use in Relapse Prevention in Heroin Addicts: A Controlled Clinical Trials with up to Six Months Follow-Up," *Proceedings of the 10th Annual Meeting and Symposium of American Academy of Addiction Psychiatry*, Nassau, Bahamas, 1999. Reason for exclusion: Wrong study design.
- PektaSno, Ö. A. Kalyonko, H. Mirsal, A. Pektas, L. L. Gooberman, and M. Beyazyrek, "Different Forms (Oral or Implant) of Naltrexone Use in Relapse Prevention on Heroin Addicts: A Controlled Clinical Trial up to 6 Months Follow-Up," *XXIst Collegium Internationale Neuro-psychopharmacologicum*, Glasgow, Scotland, July 12–16, 1998. Reason for exclusion: No functional outcomes.
- Peles, E., S. Linzy, M. J. Kreek, and M. Adelson, "Prospective Study of Qtc Changes Among Former Opiate Addicts Since Admission to Methadone Maintenance Treatment: Benzodiazepine Risk," *Journal of Addiction Medicine*, Vol. 7, No. 6, November–December 2013, pp. 428–434. Reason for exclusion: Wrong study design.
- Peles, E., S. Schreiber, J. Gordon, and M. Adelson, "Significantly Higher Methadone Dose for Methadone Maintenance Treatment (MMT) Patients with Chronic Pain," *Pain*, Vol. 113, No. 3, February 2005, pp. 340–346. Reason for exclusion: Wrong study design.

- Peles, E., S. Schreiber, T. Hetzroni, M. Adelson, and R. Defrin, "The Differential Effect of Methadone Dose and of Chronic Pain on Pain Perception of Former Heroin Addicts Receiving Methadone Maintenance Treatment," *Journal of Pain*, Vol. 12, No. 1, January 2011, pp. 41–50. Reason for exclusion: Wrong study design.
- Peles, E., A. Weinstein, A. Sason, M. Adelson, and S. Schreiber, "Stroop Task Among Patients With Obsessive-Compulsive Disorder (OCD) and Pathological Gambling (PG) in Methadone Maintenance Treatment (MMT)," *CNS Spectrums*, Vol. 19, No. 6, December 2014, pp. 509–518. Reason for exclusion: Wrong study design.
- "People in CJ System Must Have Access to MAT: Legal Action Center . . . Medication-Assisted Treatment (MAT)," *Alcoholism and Drug Abuse Weekly*, Vol. 23, No. 47, 2011, pp. 5–6. Reason for exclusion: Wrong study design.
- Perea-Milla, E., L. C. Aycaguer, J. C. Cerda, F. G. Saiz, F. Rivas-Ruiz, A. Danet, M. R. Vallecillo, and E. Oviedo-Joekes, "Efficacy of Prescribed Injectable Diacetylmorphine in the Andalusian Trial: Bayesian Analysis of Responders and Non-Responders According to a Multi Domain Outcome Index," *Trials*, Vol. 10, 2009, p. 70. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Perez de los Cobos, J., S. Martin, A. Etcheberrigaray, J. Trujols, F. Batlle, A. Tejero, J. M. Queralto, and M. Casas, "A Controlled Trial of Daily Versus Thrice-Weekly Buprenorphine Administration for the Treatment of Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 59, No. 3, June 1, 2000, pp. 223–233. Reason for exclusion: No functional outcomes.
- Perez de Los Cobos, J., J. Trujols, N. Sinol, S. Duran-Sindreu, and F. Batlle, "Satisfaction with Methadone Among Heroin-Dependent Patients with Current Substance Use Disorders During Methadone Maintenance Treatment," *Journal of Clinical Psychopharmacology*, Vol. 36, No. 2, April 2016, pp. 157–162. Reason for exclusion: Wrong study design.
- Petitjean, S., R. Stohler, J. J. Deglon, S. Livoti, D. Waldvogel, C. Uehlinger, and D. Ladewig, "Double-Blind Randomized Trial of Buprenorphine and Methadone in Opiate Dependence," *Drug and Alcohol Dependence*, Vol. 62, No. 1, March 1, 2001, pp. 97–104. Reason for exclusion: No functional outcomes.
- Petrakis, I., K. M. Carroll, C. Nich, L. Gordon, T. Kosten, and B. Rounsaville, "Fluoxetine Treatment of Depressive Disorders in Methadone-Maintained Opioid Addicts," *Drug and Alcohol Dependence*, Vol. 50, No. 3, May 1, 1998, pp. 221–226. Reason for exclusion: Only measures effects of a co-intervention.
- Petrov, S. V., M. M. Orbetzova, Y. T. Iliev, and D. T. Boyadzhiev, "Chronic Treatment with Opiate Agonists in Bulgaria—Assessing the Quality of Life Using SF 36 v. 2," *Folia Med (Plovdiv)*, Vol. 58, No. 2, April–June 2016, pp. 115–121. Reason for exclusion: No functional outcomes.
- Petry, N. M., "Methadone Plus Contingency Management of Performance Feedback Reduces Cocaine and Opiate Use in People with Drug Addiction," *Evidence Based Mental Health*, Vol. 8, No. 4, 2005, p. 112. Reason for exclusion: Only measures effects of a co-intervention.

- Petry, N. M., W. K. Bickel, and G. J. Badger, "A Comparison of Four Buprenorphine Dosing Regimens in the Treatment of Opioid Dependence," *Clinical Pharmacology and Therapeutics*, Vol. 66, No. 3, September 1999, pp. 306–314. Reason for exclusion: No functional outcomes.
- Petry, N. M., W. K. Bickel, and G. J. Badger, "A Comparison of Four Buprenorphine Dosing Regimens Using Open-Dosing Procedures: Is Twice-Weekly Dosing Possible?" *Addiction*, Vol. 95, No. 7, July 2000, pp. 1069–1077. Reason for exclusion: Wrong study design.
- Petry, N. M., W. K. Bickel, and G. J. Badger, "Examining the Limits of the Buprenorphine Interdosing Interval: Daily, Every-Third-Day and Every-Fifth-Day Dosing Regimens," *Addiction*, Vol. 96, No. 6, June 2001, pp. 823–834. Reason for exclusion: No functional outcomes.
- Pezzarossa, A., N. Orlandi, G. Gerra, and P. P. Vescovi, "Effect of Methadone on Glucose Tolerance and Arginine-Induced Insulin and Glucagon Release in Man," *Diabetes, Nutrition, and Metabolism—Clinical and Experimental*, Vol. 3, No. 4, 1990. Reason for exclusion: Wrong setting (i.e., inpatient).
- Pfohl, D. N., J. I. Allen, R. L. Atkinson, D. S. Knopman, R. J. Malcolm, J. E. Mitchell, and J. E. Morley, "Naltrexone Hydrochloride (Trexan): A Review of Serum Transaminase Elevations at High Dosage," *NIDA Research Monograph*, Vol. 67, 1986, pp. 66–72. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Phillips, J., K. M. Carpenter, and E. V. Nunes, "Suicide Risk in Depressed Methadone-Maintained Patients: Associations with Clinical and Demographic Characteristics," *The American Journal on Addictions*, Vol. 13, No. 4, July–September 2004, pp. 327–332. Reason for exclusion: Wrong study design.
- Pinto, H., V. Maskrey, L. Swift, D. Rumball, A. Wagle, and R. Holland, "The SUMMIT Trial: A Field Comparison of Buprenorphine Versus Methadone Maintenance Treatment," *Journal of Substance Abuse Treatment*, Vol. 39, No. 4, 2010. Reason for exclusion: No functional outcomes.
- Pinto, Hayley, Daphne Rumball, Vivienne Maskrey, and Richard Holland, "A Pilot Study for a Randomized Controlled and Patient Preference Trial of Buprenorphine Versus Methadone Maintenance Treatment in the Management of Opiate Dependent Patients," *Journal of Substance Use*, Vol. 13, No. 2, April 2008, pp. 73–82. Reason for exclusion: No functional outcomes.
- Piralishvili, G., I. Gamkrelidze, N. Nikolaishvili, and M. Chavchanidze, "Evaluation of the Quality of Life (Whoqol-Bref) Among Methadone and Suboxone Substitution State Program Patients and Healthy Volunteers in Georgia," *Georgian Medical News*, No. 213, December 2012, pp. 44–47. Reason for exclusion: Not English.

- Piralishvili, Gvantsa, David Otiashvili, Zura Sikharulidze, George Kamkamidze, Sabrina Poole, and George E. Woody, "Opioid Addicted Buprenorphine Injectors: Drug Use During and After 12-Weeks of Buprenorphine–Naloxone or Methadone in the Republic of Georgia," *Journal of Substance Abuse Treatment*, Vol. 50, March 2015, pp. 32–37. Reason for exclusion: No functional outcomes.
- Pirastu, R., R. Fais, M. Messina, V. Bini, S. Spiga, D. Falconieri, and M. Diana, "Impaired Decision-Making in Opiate-Dependent Subjects: Effect of Pharmacological Therapies," *Drug and Alcohol Dependence*, Vol. 83, No. 2, June 28, 2006, pp. 163–168. Reason for exclusion: Wrong study design.
- Pluck, G., K. H. Lee, R. Rele, S. A. Spence, S. Sarkar, O. Lagundoye, and R. W. Parks, "Premorbid and Current Neuropsychological Function in Opiate Abusers Receiving Treatment," *Drug and Alcohol Dependence*, Vol. 124, No. 1–2, July 1, 2012, pp. 181–184. Reason for exclusion: Wrong study design.
- Pollack, H. A., and T. D'Aunno, "Dosage Patterns in Methadone Treatment: Results from a National Survey, 1988–2005," *Health Services Research*, Vol. 43, No. 6, December 2008, pp. 2143–2163. Reason for exclusion: Wrong study design.
- Polsky, D., H. A. Glick, J. Yang, G. A. Subramaniam, S. A. Poole, and G. E. Woody, "Cost-Effectiveness of Extended Buprenorphine-Naloxone Treatment for Opioid-Dependent Youth: Data from a Randomized Trial," *Addiction*, Vol. 105, No. 9, September 2010, pp. 1616–1624. Reason for exclusion: No functional outcomes.
- Poole, S. A., A. Pecoraro, G. Subramaniam, G. Woody, and V. L. Vetter, "Presence or Absence of QTc Prolongation in Buprenorphine-Naloxone Among Youth with Opioid Dependence," *Journal of Addiction Medicine*, Vol. 10, No. 1, 2016. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Potter, J. S., J. A. Dreifuss, E. N. Marino, S. E. Provost, D. R. Dodd, L. S. Rice, G. M. Fitzmaurice, M. L. Griffin, and R. D. Weiss, "The Multi-Site Prescription Opioid Addiction Treatment Study: 18-Month Outcomes," *Journal of Substance Abuse Treatment*, Vol. 48, No. 1, January 2015, pp. 62–69. Reason for exclusion: No functional outcomes.
- Potter, J. S., E. N. Marino, M. P. Hillhouse, S. Nielsen, K. Wiest, C. P. Canamar, J. A. Martin, A. Ang, R. Baker, A. J. Saxon, and W. Ling, "Buprenorphine/naloxone and Methadone Maintenance Treatment Outcomes for Opioid Analgesic, Heroin, and Combined Users: Findings from Starting Treatment with Agonist Replacement Therapies (START)," *Journal of Studies on Alcohol and Drugs*, Vol. 74, No. 4, July 2013, pp. 605–613. Reason for exclusion: No functional outcomes.
- Prakash, S., and Y. Balhara, "Perceptions Related to Pharmacological Treatment of Opioid Dependence Among Individuals Seeking Treatment at a Tertiary Care Center in Northern India: A Descriptive Study," *Substance Use and Misuse*, Vol. 51, No. 7, June 6, 2016, pp. 861–869. Reason for exclusion: No functional outcomes.

- Praveen, K. T., F. Law, J. O'Shea, and J. Melichar, "Opioid Dependence," *American Family Physician*, Vol. 86, No. 6, September 15, 2012, pp. 565–566. Reason for exclusion: Wrong study design.
- "President Calls for More MAT to Fight Opioid Epidemic . . . Medication-Assisted Treatment," *Alcoholism and Drug Abuse Weekly*, Vol. 27, No. 42, 2015, pp. 1–3. Reason for exclusion: Wrong study design.
- Preston, K., W. Kowalczyk, K. Phillips, M. Jobes, M. Furnari, U. Ghitza, and D. Epstein, "Clonidine Increases the Likelihood That Abstinence Can Withstand Leisure Time in Buprenorphine-Maintained Outpatients," *Neuropsychopharmacology*, Vol. 40, 2015. Reason for exclusion: Only measures effects of a co-intervention.
- Preston, K. L., G. E. Bigelow, and I. A. Liebson, "Buprenorphine and Naloxone Alone and in Combination in Opioid-Dependent Humans," *Psychopharmacology (Berl)*, Vol. 94, No. 4, 1988, pp. 484–490. Reason for exclusion: Wrong setting (i.e., inpatient).
- Preston, K. L., U. E. Ghitza, J. P. Schmittner, J. R. Schroeder, and D. H. Epstein, "Randomized Trial Comparing Two Treatment Strategies Using Prize-Based Reinforcement of Abstinence in Cocaine and Opiate Users," *Journal of Applied Behavior Analysis*, Vol. 41, No. 4, Winter 2008, pp. 551–563. Reason for exclusion: Only measures effects of a co-intervention.
- Preston, K. L., K. Silverman, A. Umbricht, A. DeJesus, I. D. Montoya, and C. R. Schuster, "Improvement in Naltrexone Treatment Compliance with Contingency Management," *Drug and Alcohol Dependence*, Vol. 54, No. 2, April 1, 1999, pp. 127–135. Reason for exclusion: Only measures effects of a co-intervention.
- Preston, K. L., A. Umbricht, and D. H. Epstein, "Methadone Dose Increase and Abstinence Reinforcement for Treatment of Continued Heroin Use During Methadone Maintenance," *Archives of General Psychiatry*, Vol. 57, No. 4, April 2000, pp. 395–404. Reason for exclusion: Only measures effects of a co-intervention.
- Preston, K. L., A. Umbricht, and D. H. Epstein, "Abstinence Reinforcement Maintenance Contingency and One-Year Follow-Up," *Drug and Alcohol Dependence*, Vol. 67, No. 2, July 1, 2002, pp. 125–137. Reason for exclusion: Only measures effects of a co-intervention.
- "Prevention. Opiate Substitute Therapy Reduces HIV Transmission," *AIDS Policy Law*, Vol. 28, No. 2, December 2012, p. 1. Reason for exclusion: Only measures effects of a co-intervention.
- Prosser, J., L. J. Cohen, M. Steinfeld, D. Eisenberg, E. D. London, and I. I. Galynker, "Neuropsychological Functioning in Opiate-Dependent Subjects Receiving and Following Methadone Maintenance Treatment," *Drug and Alcohol Dependence*, Vol. 84, No. 3, October 1, 2006, pp. 240–247. Reason for exclusion: Wrong study design.
- Prosser, J., E. D. London, and I. I. Galynker, "Sustained Attention in Patients Receiving and Abstinent Following Methadone Maintenance Treatment for Opiate Dependence: Performance and Neuroimaging Results," *Drug and Alcohol Dependence*, Vol. 104, No. 3, October 1, 2009, pp. 228–240. Reason for exclusion: Wrong study design.

- Prosser, J. M., D. Eisenberg, E. E. Davey, M. Steinfeld, L. J. Cohen, E. D. London, and I. I. Galyunker, "Character Pathology and Neuropsychological Test Performance in Remitted Opiate Dependence," *Substance Abuse, Treatment, and Prevention Policy*, Vol. 3, November 19, 2008, p. 23. Reason for exclusion: Wrong study design.
- Prosser, J. M., M. Steinfeld, L. J. Cohen, S. Derbyshire, D. P. Eisenberg, R. A. Cruciani, and I. I. Galyunker, "Abnormal Heat and Pain Perception in Remitted Heroin Dependence Months After Detoxification from Methadone-Maintenance," *Drug and Alcohol Dependence*, Vol. 95, No. 3, June 1, 2008, pp. 237–244. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- "Public Policy Statement on Buprenorphine for Opiate Dependence and Withdrawal," *Journal of Addictive Diseases*, Vol. 22, No. 3, 2003, pp. 113–114. Reason for exclusion: Wrong study design.
- Pud, D., D. Cohen, E. Lawental, and E. Eisenberg, "Opioids and Abnormal Pain Perception: New Evidence from a Study of Chronic Opioid Addicts and Healthy Subjects," *Drug and Alcohol Dependence*, Vol. 82, No. 3, May 20, 2006, pp. 218–223. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Qiu, Y. W., L. J. Han, X. F. Lv, G. H. Jiang, J. Z. Tian, F. Z. Zhuo, H. H. Su, C. L. Lin, and X. L. Zhang, "Regional Homogeneity Changes in Heroin-Dependent Individuals: Resting-State Functional MR Imaging Study," *Radiology*, Vol. 261, No. 2, November 2011, pp. 551–559. Reason for exclusion: Wrong study design.
- Rabinowitz, A., "Enhancing Medication-Assisted Treatment: Success Beyond Harm Reduction," *Journal of Social Work Practice in the Addictions*, Vol. 9, No. 2, 2009, pp. 240–243. Reason for exclusion: Wrong study design.
- Rahimi-Movaghar, A., M. Amin-Esmaili, M. Hefazi, and R. Yousefi-Nooraie, "Pharmacological Therapies for Maintenance Treatments of Opium Dependence," *The Cochrane Database of Systematic Reviews*, No. 1, 2013, p. Cd007775. Reason for exclusion: No functional outcomes.
- Raisch, D. W., H. M. Campbell, D. A. Garnand, M. A. Jones, M. R. Sather, R. Naik, and W. Ling, "Health-Related Quality of Life Changes Associated with Buprenorphine Treatment for Opioid Dependence," *Quality of Life Research*, Vol. 21, No. 7, September 2012, pp. 1177–1183. Reason for exclusion: Wrong study design.
- Rao, Ravindra, Atul Ambekar, Seema Yadav, Hem Sethi, and Anju Dhawan, "Slow-Release Oral Morphine as a Maintenance Agent in Opioid Dependence Syndrome: An Exploratory Study from India," *Journal of Substance Use*, Vol. 17, No. 3, June 2012, pp. 294–300. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Rapeli, P., C. Fabritius, H. Alho, M. Salaspuro, K. Wahlbeck, and H. Kalska, "Methadone Vs. Buprenorphine/Naloxone During Early Opioid Substitution Treatment: A Naturalistic Comparison of Cognitive Performance Relative to Healthy Controls," *BMC Clinical Pharmacology*, Vol. 7, June 12, 2007, p. 5. Reason for exclusion: Wrong study design.

- Rapeli, P., C. Fabritius, H. Kalska, and H. Alho, "Cognitive Functioning in Opioid-Dependent Patients Treated with Buprenorphine, Methadone, and Other Psychoactive Medications: Stability and Correlates," *BMC Clinical Pharmacology*, Vol. 11, 2011, p. 13. Reason for exclusion: Article reporting follow up of included studies.
- Rass, O., B. A. Kleykamp, R. G. Vandrey, G. E. Bigelow, J. M. Leoutsakos, M. L. Stitzer, E. C. Strain, M. L. Copersino, and M. Z. Mintzer, "Cognitive Performance in Methadone Maintenance Patients: Effects of Time Relative to Dosing and Maintenance Dose Level," *Experimental and Clinical Psychopharmacology*, Vol. 22, No. 3, June 2014, pp. 248–256. Reason for exclusion: Wrong study design.
- Rass, O., R. L. Schacht, K. Buckheit, M. W. Johnson, E. C. Strain, and M. Z. Mintzer, "A Randomized Controlled Trial of the Effects of Working Memory Training in Methadone Maintenance Patients," *Drug and Alcohol Dependence*, Vol. 156, November 1, 2015, pp. 38–46. Reason for exclusion: Only measures effects of a co-intervention.
- Rawson, R. A., M. Glazer, E. J. Callahan, and R. P. Liberman, "Naltrexone and Behaviour Therapy for Heroin Addiction," *NIDA Research Monograph*, Vol. 25, 1979. Reason for exclusion: Only measures effects of a co-intervention.
- Rawson, R. A., and F. S. Tennant Jr., "Five-Year Follow-Up of Opiate Addicts with Naltrexone and Behavior Therapy," *NIDA Research Monograph*, Vol. 49, 1984. Reason for exclusion: Wrong study design.
- Rea, F., J. R. Bell, M. R. Young, and R. P. Mattick, "A Randomised, Controlled Trial of Low Dose Naltrexone for the Treatment of Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 75, No. 1, July 15, 2004, pp. 79–88. Reason for exclusion: No functional outcomes.
- Redding, S. E., S. Liu, W. W. Hung, and K. S. Boockvar, "Opioid Interruptions, Pain, and Withdrawal Symptoms in Nursing Home Residents," *Clinical Therapeutics*, Vol. 36, No. 11, 2014, pp. 1555–1563. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Reece, A. S., "Comparative Treatment and Mortality Correlates and Adverse Event Profile of Implant Naltrexone and Sublingual Buprenorphine," *Journal of Substance Abuse Treatment*, Vol. 37, No. 3, October 2009, pp. 256–265. Reason for exclusion: No functional outcomes.
- Reece, A. S., "Favorable Mortality Profile of Naltrexone Implants for Opiate Addiction," *Journal of Addictive Diseases*, Vol. 29, No. 1, January 2010, pp. 30–50. Reason for exclusion: No functional outcomes.
- Reece, A. S., and P. Davidson, "Deficit of Circulating Stem—Progenitor Cells in Opiate Addiction: A Pilot Study," *Substance Abuse, Treatment, and Prevention Policy*, Vol. 2, July 5, 2007, p. 19. Reason for exclusion: No functional outcomes.
- Reece, A. S., and G. K. Hulse, "Impact of Opioid Pharmacotherapy on Arterial Stiffness and Vascular Ageing: Cross-Sectional and Longitudinal Studies," *Cardiovascular Toxicology*, Vol. 13, No. 3, September 2013, pp. 254–266. Reason for exclusion: No functional outcomes.

- Reece, A. S., and G. K. Hulse, "Lifetime Opiate Exposure as an Independent and Interactive Cardiovascular Risk Factor in Males: A Cross-Sectional Clinical Study," *Vascular Health Risk Management*, Vol. 9, 2013, pp. 551–561. Reason for exclusion: No functional outcomes.
- Reimer, J., U. Verthein, A. Karow, I. Schafer, D. Naber, and C. Haasen, "Physical and Mental Health in Severe Opioid-Dependent Patients Within a Randomized Controlled Maintenance Treatment Trial," *Addiction*, Vol. 106, No. 9, September 2011, pp. 1647–1655. Reason for exclusion: No functional outcomes.
- Reimer, Jens, Uwe Verthein, Anne Karow, Ingo Schäfer, Dieter Naber, and Christian Haasen, "Physical and Mental Health in Severe Opioid-Dependent Patients Within a Randomized Controlled Maintenance Treatment Trial," *Addiction*, Vol. 106, No. 9, September 2011, pp. 1647–1655. Reason for exclusion: No functional outcomes.
- Renault, Pierre F., "Methadone Maintenance: The Effect of Knowledge of Dosage," *The International Journal of the Addictions*, Vol. 8, No. 1, 1973, pp. 41–47. Reason for exclusion: Only measures effects of a co-intervention.
- "Research in Brief," *Nida Notes*, Vol. 22, No. 5, 2009, p. 3. Reason for exclusion: Wrong study design.
- Resnick, R. B., M. Galanter, C. Pycha, A. Cohen, P. Grandison, and N. Flood, "Buprenorphine: An Alternative to Methadone for Heroin Dependence Treatment," *Psychopharmacology Bulletin*, Vol. 28, No. 1, 1992, pp. 109–113. Reason for exclusion: Wrong study design.
- Reynaud-Maurupt, C., M. P. Carrieri, J. A. Gastaud, C. Pradier, Y. Obadia, and J. P. Moatti, "Impact of Drug Maintenance Treatment on Injection Practices Among French HIV-Infected IDUs. The MANIF 2000 Study Group," *AIDS Care*, Vol. 12, No. 4, August 2000, pp. 461–470. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Rezaiyan, M. K., H. K. Moghadam, H. Khosrojerdi, and R. Afshari, "Very Low-Dose Naltrexone Versus Placebo in Alleviating Withdrawal Manifestation," *Clinical Toxicology*, Vol. 52, Supplement 1, May 2014, p. 368. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Rhoades, H. M., D. Creson, R. Elk, J. Schmitz, and J. Grabowski, "Retention, HIV Risk, and Illicit Drug Use During Treatment: Methadone Dose and Visit Frequency," *American Journal of Public Health*, Vol. 88, No. 1, January 1998, pp. 34–39. Reason for exclusion: Only measures effects of a co-intervention.
- Rich, J. D., M. McKenzie, S. Larney, J. B. Wong, L. Tran, J. Clarke, A. Noska, M. Reddy, and N. Zaller, "Methadone Continuation Versus Forced Withdrawal on Incarceration in a Combined U.S. Prison and Jail: A Randomised, Open-Label Trial," *Lancet*, Vol. 386, No. 9991, July 25, 2015, pp. 350–359. Reason for exclusion: Wrong setting (i.e., inpatient).

- Rich, J. D., M. McKenzie, D. C. Shield, F. A. Wolf, R. G. Key, M. Poshkus, and J. Clarke, "Linkage with Methadone Treatment Upon Release from Incarceration: A Promising Opportunity," *Journal of Addictive Diseases*, Vol. 24, No. 3, 2005, pp. 49–59. Reason for exclusion: Only measures effects of a co-intervention.
- Rieckmann, T. R., A. J. Abraham, A. E. Kovas, B. H. McFarland, and P. M. Roman, "Impact of Research Network Participation on the Adoption of Buprenorphine for Substance Abuse Treatment," *Addictive Behaviors*, Vol. 39, No. 5, May 2014, pp. 889–896. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Riksheim, M., M. Gossop, and T. Clausen, "From Methadone to Buprenorphine: Changes During a 10-Year Period Within a National Opioid Maintenance Treatment Programme," *Journal of Substance Abuse Treatment*, Vol. 46, No. 3, March 2014, pp. 291–294. Reason for exclusion: Wrong study design.
- Riss, G. L., D. I. Chang, C. Wevers, A. M. Westendorf, J. Buer, N. Scherbaum, and W. Hansen, "Opioid Maintenance Therapy Restores CD4+ T Cell Function by Normalizing CD4+CD25(High) Regulatory T Cell Frequencies in Heroin User," *Brain, Behavior, and Immunity*, Vol. 26, No. 6, August 2012, pp. 972–978. Reason for exclusion: No functional outcomes.
- Ritter, A. J., N. Lintzeris, N. Clark, J. J. Kutin, G. Bammer, and M. Panjari, "A Randomized Trial Comparing Levo-Alpha Acetylmethadol with Methadone Maintenance for Patients in Primary Care Settings in Australia," *Addiction*, Vol. 98, No. 11, November 2003, pp. 1605–1613. Reason for exclusion: No functional outcomes.
- Roberson, C. M., "Outpatient Opioid Addiction Treatment Using Buprenorphine," *Ala Nurse*, Vol. 37, No. 2, June–August 2010, pp. 13–16; quiz 17. Reason for exclusion: Wrong study design.
- Robertson, J. R., G. M. Raab, M. Bruce, J. S. McKenzie, H. R. Storkey, and A. Salter, "Addressing the Efficacy of Dihydrocodeine Versus Methadone as an Alternative Maintenance Treatment for Opiate Dependence: A Randomized Controlled Trial," *Addiction*, Vol. 101, No. 12, December 2006, pp. 1752–1759. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Robles, E., B. E. Huang, P. M. Simpson, and D. E. McMillan, "Delay Discounting, Impulsiveness, and Addiction Severity in Opioid-Dependent Patients," *Journal of Substance Abuse Treat*, Vol. 41, No. 4, December 2011, pp. 354–362. Reason for exclusion: Wrong study design.
- Rong, C., H. F. Jiang, R. W. Zhang, L. J. Zhang, J. C. Zhang, J. Zhang, and X. S. Feng, "Factors Associated with Relapse Among Heroin Addicts: Evidence from a Two-Year Community-Based Follow-Up Study in China," *International Journal of Environmental Research and Public Health*, Vol. 13, No. 2, January 28, 2016, p. 177. Reason for exclusion: No functional outcomes.

- Roozafzoon, R., A. Goodarzi, N. Vouseoghi, M. Sedaghati, P. Yaghmaei, and M. R. Zarrindast, "Expression of NMDA Receptor Subunits in Human Peripheral Blood Lymphocytes in Opioid Addiction," *European Journal of Pharmacology*, Vol. 638, No. 1–3, July 25, 2010, pp. 29–32. Reason for exclusion: No functional outcomes.
- Rosen, M. I., T. J. McMahon, S. W. Woods, H. R. Pearsall, and T. R. Kosten, "A Pilot Study of Dextromethorphan in Naloxone-Precipitated Opiate Withdrawal," *European Journal of Pharmacology*, Vol. 307, No. 3, July 4, 1996, pp. 251–257. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Rosen, M. I., E. A. Wallace, T. J. McMahon, H. R. Pearsall, S. W. Woods, L. H. Price, and T. R. Kosten, "Buprenorphine: Duration of Blockade of Effects of Intramuscular Hydromorphone," *Drug and Alcohol Dependence*, Vol. 35, No. 2, April 1994, pp. 141–149. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Rosenblum, A., H. Joseph, C. Fong, S. Kipnis, C. Cleland, R. K. Portenoy, Andrew Rosenblum, Herman Joseph, Chunki Fong, Steven Kipnis, Charles Cleland, and Russell K. Portenoy, "Prevalence and Characteristics of Chronic Pain Among Chemically Dependent Patients in Methadone Maintenance and Residential Treatment Facilities," *JAMA: Journal of the American Medical Association*, Vol. 289, No. 18, 2003, pp. 2370–2378. Reason for exclusion: No functional outcomes.
- Rosenthal, R. N., M. R. Lofwall, S. Kim, M. Chen, K. L. Beebe, and F. J. Vocci, "Effect of Buprenorphine Implants on Illicit Opioid Use Among Abstinent Adults with Opioid Dependence Treated with Sublingual Buprenorphine: A Randomized Clinical Trial," *JAMA*, Vol. 316, No. 3, July 19, 2016, pp. 282–290. Reason for exclusion: No functional outcomes.
- Rothenberg, J. L., M. A. Sullivan, S. H. Church, A. Seracini, and E. V. Nunes, "Behavioral Naltrexone Therapy: An Integrated Treatment for Opiate Addiction and Naltrexone Maintenance," *Proceedings of the 65th Annual Scientific Meeting of the College on Problems of Drug Dependence*, Bal Harbour, Fla., June 2000. Reason for exclusion: Only measures effects of a co-intervention.
- Rotheram-Fuller, E., S. Shoptaw, S. M. Berman, and E. D. London, "Impaired Performance in a Test of Decision-Making by Opiate-Dependent Tobacco Smokers," *Drug and Alcohol Dependence*, Vol. 73, No. 1, 2004. Reason for exclusion: Wrong study design.
- Rounsaville, Bruce J., Thomas R. Kosten, Myrna M. Weissman, and Herbert D. Kleber, "A 25 Year Follow-Up of Short-Term Interpersonal Psychotherapy in Methadone-Maintained Opiate Addicts," *Comprehensive Psychiatry*, Vol. 27, No. 3, May–June 1986, pp. 201–210. Reason for exclusion: Only measures effects of a co-intervention.
- Roux, P., L. Michel, J. Cohen, M. Mora, A. Morel, J. F. Aubertin, J. C. Desenclos, B. Spire, and P. M. Carrieri, "Methadone Induction in Primary Care (ANRS-Methaville): A Phase III Randomized Intervention Trial," *BMC Public Health*, Vol. 12, 2012, p. 488. Reason for exclusion: Only measures effects of a co-intervention.

- Roux, P., M. A. Sullivan, J. Cohen, L. Fugon, J. D. Jones, S. K. Vosburg, Z. D. Cooper, J. M. Manubay, S. Mogali, and S. D. Comer, "Buprenorphine/Naloxone as a Promising Therapeutic Option for Opioid Abusing Patients with Chronic Pain: Reduction of Pain, Opioid Withdrawal Symptoms, and Abuse Liability of Oral Oxycodone," *Pain*, Vol. 154, No. 8, August 2013, pp. 1442–1448. Reason for exclusion: Wrong setting (i.e., inpatient).
- Ruetsch, C., "Treating Prescription Opioid Dependence," *JAMA*, Vol. 312, No. 11, 2014. Reason for exclusion: Wrong study design.
- Ruetsch, C., and J. Tkacz, "The Effect of a Telephonic Patient Support Program on Treatment Outcomes Among a Sample of Opioid Dependence Patients: One Year Follow-Up," *Value in Health*, Vol. 13, No. 3, 2010. Reason for exclusion: Only measures effects of a co-intervention.
- Ruetsch, C., and J. Tkacz, "A Longitudinal Analysis of the Effect of Buprenorphine–Medication Assisted Treatment (B-MAT) and a Structured Patient Support Program on B-MAT Adherence in a National Sample of Opioid Dependent Patients," *Value in Health*, Vol. 13, No. 3, 2010. Reason for exclusion: Only measures effects of a co-intervention.
- Ruetsch, C., J. Tkacz, T. L. McPherson, and J. Cacciola, "The Effect of Telephonic Patient Support on Treatment for Opioid Dependence: Outcomes at One Year Follow-Up," *Addictive Behavior*, Vol. 37, No. 5, May 2012, pp. 686–689. Reason for exclusion: Only measures effects of a co-intervention.
- Ruger, J. P., M. Chawarski, M. Mazlan, N. Ng, and R. Schottenfeld, "Cost-Effectiveness of Buprenorphine and Naltrexone Treatments for Heroin Dependence in Malaysia," *PLoS One*, Vol. 7, No. 12, 2012, p. e50673. Reason for exclusion: No functional outcomes.
- Rundio, A., "Buprenorphine Prescribing by APRNs," *Journal of Addictions Nursing*, Vol. 23, No. 1, February 2012, pp. 80–81. Reason for exclusion: Wrong study design.
- Rush, C. R., and J. A. Ali, "Naltrexone Does Not Attenuate the Acute Behavioral Effects of Ethanol or Pentobarbital in Humans," *Behavioural Pharmacology*, Vol. 10, No. 4, July 1999, pp. 401–413. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Sacerdote, P., S. Franchi, G. Gerra, V. Leccese, A. E. Panerai, and L. Somaini, "Buprenorphine and Methadone Maintenance Treatment of Heroin Addicts Preserves Immune Function," *Brain, Behavior, and Immunity*, Vol. 22, No. 4, May 2008, pp. 606–613. Reason for exclusion: No functional outcomes.
- Sajid, A., A. Whiteman, R. L. Bell, M. S. Greene, E. A. Engleman, and R. A. Chambers, "Prescription Drug Monitoring Program Data Tracking of Opioid Addiction Treatment Outcomes in Integrated Dual Diagnosis Care Involving Injectable Naltrexone," *American Journal on Addictions*, Vol. 25, No. 7, October 2016, pp. 557–564. Reason for exclusion: Wrong study design.

- Saleh, M. I., “Modeling Longitudinal Changes in Buprenorphine Treatment Outcome for Opioid Dependence,” *Pharmacopsychiatry*, Vol. 47, No. 7, November 2014, pp. 251–258. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Salehi, M., A. Zargar, and M. A. Ramezani, “Effects of Dextromethorphan on Reducing Methadone Dosage in Opium Addicts Undergoing Methadone Maintenance Therapy: A Double Blind Randomized Clinical Trial,” *Journal of Research in Medical Sciences*, Vol. 16, No. 10, October 2011, pp. 1354–1360. Reason for exclusion: Only measures effects of a co-intervention.
- Salisbury-Afshar, E., “Buprenorphine Maintenance vs. Methadone Maintenance or Placebo for Opioid Use Disorder,” *American Family Physician*, Vol. 91, No. 3, 2015. Reason for exclusion: Wrong study design.
- Sam, L., J. Cami, T. Fernandez, J. M. Olle, J. M. Peri, and M. Torrens, “Double Blind Assessment of Buprenorphine Withdrawal in Opiate-Addicts,” *NIDA Research Monograph*, Vol. 105, 1990, p. 455. Reason for exclusion: Conference abstract.
- Samet, J. H., A. Y. Walley, and C. Bridden, “Illicit Drugs, Alcohol, and Addiction in Human Immunodeficiency Virus,” *Panminerva Medica*, Vol. 49, No. 2, June 2007, pp. 67–77. Reason for exclusion: Wrong study design.
- San, L., G. Pomarol, J. M. Peri, J. M. Olle, and J. Cami, “Follow-Up After a Six-Month Maintenance Period on Naltrexone Versus Placebo in Heroin Addicts,” *British Journal of Addiction*, Vol. 86, No. 8, August 1991, pp. 983–990. Reason for exclusion: No functional outcomes.
- Santos, E. F., “A Naturalistic Analytic Study of Veterans Using Naltrexone in Puerto Rico,” *Boletin De La Asociacion Medica De Puerto Rico*, Vol. 81, No. 4, April 1989, pp. 134–136. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Saunders, Bill, Wilkinson, Celia, Phillips, and Mike, “The Impact of Brief Motivational Intervention with Opiate Users Attending a Methadone Programme,” *Addiction*, Vol. 90, No. 3, 1995. Reason for exclusion: Only measures effects of a co-intervention.
- Saunders, B., C. Wilkinson, and M. Phillips, “The Impact of a Brief Motivational Intervention with Opiate Users Attending a Methadone Programme,” *Addiction*, Vol. 90, No. 3, March 1995, pp. 415–424. Reason for exclusion: Only measures effects of a co-intervention.
- Saunders, E. C., M. P. McGovern, C. Lambert-Harris, A. Meier, B. McLeman, and H. Xie, “The Impact of Addiction Medications on Treatment Outcomes for Persons with Co-Occurring PTSD and Opioid Use Disorders,” *The American Journal on Addictions*, Vol. 24, No. 8, December 2015, pp. 722–731. Reason for exclusion: Only measures effects of a co-intervention.
- Saunders, J. B., and A. H. Richards, “Getting to Grips with Heroin and Other Opioid Use,” *The Medical Journal of Australia*, Vol. 173, No. 10, November 20, 2000, pp. 509–510. Reason for exclusion: Wrong study design.

- Savage, C., E. G. Karp, S. F. Curran, T. E. Hanlon, and O. L. McCabe, "Methadone/LAAM Maintenance: A Comparison Study," *Comprehensive Psychiatry*, Vol. 17, No. 3, 1976. Reason for exclusion: No functional outcomes.
- Savvas, S. M., A. A. Somogyi, and J. M. White, "The Effect of Methadone on Emotional Reactivity," *Addiction*, Vol. 107, No. 2, February 2012, pp. 388–392. Reason for exclusion: Wrong study design.
- Saxon, A. J., Y. I. Hser, G. Woody, and W. Ling, "Medication-Assisted Treatment for Opioid Addiction: Methadone and Buprenorphine," *Journal of Food and Drug Analysis*, Vol. 21, No. 4, December 2013, pp. S69–S72. Reason for exclusion: No functional outcomes.
- Saxon, A. J., W. Ling, M. Hillhouse, C. Thomas, A. Hasson, A. Ang, G. Doraimani, G. Tasissa, Y. Lokhnygina, J. Leimberger, R. D. Bruce, J. McCarthy, K. Wiest, P. McLaughlin, R. Bilangi, A. Cohen, G. Woody, and P. Jacobs, "Buprenorphine/Naloxone and Methadone Effects on Laboratory Indices of Liver Health: A Randomized Trial," *Drug and Alcohol Dependence*, Vol. 128, No. 1–2, February 1, 2013, pp. 71–76. Reason for exclusion: No functional outcomes.
- Saxon, A. J., E. A. Wells, C. Fleming, T. R. Jackson, and D. A. Calsyn, "Pre-Treatment Characteristics, Program Philosophy and Level of Ancillary Services as Predictors of Methadone Maintenance Treatment Outcome," *Addiction*, Vol. 91, No. 8, August 1996, pp. 1197–1209. Reason for exclusion: Only measures effects of a co-intervention.
- Schafer, I., F. J. Eiroa-Orosa, U. Verthein, C. Dilg, C. Haasen, and J. Reimer, "Effects of Psychiatric Comorbidity on Treatment Outcome in Patients Undergoing Diamorphine or Methadone Maintenance Treatment," *Psychopathology*, Vol. 43, No. 2, 2010, pp. 88–95. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Schechter, M. T., "NAOMI—Her Time Has Come. North American Opiate Medication Initiative," *Journal of Urban Health*, Vol. 79, No. 2, June 2002, pp. 164–165. Reason for exclusion: Wrong study design.
- Schechter, A., "The Role of Narcotic Antagonists in the Rehabilitation of Opiate Addicts: A Review of Naltrexone," *American Journal of Drug and Alcohol Abuse*, Vol. 7, No. 1, 1980, pp. 1–18. Reason for exclusion: Wrong study design.
- Scherbaum, N., T. Finkbeiner, K. Leifert, and M. Gastpar, "The Efficacy of L-Methadone and Racemic Methadone in Substitution Treatment for Opiate Addicts—A Double-Blind Comparison," *Pharmacopsychiatry*, Vol. 29, No. 6, 1996, pp. 212–215. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Scherbaum, N., J. Kluwig, M. Specka, D. Krause, B. Merget, T. Finkbeiner, and M. Gastpar, "Group psychotherapy for Opiate Addicts in Methadone Maintenance Treatment—A Controlled Trial," *European Addiction Research*, Vol. 11, No. 4, 2005, pp. 163–171. Reason for exclusion: Only measures effects of a co-intervention.

- Schiff, M., S. Levit, and R. C. Moreno, "Retention and Illicit Drug Use Among Methadone Patients in Israel: A Gender Comparison," *Addictive Behaviors*, Vol. 32, No. 10, October 2007, pp. 2108–2119. Reason for exclusion: Wrong study design.
- Schindler, S. D., R. Ortner, A. Peternell, H. Eder, E. Opgenoorth, and G. Fischer, "Maintenance Therapy with Synthetic Opioids and Driving Aptitude," *European Addiction Research*, Vol. 10, No. 2, 2004, pp. 80–87. Reason for exclusion: Wrong study design.
- Schluger, J. H., G. Bart, M. Green, A. Ho, and M. J. Kreek, "Corticotropin-Releasing Factor Testing Reveals a Dose-Dependent Difference in Methadone Maintained vs. Control Subjects," *Neuropsychopharmacology*, Vol. 28, No. 5, May 2003, pp. 985–994. Reason for exclusion: No functional outcomes.
- Schmittner, John, Jennifer R. Schroeder, David H. Epstein, and Kenzie L. Preston, "Menstrual Cycle Length During Methadone Maintenance," *Addiction*, Vol. 100, No. 6, June 2005, pp. 829–836. Reason for exclusion: Wrong study design.
- Schmitz, J. M., H. M. Rhoades, R. Elk, D. Creson, I. Hussein, and J. Grabowski, "Medication Take-Home Doses and Contingency Management," *Experimental and Clinical Psychopharmacology*, Vol. 6, No. 2, May 1998, pp. 162–168. Reason for exclusion: Only measures effects of a co-intervention.
- Schottenfeld, R. S., M. C. Chawarski, and M. Mazlan, "Maintenance Treatment with Buprenorphine and Naltrexone for Heroin Dependence in Malaysia: A Randomised, Double-Blind, Placebo-Controlled Trial," *Lancet*, Vol. 371, No. 9631, June 28, 2008, pp. 2192–2200. Reason for exclusion: No functional outcomes.
- Schottenfeld, R. S., M. C. Chawarski, J. R. Pakes, M. V. Pantalon, K. M. Carroll, and T. R. Kosten, "Methadone Versus Buprenorphine with Contingency Management or Performance Feedback for Cocaine and Opioid Dependence," *American Journal of Psychiatry*, Vol. 162, No. 2, February 2005, pp. 340–349. Reason for exclusion: Only measures effects of a co-intervention.
- Schottenfeld, R. S., J. Pakes, P. O'Connor, M. Chawarski, A. Oliveto, and T. R. Kosten, "Thrice-Weekly Versus Daily Buprenorphine Maintenance," *Biological Psychiatry*, Vol. 47, No. 12, June 15, 2000, pp. 1072–1079. Reason for exclusion: No functional outcomes.
- Schottenfeld, R. S., J. R. Pakes, and T. R. Kosten, "Prognostic Factors in Buprenorphine-Versus Methadone-Maintained Patients," *The Journal of Nervous and Mental Disease*, Vol. 186, No. 1, January 1998, pp. 35–43. Reason for exclusion: No functional outcomes.
- Schottenfeld, R. S., J. R. Pakes, A. Oliveto, D. Ziedonis, and T. R. Kosten, "Buprenorphine vs. Methadone Maintenance Treatment for Concurrent Opioid Dependence and Cocaine Abuse," *Archives of General Psychiatry*, Vol. 54, No. 8, August 1997, pp. 713–720. Reason for exclusion: No functional outcomes.

- Schroeder, J. R., D. H. Epstein, A. Umbricht, and K. L. Preston, "Changes in HIV Risk Behaviors Among Patients Receiving Combined Pharmacological and Behavioral Interventions for Heroin and Cocaine Dependence," *Addictive Behaviors*, Vol. 31, No. 5, May 2006, pp. 868–879. Reason for exclusion: Only measures effects of a co-intervention.
- Schuh, K. J., S. L. Walsh, G. E. Bigelow, K. L. Preston, and M. L. Stitzer, "Buprenorphine, Morphine and Naloxone Effects During Ascending Morphine Maintenance in Humans," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 278, No. 2, 1996. Reason for exclusion: Wrong setting (i.e., inpatient).
- Schuh, Kory J., Sharon L. Walsh, and Maxine L. Stitzer, "Onset, Magnitude and Duration of Opioid Blockade Produced by Buprenorphine and Naltrexone in Humans," *Psychopharmacology*, Vol. 145, No. 2, July 1999, pp. 162–174. Reason for exclusion: Wrong setting (i.e., inpatient).
- Schulte, M., Y. Hser, A. Saxon, E. Evans, L. Li, D. Huang, M. Hillhouse, C. Thomas, and W. Ling, "Risk Factors Associated with HCV Among Opioid-Dependent Patients in a Multisite Study," *Journal of Community Health*, Vol. 40, No. 5, October 2015, pp. 940–947. Reason for exclusion: No functional outcomes.
- Schuman-Olivier, Zev, M. Claire Greene, Brandon G. Bergman, and John F. Kelly, "Is Residential Treatment Effective for Opioid Use Disorders? A Longitudinal Comparison of Treatment Outcomes Among Opioid Dependent, Opioid Misusing, and Non-Opioid Using Emerging Adults with Substance Use Disorder," *Drug and Alcohol Dependence*, Vol. 144, 2014, pp. 178–185. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Schuman-Olivier, Zev, Roger D. Weiss, Bettina B. Hoepfner, Jacob Borodovsky, and Mark J. Albanese, "Emerging Adult Age Status Predicts Poor Buprenorphine Treatment Retention," *Journal of Substance Abuse Treatment*, Vol. 47, No. 3, September 2014, pp. 202–212. Reason for exclusion: Wrong study design.
- Schuster, C. R., B. B. Smith, and J. H. Jaffe, "Drug Abuse in Heroin Users. An Experimental Study of Self-Administration of Methadone, Codeine, and Pentazocine," *Archives of General Psychiatry*, Vol. 24, No. 4, 1971. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Schwartz, K., I. Herman, G. Peer, A. Weizman, and M. Rehavi, "Elevated Platelet Vesicular Monoamine Transporter 2 in Former Heroin Addicts Maintained on Methadone," *Journal of Neural Transmission*, Vol. 114, No. 2, February 2007, pp. 281–284. Reason for exclusion: No functional outcomes.
- Schwartz, R. P., P. K. Alexandre, S. M. Kelly, K. E. O'Grady, J. Gryczynski, and J. H. Jaffe, "Interim Versus Standard Methadone Treatment: A Benefit-Cost Analysis," *Journal of Substance Abuse Treatment*, Vol. 46, No. 3, March 2014, pp. 306–314. Reason for exclusion: Only measures effects of a co-intervention.

- Schwartz, R. P., D. A. Highfield, J. H. Jaffe, J. V. Brady, C. B. Butler, C. O. Rouse, J. M. Callaman, K. E. O'Grady, and R. J. Battjes, "A Randomized Controlled Trial of Interim Methadone Maintenance," *Archives of General Psychiatry*, Vol. 63, No. 1, January 2006, pp. 102–109. Reason for exclusion: No functional outcomes.
- Schwartz, R. P., J. H. Jaffe, D. A. Highfield, J. M. Callaman, and K. E. O'Grady, "A Randomized Controlled Trial of Interim Methadone Maintenance: 10-Month Follow-Up," *Drug and Alcohol Dependence*, Vol. 86, No. 1, January 5, 2007, pp. 30–36. Reason for exclusion: Article reporting follow up of included studies.
- Schwartz, R. P., S. M. Kelly, K. E. O'Grady, D. Gandhi, and J. H. Jaffe, "Interim Methadone Treatment Compared to Standard Methadone Treatment: 4-Month Findings," *Journal of Substance Abuse Treatment*, Vol. 41, No. 1, July 2011, pp. 21–29. Reason for exclusion: Only measures effects of a co-intervention.
- Schwartz, R. P., S. M. Kelly, K. E. O'Grady, D. Gandhi, and J. H. Jaffe, "Randomized Trial of Standard Methadone Treatment Compared to Initiating Methadone Without Counseling: 12-Month Findings," *Addiction*, Vol. 107, No. 5, May 2012, pp. 943–952. Reason for exclusion: Only measures effects of a co-intervention.
- "Seattle OTP Emphasizes Safety Amid Policy Responses to Rx Abuse . . . Opioid Treatment Program," *Alcoholism and Drug Abuse Weekly*, Vol. 24, No. 8, 2012, pp. 1–8. Reason for exclusion: Wrong study design.
- Sedaghati, M., N. Vouseoghi, A. Goodarzi, P. Yaghmaei, A. Mokri, and M. R. Zarrindast, "Expression of NR3B but Not NR2D Subunit of NMDA Receptor in Human Blood Lymphocytes Can Serve as a Suitable Peripheral Marker for Opioid Addiction Studies," *European Journal of Pharmacology*, Vol. 633, No. 1–3, May 10, 2010, pp. 50–54. Reason for exclusion: No functional outcomes.
- Sell, L. A., P. J. Cowen, and P. J. Robson, "Ondansetron and Opiate Craving. A Novel Pharmacological Approach to Addiction," *British Journal of Psychiatry*, Vol. 166, No. 4, April 1995, pp. 511–514. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Senay, E. C., W. Dorus, F. Goldberg, and W. Thornton, "Withdrawal from Methadone Maintenance. Rate of Withdrawal and Expectation," *Archives of General Psychiatry*, Vol. 34, No. 3, March 1977, pp. 361–367. Reason for exclusion: No functional outcomes.
- Senay, E. C., W. Dorus, and P. F. Renault, "Methadyl Acetate and Methadone. An Open Comparison," *JAMA*, Vol. 237, No. 2, January 10, 1977, pp. 138–142. Reason for exclusion: No functional outcomes.
- Seow, S. S. W., G. Swensen, and D. Willis, "Extraneous Drug Use in Methadone-Supported Patients," *Medical Journal of Australia*, Vol. 1, No. 6, 1980. Reason for exclusion: No functional outcomes.

- Seow, S. S., A. J. Quigley, K. F. Ilett, L. J. Dusci, G. Swensen, A. Harrison-Stewart, and L. Rapoport, "Buprenorphine: A New Maintenance Opiate?" *The Medical Journal of Australia*, Vol. 144, No. 8, April 14, 1986, pp. 407–411. Reason for exclusion: No functional outcomes.
- Sha, L. J., Z. Y. Liu, Z. X. Zhang, L. X. Cheng, and H. Chen, "Treatment of 100 Heroin Addicts with Dihydroetorphine, Methadone and Buprenorphine," *Medical Journal of National Defending Force in Southwest China*, Vol. 7, No. 5, 1997. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Shearer, James, Alex D. Wodak, and Kate A. Dolan, "Evaluation of a Prison-Based Naltrexone Program," *International Journal of Prisoner Health*, Vol. 3, No. 3, September 2007, pp. 214–224. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Sherman, J., "Women on Methadone," *Australian Family Physician*, Vol. 35, No. 11, November 2006, p. 839; author reply 839. Reason for exclusion: Wrong study design.
- Sheynin, J., A. A. Moustafa, K. D. Beck, R. J. Servatius, P. A. Casbolt, P. Haber, M. Elsayed, L. Hogarth, and C. E. Myers, "Exaggerated Acquisition and Resistance to Extinction of Avoidance Behavior in Treated Heroin-Dependent Men," *Journal of Clinical Psychiatry*, Vol. 77, No. 3, March 2016, pp. 386–394. Reason for exclusion: Wrong study design.
- Shi, J., L. Y. Zhao, M. L. Copersino, Y. X. Fang, Y. Chen, J. Tian, Y. Deng, Y. Shuai, J. Jin, and L. Lu, "PET Imaging of Dopamine Transporter and Drug Craving During Methadone Maintenance Treatment and After Prolonged Abstinence in Heroin Users," *European Journal of Pharmacology*, Vol. 579, No. 1–3, 2008. Reason for exclusion: Wrong study design.
- Shmygalev, S., M. Damm, K. Weckbecker, G. Berghaus, F. Petzke, and R. Sabatowski, "The Impact of Long-Term Maintenance Treatment with Buprenorphine on Complex Psychomotor and Cognitive Function," *Drug and Alcohol Dependence*, Vol. 117, No. 2–3, September 1, 2011, pp. 190–197. Reason for exclusion: Wrong study design.
- Shufman, E. N., S. Porat, E. Witztum, D. Gandacu, R. Bar-Hamburger, and Y. Ginath, "The Efficacy of Naltrexone in Preventing Reabuse of Heroin After Detoxification," *Biological Psychiatry*, Vol. 35, No. 12, June 15, 1994, pp. 935–945. Reason for exclusion: No functional outcomes.
- Shuster, Alyson, "Comparison of Buprenorphine Versus Methadone Treatment in a Medicaid Population," *Dissertation Abstracts International: Section B: The Sciences and Engineering*, Vol. 74, No. 3-B(E), 2013. Reason for exclusion: Dissertation.
- Sibbald, B., "Methadone Maintenance Expands Inside Federal Prisons," *Canadian Medical Association Journal*, Vol. 167, No. 10, November 12, 2002, p. 1154. Reason for exclusion: Wrong study design.

- Sigmon, S. C., K. Dunn, K. Saulsgiver, S. Heil, and S. Higgins, "Randomized, Double-Blind Trial Evaluating Buprenorphine Tapper for Prescription Opioid Abuse," *Proceedings of the 71th Annual Scientific Meeting of the College on Problems of Drug Dependence*, Reno/Sparks, Nev., June 20–25, 2009. Reason for exclusion: Conference abstract.
- Sigmon, S. C., K. E. Dunn, K. Saulsgiver, M. E. Patrick, G. J. Badger, S. H. Heil, J. R. Brooklyn, and S. T. Higgins, "A Randomized, Double-Blind Evaluation of Buprenorphine Taper Duration in Primary Prescription Opioid Abusers," *JAMA Psychiatry*, Vol. 70, No. 12, December 2013, pp. 1347–1354. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Sikowitz, D., D. M. Ziedonis, M. Hayes, and T. R. Kosten, "Prolonged Adjunctive Clonidine Usage Improves Treatment Outcomes for Opioid Addicts Given Naltrexone," *NIDA Research Monograph*, Vol. 141, 1994. Reason for exclusion: Only measures effects of a co-intervention.
- Simojoki, K., P. Lillsunde, N. Lintzeris, and H. Alho, "Bioavailability of Buprenorphine from Crushed and Whole Buprenorphine (Subutex) Tablets," *European Addiction Research*, Vol. 16, No. 2, 2010, pp. 85–90. Reason for exclusion: No functional outcomes.
- Simpson, D. D., G. W. Joe, J. M. Greener, and G. A. Rowan-Szal, "Modeling Year 1 Outcomes with Treatment Process and Post-Treatment Social Influences," *Substance Use and Misuse*, Vol. 35, No. 12–14, October–December 2000, pp. 1911–1930. Reason for exclusion: Wrong study design.
- Simpson, D. D., G. W. Joe, G. A. Rowan-Szal, and J. M. Greener, "Drug Abuse Treatment Process Components That Improve Retention," *Journal of Substance Abuse Treatment*, Vol. 14, No. 6, November–December 1997, pp. 565–572. Reason for exclusion: Only measures effects of a co-intervention.
- Singhal, A., B. M. Tripathi, H. R. Pal, R. Jena, and R. Jain, "Effect of Buprenorphine on Psychomotor Functions in Patients on Buprenorphine Maintenance," *Journal of Opioid Management*, Vol. 4, No. 1, January–February 2008, pp. 41–47. Reason for exclusion: Wrong study design.
- Sinha, R., A. Kimmerling, C. Doebrick, and T. R. Kosten, "Effects of Lofexidine on Stress-Induced and Cue-Induced Opioid Craving and Opioid Abstinence Rates: Preliminary Findings," *Psychopharmacology (Berl)*, Vol. 190, No. 4, March 2007, pp. 569–574. Reason for exclusion: Only measures effects of a co-intervention.
- Skeie, I., M. Brekke, M. Lindbaek, and H. Waal, "Somatic Health Among Heroin Addicts Before and During Opioid Maintenance Treatment: A Retrospective Cohort Study," *BMC Public Health*, Vol. 8, January 31, 2008, p. 43. Reason for exclusion: Wrong study design.
- Song, S., J. Tkacz, and C. Ruetsch, "Medication Compliance Enhancement Program for Opioid Dependence: Effectiveness and Perspectives After 6 Months," *Journal of the American Pharmacists Association*, Vol. 50, No. 2, March–April 2010, p. 312. Reason for exclusion: Only measures effects of a co-intervention.

- Sorensen, J. L., C. L. Masson, K. Delucchi, K. Sporer, P. G. Barnett, F. Mitsuishi, C. Lin, Y. Song, T. Chen, and S. M. Hall, "Randomized Trial of Drug Abuse Treatment-Linkage Strategies," *Journal of Consulting and Clinical Psychology*, Vol. 73, No. 6, December 2005, pp. 1026–1035. Reason for exclusion: Only measures effects of a co-intervention.
- Soyka, M., C. Limmer, R. Lehnert, G. Koller, G. Martin, H. Kufner, S. Kagerer, and A. Haberthur, "A Comparison of Cognitive Function in Patients Under Maintenance Treatment with Heroin, Methadone, or Buprenorphine and Healthy Controls: An Open Pilot Study," *American Journal of Drug and Alcohol Abuse*, Vol. 37, No. 6, November 2011, pp. 497–508. Reason for exclusion: Wrong study design.
- Soyka, M., A. Träder, J. Klotsche, M. Backmund, G. Buhringer, J. T. Rehm, and H.-U. Wittchen, "Mortality in Long-Term Opioid Maintenance Treatment in Germany: Frequency, Causes and Predictors. [German]," *Suchtmedizin in Forschung und Praxis*, Vol. 13, No. 5, 2011. Reason for exclusion: Wrong study design.
- Soyka, Michael, Anna Träder, Jens Klotsche, Markus Backmund, Gerhard Bühringer, Jürgen Rehm, and Hans-Ulrich Wittchen, "Six-Year Mortality Rates of Patients in Methadone and Buprenorphine Maintenance Therapy: Results from a Nationally Representative Cohort Study," *Journal of Clinical Psychopharmacology*, Vol. 31, No. 5, October 2011, pp. 678–680. Reason for exclusion: Wrong study design.
- Soyka, M., C. Zingg, G. Koller, and K. Hennig-Fast, "Cognitive Function in Short- and Long-Term Substitution Treatment: Are There Differences?" *The World Journal of Biological Psychiatry*, Vol. 11, No. 2, Part 2, March 2010, pp. 400–408. Reason for exclusion: Wrong study design.
- Soyka, M., C. Zingg, G. Koller, and H. Kuefner, "Retention Rate and Substance Use in Methadone and Buprenorphine Maintenance Therapy and Predictors of Outcome: Results from a Randomized Study," *International Journal of Neuropsychopharmacology*, Vol. 11, No. 5, August 2008, pp. 641–653. Reason for exclusion: No functional outcomes.
- Specka, M., T. Finkbeiner, E. Lodemann, K. Leifert, J. Kluwig, and M. Gastpar, "Cognitive-Motor Performance of Methadone-Maintained Patients," *European Addiction Research*, Vol. 6, No. 1, March 2000, pp. 8–19. Reason for exclusion: Wrong study design.
- Spiga, R., D. B. Huang, R. A. Meisch, and J. Grabowski, "Human Methadone Self-Administration: Effects of Diazepam Pretreatment," *Experimental and Clinical Psychopharmacology*, Vol. 9, No. 1, February 2001, pp. 40–46. Reason for exclusion: Wrong study design.
- Sporer, K. A., "Strategies for Preventing Heroin Overdose," *BMJ*, Vol. 326, No. 7386, February 22, 2003, pp. 442–444. Reason for exclusion: Wrong study design.

- Springer, S. A., S. E. Brown, A. Di Paola, and F. L. Altice, "Correlates of Retention on Extended-Release Naltrexone Among Persons Living with HIV Infection Transitioning to the Community from the Criminal Justice System," *Drug and Alcohol Dependence*, Vol. 157, December 1, 2015, pp. 158–165. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Springer, S. A., S. Chen, and F. L. Altice, "Improved HIV and Substance Abuse Treatment Outcomes for Released HIV-Infected Prisoners: The Impact of Buprenorphine Treatment," *Journal of Urban Health*, Vol. 87, No. 4, July 2010, pp. 592–602. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Spruit, I. P., "The Effectiveness of the Medical Prescription of Heroin Studied by Randomized Trials in the Netherlands, Watched Suspiciously by Parliament and Neighborhoods," *Substance Use and Misuse*, Vol. 37, No. 4, March 2002, pp. 555–563. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Staab, C., "My Trauma Patient Takes Methadone: How Do I Manage His Pain?" *Journal of Emergency Nursing*, Vol. 40, No. 5, September 2014, pp. 509–511. Reason for exclusion: Only measures effects of a co-intervention.
- Staak, M., G. Berghaus, R. Glazinski, K. Hoher, S. Joo, and B. Friedel, "Experimental Research on Driver Fitness of Methadone-Substituted Heroin Addicts," *Blutalkohol*, Vol. 30, No. 6, 1993. Reason for exclusion: Not English.
- Stallvik, M., B. Nordstrand, T. Kristensen, J. Bathen, E. Skogvoll, and O. Spigset, "Corrected QT Interval During Treatment with Methadone and Buprenorphine-Relation to Doses and Serum Concentrations," *Drug and Alcohol Dependence*, Vol. 129, No. 1–2, 2013. Reason for exclusion: No functional outcomes.
- Steffen, T., R. Blattler, F. Gutzwiller, and M. Zwahlen, "HIV and Hepatitis Virus Infections Among Injecting Drug Users in a Medically Controlled Heroin Prescription Programme," *European Journal of Public Health*, Vol. 11, No. 4, December 2001, pp. 425–430. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Steffen, T., S. Christen, R. Blattler, and F. Gutzwiller, "Infectious Diseases and Public Health: Risk-Taking Behavior During Participation in the Swiss Program for a Medical Prescription of Narcotics (PROVE)," *Substance Use and Misuse*, Vol. 36, No. 1–2, January 2001, pp. 71–89. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Steigman, I., "Avatar Based Recovery Platform: New Directions in the Treatment of Chemical Dependency," *European Psychiatry*, Vol. 27, Supplement 1, 2012, p. P-92. Reason for exclusion: Only measures effects of a co-intervention.
- Stein, M. D., D. S. Herman, M. Kettavong, P. A. Cioe, P. D. Friedmann, T. Tellioglu, and B. J. Anderson, "Antidepressant Treatment Does Not Improve Buprenorphine Retention Among Opioid-Dependent Persons," *Journal of Substance Abuse Treatment*, Vol. 39, No. 2, September 2010, pp. 157–166. Reason for exclusion: Only measures effects of a co-intervention.

- Stein, M. D., D. S. Herman, E. Moitra, J. Hecht, R. Lopez, B. J. Anderson, and R. A. Brown, "A Preliminary Randomized Controlled Trial of a Distress Tolerance Treatment for Opioid Dependent Persons Initiating Buprenorphine," *Drug and Alcohol Dependence*, Vol. 147, February 1, 2015, pp. 243–250. Reason for exclusion: Only measures effects of a co-intervention.
- Stein, M. D., M. E. Kurth, K. M. Sharkey, B. J. Anderson, R. P. Corso, and R. P. Millman, "Trazodone for Sleep Disturbance During Methadone Maintenance: A Double-Blind, Placebo-Controlled Trial," *Drug and Alcohol Dependence*, Vol. 120, No. 1, 2012, pp. 65–73. Reason for exclusion: Only measures effects of a co-intervention.
- Stella, L., C. D'Ambra, F. Mazzeo, A. Capuano, F. Del Franco, A. Avolio, and F. Ambrosino, "Naltrexone Plus Benzodiazepine Aids Abstinence in Opioid-Dependent Patients," *Life Sciences*, Vol. 77, No. 21, October 7, 2005, pp. 2717–2722. Reason for exclusion: Only measures effects of a co-intervention.
- Stenbacka, M., L. Brandt, and L. Lettholm, "Leaving Methadone Maintenance Treatment: The Role of Personality Traits and Psychiatric Status," *Journal of Psychoactive Drugs*, Vol. 36, No. 2, June 2004, pp. 227–234. Reason for exclusion: No functional outcomes.
- Stimmel, B., "QT or Not QT, That Is the Question: Routine Electrocardiograms for Individuals in Methadone Maintenance Treatment," *Journal of Addictive Diseases*, Vol. 30, No. 4, October 2011, pp. 307–308. Reason for exclusion: Wrong study design.
- Stimmel, B., R. Hanbury, V. Sturiano, D. Korts, G. Jackson, and M. Cohen, "Alcoholism as a Risk Factor in Methadone Maintenance. A Randomized Controlled Trial," *American Journal of Medicine*, Vol. 73, No. 5, November 1982, pp. 631–636. Reason for exclusion: Wrong study design.
- Stimmel, B., D. Korts, M. Cohen, G. Jackson, V. Sturiano, and R. Hanbury, "Opiate Addiction and Alcoholism: The Feasibility of Combined Treatment Approaches," *Annals of the New York Academy of Sciences*, Vol. 362, 1981, pp. 50–56. Reason for exclusion: Wrong study design.
- Stimolo, C., V. D. Favero, G. Zecchinato, R. Buson, D. Cusin, P. Pellachin, and P. Simonetto, "Safety and Tolerability of the Switch from Buprenorphine to Buprenorphine/Naloxone in an Italian Addiction Treatment Centre," *Clinical Drug Investigation*, Vol. 30, Supplement 1, 2010, pp. 27–31. Reason for exclusion: Wrong study design.
- Stine, S. M., and T. R. Kosten, "Opiate Withdrawal Symptoms and Cocaine Abuse: Differential Response at High Vs. Low Dose Methadone or Buprenorphine Maintenance," *NIDA Research Monograph*, Vol. 141, 1994. Reason for exclusion: Conference abstract.
- Stine, Susan M., Steven M. Southwick, Ismene L. Petrakis, Thomas R. Kosten, Dennis S. Charney, and John H. Krystal, "Yohimbine-Induced Withdrawal and Anxiety Symptoms in Opioid-Dependent Patients," *Biological Psychiatry*, Vol. 51, No. 8, April 2002, pp. 642–651. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Stitzer, M. L., M. Y. Iguchi, and L. J. Felch, "Contingent Take-Home Incentive: Effects on Drug Use of Methadone Maintenance Patients," *Journal of Consulting and Clinical Psychology*, Vol. 60, No. 6, December 1992, pp. 927–934. Reason for exclusion: Only measures effects of a co-intervention.
- Stitzer, Maxine L., Nancy M. Petry, and Jessica Peirce, "Motivational Incentives Research in the National Drug Abuse Treatment Clinical Trials Network," *Journal of Substance Abuse Treatment*, Vol. 38, Supplement 1, June 2010, pp. S61–S69. Reason for exclusion: Only measures effects of a co-intervention.
- Stoermer, R., J. Drewe, K. M. Dursteler-Mac Farland, C. Hock, F. Mueller-Spahn, D. Ladewig, R. Stohler, and R. Mager, "Safety of Injectable Opioid Maintenance Treatment for Heroin Dependence," *Biological Psychiatry*, Vol. 54, No. 8, October 15, 2003, pp. 854–861. Reason for exclusion: No functional outcomes.
- Stoller, K. B., G. E. Bigelow, S. L. Walsh, and E. C. Strain, "Effects of Buprenorphine/Naloxone in Opioid-Dependent Humans," *Psychopharmacology (Berl)*, Vol. 154, No. 3, March 2001, pp. 230–242. Reason for exclusion: Wrong setting (i.e., inpatient).
- Stoychev, K. R., K. M. Ivanov, M. G. Alexandrova, R. D. Stoychev, L. Z. Tumbev, and D. K. Kirov, "Efficacy of Melatonin in Methadone Maintained Patients with Insomnia: Results from an Open Comparison Study," *European Neuropsychopharmacology*, Vol. 19, 2009. Reason for exclusion: Only measures effects of a co-intervention.
- Strain, E. C., K. B. Stoller, S. L. Walsh, and G. E. Bigelow, "Buprenorphine/Naloxone Combination Tablet: Effects in Non-Dependent Opioid Abusers," *NIDA Research Monograph*, 1999. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Strain, E. C., K. Stoller, S. L. Walsh, and G. E. Bigelow, "Effects of Buprenorphine Versus Buprenorphine/Naloxone Tablets in Non-Dependent Opioid Abusers," *Psychopharmacology*, Vol. 148, No. 4, 2000. Reason for exclusion: Wrong setting (i.e., inpatient).
- Strain, E. C., S. L. Walsh, and G. E. Bigelow, "Blockade of Hydromorphone Effects by Buprenorphine/Naloxone and Buprenorphine," *Psychopharmacology*, Vol. 159, No. 2, 2002. Reason for exclusion: Wrong setting (i.e., inpatient).
- Strain, E. C., "Review: There Is Insufficient Evidence for Naltrexone Maintenance Treatment in Opioid Dependence," *Evidence Based Mental Health*, Vol. 6, No. 2, 2003, p. 57. Reason for exclusion: Wrong setting (i.e., inpatient).
- Strain, E. C., G. E. Bigelow, I. A. Liebson, and M. L. Stitzer, "Moderate- vs. High-Dose Methadone in the Treatment of Opioid Dependence: A Randomized Trial," *JAMA*, Vol. 281, No. 11, March 17, 1999, pp. 1000–1005. Reason for exclusion: No functional outcomes.
- Strain, E. C., G. E. Bigelow, K. L. Preston, and I. Liebson, "The Effects of Buprenorphine in Methadone-Dependent Volunteers," *NIDA Research Monograph*, Vol. 105, 1990, p. 584. Reason for exclusion: Conference abstract.

- Strain, E. C., J. A. Harrison, and G. E. Bigelow, "Induction of Opioid-Dependent Individuals onto Buprenorphine and Buprenorphine/Naloxone Soluble-Films," *Clinical Pharmacology and Therapeutics*, Vol. 89, No. 3, March 2011, pp. 443–449. Reason for exclusion: No functional outcomes.
- Strain, E. C., K. L. Preston, I. A. Liebson, and G. E. Bigelow, "Acute Effects of Buprenorphine, Hydromorphone and Naloxone in Methadone-Maintained Volunteers," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 261, No. 3, June 1992, pp. 985–993. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Strain, E. C., K. L. Preston, I. A. Liebson, and G. E. Bigelow, "Buprenorphine Effects in Methadone-Maintained Volunteers: Effects at Two Hours After Methadone," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 272, No. 2, February 1995, pp. 628–638. Reason for exclusion: Wrong setting (i.e., inpatient).
- Strain, E. C., M. L. Stitzer, I. A. Liebson, and G. E. Bigelow, "Dose-Response Effects of Methadone in the Treatment of Opioid Dependence," *Annals of Internal Medicine*, Vol. 119, No. 1, July 1, 1993, pp. 23–27. Reason for exclusion: No functional outcomes.
- Strain, E. C., M. L. Stitzer, I. A. Liebson, and G. E. Bigelow, "Buprenorphine Versus Methadone in the Treatment of Opioid-Dependent Cocaine Users," *Psychopharmacology (Berl)*, Vol. 116, No. 4, December 1994, pp. 401–406. Reason for exclusion: Only measures effects of a co-intervention.
- Strain, E. C., M. L. Stitzer, I. A. Liebson, and G. E. Bigelow, "Comparison of Buprenorphine and Methadone in the Treatment of Opioid Dependence," *American Journal of Psychiatry*, Vol. 151, No. 7, July 1994, pp. 1025–1030. Reason for exclusion: No functional outcomes.
- Strain, E. C., M. L. Stitzer, I. A. Liebson, and G. E. Bigelow, "Buprenorphine Versus Methadone in the Treatment of Opioid Dependence: Self-Reports, Urinalysis, and Addiction Severity Index," *Journal of Clinical Psychopharmacology*, Vol. 16, No. 1, February 1996, pp. 58–67. Reason for exclusion: No functional outcomes.
- Strain, E. C., S. L. Walsh, K. L. Preston, I. A. Liebson, and G. E. Bigelow, "The Effects of Buprenorphine in Buprenorphine-Maintained Volunteers," *Psychopharmacology (Berl)*, Vol. 129, No. 4, February 1997, pp. 329–338. Reason for exclusion: Wrong setting (i.e., inpatient).
- Strang, J., N. Metrebian, N. Lintzeris, L. Potts, T. Carnwath, S. Mayet, H. Williams, D. Zador, R. Evers, T. Groshkova, V. Charles, A. Martin, and L. Forzisi, "Supervised Injectable Heroin or Injectable Methadone Versus Optimised Oral Methadone as Treatment for Chronic Heroin Addicts in England After Persistent Failure in Orthodox Treatment (RIOTT): A Randomised Trial," *Lancet*, Vol. 375, No. 9729, May 29, 2010, pp. 1885–1895. Reason for exclusion: No functional outcomes.

- Strasser, J., G. A. Wiesbeck, N. Meier, R. Stohler, and K. M. Dursteler-Macfarland, "Effects of a Single 50% Extra Dose of Methadone on Heroin Craving and Mood in Lower- Versus Higher-Dose Methadone Patients," *Journal of Clinical Psychopharmacology*, Vol. 30, No. 4, August 2010, pp. 450–454. Reason for exclusion: No functional outcomes.
- Streltzer, J., R. Davidson, and D. Goebert, "An Observational Study of Buprenorphine Treatment of the Prescription Opioid Dependent Pain Patient," *The American Journal on Addiction*, Vol. 24, No. 4, June 2015, pp. 357–361. Reason for exclusion: Wrong study design.
- Strong, D. R., R. A. Brown, M. Sims, D. S. Herman, B. J. Anderson, and M. D. Stein, "Persistence on a Stress-Challenge Task Before Initiating Buprenorphine Treatment Was Associated with Successful Transition from Opioid Use to Early Abstinence," *Journal of Addiction Medicine*, Vol. 6, No. 3, September 2012, pp. 219–225. Reason for exclusion: Wrong study design.
- "Study Finds Methadone Causes Testosterone Deficiency in Men," *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 35, 2014, pp. 4–5. Reason for exclusion: Wrong study design.
- "Study Finds Some Benefit of Methadone over Buprenorphine," *Alcoholism and Drug Abuse Weekly*, Vol. 26, No. 21, 2014, pp. 3–4. Reason for exclusion: Wrong study design.
- "Study Indicates Benefits of Extended Treatment with Suboxone for Opioid-Addicted Youths," *Brown University Child and Adolescent Psychopharmacology Update*, Vol. 11, No. 1, 2009, pp. 1–3. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Subramaniam, G. A., D. Warden, A. Minhajuddin, M. J. Fishman, M. L. Stitzer, B. Adinoff, M. Trivedi, R. Weiss, J. Potter, S. A. Poole, and G. E. Woody, "Predictors of Abstinence: National Institute of Drug Abuse Multisite Buprenorphine/Naloxone Treatment Trial in Opioid-Dependent Youth," *Journal of American Academy of Child and Adolescent Psychiatry*, Vol. 50, No. 11, November 2011, pp. 1120–1128. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Suh, J. J., D. D. Langleben, R. N. Ehrman, J. G. Hakun, Z. Wang, Y. Li, S. I. Busch, C. P. O'Brien, and A. R. Childress, "Low Prefrontal Perfusion Linked to Depression Symptoms in Methadone-Maintained Opiate-Dependent Patients," *Drug and Alcohol Dependence*, Vol. 99, No. 1–3, January 1, 2009, pp. 11–17. Reason for exclusion: Wrong study design.
- Sullivan, J. G., and L. Webster, "Novel Buccal Film Formulation of Buprenorphine-Naloxone for the Maintenance Treatment of Opioid Dependence: A 12-Week Conversion Study," *Clinical Therapy*, Vol. 37, No. 5, May 1, 2015, pp. 1064–1075. Reason for exclusion: Wrong study design.
- Sullivan, L. E., D. Barry, B. A. Moore, M. C. Chawarski, J. M. Tetrault, M. V. Pantalon, P. G. O'Connor, R. S. Schottenfeld, and D. A. Fiellin, "A Trial of Integrated Buprenorphine/Naloxone and HIV Clinical Care," *Clinical Infectious Diseases*, Vol. 43, Supplement 4, December 15, 2006, pp. S184–190. Reason for exclusion: Only measures effects of a co-intervention.

- Sullivan, L. E., M. Chawarski, P. G. O'Connor, R. S. Schottenfeld, and D. A. Fiellin, "The Practice of Office-Based Buprenorphine Treatment of Opioid Dependence: Is It Associated with New Patients Entering into Treatment?" *Drug and Alcohol Dependence*, Vol. 79, No. 1, July 2005, pp. 113–116. Reason for exclusion: Wrong study design.
- Sullivan, M. A., A. Bisaga, A. Glass, K. Mishlen, M. Pavlicova, K. M. Carpenter, J. J. Mariani, F. R. Levin, and E. V. Nunes, "Opioid Use and Dropout in Patients Receiving Oral Naltrexone with or Without Single Administration of Injection Naltrexone," *Drug and Alcohol Dependence*, Vol. 147, February 1, 2015, pp. 122–129. Reason for exclusion: No functional outcomes.
- Sullivan, M. A., A. Bisaga, J. J. Mariani, A. Glass, F. R. Levin, S. D. Comer, and E. V. Nunes, "Naltrexone Treatment for Opioid Dependence: Does Its Effectiveness Depend on Testing the Blockade?" *Drug and Alcohol Dependence*, Vol. 133, No. 1, November 1, 2013, pp. 80–85. Reason for exclusion: No functional outcomes.
- Sullivan, S., "Opioid Substitution: Improving Cost-Efficiency," *Bulletin of the World Health Organization*, Vol. 91, No. 2, February 1, 2013, p. 83. Reason for exclusion: Wrong study design.
- Svikis, D., K. Silverman, N. Haug, H. Jones, R. Mejia, and M. Stitzer, "Clinical Efficacy of Alternative Voucher Incentive Programs for Pregnant Drug-Dependent Women," *62nd Annual Scientific Meeting of the College on Problems of Drug Dependence*, San Juan, Puerto Rico, June 17–22, 2000. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Swift, R. M., "Naltrexone and Nalmefene: Any Meaningful Difference?" *Biological Psychiatry*, Vol. 73, No. 8, April 15, 2013, pp. 700–701. Reason for exclusion: Wrong study design.
- Syed, Y. Y., and G. M. Keating, "Extended-Release Intramuscular Naltrexone (VIVITROL[R]): A Review of Its Use in the Prevention of Relapse to Opioid Dependence in Detoxified Patients," *CNS Drugs*, Vol. 27, No. 10, October 2013, pp. 851–861. Reason for exclusion: Wrong study design.
- Tabatabaei-Jafari, H., H. Ekhtiari, H. Ganjgahi, P. Hassani-Abharian, M. A. Oghabian, A. Moradi, N. Sadighi, and M. Zarei, "Patterns of Brain Activation During Craving in Heroin Dependents Successfully Treated by Methadone Maintenance and Abstinence-Based Treatments," *Journal of Addiction Medicine*, Vol. 8, No. 2, March–April 2014, pp. 123–129. Reason for exclusion: No functional outcomes.
- Tacke, U., "Abuse of Buprenorphine by Intravenous Injection—The French Connection," *Addiction*, Vol. 97, No. 10, October 2002, p. 1355. Reason for exclusion: Wrong study design.
- Tang, Y. L., D. Zhao, C. Zhao, and J. F. Cubells, "Opiate Addiction in China: Current Situation and Treatments," *Addiction*, Vol. 101, No. 5, May 2006, pp. 657–665. Reason for exclusion: No functional outcomes.

- Tanner, G. R., N. Bordon, S. Conroy, and D. Best, "Comparing Methadone and Suboxone in Applied Treatment Settings: The Experiences of Maintenance Patients in Lanarkshire," *Journal of Substance Use*, Vol. 16, No. 3, 2011. Reason for exclusion: Wrong study design.
- Tanum, L., K. Solli, A. Opheim, K. Sharma-Haase, H. Latif, and N. Kunoe, "Optimal Prevention of Relapse Among Opioid Users: A 12-Week Randomized Controlled Trial of Extended-Release Naltrexone Injections Versus Daily Buprenorphine-Naloxone," *Neuropsychopharmacology*, Vol. 41, Supplement 1, 2016, pp. S265–S266. Reason for exclusion: Conference abstract.
- Targosz, D., E. Kolarzyk, and D. Pach, "The Changes in Spirometric Measurements During 6 Months' Methadone Maintenance Treatment in Opiate Dependent Patients," *Przegl Lek*, Vol. 58, No. 4, 2001, pp. 254–257. Reason for exclusion: Wrong study design.
- Teagle, S., "Depot Naltrexone Appears Safe and Effective for Heroin Addiction," *Nida Notes*, Vol. 21, No. 3, 2007, p. 7. Reason for exclusion: Wrong study design.
- Teichtahl, H., A. Prodromidis, B. Miller, G. Cherry, and I. Kronborg, "Sleep-Disordered Breathing in Stable Methadone Programme Patients: A Pilot Study," *Addiction*, Vol. 96, No. 3, March 2001, pp. 395–403. Reason for exclusion: Wrong study design.
- Termorshuizen, F., A. Krol, M. Prins, R. Geskus, W. van den Brink, and E. J. van Ameijden, "Prediction of Relapse to Frequent Heroin Use and the Role of Methadone Prescription: An Analysis of the Amsterdam Cohort Study Among Drug Users," *Drug and Alcohol Dependence*, Vol. 79, No. 2, August 1, 2005, pp. 231–240. Reason for exclusion: Wrong study design.
- Teruya, Cheryl, Robert P. Schwartz, Shannon Gwin Mitchell, Albert L. Hasson, Christie Thomas, Samantha H. Buoncrisiani, Yih-Ing Hser, Katharina Wiest, Allan J. Cohen, Naomi Glick, Petra Jacobs, Paul McLaughlin, and Walter Ling, "Patient Perspectives on Buprenorphine/Naloxone: A Qualitative Study of Retention During the Starting Treatment with Agonist Replacement Therapies (START) Study," *Journal of Psychoactive Drugs*, Vol. 46, No. 5, October 2014, pp. 412–426. Reason for exclusion: Wrong study design.
- Tetrault, J., B. Moore, D. Barry, P. O'Connor, R. Schottenfeld, D. Fiellin, and L. Sullivan, "Brief Versus Extended Counseling for HIV Clinic Based Buprenorphine Treatment of Opioid Dependence," *Journal of General Internal Medicine*, Vol. 26, Supplement 1, May 2011, pp. S133–S134. Reason for exclusion: Only measures effects of a co-intervention.
- Tetrault, J. M., B. A. Moore, D. T. Barry, P. G. O'Connor, R. Schottenfeld, D. A. Fiellin, and L. E. Fiellin, "Brief Versus Extended Counseling Along with Buprenorphine/Naloxone for HIV-Infected Opioid Dependent Patients," *Journal of Substance Abuse Treatment*, Vol. 43, No. 4, December 2012, pp. 433–439. Reason for exclusion: Only measures effects of a co-intervention.
- Thomas, M., F. Kauders, M. Harris, J. Cooperstein, G. Hough, and R. Resnick, "Clinical Experiences with Naltrexone in 370 Detoxified Addicts," *NIDA Research Monograph*, No. 9, September 1976, pp. 88–92. Reason for exclusion: Wrong study design.

- Tiurina, A., E. Krupitsky, E. Blokhina, N. Bushara, V. Palatkin, and E. Zvartay, "Analysis of the Impact of Different Forms of Naltrexone on the Syndrome of Anhedonia and Depression Among Heroin Addicts [Conference Abstract]," *European Neuropsychopharmacology [abstracts from the 11th ECNP Regional Meeting St. Petersburg Russian Federation. April 14–16, 2011]*, Vol. 21, No. 2, 2011. Reason for exclusion: Only measures effects of a co-intervention.
- Tiurina, A., E. Krupitsky, E. Zvartay, and G. Woody, "Analysis of Impact of Different Forms of Naltrexone on the Syndrome of Anhedonia and Depression Among Heroin Addicts," *European Neuropsychopharmacology*, Vol. 21, 2011. Reason for exclusion: Only measures effects of a co-intervention.
- Tjagvad, C., S. Skurtveit, K. Linnet, L. V. Andersen, D. J. Christoffersen, and T. Clausen, "Methadone-Related Overdose Deaths in a Liberal Opioid Maintenance Treatment Programme," *European Addiction Research*, Vol. 22, No. 5, 2016, pp. 249–258. Reason for exclusion: Wrong study design.
- Tolomeo, S., S. Gray, K. Matthews, J. D. Steele, and A. Baldacchino, "Multifaceted Impairments in Impulsivity and Brain Structural Abnormalities in Opioid Dependence and Abstinence," *Psychological Medicine*, Vol. 46, No. 13, October 2016, pp. 2841–2853. Reason for exclusion: Wrong study design.
- Touzeau, D, "P-56 Efficacy of Buprenorphine/Naloxone Combination in Reducing Intravenous Misuse in Opioid-Dependent Patients (Rime Study)," *Alcohol and Alcoholism*, Vol. 49, Supplement 1, 2014. Reason for exclusion: Conference abstract.
- Tran, B. X., and L. T. Nguyen, "Impact of Methadone Maintenance on Health Utility, Health Care Utilization and Expenditure in Drug Users with HIV/AIDS," *International Journal of Drug Policy*, Vol. 24, No. 6, November 2013, pp. e105–110. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Tran, B. X., A. Ohinmaa, A. T. Duong, L. T. Nguyen, P. X. Vu, S. Mills, S. Houston, and P. Jacobs, "The Cost-Effectiveness and Budget Impact of Vietnam's Methadone Maintenance Treatment Programme in HIV Prevention and Treatment Among Injection Drug Users," *Global Public Health*, Vol. 7, No. 10, 2012, pp. 1080–1094. Reason for exclusion: Only measures effects of a co-intervention.
- Traynor, K., "Pharmacists May Dispense New Opioid Addiction Treatment," *American Journal of Health-System Pharmacy*, Vol. 59, No. 22, November 15, 2002, pp. 2150, 2152. Reason for exclusion: Wrong study design.
- "Treatment Options for Opiate Addiction," *Journal of the Kentucky Medical Association*, Vol. 102, No. 5, May 2004, pp. 221–224. Reason for exclusion: Wrong study design.
- Trueblood, B., B. A. Judson, and A. Goldstein, "Acceptability of Methadyl Acetate (LAAM) as Compared with Methadone in a Treatment Program for Heroin Addicts," *Drug and Alcohol Dependence*, Vol. 3, No. 2, 1978. Reason for exclusion: No functional outcomes.

- Tsui, J. I., M. C. Lira, D. M. Cheng, M. R. Winter, D. P. Alford, J. M. Liebschutz, J. Mao, R. R. Edwards, and J. H. Samet, "Hepatitis C Virus Infection and Pain Sensitivity in Patients on Methadone or Buprenorphine Maintenance Therapy for Opioid Use Disorders," *Drug and Alcohol Dependence*, Vol. 153, August 1, 2015, pp. 286–292. Reason for exclusion: Wrong study design.
- Tuchman, E., C. Gregory, M. Simson, and E. Drucker, "Safety, Efficacy, and Feasibility of Office-Based Prescribing and Community Pharmacy Dispensing of Methadone: Results of a Pilot Study in New Mexico," *Addictive Disorders and Their Treatment*, Vol. 5, No. 2, 2006, pp. 43–51. Reason for exclusion: Only measures effects of a co-intervention.
- Tucker, T., A. Ritter, C. Maher, and H. Jackson, "A Randomized Control Trial of Group Counseling in a Naltrexone Treatment Program," *Journal of Substance Abuse Treatment*, Vol. 27, No. 4, December 2004, pp. 277–288. Reason for exclusion: Only measures effects of a co-intervention.
- Tuma, A. Hussain, Carole Siegel, Mary J. Alexander, and Joseph Wanderling, "Effects of Compliance on Outcome Independent of Pharmacological Efficacy in the Treatment of Opioid Dependence: A Post Hoc Regression Analysis," *The American Journal on Addictions*, Vol. 2, No. 3, Summer 1993, pp. 238–249. Reason for exclusion: No functional outcomes.
- Tusel, D. J., N. A. Piotrowski, K. L. Sees, P. M. Reilly, P. Banys, P. Meek, and S. M. Hall, "Contingency Contracting for Illicit Drug Use with Opioid Addicts in Methadone Treatment," *NIDA Research Monograph*, Vol. 153, 1994. Reason for exclusion: Only measures effects of a co-intervention.
- Tuten, M., D. S. Svikis, L. Keyser-Marcus, K. E. O'Grady, and H. E. Jones, "Lessons Learned from a Randomized Trial of Fixed and Escalating Contingency Management Schedules in Opioid-Dependent Pregnant Women," *American Journal of Drug and Alcohol Abuse*, Vol. 38, No. 4, July 2012, pp. 286–292. Reason for exclusion: Only measures effects of a co-intervention.
- Tyndale, R., "Drug Addiction: A Critical Problem Calling for Novel Solutions," *Clinical Pharmacology and Therapeutics*, Vol. 83, No. 4, April 2008, pp. 503–506. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Uchtenhagen, A., F. Gutzwiller, A. Dobler-Mikola, and T. Steffen, "Programme for a Medical Prescription of Narcotics. A Synthesis of Results," *European Addiction Research*, Vol. 3, No. 4, December 1997, pp. 160–163. Reason for exclusion: Wrong study design.
- Uehlinger, C., J. Deglon, S. Livoti, S. Petitjean, D. Waldvogel, and D. Ladewig, "Comparison of Buprenorphine and Methadone in the Treatment of Opioid Dependence. Swiss Multicentre Study," *European Addiction Research*, Vol. 4, Supplement 1, 1998, pp. 13–18. Reason for exclusion: No functional outcomes.

- Umbricht-Schneiter, A., D. H. Ginn, K. M. Pabst, and G. E. Bigelow, "Providing Medical Care to Methadone Clinic Patients: Referral vs. On-Site Care," *American Journal of Public Health*, Vol. 84, No. 2, February 1994, pp. 207–210. Reason for exclusion: Only measures effects of a co-intervention.
- "Underused Medications Key to Addiction Relapse Prevention," *Brown University Psychopharmacology Update*, Vol. 16, No. 10, 2005, pp. 1–6. Reason for exclusion: Wrong study design.
- Vagenas, P., A. Di Paola, M. Herme, T. Lincoln, D. J. Skiest, F. L. Altice, and S. A. Springer, "An Evaluation of Hepatic Enzyme Elevations Among HIV-Infected Released Prisoners Enrolled in Two Randomized Placebo-Controlled Trials of Extended Release Naltrexone," *Journal of Substance Abuse Treatment*, Vol. 47, No. 1, July 2014, pp. 35–40. Reason for exclusion: No functional outcomes.
- van den Brink, W., "Science-Based Practice for Methadone Treatment: Another Step Forward," *Soz Präventivmed*, Vol. 48, Supplement 1, 2003, pp. S15–16; discussion on pp. 25–27. Reason for exclusion: Wrong study design.
- van den Brink, W., V. M. Hendriks, P. Blanken, M. W. Koeter, B. J. van Zwieten, and J. M. van Ree, "Medical Prescription of Heroin to Treatment Resistant Heroin Addicts: Two Randomised Controlled Trials," *BMJ*, Vol. 327, No. 7410, August 9, 2003, p. 310. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Vanichseni, S., B. Wongsuwan, K. Choopanya, and K. Wongpanich, "A Controlled Trial of Methadone Maintenance in a Population of Intravenous Drug Users in Bangkok: Implications for Prevention of HIV," *The International Journal of the Addictions*, Vol. 26, No. 12, December 1991, pp. 1313–1320. Reason for exclusion: No functional outcomes.
- Vastag, B., "In-Office Opiate Treatment "Not a Panacea": Physicians Slow to Embrace Therapeutic Option," *JAMA*, Vol. 290, No. 6, August 13, 2003, pp. 731–735. Reason for exclusion: Wrong study design.
- Verdejo-García, A., D. I. Lubman, K. Roffel, R. Vilar-López, E. Bora, T. MacKenzie, and M. Yücel, "Cingulate Biochemistry in Heroin Users on Substitution Pharmacotherapy," *The Australian and New Zealand Journal of Psychiatry*, Vol. 47, No. 3, 2013. Reason for exclusion: Wrong study design.
- Verthein, U., T. Beck, C. Haasen, and J. Reimer, "Mental Symptoms and Drug Use in Maintenance Treatment with Slow-Release Oral Morphine Compared to Methadone: Results of a Randomized Crossover Study," *European Addiction Research*, Vol. 21, No. 2, 2015, pp. 97–104. Reason for exclusion: Only measures effects of a co-intervention.
- Verthein, U., C. Haasen, and D. Naber, "Appropriateness of Methadone Maintenance Treatment—from Individual Practice to Evidence-Based Medicine," *Soz Präventivmed*, Vol. 48, Supplement 1, 2003, pp. S19–20; discussion on pp. S25–S27. Reason for exclusion: Wrong study design.

- Verthein, U., C. Haasen, and J. Reimer, "Switching from Methadone to Diamorphine: 2-Year Results of the German Heroin-Assisted Treatment Trial," *Substance Use Misuse*, Vol. 46, No. 8, 2011, pp. 980–991. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Verthein, U, J Reimer, R Ullmann, and C Haasen, "Psychological Well-Being in Substitution Therapy with Levomethadone and D, L-Methadone: A Double-Blind, Randomised, Cross-Over Study. [German]," *Sucht*, Vol. 53, No. 1, 2007. Reason for exclusion: Not English.
- Verthein, U., R. Ullmann, A. Lachmann, A. Düring, B. Koch, H. G. Meyer-Thompson, R. Schmidt, J. Reimer, and C. Haasen, "The Effects of Racemic D,L-Methadone and L-Methadone in Substituted Patients—A Randomized Controlled Study," *Drug and Alcohol Dependence*, Vol. 80, No. 2, November 1, 2005, pp. 267–271. Reason for exclusion: No functional outcomes.
- Verthein, U., R. Ullmann, A. Lachmann, A. Düring, B. Koch, H. G. Meyer-Thompson, R. Schmidt, J. Reimer, and C. Haasen, "The Effects of Racemic D,L-Methadone and L-Methadone in Substituted Patients—A Randomized Controlled Study," *Drug and Alcohol Dependence*, Vol. 80, No. 2, 1, 2005, pp. 267–271. Reason for exclusion: No functional outcomes.
- Vidjak, N., "Treating Heroin Addiction: Comparison of Methadone Therapy, Hospital Therapy Without Methadone, and Therapeutic Community," *Croatian Medical Journal*, Vol. 44, No. 1, 2003. Reason for exclusion: No functional outcomes.
- Vigezzi, P., L. Guglielmino, P. Marzorati, R. Silenzio, M. De Chiara, F. Corrado, L. Cocchi, and E. Cozzolino, "Multimodal Drug Addiction Treatment: A Field Comparison of Methadone and Buprenorphine Among Heroin- and Cocaine-Dependent Patients," *Journal of Substance Abuse Treatment*, Vol. 31, No. 1, July 2006, pp. 3–7. Reason for exclusion: No functional outcomes.
- Vilalta, J., J. Treserra, L. García-Esteve, M. García-Giralt, and E. Cirera, "Methadone, Clonidine and Levomepromazine in the Treatment of Opiate Abstinence Syndrome: Double-Blind Clinical Trial in Heroin-Addicted Patients Admitted to a General Hospital for Organic Pathology," *Medicina Clínica*, Vol. 88, No. 17, 1987. Reason for exclusion: Not English.
- Vinson, D. C., "Patient Abusing Alcohol or Drugs? Help Starts with a Single Question," *Journal of Family Practice*, Vol. 62, No. 2, February 2013, pp. 63–69. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Vo, Hoa T., Erika Robbins, Meghan Westwood, Debra Lezama, and Marc Fishman, "Relapse Prevention Medications in Community Treatment for Young Adults with Opioid Addiction," *Substance Abuse*, Vol. 37, No. 3, 2016, pp. 392–397. Reason for exclusion: Wrong study design.

- Vocci, F. J., R. P. Schwartz, M. E. Wilson, M. S. Gordon, T. W. Kinlock, T. T. Fitzgerald, K. E. OGrady, and J. H. Jaffe, "Buprenorphine Dose Induction in Non-Opioid-Tolerant Pre-Release Prisoners," *Drug and Alcohol Dependence*, Vol. 156, November 1, 2015, pp. 133–138. Reason for exclusion: Wrong setting (i.e., inpatient).
- Volavka, J., B. James, D. Reker, A. Mallya, D. Cho, and J. Pevnick, "EEG and Other Effects of Naltrexone and Heroin in Man," *Pharmakopsychiatrie, Neuro-Psychopharmakologie*, Vol. 12, No. 1, 1979. Reason for exclusion: Wrong study design.
- Volavka, J., R. B. Resnick, R. S. Kestenbaum, and A. M. Freedman, "Short Term Effects of Naltrexone in 155 Heroin Ex Addicts," *Biological Psychiatry*, Vol. 11, No. 6, 1976. Reason for exclusion: Wrong study design.
- Volavka, J., K. Verebey, R. Resnick, and S. Mulé, "Methadone Dose, Plasma Level, and Cross-Tolerance to Heroin in Man," *The Journal of Nervous and Mental Disease*, Vol. 166, No. 2, 1978. Reason for exclusion: No functional outcomes.
- Waal, H., "Is Sustained Release Naltrexone an Option for Heroin-Dependent Pregnant Women?" *Addiction*, Vol. 108, No. 2, February 2013, pp. 252–253. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Wachholtz, A., and G. Gonzalez, "Co-Morbid Pain and Opioid Addiction: Long Term Effect of Opioid Maintenance on Acute Pain," *Drug and Alcohol Dependence*, Vol. 145, December 1, 2014, pp. 143–149. Reason for exclusion: Wrong study design.
- Wachtel, S. R., and H. de Wit, "Naltrexone Does Not Block the Subjective Effects of Oral Delta(9)-Tetrahydrocannabinol in Humans," *Drug and Alcohol Dependence*, Vol. 59, No. 3, June 1, 2000, pp. 251–260. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Walsh, S. L., H. L. June, K. J. Schuh, K. L. Preston, G. E. Bigelow, and M. L. Stitzer, "Effects of Buprenorphine and Methadone in Methadone-Maintained Subjects," *Psychopharmacology*, Vol. 119, No. 3, 1995. Reason for exclusion: Wrong study design.
- Walsh, S. L., P. A. Nuzzo, S. Babalonis, V. Casselton, and M. R. Lofwall, "Intranasal Buprenorphine Alone and in Combination with Naloxone: Abuse Liability and Reinforcing Efficacy in Physically Dependent Opioid Abusers," *Drug and Alcohol Dependence*, Vol. 162, 2016. Reason for exclusion: No functional outcomes.
- Walsh, S. L., P. A. Nuzzo, C. Elayi, and M. R. Lofwall, "Intranasal Buprenorphine Alone and in Combination with Naloxone: Reinforcing Efficacy and Abuse Liability in Physically Dependent Opioid Abusers," *Drug and Alcohol Dependence*, Vol. 156, 2015. Reason for exclusion: Conference abstract.
- Walsh, S. L., A. E. Chausmer, E. C. Strain, and G. E. Bigelow, "Evaluation of the Mu and Kappa Opioid Actions of Butorphanol in Humans Through Differential Naltrexone Blockade," *Psychopharmacology (Berl)*, Vol. 196, No. 1, January 2008, pp. 143–155. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Walsh, S. L., S. F. Gilson, D. R. Jasinski, J. M. Stapleton, R. L. Phillips, R. F. Dannals, J. Schmidt, K. L. Preston, R. Grayson, G. E. Bigelow, et al., "Buprenorphine Reduces Cerebral Glucose Metabolism in Polydrug Abusers," *Neuropsychopharmacology*, Vol. 10, No. 3, May 1994, pp. 157–170. Reason for exclusion: Wrong setting (i.e., inpatient).
- Walsh, S. L., K. L. Preston, G. E. Bigelow, and M. L. Stitzer, "Acute Administration of Buprenorphine in Humans: Partial Agonist and Blockade Effects," *Journal of Pharmacology and Experimental Therapeutics*, Vol. 274, No. 1, July 1995, pp. 361–372. Reason for exclusion: No functional outcomes.
- Walter, M., G. A. Wiesbeck, B. Degen, J. Albrich, M. Oppel, A. Schulz, H. Schächinger, and K. M. Dürsteler-MacFarland, "Heroin Reduces Startle and Cortisol Response in Opioid-Maintained Heroin-Dependent Patients," *Addiction Biology*, Vol. 16, No. 1, 2011. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Walter, Marc, Gerhard A. Wiesbeck, Bigna Degen, Jürgen Albrich, Monika Oppel, André Schulz, Hartmut Schächinger, and Kenneth M. Dürsteler-MacFarland, "Heroin Reduces Startle and Cortisol Response in Opioid-Maintained Heroin-Dependent Patients," *Addiction Biology*, Vol. 16, No. 1, January 2011, pp. 145–151. Reason for exclusion: Wrong study design.
- Wang, D., H. Teichtahl, C. Goodman, O. Drummer, R. R. Grunstein, and I. Kronborg, "Subjective Daytime Sleepiness and Daytime Function in Patients on Stable Methadone Maintenance Treatment: Possible Mechanisms," *Journal of Clinical Sleep Medicine*, Vol. 4, No. 6, December 15, 2008, pp. 557–562. Reason for exclusion: Wrong study design.
- Wang, E. A., B. A. Moore, L. E. Sullivan, and D. A. Fiellin, "Effect of Incarceration History on Outcomes of Primary Care Office-Based Buprenorphine/Naloxone," *Journal of General Internal Medicine*, Vol. 25, No. 7, July 2010, pp. 670–674. Reason for exclusion: Wrong study design.
- Wang, G. Y., R. Kydd, and B. R. Russell, "Auditory Event-Related Potentials in Methadone Substituted Opiate Users," *Journal of Psychopharmacology*, Vol. 29, No. 9, September 2015, pp. 983–995. Reason for exclusion: Wrong study design.
- Wang, G. Y., R. Kydd, T. A. Woules, M. Jensen, and B. R. Russell, "Changes in Resting EEG Following Methadone Treatment in Opiate Addicts," *Clinical Neurophysiology*, Vol. 126, No. 5, May 2015, pp. 943–950. Reason for exclusion: No functional outcomes.
- Wang, G. Y., R. R. Kydd, and B. R. Russell, "Quantitative EEG and Low-Resolution Electromagnetic Tomography (LORETA) Imaging of Patients Undergoing Methadone Treatment for Opiate Addiction," *Clinical EEG Neuroscience*, Vol. 47, No. 3, July 2016, pp. 180–187. Reason for exclusion: No functional outcomes.
- Wang, G. Y., T. A. Woules, R. Kydd, M. Jensen, and B. R. Russell, "Neuropsychological Performance of Methadone-Maintained Opiate Users," *Journal of Psychopharmacology*, Vol. 28, No. 8, August 2014, pp. 789–799. Reason for exclusion: Wrong study design.

- Wang, L., X. Wei, X. Wang, J. Li, H. Li, and W. Jia, "Long-Term Effects of Methadone Maintenance Treatment with Different Psychosocial Intervention Models," *PLoS One*, Vol. 9, No. 2, 2014. Reason for exclusion: Only measures effects of a co-intervention.
- Wang, R. I. H., C. Kochar, T. Hasegawa, and B. L. Roh, "Clinical Comparison of Propoxyphene Napsylate and Methadone in the Treatment of Opiate Dependence," *NIDA Research Monograph Series*, Vol. 41, 1981. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wang, R. I., C. Kochar, A. T. Hasegawa, and B. L. Roh, "Propoxyphene Napsylate Compared to Methadone for Opiate Dependence," *Psychopharmacology (Berl)*, Vol. 75, No. 4, 1981, pp. 335–338. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wang, R. I., C. Kochar, A. T. Hasegawa, and B. L. Roh, "Initial Methadone Dose in Treating Opiate Addiction," *The International Journal of the Addictions*, Vol. 17, No. 2, February 1982, pp. 357–363. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wang, X., L. Tan, Y. Li, Y. Zhang, D. Zhou, T. Liu, and W. Hao, "HCV and HIV Infection Among Heroin Addicts in Methadone Maintenance Treatment (MMT) and Not in MMT in Changsha and Wuhan, China," *PLoS One*, Vol. 7, No. 9, 2012, p. e45632. Reason for exclusion: No functional outcomes.
- Wang, Y., W. Li, Q. Li, W. Yang, J. Zhu, and W. Wang, "White Matter Impairment in Heroin Addicts Undergoing Methadone Maintenance Treatment and Prolonged Abstinence: A Preliminary DTI Study," *Neuroscience Letters*, Vol. 494, No. 1, April 20, 2011, pp. 49–53. Reason for exclusion: No functional outcomes.
- Wang, Y., H. Wang, W. Li, J. Zhu, M. S. Gold, D. Zhang, L. Wang, Y. Li, X. Yan, J. Cheng, Q. Li, and W. Wang, "Reduced Responses to Heroin-Cue-Induced Craving in the Dorsal Striatum: Effects of Long-Term Methadone Maintenance Treatment," *Neuroscience Letters*, Vol. 581, October 3, 2014, pp. 120–124. Reason for exclusion: No functional outcomes.
- Warden, D., G. A. Subramaniam, T. Carmody, G. E. Woody, A. Minhajuddin, S. A. Poole, J. Potter, M. Fishman, M. Bogenschutz, A. Patkar, and M. H. Trivedi, "Predictors of Attrition with Buprenorphine/Naloxone Treatment in Opioid Dependent Youth," *Addictive Behaviors*, Vol. 37, No. 9, September 2012, pp. 1046–1053. Reason for exclusion: No functional outcomes.
- Wardle, M., A. Bershad, K. Yan, and H. Wit, "Opioid Antagonism Alters Attention to and Recognition of Emotional Expressions in Healthy Adults," *Neuropsychopharmacology*, Vol. 39, 2014. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Wardle, M. C., A. K. Bershad, M. Pulaski, and H. De Wit, "Opioid Antagonism Decreases Hedonic Responses to Social Stimuli in Healthy Adults," *Neuropsychopharmacology*, Vol. 38, Supplement 2, December 2013, p. S504. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wasserman, D. A., A. L. Stewart, and K. L. Delucchi, "Social Support and Abstinence from Opiates and Cocaine During Opioid Maintenance Treatment," *Drug and Alcohol Dependence*, Vol. 65, No. 1, December 1, 2001, pp. 65–75. Reason for exclusion: Only measures effects of a co-intervention.
- Waters, N., T. J. Gaha, and I. Reynolds, "Random Urine Analyses from Drug Addicts in a Methadone Treatment Programme," *The Medical Journal of Australia*, Vol. 2, No. 5, August 2, 1975, pp. 170–172. Reason for exclusion: Only measures effects of a co-intervention.
- Webster, Lynn, Daniel Gruener, Todd Kirby, Xiang Qinfang, Evan Tzanis, and Andrew Finn, "Evaluation of the Tolerability of Switching Patients on Chronic Full μ -Opioid Agonist Therapy to Buccal Buprenorphine," *Pain Medicine*, Vol. 17, No. 5, 2016, pp. 899–907. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Webster, L., P. Hjelmstrom, M. Sumner, and E. W. Gunderson, "Efficacy and Safety of a Sublingual Buprenorphine/Naloxone Rapidly Dissolving Tablet for the Treatment of Adults with Opioid Dependence: A Randomized Trial," *Journal of Addictive Diseases*, Vol. 35, No. 4, October–December 2016, pp. 325–338. Reason for exclusion: Wrong study design.
- Webster, L., P. Hjelmstrom, M. Sumner, and C. Welsh, "Titration to Effective Dose of Zubsolv, an Advanced Formulation of Sublingual Buprenorphine/Naloxone, in Adults with Opioid Dependence," *Journal of the American Pharmacists Association*, Vol. 55, No. 2, 2015. Reason for exclusion: Conference abstract.
- Webster, L. R., M. D. Smith, C. Unal, and A. Finn, "Low-Dose Naloxone Provides an Abuse-Deterrent Effect to Buprenorphine," *Journal of Pain Research*, Vol. 8, 2015. Reason for exclusion: Wrong setting (i.e., inpatient).
- Wedam, E. F., G. E. Bigelow, R. E. Johnson, P. A. Nuzzo, and M. C. P. Haigney, "QT-Interval Effects of Methadone, Levomethadyl, and Buprenorphine in a Randomized Trial," *Archives of Internal Medicine*, Vol. 167, No. 22, 2007. Reason for exclusion: No functional outcomes.
- Weich, L., "Medical Treatment of Opioid Dependence Within the South African Context," *The South African Dental Journal*, Vol. 65, No. 4, May 2010, pp. 168–171. Reason for exclusion: Wrong study design.
- Weich, L., C. Perkel, N. van Zyl, Rataemane, and L. Naidoo, "South African Guidelines for the Management of Opioid Dependence: Updated 2009," *The South African Medical Journal*, Vol. 100, No. 1, January 2010, pp. 1–18. Reason for exclusion: Wrong study design.

- Weinrieb, R. M., R. Barnett, K. G. Lynch, M. DePiano, A. Atanda, and K. M. Olthoff, “A Matched Comparison Study of Medical and Psychiatric Complications and Anesthesia and Analgesia Requirements in Methadone-Maintained Liver Transplant Recipients,” *Liver Transplant*, Vol. 10, No. 1, January 2004, pp. 97–106. Reason for exclusion: Wrong study design.
- Weinstein, A. M., B. T. Feldtkeller, F. Law, J. Myles, and D. J. Nutt, “The Processing of Automatic Thoughts of Drug Use and Craving in Opiate-Dependent Individuals,” *Experimental and Clinical Psychopharmacology*, Vol. 8, No. 4, November 2000, pp. 549–553. Reason for exclusion: No functional outcomes.
- Weiss, L., J. E. Egan, M. Botsko, J. Netherland, D. A. Fiellin, and R. Finkelstein, “The BHIVES Collaborative: Organization and Evaluation of a Multisite Demonstration of Integrated Buprenorphine/Naloxone and HIV Treatment,” *Journal of Acquired Immune Deficiency Syndromes*, Vol. 56, Supplement 1, 2011. Reason for exclusion: Wrong study design.
- Weiss, R. D., J. S. Potter, D. A. Fiellin, M. Byrne, H. S. Connery, W. Dickinson, J. Gardin, M. L. Griffin, M. N. Gourevitch, D. L. Haller, A. L. Hasson, Z. Huang, P. Jacobs, A. S. Kosinski, R. Lindblad, E. F. McCance-Katz, S. E. Provost, J. Selzer, E. C. Somoza, S. C. Sonne, and W. Ling, “Adjunctive Counseling During Brief and Extended Buprenorphine-Naloxone Treatment for Prescription Opioid Dependence: A 2-Phase Randomized Controlled Trial,” *Archives of General Psychiatry*, Vol. 68, No. 12, December 2011, pp. 1238–1246. Reason for exclusion: Only measures effects of a co-intervention.
- Weiss, R. D., J. S. Potter, M. L. Griffin, S. E. Provost, G. M. Fitzmaurice, K. A. McDermott, E. N. Srisarajivakul, D. R. Dodd, J. A. Dreifuss, R. K. McHugh, and K. M. Carroll, “Long-Term Outcomes from the National Drug Abuse Treatment Clinical Trials Network Prescription Opioid Addiction Treatment Study,” *Drug and Alcohol Dependence*, Vol. 150, May 1, 2015, pp. 112–119. Reason for exclusion: Wrong study design.
- Weizman, T., M. Gelkopf, Y. Melamed, M. Adelson, and A. Bleich, “Treatment of Benzodiazepine Dependence in Methadone Maintenance Treatment Patients: A Comparison of Two Therapeutic Modalities and the Role of Psychiatric Comorbidity,” *Australian and New Zealand Journal of Psychiatry*, Vol. 37, No. 4, August 2003, pp. 458–463. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wells, Elizabeth A., Ron Jackson, O. Rachel Diaz, Viki Stanton, Andrew J. Saxon, and Antoinette Krupski, “Acupuncture as an Adjunct to Methadone Treatment Services,” *The American Journal on Addictions*, Vol. 4, No. 3, Summer 1995, pp. 198–214. Reason for exclusion: Only measures effects of a co-intervention.
- West, S. L., K. K. O’Neal, and C. W. Graham, “A Meta-Analysis Comparing the Effectiveness of Buprenorphine and Methadone,” *Journal of Substance Abuse*, Vol. 12, No. 4, 2000, pp. 405–414. Reason for exclusion: No functional outcomes.
- “When Is It Time to Stop Medication-Assisted Opioid Treatment?” *Alcoholism and Drug Abuse Weekly*, Vol. 22, No. 21, 2010, pp. 3–4. Reason for exclusion: Wrong study design.

- White, Jason, James Bell, John B. Saunders, Paul Williamson, Maria Makowska, Aaron Farquharson, and Katherine L. Beebe, “Open-Label Dose-Finding Trial of Buprenorphine Implants (Probuphine)® for Treatment of Heroin Dependence,” *Drug and Alcohol Dependence*, Vol. 103, No. 1–2, July 2009, pp. 37–43. Reason for exclusion: No functional outcomes.
- White, J. M., C. Danz, J. Kneebone, S. F. La Vincente, D. A. Newcombe, and R. L. Ali, “Relationship Between LAAM-Methadone Preference and Treatment Outcomes,” *Drug and Alcohol Dependence*, Vol. 66, No. 3, May 1, 2002, pp. 295–301. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Whysner, J. A., D. B. Thomas, W. Ling, and C. Charuvastra, “On the Relative Efficacy of LAAM and Methadone,” *NIDA Research Monograph*, Vol. 27, 1979, pp. 429–433. Reason for exclusion: Wrong study design.
- Wiest, K. L., J. B. Colditz, K. Carr, V. J. Asphaug, D. McCarty, and P. A. Pilkonis, “Pain and Emotional Distress Among Substance-Use Patients Beginning Treatment Relative to a Representative Comparison Group,” *Journal of Addiction Medicine*, Vol. 8, No. 6, November–December 2014, pp. 407–414. Reason for exclusion: No functional outcomes.
- Wikner, B. N., I. Ohman, T. Selden, H. Druid, L. Brandt, and H. Kieler, “Opioid-Related Mortality and Filled Prescriptions for Buprenorphine and Methadone,” *Drug and Alcohol Review*, Vol. 33, No. 5, September 2014, pp. 491–498. Reason for exclusion: No functional outcomes.
- Wilder, C. M., S. C. Miller, E. Tiffany, T. Winhusen, E. L. Winstanley, and M. D. Stein, “Risk Factors for Opioid Overdose and Awareness of Overdose Risk Among Veterans Prescribed Chronic Opioids for Addiction or Pain,” *Journal of Addictive Diseases*, Vol. 35, No. 1, 2016, pp. 42–51. Reason for exclusion: Wrong study design.
- Williams, M. D., S. B. Hicks, and K. R. Williams, “Physical Versus Psychological Dependency During Detoxification and Maintenance,” *Proceedings. National Conference on Methadone Treatment*, Vol. 2, 1973. Reason for exclusion: Conference abstract.
- Wilson, M. E., T. W. Kinlock, M. S. Gordon, K. E. O’Grady, and R. P. Schwartz, “Postprison Release HIV-Risk Behaviors in a Randomized Trial of Methadone Treatment for Prisoners,” *The American Journal on Addictions*, Vol. 21, No. 5, September–October 2012, pp. 476–487. Reason for exclusion: No functional outcomes.
- Wilson, M. E., R. P. Schwartz, K. E. O’Grady, and J. H. Jaffe, “Impact of Interim Methadone Maintenance on HIV Risk Behaviors,” *Journal of Urban Health*, Vol. 87, No. 4, July 2010, pp. 586–591. Reason for exclusion: No functional outcomes.
- Winklbaaur, B., R. Jagsch, N. Ebner, K. Thau, and G. Fischer, “Quality of Life in Patients Receiving Opioid Maintenance Therapy. A Comparative Study of Slow-Release Morphine Versus Methadone Treatment,” *European Addiction Research*, Vol. 14, No. 2, 2008, pp. 99–105. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Wittchen, H. U., S. M. Apelt, M. Soyka, M. Gastpar, M. Backmund, J. Golz, M. R. Kraus, F. Tretter, M. Schafer, J. Siegert, N. Scherbaum, J. Rehm, and G. Buhringer, "Feasibility and Outcome of Substitution Treatment of Heroin-Dependent Patients in Specialized Substitution Centers and Primary Care Facilities in Germany: A Naturalistic Study in 2694 Patients," *Drug and Alcohol Dependence*, Vol. 95, No. 3, June 1, 2008, pp. 245–257. Reason for exclusion: Only measures effects of a co-intervention.
- Wodak, A., and R. Graham, "Naltrexone Implants as Treatment for Heroin Dependence: Part II," *Addiction Biology*, Vol. 10, No. 2, June 2005, pp. 202; author reply 202–204. Reason for exclusion: Wrong study design.
- Wodak, A. D., and W. D. Hall, "Buprenorphine: Better Late than Never," *The Medical Journal of Australia*, Vol. 175, No. 7, October 1, 2001, pp. 349–350. Reason for exclusion: Wrong study design.
- Wolfe, R. C., "Methadone Maintenance as Medical Treatment: The Continuing Need for Controlled Clinical Study," *Proceedings. National Conference on Methadone Treatment*, Vol. 1, 1973. Reason for exclusion: Wrong study design.
- Wolff, R. F., K. Reid, M. Di Nisio, D. Aune, C. Truysers, A. V. Hernandez, K. Misso, R. Riemsma, and J. Kleijnen, "Systematic Review of Adverse Events of Buprenorphine Patch Versus Fentanyl Patch in Patients with Chronic Moderate-to-Severe Pain," *Pain Management*, Vol. 2, No. 4, July 2012, pp. 351–362. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Wolstein, J., M. Gastpar, T. Finkbeiner, C. Heinrich, R. Heitkamp, T. Poehlke, and N. Scherbaum, "A Randomized, Open-Label Trial Comparing Methadone and Levo-Alpha-Acetylmethadol (LAAM) in Maintenance Treatment of Opioid Addiction," *Pharmacopsychiatry*, Vol. 42, No. 1, January 2009, pp. 1–8. Reason for exclusion: No functional outcomes.
- Woody, G., D. Bruce, P. T. Korthuis, S. Chhatre, M. Hillhouse, P. Jacobs, J. Sorensen, A. J. Saxon, S. Poole, D. Metzger, and W. Ling, "Authors' Reply: "Risk Reduction with Buprenorphine-Naloxone and Methadone: Patient's Choice," *Journal of Acquired Immune Deficiency Syndromes*, Vol. 67, No. 5, December 15, 2014, pp. e142–143. Reason for exclusion: Wrong study design.
- Woody, G. E., D. Otiashvili, G. Piralishvili, Z. Sikharulidze, G. Kamkamidze, and S. Poole, "Methadone and Suboxone for Subutex Injectors: Primary Outcomes of Pilot RCT," *Neuropsychopharmacology*, Vol. 38, 2013. Reason for exclusion: Conference abstract.
- Woody, G. E., D. Bruce, P. T. Korthuis, S. Chhatre, S. Poole, M. Hillhouse, P. Jacobs, J. Sorensen, A. J. Saxon, D. Metzger, and W. Ling, "HIV Risk Reduction with Buprenorphine-Naloxone or Methadone: Findings from a Randomized Trial," *Journal of Acquired Immune Deficiency Syndromes*, Vol. 66, No. 3, July 1, 2014, pp. 288–293. Reason for exclusion: No functional outcomes.

- Woody, G. E., A. T. McLellan, L. Luborsky, and C. P. O'Brien, "Psychotherapy in Community Methadone Programs: A Validation Study," *American Journal of Psychiatry*, Vol. 152, No. 9, September 1995, pp. 1302–1308. Reason for exclusion: Only measures effects of a co-intervention.
- Woody, G. E., S. A. Poole, G. Subramaniam, K. Dugosh, M. Bogenschutz, P. Abbott, A. Patkar, M. Publicker, K. McCain, J. S. Potter, R. Forman, V. Vetter, L. McNicholas, J. Blaine, K. G. Lynch, and P. Fudala, "Extended vs. Short-Term Buprenorphine-Naloxone for Treatment of Opioid-Addicted Youth: A Randomized Trial," *JAMA*, Vol. 300, No. 17, November 5, 2008, pp. 2003–2011. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Wouldes, T. A., V. E. Pritchard, and L. J. Woodward, "Neurobehaviour of Neonates Exposed Prenatally to Methadone Maintenance Treatment," *Journal of Paediatrics and Child Health*, Vol. 48, Supplement 1, March, 2012, p. 72. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Wu, Yafei, Shiyan Yan, Yanping Bao, Zhi Lian, Zhi Qu, and Zhimin Liu, "Cross-Sectional Study of the Severity of Self-Reported Depressive Symptoms in Heroin Users Who Participate in a Methadone Maintenance Treatment Program," *Shanghai Archives of Psychiatry*, Vol. 28, No. 1, February 2016, pp. 35–41. Reason for exclusion: Wrong study design.
- Wu, Z., and R. Detels, "Methadone Treatment in Primary Care," *Lancet HIV*, Vol. 2, No. 10, October 2015, pp. e408–409. Reason for exclusion: Wrong study design.
- Xiao, G. C., L. L. Xiao, C. D. Lai, and L. X. Li, "Clinical Study of Methadone Hydrochloride Combined with Buprenorphine Hydrochloride in the Treatment with Gradually Reducing Dosage of Heroin Addicts, Compared with Ordinary Dosage of Buprenorphine Hydrochloride," *Hebei Medical Journal*, Vol. 23, No. 7, 2001. Reason for exclusion: Not English.
- Xiao, L., Y. L. Tang, A. K. Smith, Y. T. Xiang, L. X. Sheng, Y. Chi, W. J. Du, S. Guo, Z. N. Jiang, G. F. Zhang, and X. N. Luo, "Nocturnal Sleep Architecture Disturbances in Early Methadone Treatment Patients," *Psychiatry Research*, Vol. 179, No. 1, August 30, 2010, pp. 91–95. Reason for exclusion: Wrong study design.
- Xiao, L., Z. Wu, W. Luo, and X. Wei, "Quality of Life of Outpatients in Methadone Maintenance Treatment Clinics," *Journal of Acquired Immune Deficiency Syndromes*, Vol. 53, Supplement 1, February 2010, pp. S116–120. Reason for exclusion: Wrong study design.
- Yan, L., E. Liu, J. M. McGoogan, S. Duan, L. T. Wu, S. Comulada, and Z. Wu, "Referring Heroin Users from Compulsory Detoxification Centers to Community Methadone Maintenance Treatment: A Comparison of Three Models," *BMC Public Health*, Vol. 13, August 13, 2013, p. 747. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Yancovitz, S. R., D. C. Des Jarlais, N. P. Peyser, E. Drew, P. Friedmann, H. L. Trigg, and J. W. Robinson, "A Randomized Trial of an Interim Methadone Maintenance Clinic," *American Journal of Public Health*, Vol. 81, No. 9, September 1991, pp. 1185–1191. Reason for exclusion: No functional outcomes.
- Yang, J., J. Li, G. Xu, J. Zhang, Z. Chen, Z. Lu, and H. Deng, "Elevated Hair Cortisol Levels among Heroin Addicts on Current Methadone Maintenance Compared to Controls," *PLoS One*, Vol. 11, No. 3, 2016, p. e0150729. Reason for exclusion: No functional outcomes.
- Yarlas, A., K. Miller, W. Wen, M. Kowalski, S. Lynch, B. Dain, C. Munera, and S. Ripa, "Long-Term Maintenance of Improvements in Patient-Reported Functioning, Quality-of-Life, and Pain Outcomes for Moderate-to-Severe Chronic Pain Patients Receiving Continuous Treatment of Butrans® (Buprenorphine) Transdermal System (BTDS)," *Journal of Pain*, Vol. 14, No. 4, Supplement 1, April 2013, p. S80. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Yaroslavtseva, T., E. Krupitsky, E. Zvartau, E. Blokhina, and G. Woody, "The Impact of Oral and Implantable Formulations of Naltrexone on Overall Functioning and Social Adjustment in Opioid Dependent Patients," *European Neuropsychopharmacology*, Vol. 24, 2014. Reason for exclusion: Conference abstract.
- Yee, A., M. Danaee, H. S. Loh, A. H. Sulaiman, and C. G. Ng, "Sexual Dysfunction in Heroin Dependents: A Comparison Between Methadone and Buprenorphine Maintenance Treatment," *PLoS One*, Vol. 11, No. 1, 2016, p. e0147852. Reason for exclusion: No functional outcomes.
- Yeh, T. L., K. C. Chen, S. H. Lin, I. H. Lee, P. S. Chen, W. J. Yao, S. Y. Lee, Y. K. Yang, R. B. Lu, M. H. Liao, and N. T. Chiu, "Availability of Dopamine and Serotonin Transporters in Opioid-Dependent Users—A Two-Isotope SPECT Study," *Psychopharmacology (Berl)*, Vol. 220, No. 1, March 2012, pp. 55–64. Reason for exclusion: No functional outcomes.
- Yen, C. N., C. S. Wang, T. Y. Wang, H. F. Chen, and H. C. Chang, "Quality of Life and Its Correlates Among Heroin Users in Taiwan," *Kaohsiung Journal of Medical Sciences*, Vol. 27, No. 5, May 2011, pp. 177–183. Reason for exclusion: No functional outcomes.
- Yucel, M., D. I. Lubman, B. J. Harrison, A. Fornito, N. B. Allen, R. M. Wellard, K. Roffel, K. Clarke, S. J. Wood, S. D. Forman, and C. Pantelis, "A Combined Spectroscopic and Functional MRI Investigation of the Dorsal Anterior Cingulate Region in Opiate Addiction," *Molecular Psychiatry*, Vol. 12, No. 7, July 2007, pp. 611, 691–702. Reason for exclusion: Wrong study design.
- Zaaijer, E. R., J. Bruijtel, P. Blanken, V. Hendriks, M. W. Koeter, M. J. Kreek, J. Booij, A. E. Goudriaan, J. M. van Ree, and W. van den Brink, "Personality as a Risk Factor for Illicit Opioid Use and a Protective Factor for Illicit Opioid Dependence," *Drug and Alcohol Dependence*, Vol. 145, December 1, 2014, pp. 101–105. Reason for exclusion: No functional outcomes.

- Zaaijer, E. R., L. van Dijk, K. de Bruin, A. E. Goudriaan, L. A. Lammers, M. W. Koeter, W. van den Brink, and J. Booij, "Effect of Extended-Release Naltrexone on Striatal Dopamine Transporter Availability, Depression and Anhedonia in Heroin-Dependent Patients," *Psychopharmacology (Berl)*, Vol. 232, No. 14, July 2015, pp. 2597–2607. Reason for exclusion: No functional outcomes.
- Zacny, J. P., K. M. Conley, J. Galinkin, J. M. Klasta, P. A. Klock, C. J. Young, and D. W. Coalson, "Behavioral and Physiological Effects of Buprenorphine and Morphine in Non-Drug Abusing Volunteers," *NIDA Research Monograph*, Vol. 178, 1998. Reason for exclusion: Wrong population (i.e., pregnant women, adolescents).
- Zacny, James P., Kristen Wroblewski, and Dennis W. Coalson, "Methylnaltrexone: Its Pharmacological Effects Alone and Effects on Morphine in Healthy Volunteers," *Psychopharmacology*, Vol. 232, No. 1, January 2015, pp. 63–73. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.
- Zajicova, A., H. Wilczek, and V. Holan, "The Alterations of Immunological Reactivity in Heroin Addicts and Their Normalization in Patients Maintained on Methadone," *Folia Biol (Praha)*, Vol. 50, No. 1, 2004, pp. 24–28. Reason for exclusion: No functional outcomes.
- Zaller, N. D., J. J. Fu, A. R. Bazazi, and J. D. Rich, "The Impact of Financial Discharge from Methadone Maintenance Therapy on Incarceration," *Journal of Opioid Management*, Vol. 6, No. 5, September–October 2010, pp. 365–370. Reason for exclusion: Only measures effects of a co-intervention.
- Zarghami, M., F. Sheikhmoonesi, S. Ala, J. Yazdani, and S. Farnia, "A Comparative Study of Beneficial Effects of Olanzapine and Sodium Valproate on Aggressive Behavior of Patients Who Are on Methadone Maintenance Therapy: A Randomized Triple Blind Clinical Trial," *European Review for Medical Pharmacological Sciences*, Vol. 17, No. 8, April 2013, pp. 1073–1081. Reason for exclusion: Only measures effects of a co-intervention.
- Zarghami, M., F. Sheikhmoonesi, S. Farnia, J. Yazdani, S. Ala, and N. Balilashak, "Comparison of the Effects of Oral Olanzapine and Sodium Valproate on Aggressiveness in Methadone Maintenance Patients: A Clinical Trial," *Asian Journal of Psychiatry*, Vol. 4, Supplement 1, July 2011, p. S41. Reason for exclusion: Only measures effects of a co-intervention.
- Zeller, Andru, "Program Blends Suboxone and Alternative Addiction Treatments," *Alcoholism and Drug Abuse Weekly*, Vol. 24, No. 26, 2012, pp. 1–7. Reason for exclusion: Wrong study design.
- Zeng, X., L. Lei, Y. Lu, and Z. Wang, "Treatment of Heroinism with Acupuncture at Points of the Du Channel," *Journal of Traditional Chinese Medicine*, Vol. 25, No. 3, September 2005, pp. 166–170. Reason for exclusion: Not MAT using methadone, buprenorphine, Suboxone, and naltrexone.

- Zhang, G. F., Y. P. Ren, L. X. Sheng, Y. Chi, W. J. Du, S. Guo, Z. N. Jiang, L. Xiao, X. N. Luo, Y. L. Tang, A. K. Smith, Z. Q. Liu, and H. X. Zhang, "Dysfunction of the Hypothalamic-Pituitary-Adrenal Axis in Opioid Dependent Subjects: Effects of Acute and Protracted Abstinence," *American Journal of Drug and Alcohol Abuse*, Vol. 34, No. 6, 2008, pp. 760–768. Reason for exclusion: Wrong setting (i.e., inpatient).
- Zhang, L., E. P. Chow, X. Zhuang, Y. Liang, Y. Wang, C. Tang, L. Ling, J. D. Tucker, and D. P. Wilson, "Methadone Maintenance Treatment Participant Retention and Behavioural Effectiveness in China: A Systematic Review and Meta-Analysis (Provisional Abstract)," *PLoS One*, Vol. 8, No. 7, 2013. Reason for exclusion: No functional outcomes.
- Zhang, Z., P. D. Friedmann, and D. R. Gerstein, "Does Retention Matter? Treatment Duration and Improvement in Drug Use," *Addiction*, Vol. 98, No. 5, May 2003, pp. 673–684. Reason for exclusion: Wrong study design.
- Zubieta, J., M. K. Greenwald, U. Lombardi, J. H. Woods, M. R. Kilbourn, D. M. Jewett, R. A. Koeppe, C. R. Schuster, and C. E. Johanson, "Buprenorphine-Induced Changes in Mu-Opioid Receptor Availability in Male Heroin-Dependent Volunteers: A Preliminary Study," *Neuropsychopharmacology*, Vol. 23, No. 3, September 2000, pp. 326–334. Reason for exclusion: No functional outcomes.

References

- Aalto, Mauri, Jukka-Pekka Visapää, Jukka T. Halme, Carola Fabritius, and Mikko Salaspuro, "Effectiveness of Buprenorphine Maintenance Treatment as Compared to a Syringe Exchange Program Among Buprenorphine Misusing Opioid-Dependent Patients," *Nordic Journal of Psychiatry*, Vol. 65, No. 4, September 2011, pp. 238–243.
- American Society of Addiction Medicine, *National Practice Guideline for the Use of Medications in the Treatment of Addiction Involving Opioid Use*, Chevy Chase, Md., 2015. As of October 25, 2017:
<http://www.asam.org/docs/default-source/practice-support/guidelines-and-consensus-docs/asam-national-practice-guideline-supplement.pdf>
- Atkins, David, Martin Eccles, Signe Flottorp, Gordon H. Guyatt, David Henry, Suzanne Hill, Alessandro Liberati, Dianne O'Connell, Andrew D. Oxman, Bob Phillips, Holger Schünemann, Tessa Tan-Torres Edejer, Gunn E. Vist, John W. Williams, Jr., and the GRADE Working Group, "Systems for Grading the Quality of Evidence and the Strength of Recommendations I: Critical Appraisal of Existing Approaches: The GRADE Working Group," *BMC Health Services Research*, Vol. 4, No. 38, December 2004.
- Baldacchino, A., M. Armanyous, D. J. K. Balfour, G. Humphris, and K. Matthews, "Neuropsychological Functioning and Chronic Methadone Use: A Systematic Review and Meta-Analysis," *Neuroscience and Biobehavioral Reviews*, Vol. 73, February 2017, pp. 23–38.
- Bale, Richard N., William W. Van Stone, John M. Kuldau, Thomas M. J. Engelsing, Robert M. Elashoff, and Vincent P. Zarcone, "Therapeutic Communities vs. Methadone Maintenance: A Prospective Controlled Study of Narcotic Addiction Treatment—Design and One-Year Follow-Up," *Archives of General Psychiatry*, Vol. 37, No. 2, February 1980, pp. 179–193.
- Ball, John C., and Alan Ross, *The Effectiveness of Methadone Maintenance Treatment: Patients, Programs, Services, and Outcome*, New York: Springer-Verlag, 1991.
- Balshem, Howard, Mark Helfand, Holger J. Schünemann, Andrew D. Oxman, Regina Kunz, Jan Brozek, Gunn E. Vist, Yngve Falck-Ytter, Joerg Meerpohl, Susan Norris, and Gordon H. Guyatt, "GRADE Guidelines: 3. Rating the Quality Of Evidence," *Journal of Clinical Epidemiology*, Vol. 64, No. 4, April 2011, pp. 401–406.
- Biernacki, Kathryn, Skye N. McLennan, Gill Terrett, Izelle Labuschagne, and Peter G. Rendell, "Decision-Making Ability in Current and Past Users of Opiates: A Meta-Analysis," *Neuroscience and Biobehavioral Reviews*, Vol. 71, December 2016, pp. 342–351.

- Byford, Sarah, Barbara Barrett, Nicola Metrebian, Teodora Groshkova, Maria Cary, Vikki Charles, Nicholas Lintzeris, and John Strang, "Cost-Effectiveness of Injectable Opioid Treatment V. Oral Methadone for Chronic Heroin Addiction," *The British Journal of Psychiatry*, Vol. 203, No. 5, November 2013, pp. 341–349.
- Compton, Peggy, Catherine P. Canamar, Maureen Hillhouse, and Walter Ling, "Hyperalgesia in Heroin Dependent Patients and the Effects of Opioid Substitution Therapy," *The Journal of Pain*, Vol. 13, No. 4, April 2012, pp. 401–409.
- Cornish, James W., David Metzger, Gero E. Woody, David Wilson, A. Thomas McLellan, Barry Vandergrift, and Charles P. O'Brien, "Naltrexone Pharmacotherapy for Opioid Dependent Federal Probationers," *Journal of Substance Abuse Treatment*, Vol. 14, No. 6, November–December 1997, pp. 529–534.
- Corsenac, Philippe, Emmanuel Lagarde, Blandine Gadegbeku, Bernard Delorme, Aurore Tricotel, Anne Castot, Nicholas Moore, Pierre Philip, Bernard Laumon, and Ludivine Orriols, "Road Traffic Crashes and Prescribed Methadone and Buprenorphine: A French Registry-Based Case-Control Study," *Drug and Alcohol Dependence*, Vol. 123, No. 1–3, June 1, 2012, pp. 91–97.
- Coviello, Donna M., James W. Cornish, Kevin G. Lynch, Arthur I. Alterman, and Charles P. O'Brien, "A Randomized Trial of Oral Naltrexone for Treating Opioid-Dependent Offenders," *The American Journal on Addictions*, Vol. 19, No. 5, September–October 2010, pp. 422–432.
- Coviello, Donna M., Dave A. Zanis, Susan A. Wesnoski, Kevin G. Lynch, and Michelle Drapkin, "Characteristics and 9-Month Outcomes of Discharged Methadone Maintenance Clients," *Journal of Substance Abuse Treatment*, Vol. 40, No. 2, March 2011, pp. 165–174.
- Crits-Christoph, Paul, Christie Lundy, Mark Stringer, Robert Gallop, and David R. Gastfriend, "Extended-Release Naltrexone for Alcohol and Opioid Problems in Missouri Parolees and Probationers," *Journal of Substance Abuse Treatment*, Vol. 56, September 2015, pp. 54–60.
- Dole, Vincent P., J. Waymond Robinson, John Orraca, Edward Towns, Paul Searcy, and Eric Caine, "Methadone Treatment of Randomly Selected Criminal Addicts," *The New England Journal of Medicine*, Vol. 280, No. 25, June 19, 1969, pp. 1372–1375.
- Egger, Matthias, George Davey Smith, Martin Schneider, and Christoph Minder, "Bias in Meta-Analysis Detected by a Simple, Graphical Test," *BMJ*, Vol. 315, No. 7109, September 13, 1997, pp. 629–634.
- Farrell-MacDonald, S. M. A. MacSwain, M. Cheverie, M. Tiesmaki, and B. Fischer, "Impact of Methadone Maintenance Treatment on Women Offenders' Post-Release Recidivism," *European Addiction Research*, Vol. 20, No. 4, 2014, pp. 192–199.

- Fudala, Paul J., T. Peter Bridge, Susan Herbert, William O. Williford, C. Nora Chiang, Karen Jones, Joseph Collins, Dennis Raisch, Paul Casadonte, R. Jeffrey Goldsmith, Walter Ling, Usha Malkerneker, Laura McNicholas, John Renner, Susan Stine, and Donald Tusel, "Office-Based Treatment of Opiate Addiction with a Sublingual-Tablet Formulation of Buprenorphine and Naloxone," *New England Journal of Medicine*, Vol. 349, No. 10, September 4, 2003, pp. 949–958.
- Gerra, Gilberto, Amir Zaimovic, Maria Augusta Raggi, Gabriele Moi, Barbara Branchi, Mirko Moroni, and Francesca Brambilla, "Experimentally Induced Aggressiveness in Heroin-Dependent Patients Treated with Buprenorphine: Comparison of Patients Receiving Methadone and Healthy Subjects," *Psychiatry Research*, Vol. 149, No. 1–3, January 15, 2007, pp. 201–213.
- Giacomuzzi, Salvatore, Georg Kemmler, Markus Ertl, and Yvonne Riemer, "Opioid Addicts at Admission Versus Slow-Release Oral Morphine, Methadone, and Sublingual Buprenorphine Maintenance Treatment Participants," *Substance Use and Misuse*, Vol. 41, No. 2, 2006, pp. 223–244.
- Giacomuzzi, Salvatore M., Markus Ertl, Georg Kemmler, Yvonne Riemer, and Alexander Vigl, "Sublingual Buprenorphine and Methadone Maintenance Treatment: A Three-Year Follow-Up of Quality of Life Assessment," *The Scientific World Journal*, Vol. 5, No. 24, 2005, pp. 452–468.
- Giacomuzzi, Salvatore M., Yvonne Riemer, Markus Ertl, Georg Kemmler, H Rössler, Hans H. Hinterhuber, and M. Kurz, "Buprenorphine Versus Methadone Maintenance Treatment in an Ambulant Setting: A Health-Related Quality of Life Assessment," *Addiction*, Vol. 98, No. 5, May 2003, pp. 693–702.
- Gordon, Michael S., Timothy W. Kinlock, Robert P. Schwartz, and Kevin E. O'Grady, "A Randomized Clinical Trial of Methadone Maintenance for Prisoners: Findings at 6 Months Post-Release," *Addiction*, Vol. 103, No. 8, August 2008, pp. 1333–1342.
- Gossop, Michael, John Marsden, Duncan Stewart, Petra Lehmann, and John Strang, "Methadone Treatment Practices and Outcome for Opiate Addicts Treated in Drug Clinics and in General Practice: Results from the National Treatment Outcome Research Study," *British Journal of General Practice*, Vol. 49, No. 438, January 1999, pp. 31–34.
- Hartung, Joachim, "An Alternative Method for Meta-Analysis," *Biometrical Journal*, Vol. 41, No. 8, December 1999, pp. 901–916.
- Hartung, Joachim, and Guido Knapp, "A Refined Method for the Meta-Analysis of Controlled Clinical Trials with Binary Outcome," *Statistics in Medicine*, Vol. 20, No. 24, December 30, 2001, pp. 3875–3889.

- Headquarters, Department of the Army, “Army 2020: Generating Health and Discipline in the Force Ahead of the Strategic Reset,” January 2012. As of October 26, 2017:
<http://www.army.mil/e2/c/downloads/232541.pdf>
- Higgins, Julian, and Sally Green, eds., *Cochrane Handbook for Systematic Reviews of Interventions*, Version 5.1.0, March 2011. As of December 13, 2017:
<http://handbook-5-1.cochrane.org/>
- IntHout, Joanna, John P. Ioannidis, and George F. Borm, “The Hartung-Knapp-Sidik-Jonkman Method for Random Effects Meta-Analysis Is Straightforward and Considerably Outperforms the Standard Dersimonian-Laird Method,” *BMC Medical Research Methodology*, Vol. 14, No. 25, 2014, pp. 1–12.
- Jones, Christopher M., “Heroin Use and Heroin Use Risk Behaviors Among Nonmedical Users of Prescription Opioid Pain Relievers—United States, 2002–2004 and 2008–2010,” *Drug and Alcohol Dependence*, Vol. 132, No. 1–2, September 1, 2013, pp. 95–100.
- Kinlock, Timothy W., Michael S. Gordon, Robert P. Schwartz, Terrence T. Fitzgerald, and Kevin E. O’Grady, “A Randomized Clinical Trial of Methadone Maintenance for Prisoners: Results at 12 Months Postrelease,” *Journal of Substance Abuse Treatment*, Vol. 37, No. 3, October 2009, pp. 277–285.
- Lee, Joshua D., Peter D. Friedmann, Timothy W. Kinlock, Edward V. Nunes, Tamara Y. Boney, Randall A. Hoskinson, Jr., Donna Wilson, Ryan McDonald, John Rotrosen, Marc N. Gourevitch, Michael Gordon, Marc Fishman, Donna T. Chen, Richard J. Bonnie, James W. Cornish, Sean M. Murphy, and Charles P. O’Brien, “Extended-Release Naltrexone to Prevent Opioid Relapse in Criminal Justice Offenders,” *New England Journal of Medicine*, Vol. 374, No. 13, March 31, 2016, pp. 1232–1242.
- Ling, Walter, Paul Casadonte, George Bigelow, Kyle M. Kampman, Ashwin Patkar, Genie L. Bailey, Richard N. Rosenthal, and Katheirne L. Beebe, “Buprenorphine Implants for Treatment of Opioid Dependence: A Randomized Controlled Trial,” *JAMA*, Vol. 304, No. 14, October 13, 2010, pp. 1576–1583. As of October 26, 2017:
http://jama.jamanetwork.com/data/Journals/JAMA/4532/joc05127_1576_1583.pdf
- Lobmaier, Philipp P., Nikolaj Kunøe, Michael Gossop, Tormod Katevoll, and Helge Waal, “Naltrexone Implants Compared to Methadone: Outcomes Six Months After Prison Release,” *European Addiction Research*, Vol. 16, No. 3, 2010, pp. 139–145.
- Magura, Stephen, Joshua D. Lee, Jason Hersherberger, Herman Joseph, Lisa Marsch, Carol Shropshire, and Andrew Rosenblum, “Buprenorphine and Methadone Maintenance in Jail and Post-Release: A Randomized Clinical Trial,” *Drug Alcohol Dependence*, Vol. 99, No. 1–3, January 1, 2009, pp. 222–230.

- Mattick, Richard P., Robert Ali, Jason M. White, Susannah O'Brien, Seija Wolk, and Cath Danz, "Buprenorphine Versus Methadone Maintenance Therapy: A Randomized Double-Blind Trial with 405 Opioid-Dependent Patients," *Addiction*, Vol. 98, No. 4, April 2003, pp. 441–452.
- Mattick, Richard P., Courtney Breen, Jo Kimber, and Marina Davoli, "Methadone Maintenance Therapy Versus No Opioid Replacement Therapy for Opioid Dependence," *Cochrane Database of Systematic Reviews*, No. 3, 2009, p. Cd002209.
- McNicholas, Laura, *TIP 40: Clinical Guidelines for the Use of Buprenorphine in the Treatment of Opioid Addiction: Treatment Improvement Protocol (TIP) Series 40*, Rockville, Md., U.S. Department of Health and Human Services; Substance Abuse and Mental Health Services Administration; Center for Substance Abuse Treatment, DHHS Publication No. (SMA) 04-3939, 2004. As of October 26, 2017:
<http://lib.adaa.washington.edu/clearinghouse/downloads/TIP-40-Clinical-Guidelines-for-the-Use-of-Buprenorphine-in-the-Treatment-of-Opioid-Addiction-54.pdf>
- Metrebian, Nicola, Teodora Groshkova, Jennifer Hellier, Vikki Charles, Anthea Martin, Luciana Forzisi, Nicholas Lintzeris, Deborah Zador, Hugh Williams, Tom Carnwath, Soraya Mayet, and John Strang, "Drug Use, Health and Social Outcomes of Hard-to-Treat Heroin Addicts Receiving Supervised Injectable Opiate Treatment: Secondary Outcomes from the Randomized Injectable Opioid Treatment Trial (RIOTT)," *Addiction*, Vol. 110, No. 3, March 2015, pp. 479–490.
- National Institute on Drug Abuse, "Drugs of Abuse: Opioids," undated. As of October 26, 2017:
<http://www.drugabuse.gov/drugs-abuse/opioids>
- Neri, S., C. M. Bruno, D. Pulvirenti, M. Malaguarnera, C. Italiano, B. Mauceri, G. Abate, D. Cilio, S. Calvagno, A. Tsami, L. Ignaccolo, D. Interlandi, L. Prestianni, M. Ricchena, and R. Noto, "Randomized Clinical Trial to Compare the Effects of Methadone and Buprenorphine on the Immune System in Drug Abusers," *Psychopharmacology*, Vol. 179, No. 3, May 2005, pp. 700–704.
- Neumann, Anne M., Richard D. Blondell, Urmo Jaanimägi, Amanda K. Giambrone, Gregory G. Homish, Jacqueline R. Lozano, Urszula Kowalik, and Mohammadreza Azadfard, "A Preliminary Study Comparing Methadone and Buprenorphine in Patients with Chronic Pain and Coexistent Opioid Addiction," *Journal of Addictive Diseases*, Vol. 32, No. 1, January 2013, pp. 68–78.
- Newman, Robert G., and Walden B. Whitehill, "Double-Blind Comparison of Methadone and Placebo Maintenance Treatments of Narcotic Addicts in Hong Kong," *Lancet*, Vol. 2, No. 8141, September 8, 1979, pp. 485–488.

- “Patient Information Leaflet: Patient Information: Suboxone, Subutex,” undated. As of November 1, 2017:
<https://www.fda.gov/downloads/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/UCM191529.pdf>
- Rapeli, Pekka, Carola Fabritius, Hely Kalska, and Hannu Alho, “Memory Function in Opioid-Dependent Patients Treated with Methadone or Buprenorphine Along with Benzodiazepine: Longitudinal Change in Comparison to Healthy Individuals,” *Substance Abuse Treatment, Prevention, and Policy*, Vol. 4, No. 6, 2009.
- Rapeli, Pekka, Carola Fabritius, Hely Kalska, and Hannu Alho, “Cognitive Functioning in Opioid-Dependent Patients Treated with Buprenorphine, Methadone, and Other Psychoactive Medications: Stability and Correlates,” *BMC Clinical Pharmacology*, Vol. 11, 2011, p. 13.
- Reijneveld, S. A., and H. N. Plomp, “Methadone Maintenance Clients in Amsterdam After Five Years,” *The International Journal of the Addictions*, Vol. 28, No. 1, January 1993, pp. 63–72.
- Rosenthal, Richard N., Walter Ling, Paul Casadonte, Frank Vocci, Genie L. Bailey, Kyle Kampman, Ashwin Patkar, Steven Chavoustie, Christine Blasey, Stacey Sigmon, and Katherine L. Beebe, “Buprenorphine Implants for Treatment of Opioid Dependence: Randomized Comparison to Placebo and Sublingual Buprenorphine/Naloxone,” *Addiction*, Vol. 108, No. 12, December 2013, pp. 2141–2149.
- Roxane Laboratories, Inc., “Dolophine® Hydrochloride CII,” Columbus, Ohio, undated. As of November 1, 2017:
https://www.accessdata.fda.gov/drugsatfda_docs/label/2006/006134s028lbl.pdf
- Rudd, Rose A., Noah Aleshire, Jon E. Zibbell, and R. Matthew Gladden, “Increases in Drug and Opioid Overdose Deaths—United States, 2000–2014,” *Morbidity and Mortality Weekly Report (Centers for Disease Control and Prevention)*, Vol. 64, No. 50, January 1, 2016, pp. 1–5.
- Sánchez-Meca, Julio, and Fulgencio Marín-Martínez, “Confidence Intervals for the Overall Effect Size in Random-Effects Meta-Analysis,” *Psychological Methods*, Vol. 13, No. 1, March 2008, pp. 31–48.
- Schwartz, Robert P., Jerome H. Jaffe, David A. Highfield, Jason M. Callaman, and Kevin E. O’Grady, “A Randomized Controlled Trial of Interim Methadone Maintenance: 10-Month Follow-Up,” *Drug and Alcohol Dependence*, Vol. 86, No. 1, January 5, 2007, pp. 30–36.
- Schwartz, Robert P., Jerome H. Jaffe, Kevin E. O’Grady, Timothy W. Kinlock, Michael S. Gordon, Sharon M. Kelly, Monique E. Wilson, and Ashraf Ahmed, “Interim Methadone Treatment: Impact on Arrests,” *Drug Alcohol Dependence*, Vol. 103, No. 3, August 1, 2009, pp. 148–154.

- Sees, Karen L., Kevin L. Delucchi, Carmen Masson, Amy Rosen, H. Wesley Clark, Helen Robillard, Peter Banys, and Sharon M. Hall, "Methadone Maintenance Versus 180-Day Psychosocially Enriched Detoxification for Treatment of Opioid Dependence: A Randomized Controlled Trial," *JAMA*, Vol. 283, No. 10, March 8, 2000, pp. 1303–1310.
- Senay, E. C., A. G. Barthwell, R. Marks, P. Bokos, D. Gillman, and R. White, "Medical Maintenance: A Pilot Study," *Journal of Addictive Diseases*, Vol. 12, No. 4, 1993, pp. 59–76.
- Sidik, Kurex, and Jeffrey N. Jonkman, "Robust Variance Estimation for Random Effects Meta-Analysis," *Journal of Computational Statistics and Data Analysis*, Vol. 50, No. 12, August 2006, pp. 3681–3701.
- Soyka, Michael, Bettina Hock, Sabine Kagerer, Robert Lehnert, Claudia Limmer, and Heinrich Kuefner, "Less Impairment on One Portion of a Driving-Relevant Psychomotor Battery in Buprenorphine-Maintained than in Methadone-Maintained Patients: Results of a Randomized Clinical Trial," *Journal of Clinical Psychopharmacology*, Vol. 25, No. 5, October 2005, pp. 490–493.
- Soyka, Michael, Martin Lieb, Sabine Kagerer, Christina Zingg, Gabriele Koller, Peter Lehnert, Claudia Limmer, Heinrich Kuefner, and Kristina Hennig-Fast, "Cognitive Functioning During Methadone and Buprenorphine Treatment: Results of a Randomized Clinical Trial," *Journal of Clinical Psychopharmacology*, Vol. 28, No. 6, December 2008, pp. 699–703.
- Strain, Eric C., Maxine L. Stitzer, Ira A. Liebson, and George E. Bigelow, "Methadone Dose and Treatment Outcome," *Drug and Alcohol Dependence*, Vol. 33, No. 2, September 1993, pp. 105–117.
- Strand, Maren Cecilie, Bente Fjeld, Marianne Arnestad, and Jorg Morland, "Can Patients Receiving Opioid Maintenance Therapy Safely Drive? A Systematic Review of Epidemiological and Experimental Studies on Driving Ability with a Focus on Concomitant Methadone or Buprenorphine Administration," *Traffic Injury Prevention*, Vol. 14, No. 1, 2013, pp. 26–38.
- Strang, John, John Marsden, Michael Cummins, Michael Farrell, Emily Finch, Michael Gossop, Duncan Stewart, and Sarah Welch, "Randomized Trial of Supervised Injectable Versus Oral Methadone Maintenance: Report of Feasibility and 6-Month Outcome," *Addiction*, Vol. 95, No. 11, November 2000, pp. 1631–1645.
- Substance Abuse and Mental Health Services Administration, and Center for Behavioral Health Statistics and Quality, *Behavioral Health Trends in the United States: Results from the 2014 National Survey on Drug Use and Health*, Rockville, Md., 2015.
- Tiihonen, Jari, Evgeny Krupitsky, Elena Verbitskaya, Elena Blokhina, Olga Mamontova, Jaana Föhr, Pekka Tuomola, Kimmo Kuoppasalmi, Vesa Kiviniemi, and Edwin Zwartau,

- “Naltrexone Implant for the Treatment of Polydrug Dependence: A Randomized Controlled Trial,” *The American Journal of Psychiatry*, Vol. 169, No. 5, May 2012, pp. 531–536.
- Toblin, Robin L., Phillip J. Quartana, Lyndon A. Riviere, Kristina Clarke Walper, and Charles W. Hoge, “Chronic Pain and Opioid Use in U.S. Soldiers After Combat Deployment,” *JAMA Intern Medicine*, Vol. 174, No. 8, August 2014, pp. 1400–1401.
- U.S. Department of Health and Human Services, *Quick Guide for Physicians Based on TIP 40: Clinical Guidelines for the Use of Buprenorphine in the Treatment of Opioid Addiction*, Washington, D.C.: DHHS Publication No. (SMA) 05-4003, 2005. As of October 26, 2017: <http://store.samhsa.gov/shin/content//SMA05-4003/SMA05-4003.pdf>
- U.S. Department of Health and Human Services, *Medication-Assisted Treatment for Opioid Addiction in Opioid Treatment Programs. A Treatment Improvement Protocol TIP 43*, Rockville, Md.: Substance Abuse and Mental Health Services Administration, HHS Publication No. (SMA) 12-4214, revised 2012. As of October 26, 2017: https://www.ncbi.nlm.nih.gov/books/NBK64164/pdf/Bookshelf_NBK64164.pdf
- U.S. Department of Health and Human Services, *Clinical Use of Extended-Release Injectable Naltrexone in the Treatment of Opioid Use Disorder: A Brief Guide*, Rockville, Md.: Substance Abuse and Mental Health Services Administration, revised February 2, 2015. As of October 26, 2017: <http://store.samhsa.gov/shin/content/SMA14-4892/SMA14-4892.pdf>
- U.S. Department of Health and Human Services, *Incorporating Alcohol Pharmacotherapies into Medical Practice: A Treatment Improvement Protocol*, TIP 49, Rockville, Md.: Substance Abuse and Mental Health Services Administration, May 2013. As of October 26, 2017: <https://store.samhsa.gov/product/TIP-49-Incorporating-Alcohol-Pharmacotherapies-Into-Medical-Practice/SMA13-4380>
- U.S. Department of Veterans Affairs, *VA/DoD Clinical Practice Guideline for the Management of Substance Use Disorders*, version 3.0, Washington, D.C., 2015.
- VA—See U.S. Department of Veterans Affairs.
- “Vivitrol,” undated. As of November 1, 2017: https://www.accessdata.fda.gov/drugsatfda_docs/label/2006/006134s028lbl.pdf
- Wang, Grace Y., Trecia A. Wouldes, and Bruce R. Russell, “Methadone Maintenance Treatment and Cognitive Function: A Systematic Review,” *Current Drug Abuse Reviews*, Vol. 6, No. 3, September 2013, pp. 220–230.
- The White House, “Presidential Memorandum—Addressing Prescription Drug Abuse and Heroin Use,” Washington, D.C., Office of the Press Secretary, October 21, 2015. As of October 26, 2017:

<https://obamawhitehouse.archives.gov/the-press-office/2015/10/21/presidential-memorandum-addressing-prescription-drug-abuse-and-heroin>

World Health Organization, *Clinical Guidelines for Withdrawal Management and Treatment of Drug Dependence in Closed Settings*, Geneva, 2009. As of October 26, 2017:

<https://www.ncbi.nlm.nih.gov/books/NBK310658/>