

AWARD NUMBER: W81XWH-15-1-0345

TITLE: A Novel Animal Model for Investigating the Neural Basis of Focal Dystonia

PRINCIPAL INVESTIGATOR: Leslie Evinger

CONTRACTING ORGANIZATION: State University of New York
Stony Brook, NY 11794-336

REPORT DATE: September 2018

TYPE OF REPORT: Annual Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE September 2018			2. REPORT TYPE Annual Report		3. DATES COVERED 31 Aug 2017 – 30 Aug 2018	
4. TITLE AND SUBTITLE A Novel Animal Model for Investigating the Neural Basis of Focal Dystonia					5a. CONTRACT NUMBER W81XWH-15-1-0345	
					5b. GRANT NUMBER PR140382	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Leslie Evinger E-Mail: leslie.erving@stonybrook.edu					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) State University of New York Office of Sponsored Programs Stony Brook University Stony Brook, NY 11794-3362					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The overall goal of the project was to develop an animal model of the focal dystonia benign essential blepharospasm. Consistent with the widely held view that dystonia results from an interaction between a predisposing condition and an environmental trigger, we proposed to use 7 Hz deep brain stimulation of the basal ganglia as the predisposing condition and dry eye as the environmental trigger. We hypothesized that the 7 Hz deep brain stimulation would exaggerate the blink adaptations to dry eye into spasms of lid closure characteristic of benign essential blepharospasm. This reporting year we demonstrated that sex differences exist in the blink adaptations to dry eye. We are determining if the sex of the rat is important in developing our animal model of blepharospasm.						
Dystonia, benign essential blepharospasm, dry eye, motor plasticity, basal ganglia, deep brain stimulation, eyelids, blinking						
16. SECURITY CLASSIFICATION OF:				17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	19b. TELEPHONE NUMBER (include area code)			

Table of Contents

	<u>Page</u>
1. Introduction.....	4
2. Keywords.....	4
3. Accomplishments.....	4
4. Impact.....	7
5. Changes/Problems.....	7
6. Products.....	7
7. Participants & Other Collaborating Organizations.....	9
8. Special Reporting Requirements.....	9
9. Appendices.....	11

1. Introduction

The focal dystonia benign essential blepharospasm (**BEB**), arises from the convergence of a predisposing condition and an environmental trigger (Hallett et al., 2008). The overall goal of our project is to develop an animal model of BEB based on the hypothesis that hypersynchronized, 7 Hz neuronal oscillations of the basal ganglia create the predisposing condition and that eye irritation from dry eye is the environmental trigger. Our demonstration that hypersynchronized oscillations in the basal ganglia produced by 7 Hz deep brain stimulation of the subthalamic nucleus (**STN DBS**) exaggerate neural plasticity in normal rats (Kaminer et al., 2014) provides a neural mechanism by which hypersynchronized basal ganglia activity creates a predisposing condition. From our studies showing that dry eye initiates neural plasticity in blink circuits to produce compensatory adaptations in blinking (Evinger et al., 2002; Schicatano et al., 2002), we predict that combining 7 Hz STN DBS and dry eye will exaggerate neural plasticity and force the normally compensatory adaptive processes in response to dry eye to transform into the characteristics of BEB, *e.g.*, spasms of lid closure, excessive blinking, and trigeminal hyperexcitability. We propose two Specific Aims to test this hypothesis. The goal of the first Specific Aim is to show that synchronized theta oscillations in the basal ganglia exaggerate plasticity in the cerebellum and the excitability of trigeminal blink circuits as occurs in BEB patients. The Major Tasks to accomplish Specific Aim 1 are: 1) to investigate effects of synchronized basal ganglia oscillations on activity of the deep cerebellar nucleus neurons; and 2) to investigate the effects of synchronized basal ganglia oscillations on the activity of superior colliculus neurons. The purpose of the second Specific Aim is to demonstrate that synchronized 7 Hz oscillations established in the basal ganglia are sufficient to predispose mammals to develop BEB. The Major Tasks to accomplish Specific Aim 2 are: 1) to determine whether combining synchronized basal ganglia 7 Hz oscillations with corneal irritation is sufficient to develop spasms of lid closure and other characteristics of the focal dystonia BEB; and 2) to perform control experiments to determine that theta frequency is critical in enabling the development of spasms of lid closure.

2. Keywords

Dystonia, benign essential blepharospasm, dry eye, motor plasticity, basal ganglia, deep brain stimulation, eyelids, blinking

3. Accomplishments

Major goals of the project

The overarching goal of the planned investigations is to test our hypothesis that hypersynchronized 7 Hz oscillations in the basal ganglia create a predisposing condition that transforms the normally adaptive modifications initiated by dry eye into the spasms of lid closure, excessive blinking, and trigeminal hyperexcitability characteristic of individuals with the focal dystonia benign essential blepharospasm (**BEB**). To test this hypothesis, the 1st Specific Aim of the project is to demonstrate that synchronized 7 Hz (**theta**) oscillations in the basal ganglia exaggerate plasticity in the cerebellum and excitability of trigeminal blink circuits. Major Task 1 of Specific Aim 1 is to investigate effects of synchronized basal ganglia oscillations on the activity of the deep cerebellar nucleus neurons. Major Task 2 of Specific Aim 1 is to investigate the effects of synchronized basal ganglia oscillations on the activity of superior colliculus neurons. The 2nd Specific Aim of the project is to demonstrate that synchronized theta oscillations established in the basal ganglia are sufficient to predispose mammals to develop blepharospasm. Major Task 1 of Specific Aim 2 is to determine whether combining synchronized basal ganglia theta oscillations with corneal irritation is sufficient to develop spasms of lid closure and other characteristics of the focal dystonia BEB. Major Task 2 of Specific Aim 2 is to perform control experiments to determine that theta frequency is critical for the development of spasms of lid closure.

What was accomplished under these goals?

In this year, we began analyzing the data collected from Major Task 2 of Specific Aim 2 studies from the last reporting period and continued data collection and analysis from Major Task 2 of Specific Aim 1.

Specific Aim 2, Major Task 2: Our previous work on our blepharospasm model used exclusively male rats (Evinger, 2015). The current experiments, however, utilized both male and female rats. We anticipated that there would be no difference between the sexes on the effect of creating a dry eye condition by removal of the exorbital gland (Project Narrative, page 7). This assumption, however, was incorrect. The blink adaptations initiated by dry eye were significantly stronger in males than in females. Blink adapta-




Figure 1. Normalized change in number of extra blinks with each trigeminal stimulus for all rats and separated by sex. ** < .01, *** < .001




Figure 2. Normalized changes in trigeminal reflex blink excitability following dry eye. * < 0.05, *** < 0.001, n.s = not significant




Figure 3. Normalized change in spontaneous blink rate (blinks/min) for all rats and separated by sex. * < 0.05, *** < 0.005, n.s. = non-significant

tions initiated by dry eye act to minimize the breaks in the corneal tear film that create the discomfort of dry eye. Because each blink reforms the corneal tear film by eliminating breaks in the tear film (Himebaugh et al., 2009), the trigeminal system adaptively increases the number of blinks evoked by a trigeminal stimulus and elevates the excitability of trigeminal reflex blinks so that smaller breaks in the corneal tear film evoke a reflex blink in response to dry eye. As we previously showed in humans (Evinger et al., 2002; Schicatano et al., 2002), dry eye increases the number of blinks evoked by a single trigeminal blink evoking stimulus, termed blink oscillations, and elevates trigeminal reflex blink excitability. In our dry eye model in rodents, blink oscillations increased for all rats, but the increase was significantly larger for males than for female rats (Fig. 1). Following dry eye, male rats exhibited an increase in blink excitability, but the change for females was insignificant (Fig. 2). Studies of humans with dry eye report that both males and females exhibit an increase in the spontaneous blink rate in subjects with dry eye relative to subjects without dry eye (Nakamori et al., 1997; Himebaugh et al., 2009). In contrast, following exorbital gland removal to create dry eye, male rats exhibited the expected increase in spontaneous blink rate, whereas females showed a significant decrease in spontaneous blink rate (Fig. 3). Currently, we are reanalyzing all our data from Specific Aim 2, Major Task 2 separating animals by sex.

Specific Aim 1, Major Task 2: As described in the Project Narrative of the grant proposal (page 9), the goal of these experiments is to determine how different frequencies of STN DBS affect the activity of superior colliculus neurons and how changes in collicular activity modify trigeminal reflex blink excitability

measured with the paired stimulus paradigm. Predictions about superior colliculus neuronal activity alterations with STN DBS come from the basal ganglia circuit that regulates reflex blink excitability through the superior colliculus (Basso and Evinger, 1996; Basso et al., 1996; Gnadt et al., 1997). The substantia nigra pars reticulata inhibits superior colliculus neurons that excite nucleus raphe neurons. These raphe neurons inhibit spinal trigeminal blink circuits. In the paired stimulus paradigm, the first reflex blink stimulus (Condition) activates trigeminal blink circuits to evoke the Condition blink and trigeminal complex neurons also send an excitatory drive to intermediate/deep layer collicular neurons. This colliculus neuron activation provides a transient excitatory drive onto the nucleus raphe magnus neurons that inhibit spinal trigeminal blink circuits (Basso and Evinger, 1996). Thus, the second reflex blink stimulus (Test) in the paired stimulus paradigm occurs during trigeminal inhibition created by nucleus raphe magnus activation. Because of this inhibition, the Test blink is smaller than the Condition blink even though both are evoked by the same stimulus. We are analyzing the data collected in these experiments with emphasis on sex differences in neuronal responses.

Changes to SOW Order

The SOW order was changed with the approved request for a no cost extension. Data analysis for Specific Aim 1, Major Task 2 continues into months 30 – 36 and Specific Aim 1, Major Task 1 data collection and analysis are projected for months 33 – 48. The new approved SOW is an appendix.

What opportunities for training and professional development has the project provided?

Nothing to Report.

How were the results disseminated to communities of interest?

We are preparing a manuscript for publication on the data from Specific Aim 2 Major Task 2 and will complete a manuscript describing data from Specific Aim 1 Major Task 2 in the coming reporting period.

What are the plans for the next reporting period to accomplish the goals?

In addition to completing and submitting manuscripts from collected data, the work during the next reporting period will focus on finishing analysis of collected data on Specific Aim 2 Major Task 1 and Specific Aim 1 Major Task 2, beginning data collection on Specific Aim 1, Major Task 1 and analysis of those data.

Major Task 2 of Specific Aim 1 (Project Narrative page 9): These experiments are nearing completion and data analysis is ongoing. With completion of data analysis, we will begin writing a manuscript.

Major Task 2 of Specific Aim 2 (Project Narrative pages 11-12): These experiments are complete and reanalysis of the data based on sex of the animal is ongoing. Progress is being made on a manuscript describing these studies.

Major Task 1 of Specific Aim 2 (Project Narrative pages 8-9):

The goal of these experiments is to determine how different frequencies of basal ganglia oscillations modify cerebellar interpositus (IP) activity and to correlate these changes with shifts in blink plasticity. As our data demonstrate that 7 Hz STN DBS exaggerates blink plasticity, 16 Hz STN DBS impairs blink plasticity, and 130 Hz STN DBS has no effect on the blink plasticity of normal rats (Kaminer et al., 2014), we will compare IP neural activity across all three frequencies. We will simultaneously record unitary activity and local field potentials (LFP) from blink related IP regions during SO stimulation before, during, and after our trigeminal reflex blink gain paradigm (page 8, project narrative). We will compare the activity of individual IP neurons and LFP before the trigeminal reflex blink gain paradigm to their activity after the trigeminal reflex blink gain paradigm.

4. Impact

What was the impact on the development of the principal discipline(s) of the project?

Nothing to Report

What was the impact on other disciplines?

Nothing to Report

What was the impact on technology transfer?

Nothing to Report

What was the impact on society beyond science and technology?

Nothing to Report

5. Changes/Problems

Changes in approach and reasons for change

There were no changes in approach.

Actual or anticipated problems or delays and actions or plans to resolve them

I have been unable to attract a postdoctoral fellow or a new graduate student to replace the one who graduated. My technician, Ms. Cynthia Lowe, left to go to graduate school and my other technician, Ms. Donna Schmidt, left for another lab paying a higher salary. The data collection and analysis of the recording studies of Specific Aim 1 required more time than originally allocated. In addition, my extensive teaching load in Spring 2018 significantly reduced my research productivity, which prevented me from accomplishing the necessary experiments. I applied for and received a no cost extension because my teaching and administrative assignments should decrease in the 2018/2019 academic year so that I can focus on completing the Major Tasks of Specific Aim 1. To increase the number of neurons recorded in each session and thereby speed up data, I am now recording with tetrodes instead of multiwire electrodes in Specific Aim 1.

Changes that had a significant impact on expenditures

Nothing to Report

Significant changes in use or care of vertebrate animals

Nothing to Report

6. Products

Journal publications

Lee AS, Blankson BK, Manza P, O'Rawe JF, Evinger C, Leung H-C. "Spontaneous but not voluntary eye blink during spatial working memory are associated with successful performance" submitted to *Brain and Cognition*

Matheis T, Evinger C, Schubert R, Mazzola S, Fels M, Kemper N, Reilmann R, Muratori L. "Biological motion perception in Huntington's disease" in preparation

Books or other non-periodical, one time publications

None to Report

Other publications, conference papers, and presentations

None to Report

Website(s) or other internet site(s)

None to Report

Technologies or techniques

None to Report

Inventions, patent applications, and/or licenses

None to Report

Other Products

None to Report

7. Participants & other collaborating organizations

What individuals have worked on the project?

Name:	Leslie Craig Evinger	Cynthia Lowe	Donna Schmidt
Project Role:	PI	Technician	Technician
Research Identifier:	0000-0002-0039-3348		
Nearest Person Month Worked:	12	9	6
Contribution:	Experimental design, manuscript preparation, performing experiments	Performing experiments	Lab manager, histology
Funding Support:			

Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?

None to Report

What other organizations were involved as partners?

None to Report

8. Special reporting requirements

Not Applicable

9. Appendices

SOW

References

- Basso MA, Evinger C (1996) An explanation for reflex blink hyperexcitability in Parkinson's disease. II. Nucleus raphe magnus. *J Neurosci* 16:7318-7330.
- Basso MA, Powers AS, Evinger C (1996) An explanation for reflex blink hyperexcitability in Parkinson's disease. I. Superior colliculus. *J Neurosci* 16:7308-7317.
- Evinger C (2015) Benign Essential Blepharospasm is a Disorder of Neuroplasticity: Lessons From Animal Models. *J Neuroophthalmol* 35:374-379.
- Evinger C, Mao JB, Powers AS, Kassem IS, Schicatano EJ, Henriquez VM, Peshori KR (2002) Dry eye, blinking, and blepharospasm. *Mov Disord* 17 Suppl 2:S75-78.
- Gnadt JW, Lu SM, Breznen B, Basso MA, Henriquez VM, Evinger C (1997) Influence of the superior colliculus on the primate blink reflex. *Exp Brain Res* 116:389-398.
- Hallett M, Evinger C, Jankovic J, Stacy M (2008) Update on blepharospasm: Report from the BEBRF International Workshop. *Neurology* 71:1275-1282.
- Himebaugh NL, Begley CG, Bradley A, Wilkinson JA (2009) Blinking and tear break-up during four visual tasks. *Optom Vis Sci* 86:E106-114.

- Kaminer J, Thakur P, Evinger C (2014) Frequency matters: beta-band subthalamic nucleus deep-brain stimulation induces Parkinsonian-like blink abnormalities in normal rats. *Eur J Neurosci* 40:3237-3242.
- Nakamori K, Odawara M, Nakajima T, Mizutani T, Tsubota K (1997) Blinking is controlled primarily by ocular surface conditions. *Am J Ophthalmol* 124:24-30.
- Schicatano EJ, Mantzouranis J, Peshori KR, Partin J, Evinger C (2002) Lid restraint evokes two types of motor adaptation. *J Neurosci* 22:569-576.

**STATEMENT OF WORK – Month/Day/Year
PROPOSED START DATE September 1, 2015**

Site: Stony Brook University
Life Sciences Building
Department of Neurobiology &
Behavior
Stony Brook, NY 11794-5230
PI: Leslie Craig Evinger, PhD

Specific Aim 2 (specified in proposal)	Timeline	Site 1
Demonstrate that synchronized theta oscillations established in the basal ganglia are sufficient to predispose mammals to develop blepharospasm		
Major Task 2: Determine whether combining synchronized basal ganglia theta oscillations with corneal irritation is sufficient to develop spasms of lid closure and other characteristics of the focal dystonia BEB	Months	
Subtask 1: Monitor blink frequency, blink reflex hyperexcitability, and spasms of lid closure in rats during 4 different conditions: 1) No DBS; 2) 7 Hz STN DBS alone; 2) 7 Hz STN DBS and dry eye; and 4) dry eye alone.	1-8	PI
Subtask 2: Data analysis of blink frequency, hyperexcitability, and spasms of lid closure and statistical comparison of data obtained in the 4 conditions.	6-9	PI
Milestone(s) Achieved: Published paper Evinger <i>J. Neuro-Ophthalmology</i> , 2015;35:374–379 describing animal model of blepharospasm.	11	PI
Major Task 2: Perform control experiments to determine that theta frequency is critical in enabling the development of spasms of lid closure		
Subtask 1: Repeat Subtask 1 of Major Task 1 using 16 Hz STN DBS or 130 Hz STN DBS.	9-18	PI
Subtask 2: Data analysis of blink frequency, hyperexcitability, and spasms of lid closure and statistical comparison with data obtained in Subtask 1 of Major Task 1.	15-19	PI
Subtask 3: Monitor blink frequency, blink reflex hyperexcitability, and spasms of lid closure in rats during 3 different conditions: 1) No DBS; 2) dry eye alone; and 3) 7 Hz STN DBS and dry eye.	18-24	PI
Subtask 4: Data analysis of blink frequency, hyperexcitability, and spasms of lid closure and statistical comparison with data obtained Subtask 1 of Major Task 1.	22-24	PI
Milestone(s) Achieved: Began work on manuscript describing results of Specific Aim 2 Major Task 2	24	PI
Specific Aim 1		
Demonstrate that synchronized theta oscillations in the basal ganglia exaggerate plasticity in the cerebellum and excitability of trigeminal blink circuits		

Major Task 2: Investigate the effects of synchronized basal ganglia oscillations on activity of the superior colliculus neurons.		
Subtask 1: Record from superior colliculus neurons while rats participate in paired stimulus paradigm and receive 7 Hz, 16 Hz, 130 Hz, or no subthalamic deep brain stimulation.	18-36	PI
Subtask 2: Data analysis of single unit and local field potential data collected in Subtask 1 and correlation of neural activity with blink reflex excitability measures.	30-36	PI
Major Task 1: Investigate effects of synchronized basal ganglia oscillations on activity of the deep cerebellar nucleus neurons.		
Subtask 1: Record from interpositus neurons while rats participate in a blink plasticity paradigm and receive 7 Hz, 16 Hz, 130 Hz, or no subthalamic deep brain stimulation.	33-46	PI
Subtask 2: Data analysis of single unit and local field potential data collected in Subtask 1 and correlation of neural activity with brainstem plasticity measures.	40-47	PI
Milestone(s) Achieved: Submit a manuscript describing data from Specific Aim 1 Major Task 2 and a manuscript reporting data from Specific Aim 1 Major Task 2.	48	PI

If human subjects are involved in the proposed study, please provide the projected quarterly enrollment in the following table.

	Year 1				Year 2				Year 3
Target Enrollment (per quarter)	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1
Site 1									
Site 2									
Site 3									
Target Enrollment (cumulative)									

Note: The Government reserves the right to request a revised SOW format and/or additional information.