



Laser Cooling of Polyatomic Molecules

John Doyle
HARVARD COLLEGE PRESIDENT & FELLOWS OF

06/28/2019
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/ RTB1
Arlington, Virginia 22203
Air Force Materiel Command

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.					
1. REPORT DATE (DD-MM-YYYY) 30-04-2019		2. REPORT TYPE Final		3. DATES COVERED (From - To) 09/30/2015-09/20/2018	
4. TITLE AND SUBTITLE Laser Cooling of Polyatomic Molecules				5a. CONTRACT NUMBER FA9550-15-1-0446	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) John Doyle				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Harvard University Cambridge MA 02138				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Public Distribution					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT During this grant we achieved and characterized efficient Sisyphus laser cooling of a polyatomic molecule from 50 mK to below 1 mK in 1 dimension. We demonstrated transverse cooling (and heating) of a SrOH beam using two different electronic transitions, studied loss channels to vibrational states (including the bending mode), and highlighted extensions to more complex metal monoalkoxides with six and more atoms. Our results were published in Physical Review Letters and selected as Editor's suggestion and featured in Physics Viewpoint. To explore laser cooling of more complex molecules, we performed detailed theoretical calculations of Franck-Condon factors and vibrational branching ratios for SrOCH3 and CaOCH3, which are isoelectronic to triatomic SrOH and CaOH. We also created a 1-D MOT of CaOH, a significant step to a MOT of CaOCH3. In addition, in collaboration with Tim Steimle of ASU, we performed dispersed fluorescence of CaOCH3, our target molecule for laser cooling in the future. We determined experimentally the Franck-Condon factors for the X-A and X-B transitions, finding both suitable for laser cooling (although the X-A transition is simpler).					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)

Reset

Laser Cooling of Polyatomic Molecules
John Doyle
FA9550-15-1-0446
Annual Report Period 9/30/2017-9/29/2018

Compared to atoms, the additional rotational and vibrational degrees of freedom in molecules give rise to a wide variety of potential and implemented scientific applications, including quantum computation, precision measurements, and quantum simulation. While ultracold diatomic molecules will be extremely valuable in opening novel research frontiers, molecules with three or more atoms have unique capabilities enabled by their significantly more complicated structure for advancing fundamental physics, chemistry, and quantum technologies. Cooling molecular degrees of freedom significantly aids in realizing such applications. Yet, the desired quantum complexity that molecules provide also leads to challenges for control, detection, and cooling.

Cooling of the external motion of neutral atoms from above room temperature into the sub-millikelvin range (leading to, e.g., Bose-Einstein condensation) commonly relies on the use of velocity-dependent optical forces. Laser cooling requires reasonably closed and strong optical electronic transitions, so its use for molecules has been severely limited. Recently, laser cooling and magneto-optical trapping has been achieved for diatomic molecules SrF, CaF and YO. However, since SrOH had 3 distinct vibrational modes, including a doubly-degenerate bending mode absent in diatomics, the question of direct laser cooling for polyatomic molecules remained widely open.

Leading up to the past year of activities, we achieved and characterized efficient Sisyphus laser cooling of a polyatomic molecule from 50 mK to below 1 mK in 1 dimension. We demonstrated transverse cooling (and heating) of a SrOH beam using two different electronic transitions, studied loss channels to vibrational states (including the bending mode), and highlighted extensions to more complex metal monoalkoxides with six and more atoms. Our results were published in Physical Review Letters and selected as Editor's suggestion and featured in *Physics Viewpoint*. To explore laser cooling of more complex molecules, we performed detailed theoretical calculations of Franck-Condon factors and vibrational branching ratios for SrOCH₃ and CaOCH₃, which are isoelectronic to triatomic SrOH and CaOH.

In the past year we have accomplished experimental and theoretical work geared toward laser cooling of symmetric top molecules, which is our proposed work under our AFOSR renewal. Symmetric top molecules (STMs) like MOCH₃ (M = Ca, Sr) have first-order linear Stark shifts in small fields. We have been working on detailed analysis of a quantum computing platform based on STMs, including detailed simulations. We will submit our thorough analysis this month (October). On the experimental side, we built over the past year the infrastructure for a MOT of

STM molecules. In addition, in collaboration with Tim Steimle of ASU, we performed dispersed fluorescence of CaOCH_3 , our target molecules for laser cooling. We determined experimentally the Franck-Condon factors for the X-A and X-B transitions, finding both suitable for laser cooling (although the X-A transition is simpler). This included, of course, production of the molecule. We are therefore in the position to laser cool this species and are currently continuing our experimental build-up to that end.