



Nanoelectropulse-induced changes in cell excitability - a molecular mechanism for membrane depolarization

Thomas Vernier
OLD DOMINION UNIVERSITY RESEARCH FOUNDATION

07/26/2019
Final Report

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory
AF Office Of Scientific Research (AFOSR)/ RTB2
Arlington, Virginia 22203
Air Force Materiel Command

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.						
1. REPORT DATE (DD-MM-YYYY) 07/21/2019		2. REPORT TYPE Final Report			3. DATES COVERED (From - To) 06/15/2014 - 06-14-2019	
4. TITLE AND SUBTITLE Nanoelectropulse-induced changes in cell excitability: A molecular mechanism for membrane depolarization				5a. CONTRACT NUMBER FA9550-14-1-0123		
				5b. GRANT NUMBER FA9550-14-1-0123		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Vernier, P. Thomas				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Frank Reidy Research Center for Bioelectronics Old Dominion University 4211 Monarch Way Ste 300 Norfolk, VA 23508				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office of Scientific Research 875 N. Randolph, Ste 325 Arlington, Virginia 22203				10. SPONSOR/MONITOR'S ACRONYM(S) AFOSR		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution A - Approved for public release						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT Molecular dynamics (MD) simulations of ion transport through lipid electropores support the hypothesis that a 5 ns electric pulse enables transport of sodium ions sufficient to depolarize an adrenal chromaffin cell. To carry out this work we converted our model systems from the old GROMOS-OPLS force fields to the newer CHARMM36 force fields and implemented changes in the ion models to improve interactions with water and with the phospholipid bilayer interface). We also initiated an investigation of the effect of transmembrane peptides on lipid bilayer systems in applied electric fields, to determine whether these peptides modify the probability of conductive defect formation.						
15. SUBJECT TERMS membrane stress, ion conductance, molecular models, adrenal chromaffin cells, pulsed electric fields, membrane depolarization, ion channels, membrane proteins, transmembrane peptides						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			P. Thomas Vernier	
U	U	U	UU	12	19b. TELEPHONE NUMBER (Include area code) 757 683 7046	

AFOSR Grant: FA9550-14-1-0123**Title:** Nanoelectropulse-induced changes in cell excitability: A molecular mechanism for membrane depolarization**Principal Investigator:** P. Thomas Vernier

Abstract Molecular dynamics (MD) simulations of ion transport through lipid electropores provide theoretical support for the hypothesis that delivery of a single, 5 ns electric pulse can enable pore-mediated transport of sodium ions sufficient to depolarize the plasma membrane of an adrenal chromaffin cell, helping to explain a series of robust experimental observations. To carry out this work we converted our model systems from the old standard GROMOS-OPLS force fields to the newer and more accurate CHARMM36 force fields. For our specific studies we implemented also changes in the native ion models to improve the behavior of the calcium ion in its interactions with water (and indirectly with the atoms of the phospholipid bilayer interface). In parallel with this effort we are working with colleagues in Buenos Aires to improve the CHARMM36 Na⁺ model that they are using in our joint study of ion transport through mechanically (rather than electrically) constrained lipid pores. In another parallel effort, sparked by the pursuit of the primary objectives, we have undertaken an exploratory investigation of the effect of transmembrane peptides on lipid bilayer systems in applied electric fields, to answer the questions whether these peptides increase or decrease the likelihood of conductive defect formation, and whether they increase or decrease ion conductance after permeabilizing defects are formed. This will help us understand better the role of ion channels and other membrane proteins in membrane water defect formation, and how electrical stress leads to depolarization of adrenal chromaffin cells (and other electrically active cells) and thus to activation of voltage-gated calcium channels.

Background

Nanosecond pulsed electric fields (nanosecond electric pulses – NEPs) show undimmed promise as a new modality for neuromodulation that provides short duration (low risk of damage) and precise targeting of specific cells and tissues. The goal of this project is to provide fundamental and practical information that will guide future in vitro and in vivo investigations aimed directly at using NEPs for neuromodulation. This project is associated with and carried out in collaboration with Gale Craviso, Normand Leblanc, and Indira Chatterjee at the University of Nevada, Reno under AFOSR grant FA9550-14-1-0018. The overall project employs both experimental (electrophysiology and fluorescence imaging) and computational (two-dimensional electrical modeling (numerical) of the cell and molecular dynamics simulations) approaches to achieve our goal.

My effort centered on molecular dynamics (MD) simulations of ion transport through liquid electropores with the aim of reconciling experimentally observed ion flux with rates extrapolated from the model characteristics of individual pores and pore areal density.

Objectives

1. To project from electropore kinetics and dynamics the physical (density and area) and temporal extent of membrane depolarization. We compared what is known experimentally regarding ion transport during normal depolarization and activation with the pore formation and ion transport rates that we observe in simulations to determine whether the molecular model of the lipid nanopore is consistent with the hypothesis that a single 5 ns pulse is sufficient to depolarize an adrenal chromaffin cell membrane as a result of ion permeabilization. Further, from pore formation and re-sealing times extracted from our simulations we proposed to evaluate support for our hypothesis that NEPs induce transient membrane depolarizations that can be amplified in amplitude and time by a train of NEPs.

2. To characterize lipid nanopore conductance for Na^+ , K^+ , Ca^{2+} , and Cl^- . Through a series of simulations, we investigated specific ion effects on bilayer and lipid nanopore physical properties, to evaluate the importance of the ion models in predicting accurately the different behaviors of Na^+ , K^+ , Ca^{2+} , and Cl^- in pore transport rates and interface binding.

3. To evaluate the effects of increasing the complexity of the homogeneous phospholipid bilayer systems used in previous studies. Looking beyond the general effects of including the three major univalent inorganic ions and the divalent cation calcium in lipid bilayer systems, we planned to assess the impact of various modulating factors — phospholipid composition, cholesterol, local ionic strength, temperature, intracellular and extracellular membrane physical constraints (cytoskeletal and glycocalyx-type attachments) — on specific ion conductances and bilayer physical properties. By moving our model toward a more realistic representation of the membrane, we increase its predictive value.

Accomplishments

1. Initial correlations of ion transport for individual lipid electropores with experimentally observed activation of adrenal chromaffin cells. We previously reported that a combination of our initial MD simulations of ion transport through lipid electropores with the numerical modeling results of the Reno group provides *theoretical support for the hypothesis that delivery of a single, 5 ns pulse can enable pore-mediated transport of Na^+ sufficient to depolarize the plasma membrane of an adrenal chromaffin cell*. A more systematic and extensive set of simulations of Na^+ , K^+ , Ca^{2+} , and Cl^- transport through lipid electropores, with improved ion models for Na^+ and Ca^{2+} , is consistent with that initial conclusion. We have, furthermore, accumulated results with systems containing the more challenging Na^+ and Ca^{2+} ions that underscore the *importance of the accuracy of ion models* in their interactions with water and with the phospholipid head group interface. This is ongoing.

2. Conversion from GROMOS-OPLS to CHARMM36 force field. At the outset we determined that this project necessitated a *conversion of the molecular models in our simulations from the widely used, but ageing, GROMOS-OPLS force fields to the newer, more accurate CHARMM36 force fields*. This is essential for maintaining the continuity of our observations into the future [1]. To confirm the validity of these changes, we compared systems constructed with the old and new force fields, using several key indices of the correspondence of the behavior of model lipid bilayer systems with experimental observations. These include area per lipid, deuterium order parameter (for atoms in the hydrocarbon tails of the phospholipids), ion binding isotherms, minimum porating electric field, and pore initiation time. For a sample of one of these comparisons, see Figure 1.




Figure 1. Pore initiation time as a function of applied electric field for GROMOS-OPLS and CHARMM36 systems containing potassium ions.

considerable computational resources (and clock hours) because of the long simulation times required for equilibration and the need for replicate simulations to generate adequate statistical confidence. This factor was not accurately estimated when project objectives were established.

This transition to the CHARMM36 force field is complete. It was delayed by unexpected problems with the native CHARMM36 calcium ion model, described below.




Figure 2. Calcium ion water coordination number for CHARMM36 systems containing either the ion model distributed with CHARMM36 (red line) or the modified ion model created in this work (blue line). Experimentally determined values are in the range 6 to 8.

has addressed this problem in a different way [2]. None of these approaches is *the* correct one; we select the one most appropriate for our immediate investigations.

We are only now proceeding to simulate the impact of various modulating factors — phospholipid composition, cholesterol, local ionic strength, temperature, intracellular and extracellular membrane physical constraints (cytoskeletal and glycocalyx-type attachments) — on ion conductances (beginning with K^+ , Ca^{2+} , and Cl^-) and bilayer physical properties.

Simulations for ion binding isotherms and lipid electropore formation require

3. Modification of CHARMM36 calcium ion model. Upon initial examination of CHARMM36 systems containing Ca^{2+} , we noticed unrealistic (and unacceptable) clustering of Ca^{2+} and Cl^- ions (i.e., undissociated $CaCl_2$ molecules) in the bulk water. We found that we were not the first to notice this deficiency in the CHARMM36 Ca^{2+} model, and we have followed an approach similar to [1] to *improve the behavior of the calcium ion in its interactions with water (and indirectly with the atoms of the phospholipid bilayer interface)* (Figure 2). This work is now complete, and a manuscript for publication is underway. Recently we learned that another group, leaders (like the authors of [1]) in molecular dynamics force field development,

4. Sodium ion model (CHARMM36) and constrained pores. The CHARMM36 *sodium* ion model has deficiencies similar to those discussed above for Ca^{2+} . Although this was not part of our original plan (these problems have been reported only recently, and many workers in the field are just now responding – or are not even aware of them yet!), improving the behavior of these ion models is essential for producing reliable simulations of lipid membranes in electric fields. Our collaborators in Buenos Aires, who have been working with us for several years on ion conductance through lipid pores, have agreed to include *improving the Na^+ model in their study of ion transport through mechanically constrained lipid pores* (Figure 3). We will incorporate the relevant data from the Buenos Aires work, which P.I. Vernier is leading and coordinating under separate funding, into the K^+ and Ca^{2+} results that we are now generating in the Norfolk effort.




Figure 3. Mechanically constrained pores for ion transport studies [2]. NC – no constraint; ACC – anode and cathode sides constrained; CC – cathode side constrained; AC – anode side constrained.

5. Transmembrane peptides and the nanoelectroporation-driven nanoscale depolarization hypothesis.

To expand our perspective of cellular responses to stress-induced membrane permeabilization, we recently initiated an *exploratory investigation of the effect of transmembrane peptides on lipid bilayer systems in applied electric fields*. We want to know whether these peptides increase or decrease the likelihood of conductive defect formation, and whether they increase or decrease ion conductance after permeabilizing defects are formed. Answers to these questions will help us understand how electrical stress leads to depolarization of adrenal chromaffin cells (and other electrically active cells) and thus to activation of voltage-gated calcium channels. We are specifically interested in the hypothesis that ion channel gating may be a nanoscale phenomenon, at least under some conditions, that requires transport of only a few ions in the immediate vicinity of the ion channel, perhaps through a process we call nanoelectroporation-driven nanoscale depolarization. Initial results from these new studies (Figure 4) indicate the (not surprising) importance of specific amino acid residues in the bilayer interior and the (perhaps surprising) tendency of some transmembrane peptides to inhibit water defect formation in the immediate vicinity and of others to facilitate nearby water bridge construction.




Figure 4. Lipid bilayer (POPC) with a transmembrane peptide, a lipid pore, and a water bridge.

Collaborations

As mentioned above, we actively collaborated with Gale Craviso, Normand Leblanc, and Indira Chatterjee at the University of Nevada, Reno on the research reported here. We maintained regular e-mail and telephone contact, and I traveled to Reno at least once per year to discuss various research issues, including pulse delivery and the interconnection of molecular modeling and experimental observations of ion transport through electropermeabilized membranes. In addition, I am working with M. Laura Fernández and Marcelo Risk at the University of Buenos Aires in Argentina specifically on mechanically constrained lipid pores and on optimization of sodium ion models on molecular simulations.

Personnel supported by grant funds

Federica Castellani (partial support)

Students supported by grant funds

Federica Castellani (partial support)

Student volunteers

None at this time.

Presentations at research meetings/conferences

Castellani, F., E. B. Sözer, and P. T. Vernier, Microsecond kinetics of ion transport and interface binding in electrically stressed phospholipid bilayers, BioEM 2019, Montpellier, 2019.

Castellani, F., E. B. Sözer, and P. T. Vernier, Microsecond kinetics of ion transport and membrane interface binding in electrically stressed lipid bilayers, Biophysical Society Annual Meeting, Baltimore, 2019.

Castellani, F., E. B. Sözer, and P. T. Vernier, Microsecond kinetics of ion transport and membrane interface binding in electrically stressed phospholipid bilayers, Photonics and Electromagnetics Research Symposium (PIERS), Rome, 2019.

Sözer, E. B., and P. T. Vernier, Modulation by field reversal — Cell volume, membrane potential, and 2 ns electrostimulation of excitable cells, BioEM 2019, Montpellier, 2019.

Vernier, P. T., Nanoscale and macroscale transport across electrically stressed cell membranes — Beyond multiscale to multidimensional and metadimensional models, Photonics and Electromagnetics Research Symposium (PIERS), Rome, 2019.

Vernier, P. T., Nanoscale electrical stress and the molecular landscape of the cell membrane, Photonics and Electromagnetics Research Symposium (PIERS), Rome, 2019.

Vernier, P. T., Transport through electrically stressed cell membranes — Introduction to the electropermeome, XXV International Symposium on Bioelectrochemistry and Bioenergetics, Limerick, Ireland, May 2019.

Zaklit, J., G. L. Craviso, N. Leblanc, P. T. Vernier, and E. B. Sözer, Electrostimulation of adrenal chromaffin cells by 2 ns pulses can be modulated by an electric field reversal, BioEM 2019, Montpellier, 2019.

Castellani, F., E. B. Sözer, and P. T. Vernier, Microsecond kinetics of ion transport and membrane interface binding before, during, and after lipid electropore formation, 15th International Bioelectrics Symposium, Prague, 2018.

Castellani, F., and P. T. Vernier, Ion models and ion transport in molecular simulations of electrically stressed biological membranes, BioEM 2018, Portorož, 2018.

Castellani, F. and P. T. Vernier, Molecular simulations of lipid electropore formation and pore-mediated calcium transport with an improved Ca^{2+} model, Biophysical Society Annual Meeting, San Francisco, 2018.

Kramar, P., D. Miklavčič, and P. T. Vernier, Detecting the structural changes of planar lipid bilayer exposing to nanosecond electric field, BioEM 2018, Portorož, 2018.

Kramar, P., M. Reberšek, D. Miklavčič, and P. T. Vernier, Planar lipid bilayers exposed to nanosecond pulsed electric field, 27th International Electrotechnical and Computer Science Conference, Portorož, 2018.

Merla, C., F. Apollonio, A. Paffi, P. T. Vernier, and M. Liberti, An in vivo exposure system for wide-band electric pulses, 12th European Conference on Antennas and Propagation (EuCAP 2018), London, 2018.

Romeo, S., P. T. Vernier, and O. Zeni, Electroporation detection with THz time-domain spectroscopy, BioEM 2018, Portorož, 2018.

Semenov, I., E. B. Sözer, and P. T. Vernier, Intracellular distribution of reactive oxygen species after nanosecond electrical stimulation, 15th International Bioelectrics Symposium, Prague, 2018.

Sözer, E. B. and P. T. Vernier, Nanosecond bipolar pulse cancellation and the electropermeome, Biophysical Society Annual Meeting, San Francisco, 2018.

Sözer, E. B., and P. T. Vernier, Nanosecond bipolar pulse cancellation below 5 ns, BioEM 2018, Portorož, 2018.

Sözer, E. B. and P. T. Vernier, Observations of transport and cancellation of transport into electrically stressed cells defy “the standard model”, 15th International Bioelectrics Symposium, Prague, 2018.

Yang, L., S. Pierce, P. T. Vernier, I. Chatterjee, G. L. Craviso, and N. Leblanc, Differential effects of single and twin high-intensity nanosecond electric pulses on membrane permeabilization and Na⁺ conductance in bovine chromaffin cells, Biophysical Society Annual Meeting, San Francisco, 2018.

Yang, L., S. Pierce, P. T. Vernier, I. Chatterjee, G. L. Craviso, and N. Leblanc, Remarkable differences in the response of voltage-gated Na⁺ current to a single or pair of high intensity nanosecond electric pulses in adrenal chromaffin cells, BioEM 2018, Portorož, 2018.

Vernier, P. T., Intense and intensified electric fields in the low-permittivity interior of the biological membrane — Nanoscale reactor for hydroxyl radical formation, Workshop on Microwaving Cells: From Biological Effects to Innovative Techniques for Cell Analysis, IEEE International Microwave Symposium (IMS 2018), Philadelphia, June 2018.

Vernier, P. T., Nanoscale electromanipulation of cell membranes — Perturbation, reorganization, restoration, 2nd Wrocław Scientific Meeting, Wrocław Medical University, Wrocław, March, 2018.

Casciola, M., Z. A. Levine, P. Campomanes, F. Castellani, E. B. Sözer, M. Tarek, and P. T. Vernier, Transport of electropore dye indicators through stable lipid defects, a molecular dynamics study, BioEM 2017, Hangzhou, 2017.

Castellani, F., P. Marracino, F. Apollonio, M. Liberti, and P. T. Vernier, Molecular simulations of lipid bilayers in electric fields — The model matters!, Biophysical Society Annual Meeting, New Orleans, 2017.

Castellani, F. and P. T. Vernier, Molecular simulations of calcium transport through lipid electropores — the model matters!, BioEM 2017, Hangzhou, 2017.

Sözer, E. B. and P. T. Vernier, Quantitative small molecule transport after nanosecond electric field exposures – Experiments and models, Biophysical Society Annual Meeting, New Orleans, 2017.

Vernier, P. T. and E. B. Sözer, Beyond the “standard model” (of electroporation) — Cells fight back!, 24th International Symposium on Bioelectrochemistry and Bioenergetics of the Bioelectrochemical Society, Lyon, 2017.

Vernier, P. T., Nanoscale and multiscale models of membranes in permeabilizing electric fields — Physics, chemistry, biology, Physical Chemistry Seminar, Oscar Rice Lecture Series, University of North Carolina, Chapel Hill, March, 2017.

Wriggers, W., F. Castellani, P. T. Vernier, and J. A. Kovacs, Spatial heat maps from fast information matching of fast and slow degrees of freedom in molecular dynamics simulations, Biophysical Society Annual Meeting, New Orleans, 2017.

Wriggers, W., J. Kovacs, F. Castellani, P. T. Vernier, and D. J. Krusienski, An information theoretic approach for creating 3D spatial images from 4D time series data, Microscopy and Microanalysis 2017, St. Louis, 2017.

Castellani, F. and P. T. Vernier, Validity of molecular simulations of lipid bilayers in electric fields — the model makes a difference!, BioEM 2016, Ghent, 2016.

Levine, Z. A., T. Gowrishankar, K. Smith, R. Son, P. T. Vernier, and J. C. Weaver, Improving predictions: Insertion of molecular scale interactions into cell electroporation models, BioEM 2016, Ghent, 2016.

Marracino, P., F. Castellani, P. T. Vernier, M. Liberti, and F. Apollonio, A new methodology for the geometric characterization of an electropore, BioEM 2016, Ghent, 2016.

Marracino, P., F. Castellani, P. T. Vernier, M. Liberti, and F. Apollonio, A new methodology for the geometrical characterization of an electropore, ICEMB, Milan, 2016.

Semenova, N., G. L. Craviso, and P. T. Vernier, Spatially localized intracellular calcium concentration increases stimulated by nanosecond electric pulses in adrenal chromaffin cells, BioEM 2016, Ghent, 2016.

Sözer, E. B., C. F. Pocetti, and P. T. Vernier, Molecular transport in the electropore, 13th International Bioelectrics Symposium, Rostock, 2016.

Sözer, E. B., C. F. Pocetti, and P. T. Vernier, Transmembrane voltage and small molecule transport after 6 ns electroporeabilization, BioEM 2016, Ghent, 2016.

Vernier, P. T., Beyond the “standard model” (of electroporation) — Cells fight back!, Institut de Pharmacologie et de Biologie Structurale, Centre National de la Recherche Scientifique, Toulouse, France, November, 2016.

Vernier, P. T., E. B. Sözer, and Z. A. Levine, Beyond the “standard model” of electroporation — Physical and physiological components of the electropore, 13th International Bioelectrics Symposium, Rostock, Germany, September, 2016.

Yang, L., G. L. Craviso, R. Terhune, P. T. Vernier, I. Chatterjee, and N. Leblanc, Potential differential effects of single high-intensity 6 ns electric pulses on macroscopic inward and outward ionic currents recorded in whole-cell patch-clamped bovine chromaffin cells, BioEM 2016, Ghent, 2016.

Castellani, F., E. B. Sözer, Z. A. Levine, P. Marracino, F. Apollonio, M. Liberti, and P. T. Vernier, Molecular simulations of ion transport in electroporated phospholipid bilayers, Biophysical Society Annual Meeting, Baltimore, 2015.

Levine, Z. A., N. Ivošević DeNardis, and P. T. Vernier, Phospholipid and hydrocarbon dynamics at charged interfaces, 1st World Congress on Electroporation and Pulsed Electric Fields in Biology, Medicine, and Food and Environmental Technologies, Portorož, September, 2015.

Merla, C., A. Paffi, P. Monaco, T. Calderaro, F. Apollonio, C. Marino, P. T. Vernier, and M. Liberti, Design of an applicator for nsPEF exposure of newborn mice, 1st World Congress on Electroporation and Pulsed Electric Fields in Biology, Medicine, and Food and Environmental Technologies, Portorož, September, 2015.

Risk, M., M. L. Fernández, and P. T. Vernier, Asymmetric electropores in mechanically constrained lipid bilayers, BioEM 2015, Asilomar, 2015.

Sözer, E. B. and P. T. Vernier, Experimental quantification of molecular transport through lipid electropores induced by nanosecond pulsed electric fields, Biophysical Society Annual Meeting, Baltimore, 2015.

Vernier, P. T., Nanosecond transport through electropermeabilized lipid bilayers and living cell membranes, Workshop on Nanosecond Pulsed Electric Fields (nsPEF) – From Modeling to Applications: Biology, Medicine, Plasma, and Apparatus, IEEE International Microwave Symposium (MTT-S), Phoenix, May, 2015.

Vernier, P. T., Phospholipid bilayers in permeabilizing electric fields — Biophysical electrostatics at the atomic scale, 31st International Review of Progress in Applied Computational Electromagnetics (ACES 2015), Williamsburg, Virginia, March, 2015.

Vernier, P. T., Quantitative boundaries on nanoscale transport through electropermeabilized membranes, 1st World Congress on Electroporation and Pulsed Electric Fields in Biology, Medicine, and Food and Environmental Technologies, Portorož, September, 2015.

Yoon, J., J. Zaklit, I. Chatterjee, P. T. Vernier, N. Semenova, N. Leblanc, and G. L. Craviso, Transient versus sustained Ca^{2+} responses evoked in adrenal chromaffin cells by 5 ns pulses: pulse delivery considerations, BioEM 2015, Asilomar, 2015.

Zaklit, J., J. Yoon, I. Chatterjee, E. Evans, P. T. Vernier, N. Leblanc, and G. L. Craviso, Ca^{2+} release from internal stores of adrenal chromaffin cells is not evoked by 5 ns electric pulses, BioEM 2015, Asilomar, 2015.

Papers Published

Sözer, E. B. and P. T. Vernier, Modulation of biological responses to 2 ns electrical stimuli by field reversal. *Biochim. Biophys. Acta* 1861:1228-1239, 2019.

- Castellani, F., J. Teissié, and P. T. Vernier, Permeabilizing phospholipid bilayers with non-normal electric fields, *J. Membrane Biol.* 251:229-236, 2018.
- Fernández, M. L., M. R. Risk, and P. T. Vernier, Electropore formation in mechanically constrained phospholipid bilayers, *J. Membrane Biol.* 251:237-245, 2018.
- Romeo, S., A. Sannino, M. R. Scarfi, P. T. Vernier, R. Cadossi, J. Gehl, O. Zeni, ESOPE-equivalent pulsing protocols for calcium electroporation: An *in vitro* optimization study on 2 cancer cell models, *Technol. Cancer Res. Treatment* 17:1-10, 2018.
- Romeo, S., P. T. Vernier, and O. Zeni, Electroporation-induced cell modifications detected with THz time-domain spectroscopy, *J. Infrared, Millimeter, Terahertz Waves* 39:854-862, 2018.
- Sabuncu, A. C, M. Stacey, G. L. Craviso, N. Semenova, P. T. Vernier, N. Leblanc, I. Chatterjee, and J. Zaklit, Dielectric properties of isolated adrenal chromaffin cells determined by microfluidic impedance spectroscopy, *Bioelectrochemistry* 119:84-91, 2018.
- Sözer, E. B., C. F. Pocetti, and P. T. Vernier, Transport of charged small molecules after electropore permeabilization — Drift and diffusion, *BMC Biophysics* 11:4, 2018.
- Castellani, F., J. Teissié, and P. T. Vernier, Permeabilizing phospholipid bilayers with non-normal electric fields, *J. Membrane Biol.* 251:229-236, 2018.
- Fernández, M. L., M. R. Risk, and P. T. Vernier, Electropore formation in mechanically constrained phospholipid bilayers, *J. Membrane Biol.* 251:237-245, 2018.
- Sözer, E. B., C. F. Pocetti, and P. T. Vernier, Asymmetric patterns of small molecule transport after nanosecond and microsecond electropore permeabilization, *J. Membrane Biol.* 251:197-210, 2018.
- Marracino, P., F. Castellani, P. T. Vernier, M. Liberti, and F. Apollonio, Geometrical characterization of an electropore from water positional fluctuations, *J. Membrane Biol.* 250:11-19, 2017.
- Marracino, P., M. Liberti, P. T. Vernier, and F. Apollonio, Statistical analytical model for hydrophilic electropore characterization: A comparison study, *RSC Advances* 7:31997, 2017.
- Merla, C., F. Apollonio, A. Paffi, C. Marino, P. T. Vernier, and M. Liberti, Monopole patch antenna for in vivo exposure to nanosecond pulsed electric fields, *Med. Biol. Eng. Comp.* 55:1073-1083, 2017.
- Merla, C., A. G. Pakhomov, I. Semenov, and P. T. Vernier, Frequency spectrum of induced transmembrane potential and permeabilization efficacy of bipolar electric pulses, *Biochim. Biophys. Acta* 1859:1282-1290, 2017.
- Sözer, E. B., Z. A. Levine, and P. T. Vernier, Quantitative limits on small molecule transport via the electroporeome — Measuring and modeling single nanosecond perturbations, *Sci. Rep.* 7:57, 2017.
- Sözer, E. B., Y.-H. Wu, S. Romeo, and P. T. Vernier, Nanometer-scale permeabilization and osmotic swelling induced by 5 ns pulsed electric fields, *J. Membrane Biol.* 250:21-30, 2017.

- Wriggers, W., F. Castellani, J. Kovacs, and P. T. Vernier, Computing spatiotemporal heat maps of lipid electropore formation: A statistical approach, *Front. Mol. Biosci.* 4:22, 2017.
- Yang, L, G. L. Craviso, P. T. Vernier, I. Chatterjee, and N. Leblanc, Nanosecond electric pulses differentially affect inward and outward currents in patch clamped adrenal chromaffin cells, *PLOS ONE* 12:e0181002, 2017.
- Zaklit, J., G. L. Craviso, N. Leblanc, L. Yang, P. T. Vernier, and I. Chatterjee, Adrenal chromaffin cells exposed to 5-ns pulses require higher electric fields to porate intracellular membranes than the plasma membrane: An experimental and modeling study, *J. Membrane Biol.* 250:535-552, 2017.
- Batista Napotnik, T., M. Reberšek, P. T. Vernier, B. Mali, D. Miklavčič, Effects of high-voltage nanosecond electric pulses on eukaryotic cells (*in vitro*): A systematic review, *Bioelectrochemistry* 110:1-12, 2016.
- Buimaga-Iarinca, L., N. Ivošević DeNardis, P. T. Vernier, A. Calborean, and C. Morari, The effect of the electric field on the α -GPC interaction with Au(111) surface: A first principles study, *J. Phys. Chem. C* 120:9740-9749, 2016.
- Levine, Z. A., N. Ivošević DeNardis, and P. T. Vernier, Phospholipid and hydrocarbon interactions with a charged electrode interface, *Langmuir* 32:2809-2819, 2016.
- Mercadal, B., P. T. Vernier, and A. Ivorra, Dependence of electroporation detection threshold on cell radius: An explanation to observations non compatible with Schwan's equation model, *J. Membrane Biol.* 249:663-676, 2016.
- Yoon, J., N. Leblanc, J. Zaklit, P. T. Vernier, I. Chatterjee, and G. L. Craviso, Enhanced monitoring of nanosecond electric pulse-evoked membrane conductance changes in whole-cell patch clamp experiments, *J. Membrane Biol.* 249:633-644, 2016.
- Craviso, G. L., C. Fisher, I. Chatterjee, and P. T. Vernier, Adrenal chromaffin cells do not swell when exposed to nanosecond electric pulses, *Bioelectrochemistry* 103:98-102, 2015.
- Hansen, E. L., E. B. Sözer, S. Romeo, S. K. Frandsen, P. T. Vernier, and J. Gehl, Dose-dependent ATP depletion and cancer cell death following calcium electroporation: Relative effect of calcium concentration and electric field strength, *PLOS ONE* 10:e0122973, 2015.
- Kohler, S., Z. A. Levine, M. Á. García-Fernández, M.-C. Ho, P. T. Vernier, P. Leveque, D. Arnaud-Cormos, Electrical analysis of cell membrane poration by an intense nanosecond pulsed electric field, using an atomistic-to-continuum method, *IEEE Trans. Microwave Theory and Techniques* 63:2032-2040, 2015.
- Pakhomov, A. G., E. Gianulis, P. T. Vernier, I. Semenov, S. Xiao, and O. N. Pakhomova, Multiple nanosecond electric pulses increase the number but not the size of long-lived nanopores in the cell membrane, *Biochim. Biophys Acta* 1848:958-966, 2015.
- Vernier, P. T., Z. A. Levine, M.-C. Ho, S. Xiao, I. Semenov, and A. G. Pakhomov, Picosecond and terahertz perturbation of interfacial water and electroporabilization of biological membranes, *J. Membrane Biol.* 248:837-47, 2015.

References

1. Enkavi, G., M. Javanainen, W. Kulig, T. Rog, and I. Vattulainen. 2019. Multiscale simulations of biological membranes: The challenge to understand biological phenomena in a living substance. *Chem. Rev.* 119:5607-5774.
2. Han, K., R. M. Venable, A. M. Bryant, C. J. Legacy, R. Shen, H. Li, B. Roux, A. Gericke, and R. W. Pastor. 2018. Graph-theoretic analysis of monomethyl phosphate clustering in ionic solutions. *J Phys Chem B* 122:1484-1494.
3. Kohagen, M., P. E. Mason, and P. Jungwirth. 2014. Accurate description of calcium solvation in concentrated aqueous solutions. *J Phys Chem B* 118:7902-7909.
4. Fernández, M. L., M. R. Risk, and P. T. Vernier. 2018. Electropore formation in mechanically constrained phospholipid bilayers. *J. Membr. Biol.* 251:237-245.