

2020 TOA

Steroid Regimens in the Prevention of Irvine-Gass Syndrome

Samuel D. Hobbs, MD, CAPT, MC, USAF
Ophthalmology Resident, PGY-3
Wilford Hall Ambulatory Surgical Center
San Antonio Military Medical Center



Disclosures

- No financial disclosures.
- The view(s) expressed herein are those of the author(s) and do not reflect the official policy or position of Brooke Army Medical Center, Wilford Hall Ambulatory Surgical Center, the U.S. Army Medical Department, the U.S. Army Office of the Surgeon General, the Department of the Army, the Department of the Air Force and Department of Defense, or the U.S. Government.


Purpose

To describe the incidence of Irvine-Gass syndrome (pseudophakic cystoid macular edema, PCME) following cataract surgery with phacoemulsification in patients treated with either difluprednate, brand-name prednisolone acetate (PA), or generic PA, with or without the use of topical NSAIDs.




Methods

- Patients who underwent cataract extraction with phacoemulsification between January 1, 2018 and December 31, 2018 were selected from billing codes.
- Exam and medical history were confirmed via chart review.
- PCME defined as $<20/25$ BCVA and new cystic IRF on OCT.




Selection Criteria




Results

Incidence of PCME by Steroids



Total: 56/899



Difluprednate: 16/303



Brand Name PA: 16/357




Generic PA: 24/239

p = 0.02




Results

Pairwise Analysis: Rates of PCME by Brand Name PA



$p = 0.63$




$p = 0.009$





What about NSAIDs?



Incidence of Irvine-Gass by NSAID



$p = 0.19$



Incidence of Irvine-Gass by NSAID



Subgroup Analysis

	No PCME (N = 844)	PCME (N = 56)	p-value
Age	69.5±9.5	68.4±10.2	p = 0.78
Gender (Male)	409 (48.5%)	31 (55.4%)	p = 0.32
Resident vs. Attending	599 (93.3%) vs. 245 (95.0%)	43 (6.7%) vs. 13 (5.0%)	P = 0.34
PC Tear	13 (92.9%)	1 (7.1%)	p = 0.89
Hx of DM	298 (92.0%)	26 (8.0%)	p = 0.08
Mild NPDR	10 (76.9%)	3 (23.1%)	p = 0.04
Mod NPDR	6 (85.7%)	1 (14.3%)	p = 0.44
PDR	6 (75.0%)	2 (25.0%)	p = 0.09
ERM	38 (73.1%)	14 (26.9%)	p = <0.0001
Hx of RVO	2 (50.0%)	2 (50.0%)	p = 0.02
Hx of RD	11 (100.0%)	0 (0.0%)	p = 0.23
Hx of Uveitis	12 (100.0%)	0 (0.0%)	p = 0.21
Hx of AMD	99 (93.4%)	7 (6.6%)	p = 0.86
Prostaglandin Use	27 (96.4%)	1 (3.6%)	p = 0.52

Conclusion

- Generic PA is associated with higher rates of Irvine-Gass syndrome than either brand name PA or difluprednate, regardless of NSAID use.
- Generic PA may be inadequate after cataract extraction with phacoemulsification to prevent PCME.

Discussion

- Prior studies agree that there is no difference in PCME rates between difluprednate and brand name PA.¹
- Although anecdotal evidence suggests higher rates of PCME with generic PA, this is the first study that supports these findings.
- Similar to other studies,² PCME in our study group occurred more frequently in patients with mild NPDR, ERM, or RVO.
- Although DM, moderate NPDR, and PDR were not significantly associated with PCME, there was a trend toward significance.
- NSAID use after cataract surgery may help prevent PCME;³ our study showed a possible benefit in brand name or generic PA (31-33% relative risk reduction), although lack of significance in our study may be related to insufficient power.

Discussion

- Brand name PA particles are smaller and more uniform than generic PA, allowing them to stay in suspension longer.⁴
- Drop concentrations less predictable with generic PA (dose uniformity: difluprednate>brand name PA>generic PA).⁵



Acknowledgements

Donovan Reed, MD, Resident Ophthalmologist

James Aden, PhD, Biostatistician

Nikhil Godbole, MD, Staff Ophthalmologist

Marissa Wedel, MD, Vitreoretinal Surgeon

References

1. Baartman B, Gans R, Goshe J. Prednisolone versus dexamethasone for prevention of pseudophakic cystoid macular edema. *Can J Ophthalmol*. 2018;53(2):131-134.
2. Chu C, Johnston R, Buscombe C, Sallam A, Mohamed Q, Yang Y. Risk Factors and Incidence of Macular Edema after Cataract Surgery. *Ophthalmology*. 2016;123(2):316-323.
3. Juthani V, Clearfield E, Chuck R. Non-steroidal anti-inflammatory drugs versus corticosteroids for controlling inflammation after uncomplicated cataract surgery. Cochrane Eyes and Vision Group, ed. *Cochrane Database Syst Rev*. July 2017.
4. Roberts C, Nelson P. Comparative analysis of prednisolone acetate suspensions. *J Ocul Pharmacol Ther*. 2007;23(2):182-187.
5. Stringer W, Bryant R. Dose uniformity of topical corticosteroid preparations: difluprednate ophthalmic emulsion 0.05% versus branded and generic prednisolone acetate ophthalmic suspension 1%. *Clin Ophthalmol*. 2010;4:1119-24.