

AWARD NUMBER: W81XWH-16-1-0173

TITLE: Targeting a Novel Androgen Receptor-Repressed Pathway in Prostate Cancer Therapy

PRINCIPAL INVESTIGATOR: Qiming J. Wang, PhD

RECIPIENT: University of Pittsburgh

REPORT DATE: JUNE 2020

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited (A)

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE JUNE 2020		2. REPORT TYPE Final report		3. DATES COVERED 1 Sep 2016-28 Feb 2020	
4. TITLE AND SUBTITLE Targeting a Novel Androgen Receptor-Repressed Pathway in Prostate Cancer Therapy				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-16-1-0173	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Qiming J. Wang, PhD E-Mail: qjw1@pitt.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Pittsburgh				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The central hypothesis of the proposal is that PKD plays a crucial role in limiting the effectiveness of ADT by increasing prostate cancer survival through upregulating AURKA expression. AURKA encodes the protein Aurora A kinase, which is a master cell cycle regulator that is often dysregulated in human cancers. Our data demonstrated an important role of PKD in regulating Aurora A stability, implying that PKD's role in cancer cell survival and proliferation may be mediated through Aurora A. The primary goal of this study is to investigate the potential roles of PKD in mediating therapeutic resistance to ADT and to investigate the impact of PKD SMI-based combination therapies to curtail ADT-induced therapy resistance. This study will fill an important knowledge gap in signaling mechanisms and therapeutic targeting of PKD in the context of ADT-induced therapy resistance in prostate cancer. If successful, it will provide the basis for translating PKD-targeted agents to the clinical settings.					
15. SUBJECT TERMS Prostate cancer, protein kinase D, androgen, androgen receptor, therapeutic resistance, androgen-deprivation therapy, Aurora A, cell cycle.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
Unclassified	Unclassified	Unclassified	Unclassified	13	19b. TELEPHONE NUMBER (include area code)

TABLE OF CONTENTS

	<u>Page No.</u>
1. Introduction	4
2. Keywords	4
3. Accomplishments	4 - 6
4. Impact	7
5. Changes/Problems	7
6. Products	8
7. Participants & Other Collaborating Organizations	9
8. Special Reporting Requirements	10
9. Appendices	10

1. INTRODUCTION:

Prostate cancer progression to castration resistance associates with poor prognosis and high mortality. Androgen deprivation therapy (ADT) has been the standard therapy for advanced metastatic prostate cancer. However, ADT has been linked to the development of therapy resistance in part through inducing prosurvival adaptive responses. Increased understandings of these adaptive mechanisms will lead to discovery of new targets/therapies for mCRPC. We will investigate the roles of protein kinase D (PKD) in therapeutic resistance to ADT and the impact of PKD inhibitor-based combination therapies to curtail ADT-induced therapy resistance.

2. KEYWORDS:

Prostate cancer, androgen, androgen receptor, Protein kinase D, androgen-deprivation therapy, therapy resistance, transcriptional regulation, aurora A kinase, mitosis, cell cycle.

3. ACCOMPLISHMENTS:

What were the major goals of the project?

We seek to test **the central hypothesis is that PKD plays a crucial role in limiting the effectiveness of ADT by increasing prostate cancer survival through upregulating AURKA expression, and PKD small molecule inhibitors may enhance the efficacy of AR antagonists in prostate cancer treatment.** Our study will provide insights to the role of PKD in ADT-induced prosurvival responses relevant to the progression to CRPC. Successful completion of this study will define the role and mechanisms of PKD in treatment (ADT)-induced prostate cancer resistance. Three Specific Aims are proposed: **Aim 1.** Determine the mechanisms through which AR represses PKD1 expression in androgen-sensitive prostate cancer cells. **Aim 2.** Test the hypothesis that androgen deprivation-induced PKD1 expression promotes prostate cancer cell survival and ADT resistance through upregulating AURKA and CENPE. **Aim 3.** Determine the functional input of PKD in ADT resistance *in vivo* and assess the efficacy of PKD SMI in combination with AR antagonists in prostate cancer tumor xenografts.

What was accomplished under these goals?

MAJOR ACTIVITIES

The primary goal of this study is to investigate the potential roles of PKD in mediating therapeutic resistance to ADT and to investigate the impact of PKD SMI-based combination therapies to curtail ADT-induced therapy resistance. This remains our main focus during the third funding cycle. Major efforts were devoted to Aim 3, where we seek to determine the functional input of PKD in ADT resistance *in vivo* and assess the efficacy of PKD SMI in combination with AR antagonists in prostate cancer mouse models. The work is still on-going. A 6-month extension of this project has been approved.

SPECIFIC OBJECTIVES

Here is a summary of the major accomplishments achieved under each Specific Aim:

Specific Aim 1. Determine the mechanisms through which AR represses PKD1 expression in androgen-sensitive prostate cancer cells. [This aim has been completed]

(1) We identified PKD1 as a novel androgen-repressed gene. Our data showed that the transcription of PKD1 was repressed by androgen in androgen-sensitive prostate cancer cells. Steroid depletion caused an up-regulation of PKD1 transcript and protein, an effect that was reversed by AR agonist R1881 in a time- and concentration-dependent manner.

(2) Our data indicate that AR is required for the suppression of PKD1 expression by androgen. Specifically, our kinetic analysis indicated that the repression of PKD1 by androgen required the induction of a repressor protein. Further, inhibition or knockdown of AR reversed AR agonist-induced PKD1 repression.

(3) We discovered a novel AR/FRS2/MEK/ERK pathway through which androgen downregulated PKD1 expression. It is the first study that has identified fibroblast growth factor receptor substrate 2 (FRS2) as a downstream target of PKD1. Our study implies that the upregulation of the prosurvival PKD1 by antiandrogens may contribute to therapeutic resistance in prostate cancer treatment.

Publication: Zhang, L., Zhao, Z., Xu, S., Tandon, M., Lavalley, C. R., Deng, F., and *Wang, Q. J. (2017) Androgen suppresses protein kinase D1 expression through fibroblast growth factor receptor substrate 2 in prostate cancer cells. *Oncotarget*, 8:12800-12811.

Specific Aim 2. Test the hypothesis that androgen deprivation-induced PKD1 expression promotes prostate cancer cell survival and ADT resistance through upregulating AURKA and CENPE. [Study has been completed]

(1) Our study has identified PKD as key regulator of G2/M transition and mitotic entry. Our data indicated that PKD, particularly PKD2, was activated during G2 and M phase of cell cycle, and its activity is required for mitotic entry and progression. Inhibition of PKD resulted in mitotic catastrophe.

(2) We have discovered Aurora A kinase as a major target of PKD in cell cycle regulation. Overexpression of Aurora A reversed the effect of PKD inhibitor on cell cycle, thus Aurora A mediated the effects of PKD on G2/M transition and mitotic progression. We also found that PKD2 localizes in the centrosome with Aurora A by binding to γ -tubulin.

(3) We conclude that PKD2 modulates cell cycle by stabilizing aurora A kinase at centrosomes. Our data demonstrated that knockdown of PKD2 caused defects in centrosome separation, elongated G2 phase, mitotic catastrophe, and eventually cell death via apoptosis. Mechanistically, PKD2 interfered with Fbxw7 function to protect Aurora A from ubiquitin- and proteasome-dependent degradation.

Publication: Roy A., Veroli V., Prasad S., *Wang, Q. J. (2018) Protein Kinase D2 Modulates Cell Cycle By Stabilizing Aurora A Kinase at Centrosomes. *Mol Cancer Res*, 16: 1785-1797.

Specific Aim 3. Determine the functional input of PKD in ADT resistance *in vivo* and assess the efficacy of PKD SMI in combination with AR antagonists in prostate cancer tumor xenografts. [Study is near completion]

(1) Our data showed that knockdown of PKD sensitized PrCa cells to enzalutamide-induced cell death, while overexpression of PKD caused chemoresistance (see data presented in YR 2018-1019 progress report).

(2) We demonstrated that the PKD inhibitor, CRT0066101, inhibited tumor cell survival with an EC₅₀ of ~2 μ M in multiple prostate cancer cell lines. This inhibitor also synergized with enzalutamide in killing of

prostate cancer cell. CRT0066101 along and in combination with enzalutamide have been evaluated in vivo in tumor xenograft models and TRAMP prostate cancer mouse model. The results are consistent with the cellular study, indicating targeted inhibition of PKD by CRT0066101 enhances the efficacy of AR antagonists in vitro and in vivo.

Publication: Chao, Y., Zhang, X., Nagpal I., and *Wang, Q. J. Inhibition of PKD by CRT0066101 synergizes with enzalutamide at suppressing prostate cancer growth and metastasis. manuscript in preparation.

OTHER ACHIECEMENTS

We are investigating the role of PKD1 in prostate cancer metastasis, a key feature of mCRPC that directly contributes to the high mortality of this malignant disease. We have successfully established a bone metastasis mouse model by intracardiac injection of metastatic PC3-ML cells. We have completed the *in vivo* efficacy analysis of a PKD inhibitor - CRT101 on prostate cancer metastasis to the bone.

Publication: Roy, A., Prasad, S., Zhao, J., and *Wang, Q. J. Protein kinase D promotes prostate cancer cell bone metastasis by positively regulating Runx2 in a MAPK-ERK1/2-dependent manner. Manuscript submitted to *Mol Cancer Res*.

What opportunities for training and professional development has the project provided?

The project provided opportunities for training postdoctoral fellows.

Since October 2011, all postdoctoral trainees in the schools of the health sciences at the University of Pittsburgh have been required to complete an annual career development plan (also known as an individual development plan - IDP) as part of our institution's [Postdoctoral Career Development and Progress Assessment Process](#).

This process, overseen by the Center for Postdoctoral Affairs in the Health Sciences, requires that a postdoc work with his or her faculty mentor to establish an annual career development plan and to also identify two additional individuals to serve as members of the postdoc's mentoring team. The postdoc also completes an annual self-assessment relative to his or her career development plan which contributes in part to the faculty mentor's annual assessment of the postdoc's progress towards his or her career goals.

How were the results disseminated to communities of interest?

Nothing to Report.

What do you plan to do during the next reporting period to accomplish the goals?

This is the final report for this award.

4. IMPACT:

What was the impact on the development of the principal discipline(s) of the project?

This application fills an important knowledge gap in signaling mechanisms and therapeutic targeting of PKD in the context of ADT-induced therapy resistance in prostate cancer. (1) The successful completion of the study will provide mechanistic insights to AR-regulated prosurvival pathways centered on PKD in therapy resistance and tumor progression. (2) PKD may be targeted to enhance the therapeutic efficacy of antiandrogens and other chemotherapeutic agents, indicating the significant translational value of the study. (3) The identification of AURKA as biomarkers of PKD SMIs may facilitate the translation of these agents to the clinic. (4) AURKA may be used for identifying PKD SMI-sensitive prostate tumor subtypes and facilitate the application of PKD-targeted agents in personalized therapy.

What was the impact on other disciplines?

Nothing to Report.

What was the impact on technology transfer?

Nothing to Report.

What was the impact on society beyond science and technology?

Nothing to Report.

5. CHANGES/PROBLEMS:

Changes in approach and reasons for change

Nothing to Report.

Actual or anticipated problems or delays and actions or plans to resolve them

Nothing to Report.

Changes that had a significant impact on expenditures

Nothing to Report.

Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents

Nothing to Report.

Nothing to Report.

Significant changes in use or care of vertebrate animals

Nothing to Report.

Significant changes in use of biohazards and/or select agents

Nothing to Report.

6. PRODUCTS: List any products resulting from the project during the reporting period. If there is nothing to report under a particular item, state “Nothing to Report.”

- **Publications, conference papers, and presentations**

Report only the major publication(s) resulting from the work under this award.

Journal publications.

1. Zhang, L., Zhao, Z., Xu, S., Tandon, M., Lavalley, C. R., Deng, F., and ***Wang, Q. J.** (2017) Androgen suppresses protein kinase D1 expression through fibroblast growth factor receptor substrate 2 in prostate cancer cells. *Oncotarget*, 8:12800-12811.
2. Roy, A., Ye, J., ***Wang, Q. J.** (2017) Protein Kinase D: A Potential Therapeutic Target in Prostate Cancer. *Molecular and Cellular Pharmacology*, 9 (1), 1-4
3. Roy, A., Ye, J., Deng, F., ***Wang, Q. J.** (2017) Protein Kinase D Signaling in Cancer: A Friend or Foe? *BBA-Rev Cancer*, 1868(1):283-294.
4. Roy A., Veroli V., Prasad S., ***Wang, Q. J.** (2018) Protein Kinase D2 Modulates Cell Cycle By Stabilizing Aurora A Kinase at Centrosomes. *Mol Cancer Res*, 16: 1785-1797.
5. Xu, W., Zeng, F., Li, S., Li G., Lai X., **Wang Q. J.**, and ***Deng F.** (2018) Crosstalk of protein kinase C ϵ with Smad2/3 promotes tumor cell proliferation in prostate cancer cells by enhancing aerobic glycolysis. *Cell Mol Life Sci*, 75: 4583-4598.
6. ¹Zhang, L., ¹Li, Z., Liu, Y., Xu, S., Tandon, M., Appelboom, B., LaValle, C. R., Chiosea, S. I., Wang, L., Sen, M., Lui, V. W. Y., Grandis, J. R., and ***Wang, Q. J.** (2018) Analysis of oncogenic activities of protein kinase D1 in head and neck squamous cell carcinoma. *BMC Cancer*, 18: 1107-1123.
7. Xu, W., Qian, J., Zeng, G., Li, S., Guo, W., Chen, L., Li, G., Zhang, Z., **Wang, Q. J.**, and ***Deng, F.** (2019) PKDs promote tumor angiogenesis through mast cell recruitment and expression of angiogenic factors in prostate cancer microenvironment. *J Exp Clin Cancer Res*, 38: 114-127.

Books or other non-periodical, one-time publications.

Nothing to Report.

Other publications, conference papers, and presentations.

“Protein Kinase D 2 Promotes Cell Cycle Progression by Stabilizing Aurora A Kinase at Centrosome in Cancer Cells”, FASEB SRC: The Protein Kinases and Protein Phosphorylation Conference, poster presentation, 2019 July.

- **Website(s) or other Internet site(s)**

Nothing to Report.

- **Technologies or techniques**

Nothing to Report.

- **Inventions, patent applications, and/or licenses**

Nothing to Report.

- **Other Products**

Nothing to Report.

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

What individuals have worked on the project?

Name: Qiming Wang

Project Role: Principal Investigator

Research Identifier (e.g. ORCID ID):

Nearest Person Month Worked: 42 mon

Contribution to the project: the overall scientific management of this application, including experimental design, data collection and analysis, supervision of postdoctoral research associates, preparation and submission of reports and manuscripts.

Funding Support: DOD PC150190

Name: Manuj Tandon

Project Role: Research Instructor

Research Identifier (e.g. ORCID ID):

Nearest Person Month Worked: 12 mon

Contribution to the project: overall execution of the project, data collection, and analysis; writing of reports and papers.

Funding Support: DOD PC150190

Name: Sahdeo Prasad, Adhiraj Roy
Project Role: Postdoctoral Associate
Research Identifier (e.g. ORCID ID):
Nearest Person Month Worked: Dr. Prasad: 12 mon / Dr. Roy: 24 mon
Contribution to the project: technical assistance in execution of experiments, data analysis, and human/mouse tissue pathohistological analysis.
Funding Support: DOD PC150190

Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?

Dr. Wang has been awarded a R01 (R01CA229431) and an AHA (19TPA34850096) grant starting 7/1/2019.

What other organizations were involved as partners?

Nothing to Report.

8. SPECIAL REPORTING REQUIREMENTS

COLLABORATIVE AWARDS:

Nothing to Report.

QUAD CHARTS:

Nothing to Report.

- 9. APPENDICES:** Attach all appendices that contain information that supplements, clarifies or supports the text. Examples include original copies of journal articles, reprints of manuscripts and abstracts, a curriculum vitae, patent applications, study questionnaires, and surveys, etc.

Three publications that are directly related to the project are attached.

Research Paper

Androgen suppresses protein kinase D1 expression through fibroblast growth factor receptor substrate 2 in prostate cancer cells

Liyong Zhang^{1,*}, Zhenlong Zhao^{1,2,*}, Shuping Xu¹, Manuj Tandon¹, Courtney R. LaValle¹, Fan Deng³, Q. Jane Wang¹

¹Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

²Department of Anesthesiology, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong, China

³Department of Cell Biology, School of Basic Medical Sciences, Southern Medical University, Guangzhou, Guangdong, China

*These authors contributed equally to this work

Correspondence to: Q. Jane Wang, email: qjw1@pitt.edu

Keywords: androgen, androgen-repressed genes, androgen receptor, protein kinase D1, prostate cancer

Received: August 21, 2016

Accepted: December 27, 2016

Published: January 06, 2017

ABSTRACT

In prostate cancer, androgen/androgen receptor (AR) and their downstream targets play key roles in all stages of disease progression. The protein kinase D (PKD) family, particularly PKD1, has been implicated in prostate cancer biology. Here, we examined the cross-regulation of PKD1 by androgen signaling in prostate cancer cells. Our data showed that the transcription of PKD1 was repressed by androgen in androgen-sensitive prostate cancer cells. Steroid depletion caused up regulation of PKD1 transcript and protein, an effect that was reversed by the AR agonist R1881 in a time- and concentration-dependent manner, thus identifying PKD1 as a novel androgen-repressed gene. Kinetic analysis indicated that the repression of PKD1 by androgen required the induction of a repressor protein. Furthermore, inhibition or knockdown of AR reversed AR agonist-induced PKD1 repression, indicating that AR was required for the suppression of PKD1 expression by androgen. Downstream of AR, we identified fibroblast growth factor receptor substrate 2 (FRS2) and its downstream MEK/ERK pathway as mediators of androgen-induced PKD1 repression. In summary, PKD1 was identified as a novel androgen-suppressed gene and could be downregulated by androgen through a novel AR/FRS2/MEK/ERK pathway. The upregulation of prosurvival PKD1 by anti-androgens may contribute to therapeutic resistance in prostate cancer treatment.

INTRODUCTION

Prostate cancer is the most common noncutaneous malignancy and the second leading cause of cancer-related deaths among men in the United States. The initiation and progression of prostate cancer is uniquely dependent on androgen receptor (AR)-induced signaling. Although androgen deprivation therapy provides an initial favorable response in advanced prostate cancer, the more aggressive castration-resistant prostate cancer (CRPC) develops invariably in almost all patients, eventually leading to death. It has become increasingly clear that continuous activation of the AR in CRPC remains the main driving force of tumor progression and metastasis. Thus,

understanding the critical events associated with the AR signaling is essential for developing novel and effective therapies to treat CRPC.

The protein kinase D (PKD) family of serine/threonine kinases belongs to the Ca²⁺/calmodulin-dependent protein kinase (CAMK) superfamily [1, 2]. To date, three isoforms of PKD have been identified, PKD1 (formerly PKC μ) [3, 4], PKD2 [5], and PKD3 (formerly PKC ν) [6]. In intact cells, PKD activation involves phosphorylation of two conserved serine residues in the activation loop by DAG-responsive PKCs [7–9], and PKD activity can be maintained independently of PKC through autophosphorylation [10, 11]. Emerging evidence supports that PKD has an important role in carcinogenesis

and tumor progression [12, 13]. A recent report suggested that a hotspot activating mutation in *PRKD1*, the gene encoding PKD1, may drive polymorphous low-grade adenocarcinoma (PLGA), the second most frequent type of malignant tumor of the minor salivary glands [14]. PKD regulates a variety of tumor-associated biological processes, including tumor cell proliferation, growth, survival, migration, invasion, secretion, and angiogenesis [12, 15–20]. Aberrant PKD activity and expression have been demonstrated in tumor cell lines and tumor tissues from the pancreas [18], skin [19, 21], breast [22], and prostate [20, 23]. In particular, PKD has been shown to play an important role in the pathogenesis of prostate cancer [20, 24–26], and targeted PKD inhibition potently blocks prostate cancer cell proliferation and survival [26, 27].

Fibroblast growth factor (FGF) signaling is a highly complex signaling network that comprises 18 ligands, which bind to and activate four highly conserved transmembrane tyrosine kinase receptors (FGFR1, FGFR2, FGFR3, and FGFR4). The FGF/FGFR pathway plays an important role in cancer development and progression by modulating a variety of biological processes, including cell proliferation, survival, and migration [28, 29]. FGFR substrate 2 (FRS2/FRS2 α), also known as FGFR-signaling adaptor SNT1 (suc1-associated neurotrophic factor target 1), is regarded as the ‘conning center’ for intracellular signaling elicited by the activation of FGFRs at the cell surface. FRS2 forms complexes with Grb2-Sos and Grb2-Gab1 to activate the Ras/Raf/MEK/ERK and PI3K/Akt pathways [29, 30]. Although FRS2 expression is not regulated by androgen [31], androgen-sensitive prostate cancer cells express FGF2, and its expression is upregulated in response to androgen stimulation [32]. Thus, androgen regulates the activity of FGFR signaling in prostate cancer cells.

In this study, we report for the first time that PKD1 was tightly regulated by androgen at the transcriptional level in prostate cancer cells and was a novel androgen-repressed gene. Inhibition or knockdown of androgen receptor (AR) blocked androgen depletion-induced PKD1 expression, indicating that AR was required for the repression of *PRKD1* gene expression. Further analysis identified FRS2 as a novel mediator of androgen-induced PKD1 repression. The regulation of PKD1 by androgen and AR may have important implications in the therapeutic response to AR-targeted agents.

RESULTS

Androgen repressed PKD1 expression in androgen-sensitive prostate cancer cells

Androgen signaling plays a crucial role in prostate cancer initiation and progression. In this study, we sought to determine whether androgen modulated PKD1 expression and signaling. PKD1 was detected in androgen-sensitive LNCaP cells and two castration-resistant LNCaP-

derivative cell lines, C4-2 (androgen-hypersensitive) and C81 (androgen-insensitive), but not in androgen-sensitive LAPC4 cells. As shown in Figure 1A, a significant increase in PKD1 expression was observed upon androgen depletion (AD) in LNCaP and C4-2 cells and to a lesser extent in C81 cells. R1881, a synthetic androgen agonist, induced remarkable concentration-dependent suppression of PKD1 expression at the transcript (Figure 1B) and protein (Figure 1C) levels in LNCaP and C4-2 cells. R1881 also suppressed PKD1 expression in VCaP cells, a castration-resistant prostate cancer cell line that expresses wild-type AR, in a concentration-dependent manner (Figure 1D). Interestingly, PKD2 expression was similarly suppressed by R1881 in a concentration-dependent manner in LNCaP and VCaP cells (Supplementary Figure 1A–1B). PKD3 was also upregulated upon androgen withdraw in LNCaP cells, despite its low endogenous expression (Supplementary Figure 1A). In contrast, androgen did not affect the expression of PKD1 and PKD2 in another castration-resistant cell line, 22Rv1, which expresses both full-length AR and truncated AR variants (Supplementary Figure 1C), suggesting that the effect of androgen may be cell context-dependent. Taken together, we concluded that PKD1 was an androgen-repressed gene.

PKD1 expression was dependent on the induction of a repressor protein

The kinetics of PKD1 regulation in response to androgen deprivation or R1881 treatment was examined. As shown in Figure 2A, androgen deprivation gradually up regulated PKD1 protein expression, which peaked at 16–24 h, while R1881 suppressed PKD1 expression with similar kinetics. The induction of PKD1 transcript and its inhibition by R1881 correlated well with the time-course of protein expression (Figure 2B).

To gain insights into the regulation of PKD1 by androgen, we first examined whether R1881 affected PKD1 mRNA stability. The half-life ($t_{1/2}$) of PKD1 mRNA was determined in the presence of actinomycin D, an inhibitor of gene transcription. As shown in Figure 2C, the $t_{1/2}$ of PKD1 mRNA was about 4 h, which was not significantly altered by the addition of R1881 ($p > 0.5$), indicating that R1881 did not impact the stability of PKD1 mRNA. Next, cycloheximide (CHX) was used to inhibit protein synthesis to determine whether the regulation of PKD1 gene expression by androgen involved *de novo* protein synthesis. CHX induced a nearly 2-fold increase in PKD1 expression and completely blocked R1881-induced PKD1 downregulation, indicating that the suppression of PKD1 expression likely required the induction of a repressor protein (Figure 2D). This finding was in line with the gradual onset of PKD1 regulation by androgen, further supporting the involvement of a repressor protein. Taken together, our data indicated that androgen-regulated PKD1 expression was dependent on the presence of a repressor protein.




Figure 1: Androgen repressed PKD1 expression. (A) Effects of androgen depletion on PKD1 expression in prostate cancer cells. LNCaP, C4-2, C81, and LAPC4 cells were grown for 48 h in normal androgen-containing (AC) or androgen-depleted (AD) medium supplemented with charcoal-stripped FBS. Cells were lysed and subjected to immunoblotting for PKD1 and GAPDH (loading control). *Bottom*, quantitative measurement of band intensity by densitometry analysis. The data were expressed as % control with C81 (AC) set as 100%. Data are the mean $\pm$ SEM of four independent experiments. (B) Androgen inhibited PKD1 transcription. Total RNAs from LNCaP were extracted, and real-time RT-qPCR was conducted using specific PKD1 primers. *GAPDH* was used as internal control. Data are the mean $\pm$ SEM of three independent experiments. (C) Androgen suppressed PKD1 protein expression. LNCaP and C4-2 cells were grown in androgen-depleted medium for 48 h, following by treatment without or with increasing concentrations of androgen R1881. Cells were harvested after 24 h and subjected to immunoblotting. *Bottom*, the band intensity was quantified by densitometry analysis, and data are the mean $\pm$ SEM of ten (LNCaP) or three (C4-2) independent experiments. (D) Androgen suppressed PKD1 protein expression in castration-resistant VCaP cells. VCaP cells were grown in androgen-depleted medium for 48 h, followed by treatment without or with androgen R1881 for 24 h. Cells were harvested for immunoblotting. Data from one of three independent experiments are shown. *Right*, quantitative measurement of band intensity from three experiments is shown. ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.




Figure 2: PKD1 expression was dependent of the induction of a repressor protein. (A) Kinetics of PKD1 regulation by androgen. *Left panels*, LNCaP cells were grown in AD medium for the indicated times. *Right panels*, LNCaP cells were grown in AD medium for 48 h, followed by treatment with R1881 (1 nM) for the indicated times. Cells were harvested and subjected to immunoblotting for PKD1 and tubulin (loading control). Cells grown in AC medium were used as the control. Representative data from one of four experiments are shown. *Bottom*, quantitative measurement of band intensity by densitometry analysis. Data are the mean $\pm$ SEM of three independent experiments. (B) Kinetics of PKD1 transcript expression. LNCaP cells were treated as above in "A". Total RNAs were extracted, and the kinetics of PKD1 mRNA induction/suppression were examined by real time RT-qPCR. (C) R1881 did not affect PKD1 mRNA stability. LNCaP cells were grown either in AD medium for 48 h, followed by the addition of actinomycin D (2 ng/mL) with or without R1881 (1 nM) for the indicated times. Total RNAs were extracted and subjected to real time RT-qPCR for analysis of PKD1 transcripts. Not significant by paired t test ($p > 0.5$). (D) PKD1 expression required the induction of a repressor protein. LNCaP cells were grown in AD medium for 48 h, followed by R1881 treatment with or without cycloheximide (CHX) for 6 or 10 h. Total RNAs were extracted, and the levels of PKD1 mRNA were measured by real-time RT-qPCR. GAPDH was used as a loading control. Data are the mean $\pm$ SEM of at least three independent experiments. ns, not significant; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$.

AR mediated PKD1 repression by androgen

Androgens are important hormones for normal physiology and are responsible for certain disease conditions. Their actions are mediated by the AR, a ligand-dependent nuclear transcription factor. Androgens binds to AR after entering the cells to form an androgen-receptor complex, which then translocates to the nucleus where it binds to androgen response elements (AREs) in the promoter regions and regulates the transcription of its target genes. The actions of AR can be blocked by AR inhibitors, such as bicalutamide (Casodex) or enzalutamide (MDV3100). Bicalutamide is known to bind AR and leads to the formation of a transcriptionally inactive AR complex [33]. In this study, we sought to examine whether AR was required for the repression of PKD1 expression by R1881, and bicalutamide was used to determine whether the inhibition of AR activity affected PKD1 expression. After androgen deprivation, LNCaP and C4-2 cells were treated with R1881 at 1 nM in the presence or absence of bicalutamide (10 μ M). As shown in Figure 3A, bicalutamide significantly reversed R1881-induced PKD1 repression in LNCaP and C4-2 cells. In LNCaP cells, inhibition of AR by bicalutamide also upregulated PKD1 protein expression in a concentration-dependent manner (Figure 3B). The specific role of AR was then examined using multiple AR-targeted siRNAs. Our data showed that knockdown of AR by three siRNAs targeting different regions of the AR transcript significantly blocked R1881-induced PKD1 suppression in LNCaP (Figure 3C) and C4-2 cells (data not shown). AR knockdown was confirmed by western blotting. Taken together, these data suggested that AR was required for the transcriptional repression of PKD1 gene expression caused by androgen stimulation.

To determine whether AR directly regulated the expression of PKD1, we analyzed the promoter region of PKD1, which led to the identification of two potential AREs upstream of the transcription start site (TSS). The human PKD1 gene spans ~45.7 kb. Analysis of up to 5000 bp of the promoter region upstream from the TSS revealed two putative AREs. (ARE1, 5'-AGTACTTTAAGCTCT-3'; ARE2, 5'-AGAACAAAATAAGCT-3'; (Supplementary Figure 2A). The regions (pm1 and pm2) that contained the AREs were separately cloned into the pTA-Luc reporter. Their activities were analyzed in LNCaP cells cultured in the presence or absence of androgen depletion, followed by treatment with or without R1881. Our data indicated that no luciferase activity was detected from both reporters in LNCaP cells (Supplementary Figure 2B), implying that the AREs in PKD1 promoter did not play an active role in regulating PKD1 transcription in response to androgen stimulation.

An AR co repressor screen revealed FRS2 as the potential mediator of androgen-induced PKD1 repression

The involvement of AR and an androgen-induced repressor protein prompted us to conduct an esiRNA screen that targeted 23 AR corepressors and other related proteins. LNCaP cells were transfected individually with 23 esiRNAs, followed by androgen depletion and treatment with or without R1881. Levels of PKD1 transcript were analyzed by real time RT-qPCR. In the controls, androgen depletion induced PKD1 expression, and treatment with R1881 caused over 2-fold reduction in PKD1 mRNA. As shown in Figure 4, similar to the non targeting siRNA, R1881-induced PKD1 repression was not affected by the depletion of all target genes, with the exception of FRS2. Knockdown of FRS2 by esiRNA completely reversed the repression of PKD1 transcription by R1881 (Figures 4C, 5A). In summary, FRS2 was identified as a potential repressor of PKD1 gene expression.

Androgen repressed PKD1 expression through a FGFR/FRS2/MEK/ERK pathway

The role of FRS2 was further validated using FRS2 siRNAs (si-FRS2-1, -2). Depletion of FRS2 abolished the R1881-induced suppression of PKD1 transcription, confirming FRS2 as a potential mediator of androgen-dependent PKD1 repression (Figure 5A). Furthermore, at the protein level, knockdown of FRS2 by two different siRNAs completely abrogated the downregulation of PKD1 by R1881 (Figure 5B). *FRS2* siRNAs caused significant knockdown of *FRS2* mRNA (Figure 5C). Thus, FRS2 mediated androgen-induced PKD1 repression.

The adaptor protein FRS2 is a major mediator of the FGFR signaling in normal and malignant cells. FGFR stimulation by FGF leads to the tyrosine phosphorylation of FRS2, which then forms a complex with Grb2 and Sos to activate the downstream Ras/Raf/MEK/ERK signaling pathway. Androgen-sensitive LNCaP cells express low levels of FGF2, and its expression is upregulated in response to androgen stimulation [32]. Here, we sought to determine whether the FRS2-mediated FGFR signaling pathway was involved in the regulation of PKD1 by androgen. As shown in Figure 5D, PD173074, an inhibitor of FGFR, significantly reversed R1881-induced PKD1 repression, indicating that FGFR activity was required for the inhibition of PKD1 by androgen/AR. Next, the role of the FGF-activated MEK/ERK MAPK signaling pathway was evaluated. Our data demonstrated that R1881-induced PKD1 suppression was abrogated in a concentration-dependent manner by UO126, a MEK inhibitor. Thus, MEK/ERK activity was also required for the suppression of PKD1 by androgen (Figure 5E). Since the suppression of PKD1 is likely associated with the secretion of a

FGFR ligand, we tested the effects of inhibiting secretory pathways on the expression of PKD1 using brefeldin A (BFA), a fungal metabolite and an inhibitor of intracellular protein transport that inhibits constitutive secretion from the *trans*-Golgi network. Our data indicated that BFA at 5 and 10 μ M completely reversed androgen-induced PKD1 suppression (Figure 5F). In summary, our data implied that androgen suppressed PKD1 expression through an indirect FGFR/FRS2/MEK/ERK pathway in prostate cancer cells.

DISCUSSION

In this study, we present the novel findings demonstrating that PKD1 was repressed by androgen/AR at the mRNA and protein levels in androgen-sensitive prostate cancer cells, identifying PKD1 as a novel androgen-repressed gene. We further identified FRS2 as a novel mediator of androgen-induced repression of PKD1 expression. The cross-regulation of PKD1 by androgen/AR places PKD1 in the AR-induced signaling




Figure 3: AR mediated PKD1 repression by androgen. (A) AR inhibition led to increased PKD1 expression. LNCaP and C4-2 cells cultured in AD medium for 48 h were treated with or without R1881 (1 nM) $\pm$ bicalutamide (Bic) (10 μ M) for 16 h. *Bottom*, quantitative measurement of band intensity from two experiments. (B) Bicalutamide caused PKD1 upregulation. LNCaP cells were treated with increasing concentrations of bicalutamide for 48 h, followed by immunoblotting for PKD1. The band intensity was quantified by densitometry analysis, and data are the mean $\pm$ SEM of three independent experiments. (C) AR was required for transcriptional regulation of PKD1 by androgen. LNCaP cells were transfected with non targeting siRNA (si-NT) or AR siRNAs (si-AR-1, -2, -3). After 48 h, the medium was replenished with AD medium with or without R1881 (1 nM) for 16 h. Cells were collected and subjected to immunoblotting for PKD1, AR, and GAPDH. *Right*, quantitative measurement of band intensity for PKD1 (*top*) and AR (*bottom*) from three experiments is shown. ns, not significant; * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. Data are the mean $\pm$ SEM of six independent experiments.

network, which is critical to prostate cancer progression. The transcriptional regulation of PKD isoforms has not been studied in the past. Our study provides the first mechanistic understanding of a novel androgen-induced AR/FRS2/MEK/ERK pathway that regulates the expression of PKD1. As a well-documented prosurvival signaling protein, PKD1 upregulation in response to androgen deprivation and anti-androgen treatment may have significant implications in therapy resistance and progression to CRPC.

The class I steroidal nuclear receptor AR is a critical regulator of tumor initiation and progression in both early and advanced prostate cancer. As a transcription factor, AR exerts its actions mainly through regulating the expression of a host of target genes. Among them, AR-stimulated genes have been extensively studied, with prostate-specific antigen (PSA) being the best characterized. In contrast, AR-repressed target genes have not been well characterized. These genes constitute a large portion of AR-targeted genes, and some have been shown to play essential roles in prostate cancer progression [34, 35]. Diverse mechanisms have been proposed to account for the repression of target gene expression by AR. These include both genomic mechanisms, such as active repression via the recruitment

of corepressor complexes, and nongenomic mechanisms, such as regulation of signaling pathways [34]. Our data showed that inhibition or silencing of AR blocked the suppression of PKD1 by R1881, indicating that AR was required for the downregulation of PKD1. Initially, analysis of the 5' promoter region of the PKD1 gene led to the identification of two potential AREs, which prompted us to investigate the direct role of AR in transcriptional repression of the PKD1 gene. However, analysis of the transcriptional activity of the ARE-containing PKD1 promoter failed to detect any androgen-induced transcriptional activities associated with pm1 and pm2, suggesting that the identified potential AREs may be inactive. Although less common, inactive AREs have been demonstrated, even in the presence of AR binding, and more complex mechanisms have been suggested to be involved in the regulation of genes nearby these AREs in prostate cancer cells [36, 37]. Importantly, kinetic analysis demonstrated a slow and gradual onset of PKD1 downregulation at the protein and transcript levels, which peaked at about 16–20 h in response to androgen; this finding also provides evidence against a mechanism involving active transcriptional repression through direct interaction with AREs. The involvement of an androgen-induced repressor





Figure 4: Screening of AR corepressors. (A–C) An esiRNA screen that targeted 23 AR corepressors and other related proteins was conducted in LNCaP cells. The cells transfected with esiRNAs were subjected to androgen depletion for 48 h, followed by treatment with or without R1881. Levels of PKD1 transcript was analyzed by real time RT-qPCR. Non targeting siRNA (si-NT) was used as the control. Student's *t*-tests were used to determine the statistical significance between the untreated and R1881-treated groups within each pair of esiRNA knockdown samples. **p* < 0.05; ***p* < 0.01; ****p* < 0.001.

is supported by the data showing that CHX abolished androgen-induced PKD1 repression. Although this may also occur without the synthesis of a repressor, for example, the suppressive effects of AR could be mediated through its interaction with a pre-existing labile protein at the AR-repressed loci; CHX treatment will similarly abolish the repressive effect mediated by this labile protein. Certainly, our findings do not exclude the possibility that there may be distal ARE sites that bind to AR and contribute to AR-mediated PKD1 repression. Overall, our current data support an AR-mediated indirect mechanism involving the cell surface adaptor protein FRS2 in the repression of PKD1 by androgen. These findings were based on an unbiased RNAi screen of a library of AR corepressor proteins. Further analysis validated the role of FRS2, as well as its upstream FGFR and the downstream MEK/ERK pathway, in the regulation of PKD1 by androgen.

In androgen-sensitive prostate cancer cells, depletion of FRS2 blocked R1881-induced PKD1

suppression at both the transcriptional and protein levels. Additionally, inhibition of FGFR and MEK, as well as protein secretion, blocked R1881-induced repression of PKD1. Thus, androgen may repress PKD1 through an AR-induced FGFR/FRS2/MEK/ERK pathway to inhibit PKD1 expression in prostate cancer cells. A previous study showed that FRS2 expression is not regulated by androgen in LNCaP cells [31]. However, in androgen-sensitive LNCaP cells, low levels of FGF2 are detected, and the expression of FGF2 is upregulated in response to androgen stimulation [32]. Additionally, androgen stimulates the activity and production of FGF2 and FGF-binding protein in PC3 prostate cancer cells with stably overexpressed AR [38]. In a different study, however, Kassen et al. showed that FGF2 is not expressed, and androgen in turn acts by increasing the bioavailability of FGF2 by releasing trapped FGF2 from the extracellular matrix through activation of heparinase, which leads to activation of FGFR and stimulation of LNCaP cell proliferation






Figure 5: FRS2 was required for androgen-induced PKD1 repression. (A) Knockdown of FRS2 reversed androgen-induced repression of PKD1 transcription. Cells were transfected with FRS2 siRNA (si-FRS2-1) and a non targeting siRNA (si-NT), followed by treatment with or without R1881. PKD1 transcripts were analyzed by real-time RT-qPCR. Representative data from one of three independent experiments with triplicate measurements are shown. (B) Knockdown of FRS2 blocked the repression of PKD1 protein by androgen. LNCaP cells were transfected with two different FRS2 siRNAs (si-FRS2-1, -2), followed by treatment with R1881. *Right*, quantitative measurement of band intensity for PKD1 from three experiments is shown. (C) Real-time RT-PCR confirmed the knockdown of FRS2. Cells from "B" were subjected to RNA extraction, followed by real-time RT-PCR for levels of PKD1 transcript. (D–F) Androgen repression of PKD1 was dependent on a secretory pathway involving FGFR and MEK. LNCaP cells were grown in AC or AD medium for 48 h, followed by treatment with or without R1881 in the presence or absence of the FGFR inhibitor PD17034 (D), the MEK inhibitor UO0126 (E), and brefeldin A (F) for 16 h. Cell lysates were subjected to immunoblotting for PKD1. Representative images from one of at least three independent experiments are shown. *Bottom*, quantitative measurement of band intensity by densitometry analysis. Data are the mean $\pm$ SEM of five to seven independent experiments.

[39]. Regardless of these discrepancies, in all cases, AR promotes the activation of FGFR in prostate cancer cells, which results in phosphorylation of FRS2 and activation of the downstream Ras/Raf/MEK/ERK signaling pathway. By inhibiting MEK activity, we confirmed the requirement

for MEK/ERK signaling activity in the suppression of PKD1 by R1881. This evidence supports the notion that PKD1 is repressed by an AR-induced FGFR/FRS2/MEK/ERK pathway in androgen-sensitive prostate cancer cells. The binding of FGF to FGFR leads to the recruitment

of multiple adaptor proteins, including FRS2, Grbs, Sos, and Gab1, and induces the activation of multiple downstream signaling pathways, including MEK/ERK, PI3K/Akt, PLC γ /PKC, and Stat3 pathways. We must state that although our data demonstrated a major role of the MEK/ERK pathway in the regulation of PKD1 expression by androgen, our data did not completely exclude the potential involvement of other pathways, which will be investigated in our future studies.

Our study identified PKD1 as an androgen/AR-repressed gene and uncovered a novel indirect mechanism through which AR regulates PKD1 expression. Although the functional implication of this regulation in prostate cancer progression is still unclear, PKD1 is an important prosurvival signaling protein in normal and cancer cells that functions by regulating multiple signaling pathways, such as stimulating NF- κ B, ERK1/2, and Akt and inhibiting JNK and p38 [20, 25, 40]. This notion is further supported by our previous findings that PKD1 protects androgen-sensitive LNCaP prostate cancer cells from phorbol ester-induced apoptosis [25]. Thus, the upregulation of PKD1 as a result of inhibition or loss of AR may promote tumor cell survival and contribute to therapeutic resistance to AR-targeted agents. This further implies that PKD may represent a viable target for mitigating therapy resistance. In castration-resistant C81, 22Rv1, and VCaP cells, we observed different responses to androgen in terms of PKD1 regulation; although androgen did not affect PKD1 expression in 22Rv1 cells, VCaP cells, which express wild-type AR, did respond to androgen stimulation by downregulating PKD1 in a concentration-dependent manner, and minor effects were also observed in C81. This cell context-dependent responsiveness to androgen may be linked to the activity of the AR/FGFR/FRS2 signaling pathway and variations in the expression of its signaling components.

In summary, our study identified PKD1 as a novel androgen/AR-suppressed gene. The suppression of PKD1 was mediated through an indirect mechanism that involved FRS2, a cell surface adaptor protein that connects FGF/FGFR to the downstream MEK/ERK signaling pathway. Our findings suggested that the prosurvival function of PKD1 may have significant implications in prostate cancer progression and therapy resistance. PKD1 may be targeted to enhance the therapeutic response to anti-androgens in prostate cancer treatment.

MATERIALS AND METHODS

Reagents and antibodies

The synthetic androgen methyl trienolone (R1881) was obtained from Perkin-Elmer Life Sciences (Boston, MA), and bicalutamide was purchased from Enzo Life Sciences (Farmingdale, NY). Charcoal-treated fetal bovine serum (FBS) was from Hyclone (Logan, UT)

and Sigma (St. Louis, MO). Other cell culture reagents and media were from American Type Culture Collection (ATCC; Rockville, MD). Anti-PKD1, anti-PKD2, and anti-AR antibodies were purchased from Cell Signaling Technology (Danvers, MA). Antibodies targeting GAPDH and α -tubulin were from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Goat anti-rabbit and goat anti-mouse horseradish peroxidase-conjugated secondary antibodies were from Promega (Madison, WI).

Cell culture and siRNA transfection

LNCaP, C4-2, C81, VCaP, and PC-3 cells were obtained from ATCC (Manassas, VA) and were cultured according to the manufacturer's recommendations. LNCaP cells were discarded after 12 passages. The 23 AR corepressor esiRNAs were obtained from Sigma-Aldrich (Supplementary Table 1). The non targeting siRNA and AR and FRS2 siRNAs were obtained from Integrated DNA Technologies (Coralville, IA). The esiRNAs and siRNAs were transfected into cells using DharmaFECT reagent according to the manufacturer's instructions (GE Dharmacon, Lafayette, CO).

Western blot analysis

Cells were collected and lysed in IP lysis buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1.5 mM MgCl₂, 10% glycerol, 1% Triton X-100, 5 mM EGTA, 20 μ M leupeptin, 1 mM AEBSF, 1 mM NaVO₃, 10 mM NaF, and 1 $\times$ protease inhibitor cocktail). Lysate protein concentrations were determined using the Pierce BCA Protein Assay Kit (Thermo Fisher, Hudson, NH). Cell lysates were separated by SDS-PAGE and transferred to nitrocellulose membranes. After blocking with 5% milk, the membranes were incubated with primary antibodies in blocking buffer at 4°C overnight. After washing, the membranes were incubated with secondary antibodies at room temperature for 1 h. Protein bands were detected using an enhanced chemiluminescence (ECL) kit. Anti- α -tubulin or anti-GAPDH antibodies were used as a loading control. Densitometry analyses were performed with ImageJ software (NIH).

Real-time RT-PCR

Total RNAs were isolated from LNCaP cells using a RNeasy kit (Qiagen, Valencia, CA) according to the manufacturer's protocol. One microgram of total RNAs was used to generate cDNA using an iScript cDNA synthesis kit. Real-time PCR was subsequently performed using SsoFast EvaGreen Supermix on a CFX96 Real-Time PCR Detection System (Bio-Rad, Richmond, CA, USA). The following primers were used: PKD1, forward primer 5'-CGCACATCATCTGCTGAAC-3' and reverse primer 5'-CTTTCGGTGCA

CAACGTTTA-3'; FRS2, forward primer 5'-ATGG GAATGAGTTAGGTTCTGGC-3' and reverse primer 5'-GCGGGTGTATAAAATCAGTTCTGTG-3'. Data were normalized automatically by using GAPDH as the loading control, with the following primers: forward primer 5'-GCAAATTCATGGCACCGT-3' and reverse primer 5'-TCGCCCCACTTGATTTTGG-3'.

Statistical analysis

All statistical analyses were carried out using GraphPad Prism IV software. A *p* value of less than 0.05 was considered statistically significant (**p* < 0.05; ***p* < 0.01; ****p* < 0.001; *****p* < 0.0001).

ACKNOWLEDGMENTS

We would like to thank Pat Fawcett, Ravi Patel, and Dr. Karthik Giridhar for technical assistance on this project.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

GRANT SUPPORT

This study was supported in part by the National Institutes of Health grant R01CA129127 (Wang), R01CA142580 (Wang), Department of Defense award PC150190, and Oversea Hong Kong & Macao Scholars Collaborative Research Fund of NSFC in China (Grant# 81328020, Deng and Wang).

REFERENCES

1. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S. The protein kinase complement of the human genome. *Science*. 2002; 298:1912–1934.
2. Wang QJ. PKD at the crossroads of DAG and PKC signaling. *Trends Pharmacol Sci*. 2006; 27:317–323.
3. Johannes FJ, Prestle J, Eis S, Oberhagemann P, Pfizenmaier K. PKC α is a novel, atypical member of the protein kinase C family. *J Biol Chem*. 1994; 269:6140–6148.
4. Valverde AM, Sinnett-Smith J, Van Lint J, Rozengurt E. Molecular cloning and characterization of protein kinase D: a target for diacylglycerol and phorbol esters with a distinctive catalytic domain. *Proc Natl Acad Sci USA*. 1994; 91:8572–8576.
5. Sturany S, Van Lint J, Muller F, Wilda M, Hameister H, Hocker M, Brey A, Gern U, Vandenheede J, Gress T, Adler G, Seufferlein T. Molecular cloning and characterization of the human protein kinase D2. A novel member of the protein kinase D family of serine threonine kinases. *J Biol Chem*. 2001; 276:3310–3318.

6. Hayashi A, Seki N, Hattori A, Kozuma S, Saito T. PKC η , a new member of the protein kinase C family, composes a fourth subfamily with PKC μ . *Biochim Biophys Acta*. 1999; 1450:99–106.
7. Waldron RT, Iglesias T, Rozengurt E. Phosphorylation-dependent protein kinase D activation. *Electrophoresis*. 1999; 20:382–390.
8. Waldron RT, Rozengurt E. Protein kinase C phosphorylates protein kinase D activation loop Ser744 and Ser748 and releases autoinhibition by the pleckstrin homology domain. *J Biol Chem*. 2003; 278:154–163.
9. Zugaza JL, Sinnett-Smith J, Van Lint J, Rozengurt E. Protein kinase D (PKD) activation in intact cells through a protein kinase C-dependent signal transduction pathway. *Embo J*. 1996; 15:6220–6230.
10. Jacamo R, Sinnett-Smith J, Rey O, Waldron RT, Rozengurt E. Sequential protein kinase C (PKC)-dependent and PKC-independent protein kinase D catalytic activation via Gq-coupled receptors: differential regulation of activation loop Ser (744) and Ser (748) phosphorylation. *J Biol Chem*. 2008; 283:12877–12887.
11. Sinnett-Smith J, Jacamo R, Kui R, Wang YM, Young SH, Rey O, Waldron RT, Rozengurt E. Protein kinase D mediates mitogenic signaling by Gq-coupled receptors through protein kinase C-independent regulation of activation loop Ser744 and Ser748 phosphorylation. *J Biol Chem*. 2009; 284:13434–13445.
12. Lavalley CR, George KM, Sharlow ER, Lazo JS, Wipf P, Wang QJ. Protein kinase D as a potential new target for cancer therapy. *Biochim Biophys Acta*. 2010; 1806:183–192.
13. Rozengurt E. Protein kinase D signaling: multiple biological functions in health and disease. *Physiology (Bethesda)*. 2011; 26:23–33.
14. Weinreb I, Piscuoglio S, Martelotto LG, Waggott D, Ng CK, Perez-Ordóñez B, Harding NJ, Alfaro J, Chu KC, Viale A, Fusco N, da Cruz Paula A, Marchio C, et al. Hotspot activating PRKD1 somatic mutations in polymorphous low-grade adenocarcinomas of the salivary glands. *Nature genetics*. 2014; 46:1166–1169.
15. Eiseler T, Doppler H, Yan IK, Goodison S, Storz P. Protein kinase D1 regulates matrix metalloproteinase expression and inhibits breast cancer cell invasion. *Breast Cancer Res*. 2009; 11:R13.
16. Kim M, Jang HR, Kim JH, Noh SM, Song KS, Cho JS, Jeong HY, Norman JC, Caswell PT, Kang GH, Kim SY, Yoo HS, Kim YS. Epigenetic inactivation of protein kinase D1 in gastric cancer and its role in gastric cancer cell migration and invasion. *Carcinogenesis*. 2008; 29:629–637.
17. Azoitei N, Kleger A, Schoo N, Thal DR, Brunner C, Pusapati GV, Filatova A, Genze F, Moller P, Acker T, Kuefer R, Van Lint J, Baust H, et al. Protein kinase D2 is a novel regulator of glioblastoma growth and tumor formation. *Neuro Oncol*. 2011; 13:710–724.
18. Ochi N, Tanasavimon S, Matsuo Y, Tong Z, Sung B, Aggarwal BB, Sinnett-Smith J, Rozengurt E, Guha S.

Protein kinase D1 promotes anchorage-independent growth, invasion, and angiogenesis by human pancreatic cancer cells. *J Cell Physiol.* 2011; 226:1074–1081.

19. Ristich VL, Bowman PH, Dodd ME, Bollag WB. Protein kinase D distribution in normal human epidermis, basal cell carcinoma and psoriasis. *Br J Dermatol.* 2006; 154:586–593.
20. Chen J, Deng F, Singh SV, Wang QJ. Protein kinase D3 (PKD3) contributes to prostate cancer cell growth and survival through a PKCepsilon/PKD3 pathway downstream of Akt and ERK 1/2. *Cancer Res.* 2008; 68:3844–3853.
21. Rennecke J, Rehberger PA, Furstemberger G, Johannes FJ, Stohr M, Marks F, Richter KH. Protein-kinase-Cmu expression correlates with enhanced keratinocyte proliferation in normal and neoplastic mouse epidermis and in cell culture. *Int J Cancer.* 1999; 80:98–103.
22. Karam M, Legay C, Auclair C, Ricort JM. Protein kinase D1 stimulates proliferation and enhances tumorigenesis of MCF-7 human breast cancer cells through a MEK/ERK-dependent signaling pathway. *Exp Cell Res.* 2012; 318:558–569.
23. Biswas MH, Du C, Zhang C, Straubhaar J, Languino LR, Balaji KC. Protein kinase D1 inhibits cell proliferation through matrix metalloproteinase-2 and matrix metalloproteinase-9 secretion in prostate cancer. *Cancer Res.* 2010; 70:2095–2104.
24. Sharlow ER, Giridhar KV, LaValle CR, Chen J, Leimgruber S, Barrett R, Bravo-Altamirano K, Wipf P, Lazo JS, Wang QJ. Potent and selective disruption of protein kinase D functionality by a benzoxoloazepinone. *J Biol Chem.* 2008; 283:33516–33526.
25. Chen J, Giridhar KV, Zhang L, Xu S, Wang QJ. A protein kinase C/protein kinase D pathway protects LNCaP prostate cancer cells from phorbol ester-induced apoptosis by promoting ERK1/2 and NF- κ B activities. *Carcinogenesis.* 2011; 32:1198–1206.
26. Lavalle CR, Bravo-Altamirano K, Giridhar KV, Chen J, Sharlow E, Lazo JS, Wipf P, Wang QJ. Novel protein kinase D inhibitors cause potent arrest in prostate cancer cell growth and motility. *BMC Chem Biol.* 2010; 10:5.
27. Tandon M, Johnson J, Li Z, Xu S, Wipf P, Wang QJ. New pyrazolopyrimidine inhibitors of protein kinase d as potent anticancer agents for prostate cancer cells. *PloS One.* 2013; 8:e75601.
28. Kwabi-Addo B, Ozen M, Ittmann M. The role of fibroblast growth factors and their receptors in prostate cancer. *Endocr Relat Cancer.* 2004; 11:709–724.
29. Turner N, Grose R. Fibroblast growth factor signalling: from development to cancer. *Nat Rev Cancer.* 2010; 10:116–129.
30. Gotoh N. Regulation of growth factor signaling by FRS2 family docking/scaffold adaptor proteins. *Cancer Sci.* 2008; 99:1319–1325.
31. Valencia T, Joseph A, Kachroo N, Darby S, Meakin S, Gnanapragasam VJ. Role and expression of FRS2 and FRS3 in prostate cancer. *BMC Cancer.* 2011; 11:484.
32. Zuck B, Goepfert C, Nedlin-Chittka A, Sohr K, Voigt KD, Knabbe C. Regulation of fibroblast growth factor-like protein(s) in the androgen-responsive human prostate carcinoma cell line LNCaP. *J Steroid Biochem Mol Biol.* 1992; 41:659–663.
33. Masiello D, Cheng S, Bublely GJ, Lu ML, Balk SP. Bicalutamide functions as an androgen receptor antagonist by assembly of a transcriptionally inactive receptor. *J Biol Chem.* 2002; 277:26321–26326.
34. Grosse A, Bartsch S, Baniahmad A. Androgen receptor-mediated gene repression. *Mol Cell Endocrinol.* 2012; 352:46–56.
35. Hendriksen PJ, Dits NF, Kokame K, Veldhoven A, van Weerden WM, Bangma CH, Trapman J, Jenster G. Evolution of the androgen receptor pathway during progression of prostate cancer. *Cancer Res.* 2006; 66:5012–5020.
36. Jia L, Berman BP, Jariwala U, Yan X, Cogan JP, Walters A, Chen T, Buchanan G, Frenkel B, Coetzee GA. Genomic androgen receptor-occupied regions with different functions, defined by histone acetylation, coregulators and transcriptional capacity. *PloS One.* 2008; 3:e3645.
37. Jariwala U, Prescott J, Jia L, Barski A, Pregizer S, Cogan JP, Arasheben A, Tilley WD, Scher HI, Gerald WL, Buchanan G, Coetzee GA, Frenkel B. Identification of novel androgen receptor target genes in prostate cancer. *Mol Cancer.* 2007; 6:39.
38. Rosini P, Bonaccorsi L, Baldi E, Chiasserini C, Forti G, De Chiara G, Lucibello M, Mongiat M, Iozzo RV, Garaci E, Cuzzolino F, Torcia MG. Androgen receptor expression induces FGF2, FGF-binding protein production, and FGF2 release in prostate carcinoma cells: role of FGF2 in growth, survival, and androgen receptor down-modulation. *Prostate.* 2002; 53:310–321.
39. Kassen AE, Sensibar JA, Sintich SM, Pruden SJ, Kozlowski JM, Lee C. Autocrine effect of DHT on FGF signaling and cell proliferation in LNCaP cells: role of heparin/heparan-degrading enzymes. *Prostate.* 2000; 44:124–132.
40. Trauzold A, Schmiedel S, Sipos B, Wermann H, Westphal S, Roder C, Klapper W, Arlt A, Lehnert L, Ungefroren H, Johannes FJ, Kalthoff H. PKCmu prevents CD95-mediated apoptosis and enhances proliferation in pancreatic tumour cells. *Oncogene.* 2003; 22:8939–8947.



Review

Protein kinase D signaling in cancer: A friend or foe?

Adhiraj Roy^a, Jing Ye^b, Fan Deng^c, Qiming Jane Wang^{a,*}^a Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, 200 Lothrop Street, Pittsburgh, PA 15261, USA^b Department of Anesthesiology, School of Basic Medical Sciences, Southern Medical University, Guangzhou, China^c Department of Cell Biology, School of Basic Medical Sciences, Southern Medical University, Guangzhou, China

ARTICLE INFO

Keywords:

PKD
Cancer
Structure
Regulation
Signaling mechanisms
Function

ABSTRACT

Protein kinase D is a family of evolutionarily conserved serine/threonine kinases that belongs to the Ca^{++} /Calmodulin-dependent kinase superfamily. Signal transduction pathways mediated by PKD can be triggered by a variety of stimuli including G protein-coupled receptor agonists, growth factors, hormones, and cellular stresses. The regulatory mechanisms and physiological roles of PKD have been well documented including cell proliferation, survival, migration, angiogenesis, regulation of gene expression, and protein/membrane trafficking. However, its precise roles in disease progression, especially in cancer, remain elusive. A plethora of studies documented the cell- and tissue-specific expressions and functions of PKD in various cancer-associated biological processes, while the causes of the differential effects of PKD have not been thoroughly investigated. In this review, we have discussed the structural-functional properties, activation mechanisms, signaling pathways and physiological functions of PKD in the context of human cancer. Additionally, we have provided a comprehensive review of the reported tumor promoting or tumor suppressive functions of PKD in several major cancer types and discussed the discrepancies that have been raised on PKD as a major regulator of malignant transformation.

1. Introduction

Protein kinases are of utmost importance in maintaining a battery of cellular activities and the human genome encodes over 500 protein kinase genes which constitute about 2% of all human genes, collectively named as the human kinome [1]. The protein kinase D (PKD) family of serine/threonine kinases falls into the Ca^{++} /Calmodulin-dependent protein kinases (CaMKs) superfamily and consists of three isoforms in mammals, notably, PKD1, PKD2 and PKD3. PKD1 was the first member identified in human and mouse in 1994 [2,3], although initially it was categorized as a member of the protein kinase C (PKC) family and named PKC μ [2,4]. It was later reclassified into the CaMK family based on sequence homology in the catalytic domain. PKD3 and PKD2, two additional PKD isoforms, were discovered thereafter [5,6].

PKDs are evolutionarily highly conserved and homologs are found in several organisms including mice (*Mus musculus*), rats (*Rattus norvegicus*), flies (*Drosophila melanogaster*) and yeast (*Saccharomyces cerevisiae*) [7]. There is also high sequence homology among PKD isoforms, although structural and functional differences have been noted. For example, PKD3 lacks PDZ (PSD-95/Discs large/ZO-1) binding motif [8] and a Src family kinase phosphorylation motif [9]. Among other

organisms, *D. melanogaster* possesses only one PKD gene [10], whereas, two orthologs termed *dfk-1* and *dfk-2* are present in *C. elegans* [11–13]. In a canonical pathway, various stimuli including hormones, phorbol esters, growth factors, cellular stress converge to the activation of PKDs through diacylglycerol (DAG) and classical or novel protein kinase C (c/nPKC) via active phospholipase C (PLC) β and γ [14,15]. Activated PKD resides in diverse subcellular locations such as cytosol, Golgi apparatus, nucleus, mitochondria to regulate a plethora of cellular functions, especially those related to malignant transformation including cell proliferation, growth, migration/invasion, apoptosis, epidermal-to-mesenchymal transition (EMT) [14,16,17].

In this review, we discuss the *status quo* of PKD isoforms in terms of their modulation of different physiological activities and mechanistic role in development and progression of human diseases focusing on cancer. In light of accumulating scientific evidence, we aim to provide an updated and comprehensive review of each of the PKD isoforms, their differential expression patterns, and how they communicate with the cellular machineries in a wide variety of cell and tissue types to coordinate its biological role in oncogenesis (Table 1). Furthermore, we discuss potential tumor promoting as well as suppressive properties of PKD in different cancer types and aim to resolve the prevailing

Abbreviations: PKC, protein kinase C; PKD, protein kinase D; PLC, phospholipase C; DAG, diacylglycerol; CRD, cysteine-rich domain; GPCR, G protein-coupled receptor; TNF, tumor necrosis factor; TGN, trans-Golgi network; EMT, epithelial-to-mesenchymal transition; MMP, matrix metalloproteinase; HDAC, histone deacetylase; BMP, bone morphogenic protein; AR, androgen receptor; RUNX, Runt-related transcription factors

* Corresponding author.

E-mail address: qjw1@pitt.edu (Q.J. Wang).

<http://dx.doi.org/10.1016/j.bbcan.2017.05.008>

Received 27 April 2017; Received in revised form 26 May 2017; Accepted 27 May 2017

Available online 31 May 2017

0304-419X/ © 2017 Elsevier B.V. All rights reserved.

Table 1
Summary of differential expression and physiological roles of PKD isoforms in major cancers.

Cancer type	PKD isoform	Function in tumorigenesis	Mode of action	References
Prostate	PKD1	Anti-tumorigenic	1. Downregulates AR. 2. Represses AR-dependent transcription of target genes by modulating Hsp27. 3. phosphorylates E-cadherin and regulates β -catenin activity.	[124] [126] [125]
		Pro-oncogenic	Wnt5 signaling pathway activation of Jun-N-terminal kinase.	[91]
		Pro-oncogenic	1. Modulates Akt and ERK1/2 activation. 2. Activates NF- κ B signaling, elevates urokinase-type plasminogen activator (uPA) and MMP-9 expression.	[128,190] [90]
Breast	PKD1	Anti-tumorigenic	1. Phosphorylates and inactivates SSH1L, interacts with cortactin and paxillin. 2. Suppression of MMP expression. 3. Inhibition of EMT via inactivation of Snail function and modulation of epithelial-to-mesenchymal markers	[81,83,137,141] [81] [139]
		Pro-oncogenic	Promotes cell-cell adhesion.	[142,143]
		Pro-oncogenic	1. Promotes Hsp27 and HDAC4/5/7 phosphorylation. 2. Upregulates P-glycoprotein (P-gp) expression, multidrug resistance.	[144] [145]
Pancreas	PKD1	Pro-oncogenic	1. Activates MEK1/2 initiated by neurotensin, increases DNA replication. 2. Inhibits CD95-mediated apoptosis, cell proliferation and survival. 3. Acts downstream of TGF α /Kras and mediates formation of ductal structures through activation of the Notch pathway.	[155,156] [153] [154]
		Pro-oncogenic	1. PKD2 activation and increase in Ser ⁸² phosphorylation of Hsp27 in PANC-1 cells stimulated by neurotensin. 2. Stimulates expression and secretion of MMP-7 and 9, induces tumor angiogenesis by releasing ECM-bound VEGF-A.	[157] [162]
Skin	PKD1	Pro-oncogenic	1. Modulates ERK/MAPK pathway and promotes proliferation. 2. Reduces UVB-induced apoptosis.	[166] [167]
		Anti-tumorigenic	Correlated with upregulation of CDK4/6 inhibitor p15 ^{INK4B} and induction of p53-independent G1 cell cycle arrest.	[169]
		Pro-oncogenic		[169]
Gastric	PKD1	Anti-tumorigenic	1. Epigenetically silenced.	[170]
		Pro-oncogenic	1. Promotes AKT and ERK signaling. 2. Inhibits apoptosis. 3. Suppresses NF- κ B signaling.	[177] [177] [177]

functional discrepancies it poses as a regulator of cancer.

2. PKD structural and functional relationships, phosphorylation, activation

All 3 members of the PKD family share discrete structural and functional similarities (Fig. 1). They possess an N-terminal regulatory domain which is subdivided into 2 tandem cysteine-rich Zn-finger like domains (CRD, C1a and C1b), a plekstrin homology (PH) domain and a C-terminal catalytic domain, which are shared by all three isoforms. However, PKD3 lacks C-terminal PDZ binding (PB) domain as it is present in both PKD1 and PKD2. The serine residues shown in the catalytic domain represent the conserved activation loop amino acids phosphorylated by the members of the c/nPKCs, which leads to PKD activation. Note: Phosphorylation sites are numbered based on the murine PKD isoforms.

C-terminal catalytic domain [14,16]. The regulatory domain plays a critical role by auto-inhibiting the kinase domain and maintaining the protein in an inactivated state and thus deletion or mutation of critical residues of the C1 domains results in constitutively active PKD [14]. The C1 domains bind DAG and phorbol esters, anchor PKD to membranes and modulate the localization of the protein to Golgi, nucleus as well as plasma membrane [14,18,19]. Although the C1 domains are active in PKD, there are intrinsic differences in their activity and selectivity for ligands. Our study has demonstrated isoform-specific




Fig. 1. Diagram of protein kinase D structure. The structure of PKD contains an N-terminal regulatory domain which consists of cysteine-rich Zn-finger like motifs (C1a and C1b), a plekstrin homology (PH) domain and a C-terminal catalytic domain, which are shared by all three isoforms. However, PKD3 lacks C-terminal PDZ binding (PB) domain as it is present in both PKD1 and PKD2. The serine residues shown in the catalytic domain represent the conserved activation loop amino acids phosphorylated by the members of the c/nPKCs, which leads to PKD activation. Note: Phosphorylation sites are numbered based on the murine PKD isoforms.

differences in the ligand binding activities of PKD isoforms with PKD3 being most sensitivity to DAG/phorbol esters [20]. As for individual C1 domains, we have shown that C1a is a high affinity receptor for DAG, while C1b is low affinity for DAG but high affinity for phorbol esters. In the context of full-length protein, C1a in several studies appears to play a more crucial role in DAG binding and membrane targeting of PKD [19–22]. Furthermore, the reduced DAG binding activity of C1b is attributed to a conserved lysine residue within this domain that impairs DAG binding [20,23]. Overall, C1 domain is central to the spatial and temporal regulation of PKD localization at different subcellular locations. In addition to the intrinsic differences of the twin C1 domains [20,24,25], their ligand binding activity, selectivity, and accessibility to ligand in a holoenzyme are intricately regulated by PKD phosphorylation [22], kinase activity [19,21,22] and interactions with protein binding partners [26], and this regulation becomes more complex with embedded nuclear localization and export signals within the structure of C1 domain [27]. Whether, how, and how much they contribute to the differential biological functions of PKD isoforms remain to be determined.

The PH domain has been shown to interact with other proteins and play a role in subcellular localization as well as nuclear export of PKD [14]. The type 1 PDZ (PSD-95/Discs large/ZO-1) binding motif is found at the C-terminus of PKD1 and PKD2 which is responsible for interaction with protein substrates such as NHERF-1 and Kidins220 [8,28]. The substrate recognition motif of PKD1 (L.X.R. (Q/K/E/M).S.X.X.X.X) displays a unique preference of leucine at the –5 position [16,29,30]. There are minor structural differences in PKD2 and PKD3. For example, PKD2 contains a serine-rich region between C1a and C1b motifs and PKD3 lacks any C-terminal PDZ domain [14]. PKD can be activated by a variety of physiological factors, such as bioactive peptides [31], lipids [32], growth factors [33], tumor necrosis factor (TNF) [34], chemokines, and many of which act through binding and activating the G protein-coupled receptors (GPCR) or receptor tyrosine kinase (RTK) and further activating phospholipase Cs (PLCs) and c/nPKCs. c/nPKCs phosphorylate the conserved serine residues in the activation loop of PKDs (for example, Ser⁷⁴⁴ and Ser⁷⁴⁸ of murine PKD1, equivalent to Ser⁷³⁸ and Ser⁷⁴² of human PKD1), leading to PKD activation [14,15]. Mutation of active site serine residues in PKD (PKD1 S744A/S748A) abolishes PKD activation [16,17]. Replacement of both serine residues with glutamic acid (PKD1 S744E/S748E) results in a constitutively active PKD implying that activation loop phosphorylation is an essential mechanism for PKD activation [14]. In further analysis of the canonical PKC-dependent activation of PKD pathway governing the functional facets of PKD in cellular physiology, emerging evidence suggests that different regulatory mechanisms control the phosphorylation at the two sites in the activation loop. For instance, Gq-coupled receptor agonists such as bombesin induce biphasic PKD activation, notably, a first rapid PKC-dependent activation through phosphorylation of PKD1 at Ser⁷⁴⁴, followed by a second PKC-independent autophosphorylation at Ser⁷⁴⁸ to sustain PKD activity [35,36]. Several lines of evidence emerged for supporting the fact that PKD1 autophosphorylation at Ser⁷⁴⁸ is the major mechanism for late sustained PKD activation in cells treated with GPCR agonists [35]. Beyond the activation loop phosphorylation, there are considerable discrepancies in the understanding of an autophosphorylation site, PKD1 Ser⁹¹⁶ (equivalent to human PKD1 Ser⁹¹⁰), which has been used as a marker of PKD1 activity status in many studies [37]. The lines of evidence supporting its use as a marker for PKD activation are: 1) PKD1 Ser⁹¹⁶ phosphorylation increases when PKD1 is activated by growth factor receptors or phorbol esters, 2) constitutively active PKD1^{S744E, S748E} exhibits high levels of endogenous Ser⁹¹⁶ phosphorylation [37,38]. It was conceived that activation of PKD through phosphorylation of Ser⁷⁴⁴ and Ser⁷⁴⁸ residues is often followed by autophosphorylation of Ser⁹¹⁶ [14,37,39,40]. However, other evidences support the contrary: 1) PKD1 activation loop phosphorylation and increase in catalytic activity by agonist stimulation do not augment Ser⁹¹⁰ phosphorylation [41–43], 2)

A catalytically inactive PKD1 mutant (PKD1^{K612W}) displays Ser⁹¹⁰ phosphorylation by endogenous PKD1 and other enzymes [8,44], 3) Ser⁹¹⁰ autophosphorylation can be achieved at exceedingly low concentration of ATP that does not require PKD1 phosphorylation of Ser⁷³⁸,⁷⁴². Additionally, the phosphorylation status on Ser⁹¹⁰ does not correlate well with the inhibition of PKD by certain inhibitors, for example, Ser⁹¹⁰ phosphorylation is resistant to Gö6976, an ATP competitive inhibitor [44] and another ATP competitive inhibitor, BPKDi that inhibits HDAC5 phosphorylation, does not inhibit PKD1 Ser⁹¹⁰ autophosphorylation [45]. Further studies are necessary to define the precise role of PKD1 Ser⁹¹⁰ autophosphorylation and its implication in the regulation of PKD1 catalytic activity.

3. Protein kinase D and somatic mutations in cancer

Benign cells acquire somatic mutations which cause dysregulation of cell proliferation, migration and invasion, key phenomena for oncogenesis. Approaches through comprehensive genomics analysis have provided valuable cues to somatic aberrations that define individual cancers [46–48]. *PRKD* is generally thought to exhibit low frequency of somatic mutations in pan-cancer analysis. However, in several recent reports, high frequency somatic mutations in *PRKD* genes have been reported in at least two rare tumors, Polymorphous low-grade adenocarcinoma (PLGA) and angiolipomas [49–51].

Polymorphous low-grade adenocarcinoma (PLGA) is an intra-oral salivary gland malignancy which preferentially affects the minor salivary glands. Weinreb et al. subjected three consecutive PLGAs to massive parallel RNA sequencing and whole exome sequencing (WES) and identified two somatic heterozygous single-nucleotide variations (SNV), which are c.2130A > T and c.2130A > C, affecting highly conserved E710 amino acid at the catalytic loop, resulting in a mutant PKD1 (PKD1 E710D) [51]. To further validate the results, the authors analyzed 53 PLGAs by Sanger sequencing and targeted amplicon sequencing of *PRKD1* exon 15 and confirmed the presence of somatic c.2130A > T and c.2130A > C mutations in 41.5% and 30.2% of PLGAs, respectively and the total mutation frequency of 72.9%, representing a single high-frequency hotspot mutation that is indicative of a driver oncogene. Homology modelling of PKD1 suggested that this E710D mutation could alter coordination with Mg²⁺ ion, affect enzyme kinetics as well as interfere with ADP-binding [51]. Cell-free *in vitro* kinase assay showed significantly increased transphosphorylation of the substrate CREBtide by PKD1^{E710D} mutant protein and its elevated catalytic activity as compared to the wild type PKD1 [51]. Further analysis indicated that the expression of PKD1^{E710D} mutant protein in embryonic kidney epithelial and non-malignant breast epithelial cells caused increased phosphorylation of Ser⁷³⁸/Ser⁷⁴² and Ser⁹¹⁰ of PKD1 as compared to the wild type PKD1. It was also demonstrated that forced expression of PKD1^{E710D} mutant protein in MCF10A and MCF12A breast cells changed the hollow spheroid, acinar-like structures into large, coalescent structures filled with lumens and irregular contours, an increased invasive phenotype typically associated with the overexpression of other oncogenes in this model system [52,53]. Taken together, these results demonstrate that the somatic mutation in PKD1 encoding PKD1^{E710D} is likely activating driver of PLGA and confers a neoplastic advantage to the epithelial cells [51]. There was another report which aimed to identify somatic mutations in PLGA that affect the kinase domains of *PRKD2* or *PRKD3* gene and act as a driver of neoplasia [50]. This study found PLGAs that lack *PRKD1* somatic mutations or PKD gene family rearrangement; do not harbor somatic mutations in the kinase domains of *PRKD2* or *PRKD3* genes. These findings appear to bring up an interesting concept that PKD1 is not functionally equivalent to PKD2 and PKD3 in tumorigenesis. There is a lack of evidence for the somatic mutations and their effect on the biology of adipocytic tumors, including angiolipoma as no genetic aberrations or chromosomal rearrangements/deregulations have been reported [54]. In a recent report, Hofvander et al. analyzed a cohort of




Fig. 2. Major signaling pathways and biological functions of PKD. Several extracellular stimuli activate phospholipase C (PLC) which catalyzes the formation of diacylglycerol (DAG). DAG recruits PKD and PKC to the plasma membrane inducing the activation of PKC which then further phosphorylates PKD at two serine (Ser^{744,748}) residues resulting in the activation of PKD. PKD can also be activated by PKC-independent pathways and the activity of PKD can be sustained through autophosphorylation at Ser⁷⁴⁸ residue (dashed line). Activated PKD regulates an array of cancer-associated functions including cell proliferation, migration, survival, regulation of gene transcription, protein/vesicle trafficking and secretion through several major signaling pathways.

benign adipocytic tumors including conventional lipoma, hibernoma and angiolipoma by WES and ultra-deep sequencing and demonstrated the presence of somatic mutations (18 out of 21) in the catalytic domain of PKD2 [49]. qRT-PCR confirmed that the level of *PRKD2* but not *PRKD1* or *PRKD3* was higher in angiolipoma than in lipomas. The authors argued that the finding of *PRKD2* mutations in 80% of the tumors strongly correlate with the neoplastic origin of angiolipoma. Further studies are necessary to evaluate the significance of the mutations in the catalytic domain of *PRKD2* gene. Collectively, these studies not only highlight the significance of PKD family in oncogenesis, but also reveal its potential utility as molecular diagnostic markers or therapeutic targets for certain tumors.

4. PKD regulates major biological processes that contribute to development and progression of cancer

PKD contributes to a broad spectrum of cellular processes including cell survival, proliferation, EMT, angiogenesis, gene transcription, secretion and vesicle transport through TGN and innate immunity (Fig. 2). Once activated, PKD rapidly localizes to different subcellular locations including plasma membrane, nucleus and mitochondria. In this section we discuss several major biological events and pathways regulated by PKD that directly contribute to development and progression of cancer.

4.1. Cell growth and proliferation

A major characteristic of cancer cells is uncontrolled cellular growth and proliferation. In this section, we will review the roles of PKD in modulating the biological pathways that control cell growth, proliferation and survival in context of neoplasia.

PKD is activated by many mitogenic GPCR agonists that mediate their response through G_q, G_i and G₁₂ implying a role of PKD in cell proliferation [55–58]. G_q-coupled receptor agonists including bombesin and vasopressin-induced activation of PKD1 and PKD2 and subsequent increased DNA replication and cell proliferation has been shown in Swiss 3 T3 fibroblast [36,59–61]. Extracellular-regulated protein kinase (ERK) pathway is a major mediator of GPCR agonists-induced mitogenic effects [62]. PKD contributes to the duration and intensity of MEK/ERK/RSK activation in GPCR agonist-stimulated cells leading to the induction of c-Fos that stimulates cell cycle progression [36,59].

4.2. Cell survival

Oxidative stress can be accounted as an imbalance between the systemic production of reactive oxygen species and a cell's ability to readily detoxify the reactive intermediates or to repair the resulting damage. Abnormalities in the normal redox state of cells can cause toxic effects through the production of reactive oxygen species (ROS), such as

O₂^{•−} (superoxide radical), OH (hydroxyl radical) and H₂O₂ (hydrogen peroxide). Oxidative stress can activate PKD via nonreceptor tyrosine kinase c-Abl and Src along with PKCδ. Reactive oxygen species (ROS) produces DAG via PLD1 and phosphatidic acid phosphatase (PAP)-mediated catalysis and further recruitment of PKD and PKCδ at the outer mitochondrial membrane [63]. A colocalized c-Abl phosphorylates Tyr⁴⁶³ in the PH domain of PKD1 which causes a conformational change, allowing the YGLY domain to be released from PH domain-mediated intramolecular autoinhibition [9]. Src-dependent phosphorylation of Tyr⁹⁵ creates a priming site for the C1b domain of PKCδ. PKCδ efficiently activates PKD1 by phosphorylating Ser⁷⁴⁶/Ser⁷⁴⁸. Activated PKD in turn activates IKKα-IKKβ-Nemo complex and nuclear import of NF-κB which results in induction of antiapoptotic and, or antioxidant genes such as manganese superoxide dismutase (MnSOD) and promotes cell survival [64]. Hence, a complete understanding of the molecular mechanism underlying ROS-induced PKD-mediated cell survival and identification of PKD substrates that activate NF-κB signaling pathway will help us better decipher the mechanism behind cancer cell survival.

4.3. Epithelial-to-mesenchymal transition (EMT)

Epithelial-to-mesenchymal transition (EMT) is a prominent neoplastic characteristics, where cell-cell adhesion is disrupted leading to enhanced cell motility and invasiveness [65]. Epithelial cells undergoing EMT process show mesenchymal cell properties including expression of mesenchymal markers and ability to migrate and invade [66]. E-cadherin is a master regulator of EMT process where it binds to β-catenin to form a protein complex and maintains cell-cell adhesion by interacting with actin and microtubule cytoskeleton owing to its anti-proliferative, antimetastatic and anti-invasion properties [67,68]. Different mechanisms of E-cadherin repression in malignant tumor have been shown including mutation, transcriptional repression, epigenetic silencing [68] and many transcription factors are involved in repression of E-cadherin and induction of EMT such as Snail and Twist [4,69–71]. Regulation of Snail protein is achieved by multiple pathways. The NF-κB signaling pathway positively modulates and stabilizes Snail protein promoting cell migration [72], whereas, phosphorylation of Snail by p21-activated kinase 1 (PAK1) [73] and GSK3β [74,75] increase Snail repression activity. Snail represses E-cadherin expression in prostate and breast cancer. PKD1 phosphorylates Snail at Ser¹¹ residue triggering nuclear exclusion of Snail by 14–3–3 adaptor protein. As a result, Snail-repressed genes are de-repressed and cell migration is suppressed via production of E-cadherin and other proteins that mediate cell-cell adhesion. Hence, PKD negatively regulates the function of Snail and inhibits EMT.

4.4. Cellular motility, migration and invasion

Cellular motility and invasion are coupled to remodeling of actin cytoskeleton [76] and degradation of extracellular matrix (ECM) [77,78]. Cellular movement is achieved when Cofilin slices actin filaments at the leading edge of motile cells, generating a supply of actin monomers and orchestrating the formation of WAVE-2-cortactin-ARP2/3 complex which ultimately creates an expanded, branched network of F-actin [76]. LIM kinase suppresses cellular migration by phosphorylating Ser³ residue of cofilin [79] and motility is restored when a protein phosphatase slingshot 1 like (SSH1L) dephosphorylates cofilin [80]. The original work by the Storz group showed that PKD phosphorylates SSH1L by complexing with SSH1L and F-actin at lamellipodium [81,82]. Phosphorylated SSH1L is sequestered to cytosol by 14–3–3 adaptor protein [83], consequently, pSer³-cofilin concentration rises and cell migration is inhibited. A follow-up study by the Storz group also reported PKD2/PKD3-mediated regulation of SSH1L and p21-activated kinase 4 (PAK4) resulting in phosphorylation of cofilin and decrease in cell migration [84]. PKD inhibits cytoskeleton

remodeling by phosphorylating the Ras effector RIN1 [85]. Phosphorylated RIN1 activates tyrosine kinase c-Abl and the RIN1-c-Abl complex phosphorylates and brings conformational change in a scaffold protein CRK that recruits F-actin remodeling proteins [85,86], resulting in suppression of cell motility. Matrix metalloproteinases (MMPs) are a class of proteases (collagenases, e.g. MMP-1; gelatinases, e.g. MMP-2) that mediate cell migration through ECM degradation. It has been reported that PKD inhibits breast cancer cell invasion by negatively regulating the transcription of several MMPs including MMP-2, MMP-7, MMP-9, MMP-10, MMP-11, MMP-13, MMP-14 and MMP-15 [81]. Histone deacetylases (HDACs) have been shown to regulate MMP expression [87,88] and PKD1 is shown to be a negative regulator of HDACs [89]. Therefore, it is conceivable that PKD may negatively regulate MMPs via HDAC regulation. PKD has also been reported to be a positive regulator of cell migration. PKD2 and PKD3 have been shown to increase prostate cancer cell invasion and migration by promoting NF-κB and urokinase-type plasminogen activator (uPA) expression/activation [90]. Yamamoto et al. has shown that Wnt5a-JNK-PKD1 axis positively regulates cell proliferation and migration of prostate cancer [91]. It has also been shown that vascular endothelial growth factors (VEGFs) promote cell proliferation and migration via PKD-mediated phosphorylation of class IIa HDACs and subsequent expression of VEGF-responsive genes [92,93]. In summary, several PKD-regulated pathways converge on the promotion as well as inhibition of cell proliferation, invasion and migration.

4.5. Angiogenesis

Angiogenesis is a process where new blood capillaries are formed and it is essential for many physiological processes such as embryonic development, wound healing and many pathological processes including tumorigenesis [94]. VEGFs are prominent in angiogenesis [95,96]. There are two related receptor tyrosine kinases that bind VEGF, VEGFR-1 and VEGFR-2 [97,98] and induce downstream signal that activates a variety of proteins such as PLC-γ, PI3-Kinase and the Src family [99,100].

As one of the best characterized angiogenic factors, VEGF exerts different biological functions in endothelial cells, such as: 1) stimulation of endothelial cell proliferation and migration [96], 2) promotion of endothelial cell survival by inducing expression of anti-apoptotic proteins such as Bcl-2 and death antagonist A1 [101–105]. PKD1 has been shown to be activated downstream of VEGFR2-PLCγ-PKC to activate ERK1/2 pathway and stimulate endothelial cell proliferation [33]. Hao et al. found that PKD2 was a major PKD isoform that mediates endothelial cell proliferation and migration [106]. In mouse embryonic stem cells, PKD2 activity is required for angiogenesis [107]. Mechanistically, PKD-phosphorylation of class IIa HDACs enable them to be sequestered to the cytoplasm by 14–3–3 proteins leading to derepression of target genes. VEGF-stimulated, PKD-mediated phosphorylation at Ser^{259/498} and concomitant nuclear export of HDAC5 induces MEF2-dependent genes and endothelial cell migration [89]. HDAC7 is involved in regulating endothelial cell morphology and migration [92]. VEGF induces PKD-mediated phosphorylation of HDAC7 at Ser^{178/344/479} and its subsequent nuclear exclusion by 14–3–3 proteins. This results in the expression of VEGF-responsive genes and promotion of endothelial cell proliferation and migration, MMP expression and EMT [89,92]. Therefore, it can be concluded that PKD plays a key role in signaling pathways that regulate angiogenesis in endothelial cells.

4.6. Bone development and innate immunity

Of note, it is noteworthy to mention that PKD is also involved in bone formation and innate immunity. Bone morphogenic proteins (BMPs), a family of multifunctional growth factors belonging to the transforming growth factor β (TGFβ) superfamily maintain skeletal integrity by modulating signaling pathways that converge on Runt-

related transcription factors (RUNX), regulator of osteoblast gene transcription. It has been shown that BMP-2 induces PKD activation via PKC-independent pathway during osteoblast lineage progression [108] and PKD activation is required for the BMP-2 mediated osteoblast differentiation [42]. Our recent report using conditional PKD1-knockout mice model has shown that PKD1 positively regulates bone development and osteoblast differentiation which could be linked to the activity of the STAT3/p38 MAPK signaling pathway [109]. Cancer mortality and morbidity are mainly caused by metastasis and bone is the 3rd most common site of metastasis. Given the important role of PKD in bone homeostasis, it is conceivable that PKD may play an important role in bone metastasis of malignant tumors [110]. PKD has been shown to be involved in innate immunity in many different ways, such as: 1) functional regulator of T and B lymphocyte [14], 2) regulator of class IIa HDACs in lymphocytes [111–113], 3) modulator of β 1 integrin activity in T lymphocytes [114], 4) regulator of IL-2 via TCR stimulation [115], 5) downstream target in Toll-like receptor 9 (TLR9) signaling in macrophages [116] and TLR2 in mouse bone marrow-derived mast cells [117], 7) promoter of cell proliferation in chronic myelogenous leukemia (CML) [118] and 8) regulator of neutrophil chemotaxis [119].

5. PKD: a friend or foe in cancer development and progression?

Among other hallmarks such as sustaining proliferative signaling, enabling replicative immortality and evading cell death by apoptosis are of utmost significance in cancer development [120]. Accumulating evidence shows prominent link between tumor development and diverse signal transduction pathways that are modulated by PKD. The precise role of PKD in tumor progression remains elusive as evidence suggests that PKD plays a critical role as both the potent tumor promoter and suppressor of tumor development. In this section, we discuss the role of PKD isoforms in cancer development and progression focusing on several major cancer types and attempt to discuss the prevailing discrepancies that are associated with these tumor-specific studies (Fig. 3).

5.1. PKD in prostate cancer

Prostate cancer is one of the most common malignancies in male and accounts for 13% of cancer-related deaths in the USA [121]. Although early diagnosis and screening methods have advanced, effective

treatments of late-stage metastasized tumors are scarce [122]. Although elevated levels of PKD is observed in human prostate carcinoma tissue compared to normal prostate epithelium, differential expression and distribution of PKD isoforms have been reported and functional analysis of the PKD isoforms has revealed their different roles in prostate cancer progression.

PKD1 is highly expressed in cultured androgen-sensitive, less metastatic LNCaP cell line; whereas, in contrast, PC3 and DU145, two androgen-insensitive, highly metastatic prostate cancer cell lines fail to express PKD1 but found to have high levels of PKD2 and significant PKD3 expression [123]. PKD1 has been reported to be a negative regulator of cell proliferation of prostate cancer cells [124–126]. It has been shown that PKD1 once overexpressed blocks migration through E-cadherin phosphorylation and regulation of β -catenin activity [125–127]. PKD1 has also been shown to downregulate androgen receptor (AR) function in prostate cancer cells [124]. This study demonstrated that PKD1 physically associates with AR through its kinase domain and makes a transcriptional complex. The binding of PKD1 with AR initiates transcription of target genes and this PKD1-AR interaction might be a novel signaling mechanism responsible for prostate cancer progression. In contrast to these studies showing a tumor-suppressive function of PKD1, several lines of independent studies have shown that PKD in fact can serve as an oncogenic factor in prostate cancer [91]. Its role as an enhancer of cell migration and invasion has been linked to Wnt5a overexpression. This study has demonstrated that Wnt5a activates Jun-N-terminal kinase via PKD and PKD is required for Wnt5a-mediated induction of MMP-1 expression, cell migration and invasion.

PKD3, on the other hand, has been shown to promote cell proliferation and survival of prostate cancer by increasing prolonged activation of Akt and ERK1/2 [123,128]. It is known that hyperactive Akt in PTEN-null prostate cancer has been linked to angiogenesis, invasion and metastasis [129,130]. Moreover, our study has reported that PKD2 and PKD3 positively regulate prostate cancer cell invasion by upregulating NF- κ B signaling and HDAC1-mediated urokinase-type plasminogen activator (uPA) expression [90] and PKD2 and PKD3 were responsible for increased matrix metalloproteinase-9 (MMP-9) expression, a key player for EMT [128]. Additionally, our previous study has demonstrated that PKD3 contributes to the growth and survival of prostate cancer cells through PKC ϵ /PKD3 pathway and downstream Akt and ERK-1/2 signaling pathways [123]. Activated PKD1 and PKD2 were shown to protect LNCaP prostate cancer cells from phorbol ester PMA-induced apoptosis by promoting downstream ERK-1/2 and NF- κ B activities [131]. Using stable inducible PKD3 knockdown prostate cancer cell lines we have demonstrated that knockdown of PKD3 inhibits secretion of multiple key tumor-promoting factors including MMP-9, IL-6, IL-8, and GRO α and inducible depletion of PKD3 in a subcutaneous xenograft model suppressed tumor growth and decreased levels of intratumoral GRO α in mice. Furthermore, we have shown that androgen represses PKD expression in androgen-sensitive prostate cancer cells in an androgen receptor (AR)-dependent manner and the response is mediated by fibroblast growth factor receptor substrate 2 (FRS2) [132] and thus, we envision that upregulation of PKD as a result of loss or inhibition of AR may promote prostate cancer tumor cell survival.

In support of our view, using several classes of structurally distinct small molecule inhibitors of PKD discovered by our group, including CID755673 and its analogs kb-NB142-70, SD-208, 1-naphthyl PP1 (1-NA-PP1), Compound 139 [128,133–136], which are all nanomolar cell-active pan-PKD inhibitors, we demonstrated that targeted inhibition of PKD by these inhibitors led to reduced proliferation, migration and invasion of prostate cancer cells. Altogether, these studies have validated PKD as a potential therapeutic target for prostate cancer. In the future, detailed investigation should be carried out to decipher the precise role of each PKD isoform at different stages of prostate cancer development, and a PKD genetically engineered mouse model of prostate cancer will further validate the functional role of PKD in prostate carcinogenesis and tumor progression.




Fig. 3. Tumor-specific roles of PKD isoforms in several major cancer types. Benign cells transform into neoplasm such as carcinoma of prostate, breast, pancreas, skin and gastric. Three isoforms of PKD, namely PKD1, PKD2 and PKD3 either promote (solid arrow) or inhibit (bar-headed solid line) cancer progression in highly tumor-specific manner.

5.2. PKD in breast cancer

Breast cancer is the leading cause of cancer death in women and it is expected to account for 29% of all new cancer diagnoses in the USA by 2017 [121]. PKD was first linked to breast cancer in a study by Bowden et al. demonstrating the association of PKD1 in a complex with cortactin and paxillin in invadopodia at sites of extracellular matrix degradation [137]. Later studies have shown that PKD1 expression, but not PKD2 or PKD3, is reduced in over 95% of invasive breast cancer samples compared to benign breast tissue [81] and thus established a tumor-suppressive role of PKD1 in breast cancer. Eiseler et al. have shown that the loss of PKD1 in breast cancer is associated with higher degree of tumor invasiveness [81]. Highly aggressive breast cancer cell lines such as MDA-MB-231 and BT-20 do not express PKD1, whereas; less invasive MCF-7 and normal mammary epithelial cells such as MCF-10A show significant PKD1 expression. It has been demonstrated that hypermethylation of the *PRKD1* promoter causes loss of PKD1 expression in invasive breast cancer [138]. In MCF-7 cell, silencing of PKD1 enhanced its migration, whereas; overexpression of constitutively active form of PKD1 in MDA-MB-231 cells led to decreased cell invasion [81]. Multicellular spheroid/3D cell culture assay using MDA-MB-231 cells demonstrated that expression of active PKD1 inhibited cell invasion compared to normal breast epithelial cells [81]. At molecular level, PKD confers its role as tumor-suppressor in breast cancer by 1) suppressing the expression of many matrix metalloproteases (MMP) such as MMP-2, MMP-7, MMP-9, MMP-10, MMP-11, MMP-13 and MMP-14 [81], 2) inhibiting EMT via inactivation of Snail function by phosphorylating Ser¹¹ residue [139,140], 3) inducing the expression of epithelial-to-mesenchymal markers (vimentin, E-cadherin) [139] and 4) phosphorylating SSH1L at Ser⁹³⁷ and Ser⁹⁷⁸ residues and negatively regulating cofilin phosphorylation and interacting with cortactin and paxillin at lamellipodia, thereby inhibits cell motility and invasion [81,83,137,141]. In contrast, opposing effects of PKD has also been demonstrated in breast cancer. Kennett et al. and Palmantier et al. have demonstrated a tumor-promoting function of PKD1 in breast cancer cells [142,143].

Although accumulating evidence indicate PKD1 as a gross tumor-suppressor and a major contributor to the maintenance of epithelial phenotype in breast cancer [81,139–141], PKD2 and PKD3 have been assigned roles of tumor-promoters where they induce cell proliferation, invasiveness and chemoresistance [144–146]. Huck et al. have demonstrated that PKD3 is a tumor-promoter in triple-negative breast cancer (TNBC) [147]. PKD3 triggered the activation of S6 kinase 1 (S6K1) which is the main downstream target of the mammalian target of rapamycin complex 1 (mTORC1). The authors have also shown that PKD3 depletion inhibited cell proliferation of TNBC and hence, identifying PKD3 as a potent chemotherapeutic target. Borges et al. have demonstrated that PKD3 is highly upregulated in estrogen receptor (ER)-negative (ER[−]) invasive ductal carcinoma (IDC) which is associated with triple-negative phenotype [148]. This study showed that ER directly binds to the *PRKD3* gene promoter and inhibits PKD3 expression. Hence, loss of ER leads to upregulation of PKD3 which eventually induce increased cell proliferation, migration and invasion. Hao et al. have shown that silencing PKD2 or PKD3 significantly inhibited proliferation of HCC1806 triple-negative breast cancer cell line and PKD3 knockdown inhibited Hsp27 and HDAC4/5/7 phosphorylation [144]. Further investigation into different PKD isoforms at cellular levels as well as using mouse genetic models will help us to better understand the biology of this family of protein kinases in the development of breast neoplasia.

5.3. PKD in pancreatic cancer

Ductal adenocarcinoma of pancreas is a very aggressive, chemotherapy-resistant type of cancer [149] and according to the American Cancer Society, pancreatic cancer will account for about 3% of all

cancers in the US and about 7% of all cancer deaths by 2017. Almost 95% of all pancreatic ductal adenocarcinoma (PDAC) display somatic activating mutations of Kras [150] and increased epidermal growth factor (EGFR) signaling [151,152]. As compared to the normal pancreas, human ductal adenocarcinoma of the pancreas shows increased PKD1 expression and kinase activity [39,153]. An important study from the Storz group has demonstrated a positive role of PKD1 in the malignant transformation of PDAC. The authors have demonstrated that in a 3D explant model, using PKD inhibitors or PKD knockdown approach that PKD1 is necessary for TGF α - and Kras-mediated formation of duct-like structures originating from acinar cells, a process called acinar-to-ductal metaplasia (ADM), which converts to pancreatic intraepithelial neoplasia (PanIN), the premalignant neoplastic precursor of PDAC. Using *in vivo* mouse model (p48^{cre} Kras^{G12D} mice), the authors have knocked out PKD1 in acinar cells and demonstrated decreased progression of acinar-to-ductal metaplasia (ADM) to PanIN [154]. This study provided strong support for a role of PKD1 in the pathogenesis of PDAC. Additionally, other studies at cellular level support pro-proliferative and pro-survival effects of PKD1 in pancreatic cancer cells. Specifically, Trauzold et al. have shown that overexpression of PKD1 in pancreatic cancer cell decreased CD95-mediated apoptosis, increased cell proliferation rate and upregulated survivin [153]. PKD1 mediates the mitogenic effects of neurotensin in pancreatic cancer cells [155,156]. A study by Yuan et al. has shown that neurotensin increases Hsp27 phosphorylation of Ser⁸² via p38 MAPK-PKD2 signaling axis in pancreatic cancer cells [157]. Hsp27 level is markedly increased in many cancers and its elevated expression contributes to increased tumorigenicity and chemoresistance [158–161]. Another recent study has demonstrated a pro-oncogenic role of PKD2 in pancreatic cancer where PKD2 functions upstream of MMP-7 and -9 in pancreatic cancer cells and induces invasion and angiogenesis *in vivo* and *in vitro* [162]. PKD2 does so by stimulating expression and secretion of MMP-7 and -9 and induces invasion in 3D extracellular matrix (ECM) culture, and furthermore, PKD2-activated MMP9 induces tumor angiogenesis by releasing ECM-bound VEGF-A [162]. Although PKD1 and PKD2 have emerged as tumor-promoters in pancreatic cancer respectively, the role of PKD3 remains obscure and further research is needed to address the function of individual PKD isoform in pancreatic cancer.

5.4. PKD in skin cancer

Basal cell carcinoma (BCC) is the most common form of malignant skin cancer in the world. An estimated 83,000 new cases of skin cancer (6% of all cancer types) have been documented in the USA by 2016 [121]. In normal epidermis, PKD1 is primarily expressed in stratum basalis, the proliferative compartment of the skin which supports a notion that PKD1 may promote hyperproliferative disorders in skin [163] and PKD1 expression was shown to be upregulated in mouse carcinomas and human hyperplastic disorders including BCC [163–165]. PKD1 has been shown to repress keratinocyte differentiation and promote cellular proliferation through modulation of MEK/ERK1/2 pathway [166]. UVB radiation is a key risk factor for developing BCC and it has been reported that activation of PKD1 by Src family of tyrosine kinases in primary mouse keratinocytes exposed to UVB reduced UVB-induced apoptosis and this activation of PKD1 was PKC-independent [167]. Rashel et al. have demonstrated a pro-proliferative role of PKD1 in epidermal adaptive response, wound healing and skin carcinogenesis [168]. Using PKD1-conditional knockout (cKO) mouse model, the authors have presented evidence that: 1) keratinocyte cells in PKD1-cKO mice showed delayed wound healing and reduced proliferative response, 2) PKD1 is a positive regulator of epidermal hyperplasia and inflammation in response to phorbol esters and 3) PKD1-cKO mice are resistant to tumor formation when subjected to two-stage chemically-induced skin carcinogenesis [168]. In a recent study, Ryvkin et al. have demonstrated an opposing role of PKD2 and PKD3 isoforms in human keratinocyte proliferation and differentiation

[169]. The authors have shown that loss of PKD2 resulted in enhanced keratinocyte proliferation suggesting an anti-tumorigenic role of PKD2. Whereas, silencing of PKD3 showed proliferation defect, loss of clonogenicity and diminished tissue regenerative ability, implying a pro-oncogenic role of PKD3. It is to be noted that PKD1 is not expressed in human keratinocytes and thus, PKD2 and PKD3 play a key role in maintaining human epidermal homeostasis, loss of which results in BCC. Further studies are needed to decipher precise roles of PKD isoforms in skin cancer.

5.5. PKD in gastric cancer

PKD1 has been shown to be a negative regulator of gastric cancer [170]. Gastric carcinoma cells as well as patient tissue samples showed decreased PKD1 expression [170]. Gene silencing of PKD1 using siRNA increased cell invasion of gastric cancer and it was found that PKD1 was epigenetically silenced in this tissue type [170]. In an independent study, Shabelnik et al. have demonstrated opposing roles of PKD1 and PKD2 in gastric cancer where PKD1 acts as tumor-suppressor and PKD2 as tumor-promoter [171]. Using tumor samples of different histological variants of primary gastric cancer and gastric adenocarcinoma cell line AGS, the authors have shown that PKD1 and PKD2 are differentially expressed, i.e., lower and higher expression of PKD1 and PKD2 respectively, in both mRNA and protein levels. pcDNA3.1-mediated overexpression of PKD1 resulted in the inhibition of cell proliferation, migration and colony formation, whereas, that of PKD2 enhanced cell proliferation, migration and colony formation abilities in AGS cells [171]. The role of PKD3 isoform in gastric cancer development and progression remains elusive.

5.6. PKD and other cancer types

The role of PKD in other cancer types is poorly defined, although some reports link PKD to certain cancers. For example, activation of PKD by phorbol esters and bombesin via PKC has been shown in small cell lung cancer (SCLC) cell lines h69, H345 and H510 [172]. Using PKC and PKD inhibitors, Brenner et al. has shown that PKD1 is involved in renal cell carcinoma and it promotes tumor progression by positively regulating the adhesion of renal carcinoma cells to endothelial cells [173]. Studies on human malignant lymphoma cells did not conclude distinct roles of PKD as PKD1 levels were undetected and there was no change in PKD2 expression in the malignant cells as compared to benign tissues [174]. In a recent study, PKD2 has been demonstrated to be a potent mediator of glioblastoma where it promotes tumor progression by upregulating integrin α -2 and -4 (ITGA2 and -4), plasminogen activator urokinase (PLAU), plasminogen activator urokinase receptor (PLAUR), and matrix metalloproteinase 1 (MMP1) [175]. When overexpressed in SW480 colon cancer cells, PKD1 suppresses nuclear β -catenin accumulation and inhibits colon cancer [176]. On the contrary, in another study using RKO human colon cancer cell line, it has been demonstrated that targeted inhibition of PKD by small molecule inhibitor suppressed AKT/ERK signaling and NF- κ B activity [177]. Hence, PKD might be a potent chemotherapeutic target for the treatment of colorectal cancer.

6. Therapeutic targeting of PKD in cancer

Accumulating evidence indicate that PKD expression is deregulated in many cancers and PKD plays a crucial role in a wide range of cancer-associated cellular processes such as cell proliferation, migration, apoptosis, EMT, and angiogenesis. This makes PKD an attractive therapeutic target for cancer and has since embarked efforts in the discovery and development of novel PKD inhibitors. Starting with the discovery of a non-ATP-competitive PKD inhibitor CID755673 [128], we have subsequently reported several first-in-class structurally distinct PKD small molecule inhibitors, including the non-ATP-competitive

CID755673 and its derivatives kb-NB142-70 and KMG-NB4-23 [128,178–181], three dual PKD inhibitors (compound 139, 1-NA-PP1, and SD208), and three cell-active PKD inhibitors (CID2011756, CID5389142 and CID1893668) [134–136,181], which are all nanomolar PKD small molecule inhibitors that potently block prostate cancer cell proliferation, migration, and invasion [128,180]. CID755673, in particular, has shown *in vivo* efficacy in other disease models [182].

In addition to our efforts, several highly potent and selective PKD inhibitors have also emerged from the pharmaceutical industry [45,183–185] including an aminopyridine arene CRT0066101 (CRT101) [186] and an aminopyrimidine phenol CRT0066051 (CRT051) [187] from Cancer Research Technology Ltd. Additionally, in a recent study, Golkowski et al. have identified novel PKD inhibitors, namely compounds 1553, 1561, 1649 and 1369 using kinobead-based proteomic assay [188]. In this study, compound 1369 was found to be highly selective and potent pan-PKD inhibitor and an important tool compound to identify the roles of PKD isoforms in cellular as well as *in vivo* models. Among all available PKD SMIs, CRT101 is by far the most potent, selective, and cell-permeable PKD small molecule inhibitor with demonstrated *in vivo* antitumor activity in multiple cancer models [148,177,189]. Specifically, CRT0066101 was found to inhibit growth of pancreatic cancer tumor xenografts [189]. CRT0066101 also caused significant inhibition of tumor growth in HCT116 xenograft nude mice, supporting the therapeutic potential of this inhibitor in colon cancer [177]. CRT0066101 has been shown to be a potent anticancer agent against highly aggressive ER negative breast cancer [148]. This study showed that similar to PKD3 knockdown effect, CRT0066101 significantly reduced breast cancer cell proliferation, migration and invasion both *in vitro* and *in vivo*.

It is interesting to note that despite the differential roles of PKD isoforms in different cancer types as implicated in our discussion above, PKD inhibitors have unequivocally exhibited antitumor activities in various *in vitro* and *in vivo* cancer models. It is possible the overall selectivity profiles of these inhibitors favor anticancer activity with PKD being the primary target. Nonetheless, with the growing reports of diverse functions of PKD isoforms in different cancers, the development of isoform-selective PKD inhibitors is a well-justified direction for future studies.

7. Perspectives and concluding remarks

Growing evidence supports PKD as a key signaling molecule that orchestrates various cancer-associated biological functions such as cell proliferation, survival, EMT, migration, invasion, secretion and angiogenesis upon activation by a battery of stimuli. Despite recent advances of our knowledge about the role of PKD in various pathological conditions including cancer, cardiac hypertrophy and inflammation, the similar or opposing roles of the same PKD isoform in different cancers or of the same cancer for different PKD isoforms has raised more unanswered questions.

To summarize briefly, emerging evidence suggests that PKD1 can function as a tumor-suppressive protein in breast and gastric cancers, where PKD1 inhibits cell survival, proliferation, migration by negatively regulating several key target proteins including SSH1L, Snail and MMPs. However, in other cancers, such as pancreatic and skin cancers, PKD1 emerged as a driver of neoplasia. It does so by many mechanisms such as activating MEK1/2 pathway that increases DNA replication, inhibits apoptosis and promotes proliferation by positively regulating ERK/MAPK pathway. Compared to the opposing roles of PKD1 in cancer development and progression, majority of studies have shown that PKD2 is a tumor-promoting protein in a wide range of cancers. It activates different biological functions such as NF- κ B signaling, MMP expression, induction of angiogenesis and inhibition of apoptosis and promotes some common cancers including carcinomas of prostate, breast, pancreas, stomach and also other cancers such as glioblastoma.

The actual role of PKD3 in cancer is elusive. It shows pro-oncogenic properties in case of prostate, breast and skin cancer but what it does in pancreatic and gastric cancer, is unknown.

In this review, we have discussed the major signaling pathways involved in development and progression of neoplasm that are modulated by PKD, the molecular mechanisms of regulation of cellular phenomenon orchestrated by PKD and manifestation of each type of PKD isotypes in the context of major carcinomas. Although PKD emerged as a potential target for chemotherapeutic intervention and pan-PKD inhibitors have shown potent anti-cancer activity in multiple cancer models both *in vivo* and *in vitro*. Many cellular studies have demonstrated differential effects of PKD isoforms in different biological processes. These differences seem to be isoform- and tumor type-dependent, which raise questions on whether it is appropriate and how to target PKD for cancer treatment. As developed in this review and going forward, there remain many unanswered questions. What are the molecular cues that direct precise and selective role of PKD in disease progression? Is it the isoform-specific function that enables PKD to selectively choose the cell type to exert its biological effect? What would be the best strategy to develop effective small molecule inhibitors of PKD that will preserve the tumor-suppressing capacity of PKD while eradicating those that are tumor-promoting? Further knowledge defining the precise role of PKD isoforms in different tumor models will provide a much clearer picture for targeting this family of protein kinases for cancer therapy.

Conflict of interests

The authors declare no conflict of interest.

Transparency document

The <http://dx.doi.org/10.1016/j.bbcan.2017.05.008> associated with this article can be found in online version.

Acknowledgement

This study was supported in part by the National Institutes of Health grant R01CA142580 (Wang), Department of Defense award PC150190, and Oversea Hong Kong & Macao Scholars Collaborative Research Fund of NSFC in China (Grant#813228020, Deng and Wang).

References

- [1] G. Manning, D.B. Whyte, R. Martinez, T. Hunter, S. Sudarsanam, The protein kinase complement of the human genome, *Science* 298 (2002) 1912–1934.
- [2] A.M. Valverde, J. Sinnott-Smith, J. Van Lint, E. Rozengurt, Molecular cloning and characterization of protein kinase D: a target for diacylglycerol and phorbol esters with a distinctive catalytic domain, *Proc. Natl. Acad. Sci. U. S. A.* 91 (1994) 8572–8576.
- [3] F.J. Johannes, J. Prestle, S. Eis, P. Oberhagemann, K. Pfizenmaier, PKC α is a novel, atypical member of the protein kinase C family, *J. Biol. Chem.* 269 (1994) 6140–6148.
- [4] A. Cano, M.A. Perez-Moreno, I. Rodrigo, A. Locascio, M.J. Blanco, M.G. del Barrio, F. Portillo, M.A. Nieto, The transcription factor snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression, *Nat. Cell Biol.* 2 (2000) 76–83.
- [5] A. Hayashi, N. Seki, A. Hattori, S. Kozuma, T. Saito, PKC η , a new member of the protein kinase C family, composes a fourth subfamily with PKC μ , *Biochim. Biophys. Acta* 1450 (1999) 99–106.
- [6] S. Sturany, J. Van Lint, F. Muller, M. Wilda, H. Hameister, M. Hocker, A. Brey, U. Gern, J. Vandenheede, T. Gress, G. Adler, T. Seufferlein, Molecular cloning and characterization of the human protein kinase D2. A novel member of the protein kinase D family of serine threonine kinases, *J. Biol. Chem.* 276 (2001) 3310–3318.
- [7] K. Ellwanger, A. Hausser, Physiological functions of protein kinase D in vivo, *IUBMB Life* 65 (2013) 98–107.
- [8] L. Sanchez-Ruiloba, N. Cabrera-Poch, M. Rodriguez-Martinez, C. Lopez-Menendez, R.M. Jean-Mairet, A.M. Higuero, T. Iglesias, Protein kinase D intracellular localization and activity control kinase D-interacting substrate of 220-kDa traffic through a postsynaptic density-95/discs large/zonula occludens-1-binding motif, *J. Biol. Chem.* 281 (2006) 18888–18900.
- [9] H. Doppler, P. Storz, A novel tyrosine phosphorylation site in protein kinase D contributes to oxidative stress-mediated activation, *J. Biol. Chem.* 282 (2007) 31873–31881.
- [10] D. Maier, A. Hausser, A.C. Nagel, G. Link, S.J. Kugler, I. Wech, K. Pfizenmaier, A. Preiss, *Drosophila* protein kinase D is broadly expressed and a fraction localizes to the Golgi compartment, *Gene Expr. Patterns* 6 (2006) 849–856.
- [11] H. Feng, M. Ren, L. Chen, C.S. Rubin, Properties, regulation, and in vivo functions of a novel protein kinase D: *Caenorhabditis elegans* DKF-2 links diacylglycerol second messenger to the regulation of stress responses and life span, *J. Biol. Chem.* 282 (2007) 31273–31288.
- [12] H. Feng, M. Ren, C.S. Rubin, Conserved domains subserve novel mechanisms and functions in DKF-1, a *Caenorhabditis elegans* protein kinase D, *J. Biol. Chem.* 281 (2006) 17815–17826.
- [13] H. Feng, M. Ren, S.L. Wu, D.H. Hall, C.S. Rubin, Characterization of a novel protein kinase D: *Caenorhabditis elegans* DKF-1 is activated by translocation-phosphorylation and regulates movement and growth in vivo, *J. Biol. Chem.* 281 (2006) 17801–17814.
- [14] E. Rozengurt, O. Rey, R.T. Waldron, Protein kinase D signaling, *J. Biol. Chem.* 280 (2005) 13205–13208.
- [15] Q.J. Wang, PKD at the crossroads of DAG and PKC signaling, *Trends Pharmacol. Sci.* 27 (2006) 317–323.
- [16] A. Rykx, L. De Kimpe, S. Mikhalap, T. Vantus, T. Seufferlein, J.R. Vandenheede, J. Van Lint, Protein kinase D: a family affair, *FEBS Lett.* 546 (2003) 81–86.
- [17] J. Van Lint, A. Rykx, Y. Maeda, T. Vantus, S. Sturany, V. Malhotra, J.R. Vandenheede, T. Seufferlein, Protein kinase D: an intracellular traffic regulator on the move, *Trends Cell Biol.* 12 (2002) 193–200.
- [18] Q.J. Wang, T.W. Fang, D. Yang, N.E. Lewin, J. Van Lint, V.E. Marquez, P.M. Blumberg, Ligand structure-activity requirements and phospholipid dependence for the binding of phorbol esters to protein kinase D, *Mol. Pharmacol.* 64 (2003) 1342–1348.
- [19] G. Anderson, J. Chen, Q.J. Wang, Individual C1 domains of PKD3 in phorbol ester-induced plasma membrane translocation of PKD3 in intact cells, *Cell. Signal.* 17 (2005) 1397–1411.
- [20] J. Chen, F. Deng, J. Li, Q.J. Wang, Selective binding of phorbol esters and diacylglycerol by individual C1 domains of the PKD family, *Biochem. J.* 411 (2008) 333–342.
- [21] M. Spitaler, E. Emslie, C.D. Wood, D. Cantrell, Diacylglycerol and protein kinase D localization during T lymphocyte activation, *Immunity* 24 (2006) 535–546.
- [22] Y. Maeda, G.V. Beznoussenko, J. Van Lint, A.A. Mironov, V. Malhotra, Recruitment of protein kinase D to the trans-Golgi network via the first cysteine-rich domain, *EMBO J.* 20 (2001) 5982–5990.
- [23] Q.J. Wang, T.W. Fang, K. Nacro, V.E. Marquez, S. Wang, P.M. Blumberg, Role of hydrophobic residues in the C1b domain of protein kinase C delta on ligand and phospholipid interactions, *J. Biol. Chem.* 276 (2001) 19580–19587.
- [24] K. Irie, A. Nakahara, H. Ohgashi, H. Fukuda, P.A. Wender, H. Konishi, U. Kikkawa, Synthesis and phorbol ester-binding studies of the individual cysteine-rich motifs of protein kinase D, *Bioorg. Med. Chem. Lett.* 9 (1999) 2487–2490.
- [25] T. Iglesias, R.T. Waldron, E. Rozengurt, Identification of in vivo phosphorylation sites required for protein kinase D activation, *J. Biol. Chem.* 273 (1998) 27662–27667.
- [26] G.V. Pusapati, D. Krndija, M. Armacki, G. von Wichert, J. von Blume, V. Malhotra, G. Adler, T. Seufferlein, Role of the second cysteine-rich domain and Pro275 in protein kinase D2 interaction with ADP-ribosylation factor 1, trans-Golgi network recruitment, and protein transport, *Mol. Biol. Cell* 21 (2010) 1011–1022.
- [27] A. Auer, J. von Blume, S. Sturany, G. von Wichert, J. Van Lint, J. Vandenheede, G. Adler, T. Seufferlein, Role of the regulatory domain of protein kinase D2 in phorbol ester binding, catalytic activity, and nucleocytoplasmic shuttling, *Mol. Biol. Cell* 16 (2005) 4375–4385.
- [28] M.T. Kunkel, E.L. Garcia, T. Kajimoto, R.A. Hall, A.C. Newton, The protein scaffold NHERF-1 controls the amplitude and duration of localized protein kinase D activity, *J. Biol. Chem.* 284 (2009) 24653–24661.
- [29] H. Doppler, P. Storz, J. Li, M.J. Comb, A. Tokar, A phosphorylation state-specific antibody recognizes Hsp27, a novel substrate of protein kinase D, *J. Biol. Chem.* 280 (2005) 15013–15019.
- [30] J.E. Huttli, E.T. Jarrell, J.D. Chang, D.W. Abbott, P. Storz, A. Tokar, L.C. Cantley, B.E. Turk, A rapid method for determining protein kinase phosphorylation specificity, *Nat. Methods* 1 (2004) 27–29.
- [31] J. Guo, Z. Gertsberg, N. Ozgen, A. Sabri, S.F. Steinberg, Protein kinase D isoforms are activated in an agonist-specific manner in cardiomyocytes, *J. Biol. Chem.* 286 (2011) 6500–6509.
- [32] N. Durand, S. Borges, P. Storz, Protein kinase D enzymes as regulators of EMT and cancer cell invasion, *J. Clin. Med.* 5 (2016).
- [33] C. Wong, Z.G. Jin, Protein kinase C-dependent protein kinase D activation modulates ERK signal pathway and endothelial cell proliferation by vascular endothelial growth factor, *J. Biol. Chem.* 280 (2005) 33262–33269.
- [34] J. Yoo, C. Chung, L. Slice, J. Sinnott-Smith, E. Rozengurt, Protein kinase D mediates synergistic expression of COX-2 induced by TNF- α and bradykinin in human colonic myofibroblasts, *Am. J. Physiol. Cell Physiol.* 297 (2009) C1576–C1587.
- [35] R. Jacamo, J. Sinnott-Smith, O. Rey, R.T. Waldron, E. Rozengurt, Sequential protein kinase C (PKC)-dependent and PKC-independent protein kinase D catalytic activation via Gq-coupled receptors: differential regulation of activation loop Ser (744) and Ser(748) phosphorylation, *J. Biol. Chem.* 283 (2008) 12877–12887.
- [36] J. Sinnott-Smith, R. Jacamo, R. Kui, Y.M. Wang, S.H. Young, O. Rey, R.T. Waldron, E. Rozengurt, Protein kinase D mediates mitogenic signaling by Gq-coupled receptors through protein kinase C-independent regulation of activation loop Ser744 and Ser748 phosphorylation, *J. Biol. Chem.* 284 (2009) 13434–13445.

- [37] S.A. Matthews, E. Rozengurt, D. Cantrell, Characterization of serine 916 as an *in vivo* autophosphorylation site for protein kinase D/protein kinase Cmu, *J. Biol. Chem.* 274 (1999) 26543–26549.
- [38] S.F. Steinberg, Regulation of protein kinase D1 activity, *Mol. Pharmacol.* 81 (2012) 284–291.
- [39] S. Guha, S. Tanasavimon, J. Sinnett-Smith, E. Rozengurt, Role of protein kinase D signaling in pancreatic cancer, *Biochem. Pharmacol.* 80 (2010) 1946–1954.
- [40] M. Jaggi, C. Du, W. Zhang, K.C. Balaji, Protein kinase D1: a protein of emerging translational interest, *Front. Biosci.* 12 (2007) 3757–3767.
- [41] I. Brandlin, S. Hubner, T. Eiseler, M. Martinez-Moya, A. Horschinek, A. Hausser, G. Link, S. Rupp, P. Storz, K. Pfizenmaier, F.J. Johannes, Protein kinase C (PKC) ϵ -mediated PKC μ activation modulates ERK and JNK signal pathways, *J. Biol. Chem.* 277 (2002) 6490–6496.
- [42] A.B. Celil, P.G. Campbell, BMP-2 and insulin-like growth factor-I mediate Osterix (Ox) expression in human mesenchymal stem cells via the MAPK and protein kinase D signaling pathways, *J. Biol. Chem.* 280 (2005) 31353–31359.
- [43] P. Storz, H. Doppler, A. Toker, Activation loop phosphorylation controls protein kinase D-dependent activation of nuclear factor kappaB, *Mol. Pharmacol.* 66 (2004) 870–879.
- [44] V.O. Rybin, J. Guo, S.F. Steinberg, Protein kinase D1 autophosphorylation via distinct mechanisms at Ser744/Ser748 and Ser916, *J. Biol. Chem.* 284 (2009) 2332–2343.
- [45] E.L. Meredith, K. Beattie, R. Burgis, M. Capparelli, J. Chapo, L. Dipietro, G. Gamber, I. Enyedy, D.B. Hood, V. Hosagrahara, C. Jewell, K.A. Koch, W. Lee, D.D. Lemon, T.A. McKinsey, M. Miranda, N. Pagratis, D. Phan, C. Plato, C. Rao, O. Rozhitskaya, N. Soldermann, C. Springer, M. van Eis, R.B. Vega, W. Yan, Q. Zhu, L.G. Monovich, Identification of potent and selective amidobiprydyl inhibitors of protein kinase D, *J. Med. Chem.* 53 (2010) 5422–5438.
- [46] M.T. Chang, S. Asthana, S.P. Gao, B.H. Lee, J.S. Chapman, C. Kandath, J. Gao, N.D. Socci, D.B. Solit, A.B. Olshen, N. Schultz, B.S. Taylor, Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity, *Nat. Biotechnol.* 34 (2016) 155–163.
- [47] L.A. Garraway, E.S. Lander, Lessons from the cancer genome, *Cell* 153 (2013) 17–37.
- [48] B. Vogelstein, N. Papadopoulos, V.E. Velculescu, S. Zhou, L.A. Diaz Jr., K.W. Kinzler, Cancer genome landscapes, *Science* 339 (2013) 1546–1558.
- [49] J. Hofvander, E. Arbajian, K.G. Stenula, K. Lindkvist-Petersson, M. Larsson, J. Nilsson, L. Magnusson, F.V. von Steyern, P. Rissler, J.L. Hornick, F. Mertens, Frequent low-level mutations of protein kinase D2 in angiolipoma, *J. Pathol.* 241 (5) (2017) 578–582, <http://dx.doi.org/10.1002/path.4865> Epub 2017 Feb 27.
- [50] S. Piscuoglio, N. Fusco, C.K. Ng, L.G. Martelotto, A. da Cruz Paula, N. Katabi, B.P. Rubin, A. Skalova, I. Weinreb, B. Weigelt, J.S. Reis-Filho, Lack of PRKD2 and PRKD3 kinase domain somatic mutations in PRKD1 wild-type classic polymorphous low-grade adenocarcinomas of the salivary gland, *Histopathology* 68 (2016) 1055–1062.
- [51] I. Weinreb, S. Piscuoglio, L.G. Martelotto, D. Waggott, C.K. Ng, B. Perez-Ordóñez, N.J. Harding, J. Alfaro, K.C. Chu, A. Viale, N. Fusco, A. da Cruz Paula, C. Marchio, R.A. Sakr, R. Lim, L.D. Thompson, S.I. Chiosea, R.R. Seethala, A. Skalova, E.B. Stelow, I. Fonseca, A. Assaad, C. How, J. Wang, R. de Borja, M. Chan-Seng-Yue, C.J. Howlett, A.C. Nichols, Y.H. Wen, N. Katabi, N. Buchner, L. Mullen, T. Kislinger, B.G. Wouters, F.F. Liu, L. Norton, J.D. McPherson, B.P. Rubin, B.A. Clarke, B. Weigelt, P.C. Boutros, J.S. Reis-Filho, Hotspot activating PRKD1 somatic mutations in polymorphous low-grade adenocarcinomas of the salivary glands, *Nat. Genet.* 46 (2014) 1166–1169.
- [52] J. Debnath, S.K. Muthuswamy, J.S. Brugge, Morphogenesis and oncogenesis of MCF-10A mammary epithelial acini grown in three-dimensional basement membrane cultures, *Methods* 30 (2003) 256–268.
- [53] J. Debnath, J.S. Brugge, Modelling glandular epithelial cancers in three-dimensional cultures, *Nat. Rev. Cancer* 5 (2005) 675–688.
- [54] H. Bartuma, K.H. Nord, G. Macchia, M. Isaksson, J. Nilsson, H.A. Domanski, N. Mandahl, F. Mertens, Gene expression and single nucleotide polymorphism array analyses of spindle cell lipomas and conventional lipomas with 13q14 deletion, *Genes Chromosomes. Cancer* 50 (2011) 619–632.
- [55] J. Yuan, L. Slice, J.H. Walsh, E. Rozengurt, Activation of protein kinase D by signaling through the alpha subunit of the heterotrimeric G protein G(q), *J. Biol. Chem.* 275 (2000) 2157–2164.
- [56] J. Yuan, L.W. Slice, J. Gu, E. Rozengurt, Cooperation of Gq, Gi, and G12/13 in protein kinase D activation and phosphorylation induced by lysophosphatidic acid, *J. Biol. Chem.* 278 (2003) 4882–4891.
- [57] J. Yuan, L.W. Slice, E. Rozengurt, Activation of protein kinase D by signaling through rho and the alpha subunit of the heterotrimeric G protein G13, *J. Biol. Chem.* 276 (2001) 38619–38627.
- [58] J.L. Zugaza, R.T. Waldron, J. Sinnett-Smith, E. Rozengurt, Bombesin, vasopressin, endothelin, bradykinin, and platelet-derived growth factor rapidly activate protein kinase D through a protein kinase C-dependent signal transduction pathway, *J. Biol. Chem.* 272 (1997) 23952–23960.
- [59] J. Sinnett-Smith, E. Zhukova, N. Hsieh, X. Jiang, E. Rozengurt, Protein kinase D potentiates DNA synthesis induced by Gq-coupled receptors by increasing the duration of ERK signaling in Swiss 3T3 cells, *J. Biol. Chem.* 279 (2004) 16883–16893.
- [60] J. Sinnett-Smith, E. Zhukova, O. Rey, E. Rozengurt, Protein kinase D2 potentiates MEK/ERK/RSK signaling, c-Fos accumulation and DNA synthesis induced by bombesin in Swiss 3T3 cells, *J. Cell. Physiol.* 211 (2007) 781–790.
- [61] E. Zhukova, J. Sinnett-Smith, E. Rozengurt, Protein kinase D potentiates DNA synthesis and cell proliferation induced by bombesin, vasopressin, or phorbol esters in Swiss 3T3 cells, *J. Biol. Chem.* 276 (2001) 40298–40305.
- [62] E. Rozengurt, Mitogenic signaling pathways induced by G protein-coupled receptors, *J. Cell. Physiol.* 213 (2007) 589–602.
- [63] C.F. Cowell, H. Doppler, I.K. Yan, A. Hausser, Y. Umezawa, P. Storz, Mitochondrial diacylglycerol initiates protein-kinase D1-mediated ROS signaling, *J. Cell Sci.* 122 (2009) 919–928.
- [64] P. Storz, H. Doppler, A. Toker, Protein kinase D mediates mitochondrion-to-nucleus signaling and detoxification from mitochondrial reactive oxygen species, *Mol. Cell. Biol.* 25 (2005) 8520–8530.
- [65] R. Kalluri, R.A. Weinberg, The basics of epithelial-mesenchymal transition, *J. Clin. Invest.* 119 (2009) 1420–1428.
- [66] J. Yang, R.A. Weinberg, Epithelial-mesenchymal transition: at the crossroads of development and tumor metastasis, *Dev. Cell* 14 (2008) 818–829.
- [67] J.P. Thiery, Epithelial-mesenchymal transitions in tumour progression, *Nat. Rev. Cancer* 2 (2002) 442–454.
- [68] F. van Roy, G. Berx, The cell–cell adhesion molecule E-cadherin, *Cell. Mol. Life Sci.* 65 (2008) 3756–3788.
- [69] E. Batlle, E. Sancho, C. Franci, D. Dominguez, M. Monfar, J. Baulida, A. Garcia De Herreros, The transcription factor snail is a repressor of E-cadherin gene expression in epithelial tumour cells, *Nat. Cell Biol.* 2 (2000) 84–89.
- [70] H. Peinado, D. Olmeda, A. Cano, Snail, Zeb and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat. Rev. Cancer* 7 (2007) 415–428.
- [71] J. Yang, S.A. Mani, J.L. Donaher, S. Ramaswamy, R.A. Itzykson, C. Come, P. Savagner, I. Gitelman, A. Richardson, R.A. Weinberg, Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis, *Cell* 117 (2004) 927–939.
- [72] Y. Wu, J. Deng, P.G. Rychahou, S. Qiu, B.M. Evers, B.P. Zhou, Stabilization of snail by NF-kappaB is required for inflammation-induced cell migration and invasion, *Cancer Cell* 15 (2009) 416–428.
- [73] Z. Yang, S. Rayala, D. Nguyen, R.K. Vadlamudi, S. Chen, R. Kumar, Pak1 phosphorylation of snail, a master regulator of epithelial-to-mesenchyme transition, modulates snail's subcellular localization and functions, *Cancer Res.* 65 (2005) 3179–3184.
- [74] J.I. Yook, X.Y. Li, I. Ota, C. Hu, H.S. Kim, N.H. Kim, S.Y. Cha, J.K. Ryu, Y.J. Choi, J. Kim, E.R. Fearon, S.J. Weiss, A Wnt-Axin2-GSK3beta cascade regulates Snail1 activity in breast cancer cells, *Nat. Cell Biol.* 8 (2006) 1398–1406.
- [75] B.P. Zhou, J. Deng, W. Xia, J. Xu, Y.M. Li, M. Gunduz, M.C. Hung, Dual regulation of snail by GSK-3beta-mediated phosphorylation in control of epithelial-mesenchymal transition, *Nat. Cell Biol.* 6 (2004) 931–940.
- [76] H. Yamaguchi, J. Condeelis, Regulation of the actin cytoskeleton in cancer cell migration and invasion, *Biochim. Biophys. Acta* 1773 (2007) 642–652.
- [77] M. Egeblad, Z. Werb, New functions for the matrix metalloproteinases in cancer progression, *Nat. Rev. Cancer* 2 (2002) 161–174.
- [78] M.D. Sternlicht, A. Lochter, C.J. Sympon, B. Huey, J.P. Rougier, J.W. Gray, D. Pinkel, M.J. Bissell, Z. Werb, The stromal proteinase MMP3/stromelysin-1 promotes mammary carcinogenesis, *Cell* 98 (1999) 137–146.
- [79] R.W. Scott, M.F. Olson, LIM kinases: function, regulation and association with human disease, *J. Mol. Med. (Berl.)* 85 (2007) 555–568.
- [80] R. Niwa, K. Nagata-Ohashi, M. Takeichi, K. Mizuno, T. Uemura, Control of actin reorganization by slingshot, a family of phosphatases that dephosphorylate ADF/cofilin, *Cell* 108 (2002) 233–246.
- [81] T. Eiseler, H. Doppler, I.K. Yan, S. Goodison, P. Storz, Protein kinase D1 regulates matrix metalloproteinase expression and inhibits breast cancer cell invasion, *Breast Cancer Res.* 11 (2009) R13.
- [82] T. Eiseler, H. Doppler, I.K. Yan, K. Kitatani, K. Mizuno, P. Storz, Protein kinase D1 regulates cofilin-mediated F-actin reorganization and cell motility through slingshot, *Nat. Cell Biol.* 11 (2009) 545–556.
- [83] P. Peterburs, J. Heering, G. Link, K. Pfizenmaier, M.A. Olayioye, A. Hausser, Protein kinase D regulates cell migration by direct phosphorylation of the cofilin phosphatase slingshot 1 like, *Cancer Res.* 69 (2009) 5634–5638.
- [84] H. Doppler, L.L. Bastea, S. Borges, S.J. Spratley, S.E. Pearce, P. Storz, Protein kinase d isoforms differentially modulate cofilin-driven directed cell migration, *PLoS One* 9 (2014) e98090.
- [85] S. Ziegler, T. Eiseler, R.P. Scholz, A. Beck, G. Link, A. Hausser, A novel protein kinase D phosphorylation site in the tumor suppressor Rab interactor 1 is critical for coordination of cell migration, *Mol. Biol. Cell* 22 (2011) 570–580.
- [86] H. Hu, J.M. Bliss, Y. Wang, J. Colicelli, RIN1 is an ABL tyrosine kinase activator and a regulator of epithelial-cell adhesion and migration, *Curr. Biol.* 15 (2005) 815–823.
- [87] L. Klampfer, J. Huang, S. Shirasawa, T. Sasazuki, L. Augenlicht, Histone deacetylase inhibitors induce cell death selectively in cells that harbor activated KrasV12: the role of signal transducers and activators of transcription 1 and p21, *Cancer Res.* 67 (2007) 8477–8485.
- [88] L.T. Liu, H.C. Chang, L.C. Chiang, W.C. Hung, Histone deacetylase inhibitor up-regulates RECK to inhibit MMP-2 activation and cancer cell invasion, *Cancer Res.* 63 (2003) 3069–3072.
- [89] C.H. Ha, W. Wang, B.S. Jhun, C. Wong, A. Hausser, K. Pfizenmaier, T.A. McKinsey, E.N. Olson, Z.G. Jin, Protein kinase D-dependent phosphorylation and nuclear export of histone deacetylase 5 mediates vascular endothelial growth factor-induced gene expression and angiogenesis, *J. Biol. Chem.* 283 (2008) 14590–14599.
- [90] Z. Zou, F. Zeng, W. Xu, C. Wang, Z. Ke, Q.J. Wang, F. Deng, PKD2 and PKD3 promote prostate cancer cell invasion by modulating NF-kappaB- and HDAC1-mediated expression and activation of uPA, *J. Cell Sci.* 125 (2012) 4800–4811.
- [91] H. Yamamoto, N. Oue, A. Sato, Y. Hasegawa, H. Yamamoto, A. Matsubara, W. Yasui, A. Kikuchi, Wnt5a signaling is involved in the aggressiveness of prostate cancer and expression of metalloproteinase, *Oncogene* 29 (2010) 2036–2046.

- [92] D. Mottet, A. Bellahcene, S. Pirotte, D. Walthregny, C. Deroanne, V. Lamour, R. Lidereau, V. Castronovo, Histone deacetylase 7 silencing alters endothelial cell migration, a key step in angiogenesis, *Circ. Res.* 101 (2007) 1237–1246.
- [93] C.H. Ha, B.S. Jhun, H.Y. Kao, Z.G. Jin, VEGF stimulates HDAC7 phosphorylation and cytoplasmic accumulation modulating matrix metalloproteinase expression and angiogenesis, *Arterioscler. Thromb. Vasc. Biol.* 28 (2008) 1782–1788.
- [94] N. Ferrara, T. Davis-Smyth, The biology of vascular endothelial growth factor, *Endocr. Rev.* 18 (1997) 4–25.
- [95] J. Folkman, Angiogenesis in cancer, vascular, rheumatoid and other disease, *Nat. Med.* 1 (1995) 27–31.
- [96] N. Ferrara, VEGF and the quest for tumour angiogenesis factors, *Nat. Rev. Cancer* 2 (2002) 795–803.
- [97] H. Chen, A. Chedotal, Z. He, C.S. Goodman, M. Tessier-Lavigne, Neuropilin-2, a novel member of the neuropilin family, is a high affinity receptor for the semaphorins Sema E and Sema IV but not Sema III, *Neuron* 19 (1997) 547–559.
- [98] A.L. Kolodkin, D.V. Levensgood, E.G. Rowe, Y.T. Tai, R.J. Giger, D.D. Ginty, Neuropilin is a semaphorin III receptor, *Cell* 90 (1997) 753–762.
- [99] B.P. Eliceiri, R. Paul, P.L. Schwartzberg, J.D. Hood, J. Leng, D.A. Cheresh, Selective requirement for Src kinases during VEGF-induced angiogenesis and vascular permeability, *Mol. Cell* 4 (1999) 915–924.
- [100] D. Guo, Q. Jia, H.Y. Song, R.S. Warren, D.B. Donner, Vascular endothelial cell growth factor promotes tyrosine phosphorylation of mediators of signal transduction that contain SH2 domains. Association with endothelial cell proliferation, *J. Biol. Chem.* 270 (1995) 6729–6733.
- [101] L.E. Benjamin, D. Golijanin, A. Itin, D. Podes, E. Keshet, Selective ablation of immature blood vessels in established human tumors follows vascular endothelial growth factor withdrawal, *J. Clin. Invest.* 103 (1999) 159–165.
- [102] L.E. Benjamin, E. Keshet, Conditional switching of vascular endothelial growth factor (VEGF) expression in tumors: induction of endothelial cell shedding and regression of hemangioblastoma-like vessels by VEGF withdrawal, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 8761–8766.
- [103] H.P. Gerber, V. Dixit, N. Ferrara, Vascular endothelial growth factor induces expression of the antiapoptotic proteins Bcl-2 and A1 in vascular endothelial cells, *J. Biol. Chem.* 273 (1998) 13313–13316.
- [104] H.P. Gerber, A. McMurtry, J. Kowalski, M. Yan, B.A. Keyt, V. Dixit, N. Ferrara, Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation, *J. Biol. Chem.* 273 (1998) 30336–30343.
- [105] F. Yuan, Y. Chen, M. Dellian, N. Safabakhsh, N. Ferrara, R.K. Jain, Time-dependent vascular regression and permeability changes in established human tumor xenografts induced by an anti-vascular endothelial growth factor/vascular permeability factor antibody, *Proc. Natl. Acad. Sci. U. S. A.* 93 (1996) 14765–14770.
- [106] Q. Hao, L. Wang, Z.J. Zhao, H. Tang, Identification of protein kinase D2 as a pivotal regulator of endothelial cell proliferation, migration, and angiogenesis, *J. Biol. Chem.* 284 (2009) 799–806.
- [107] M. Muller, J. Schroer, N. Azoitei, T. Eiseler, W. Bergmann, R. Kohntop, Q. Lin, I.G. Costa, M. Zenke, F. Genze, C. Weidgang, T. Seufferlein, S. Liebau, A. Kleger, A time frame permissive for protein kinase D2 activity to direct angiogenesis in mouse embryonic stem cells, *Sci. Rep.* 5 (2015) 11742.
- [108] J. Lemonnier, C. Ghayor, J. Guicheux, J. Caverzasio, Protein kinase C-independent activation of protein kinase D is involved in BMP-2-induced activation of stress mitogen-activated protein kinases JNK and p38 and osteoblastic cell differentiation, *J. Biol. Chem.* 279 (2004) 259–264.
- [109] S. Li, W. Xu, Z. Xing, J. Qian, L. Chen, R. Gu, W. Guo, X. Lai, W. Zhao, S. Li, Y. Wang, Q.J. Wang, F. Deng, A conditional knockout mouse model reveals a critical role of PKD1 in osteoblast differentiation and bone development, *Sci. Rep.* 7 (2017) 40505.
- [110] S. Nandana, M. Tripathi, P. Duan, C.Y. Chu, R. Mishra, C. Liu, R. Jin, H. Yamashita, M. Zayzafoon, N.A. Bhowmick, H.E. Zhou, R.J. Matusik, L.W. Chung, Bone metastasis of prostate cancer can be therapeutically targeted at the TBX2-WNT signaling axis, *Cancer Res.* 77 (6) (2017) 1331–1344, <http://dx.doi.org/10.1158/0008-5472.CAN-16-0497> Epub 2017 Jan 20.
- [111] F. Dequiedt, J. Van Lint, E. Lecomte, V. Van Duppen, T. Seufferlein, J.R. Vandenheede, R. Wattiez, R. Kettmann, Phosphorylation of histone deacetylase 7 by protein kinase D mediates T cell receptor-induced Nur77 expression and apoptosis, *J. Exp. Med.* 201 (2005) 793–804.
- [112] S.A. Matthews, P. Liu, M. Spitaler, E.N. Olson, T.A. McKinsey, D.A. Cantrell, A.M. Scharenberg, Essential role for protein kinase D family kinases in the regulation of class II histone deacetylases in B lymphocytes, *Mol. Cell. Biol.* 26 (2006) 1569–1577.
- [113] M. Parra, H. Kasler, T.A. McKinsey, E.N. Olson, E. Verdin, Protein kinase D1 phosphorylates HDAC7 and induces its nuclear export after T-cell receptor activation, *J. Biol. Chem.* 280 (2005) 13762–13770.
- [114] R.B. Medeiros, D.M. Dickey, H. Chung, A.C. Quale, L.R. Nagarajan, D.D. Billadeau, Y. Shimizu, Protein kinase D1 and the beta 1 integrin cytoplasmic domain control beta 1 integrin function via regulation of Rap1 activation, *Immunity* 23 (2005) 213–226.
- [115] A. Irie, K. Harada, H. Tsukamoto, J.R. Kim, N. Araki, Y. Nishimura, Protein kinase D2 contributes to either IL-2 promoter regulation or induction of cell death upon TCR stimulation depending on its activity in Jurkat cells, *Int. Immunol.* 18 (2006) 1737–1747.
- [116] J.E. Park, Y.I. Kim, A.K. Yi, Protein kinase D1: a new component in TLR9 signaling, *J. Immunol.* 181 (2008) 2044–2055.
- [117] T.R. Murphy, H.J. Legere 3rd, H.R. Katz, Activation of protein kinase D1 in mast cells in response to innate, adaptive, and growth factor signals, *J. Immunol.* 179 (2007) 7876–7882.
- [118] N. Yoo, H.R. Lee, J.M. Son, H.B. Kang, H.G. Lee, S.R. Yoon, S.Y. Yoon, J.W. Kim, Genkwaadin promotes leukocyte migration by increasing CD44 expression via PKD1/NF-kappaB signaling pathway, *Immunol. Lett.* 173 (2016) 69–76.
- [119] X. Xu, N. Gera, H. Li, M. Yun, L. Zhang, Y. Wang, Q.J. Wang, T. Jin, GPCR-mediated PLCbetagamma/PKCbeta/PKD signaling pathway regulates the cofilin phosphatase slingshot 2 in neutrophil chemotaxis, *Mol. Biol. Cell* 26 (2015) 874–886.
- [120] D. Hanahan, R.A. Weinberg, Hallmarks of cancer: the next generation, *Cell* 144 (2011) 646–674.
- [121] R.L. Siegel, K.D. Miller, A. Jemal, Cancer statistics, 2016, *CA Cancer J. Clin.* 66 (2016) 7–30.
- [122] T. Shirai, Significance of chemoprevention for prostate cancer development: experimental in vivo approaches to chemoprevention, *Pathol. Int.* 58 (2008) 1–16.
- [123] J. Chen, F. Deng, S.V. Singh, Q.J. Wang, Protein kinase D3 (PKD3) contributes to prostate cancer cell growth and survival through a PKCepsilon/PKD3 pathway downstream of Akt and ERK 1/2, *Cancer Res.* 68 (2008) 3844–3853.
- [124] P. Mak, M. Jaggi, V. Syed, S.C. Chauhan, S. Hassan, H. Biswas, K.C. Balaji, Protein kinase D1 (PKD1) influences androgen receptor (AR) function in prostate cancer cells, *Biochem. Biophys. Res. Commun.* 373 (2008) 618–623.
- [125] M. Jaggi, S.L. Johansson, J.J. Baker, L.M. Smith, A. Galich, K.C. Balaji, Aberrant expression of E-cadherin and beta-catenin in human prostate cancer, *Urol. Oncol.* 23 (2005) 402–406.
- [126] S. Hassan, M.H. Biswas, C. Zhang, C. Du, K.C. Balaji, Heat shock protein 27 mediates repression of androgen receptor function by protein kinase D1 in prostate cancer cells, *Oncogene* 28 (2009) 4386–4396.
- [127] M. Jaggi, P.S. Rao, D.J. Smith, M.J. Wheelock, K.R. Johnson, G.P. Hemstreet, K.C. Balaji, E-cadherin phosphorylation by protein kinase D1/protein kinase C(mu) is associated with altered cellular aggregation and motility in prostate cancer, *Cancer Res.* 65 (2005) 483–492.
- [128] E.R. Sharlow, K.V. Giridhar, C.R. LaValle, J. Chen, S. Leimgruber, R. Barrett, K. Bravo-Altamirano, P. Wipf, J.S. Lazo, Q.J. Wang, Potent and selective disruption of protein kinase D functionality by a benzoxazolepinolone, *J. Biol. Chem.* 283 (2008) 33516–33526.
- [129] E.C. Nelson, C.P. Evans, P.C. Mack, R.W. Devere-White, P.N. Lara Jr., Inhibition of Akt pathways in the treatment of prostate cancer, *Prostate Cancer Prostatic Dis.* 10 (2007) 331–339.
- [130] D. Sarker, A.H. Reid, T.A. Yap, J.S. de Bono, Targeting the PI3K/AKT pathway for the treatment of prostate cancer, *Clin. Cancer Res.* 15 (2009) 4799–4805.
- [131] J. Chen, K.V. Giridhar, L. Zhang, S. Xu, Q.J. Wang, A protein kinase C/protein kinase D pathway protects LNCaP prostate cancer cells from phorbol ester-induced apoptosis by promoting ERK1/2 and NF-kappaB activities, *Carcinogenesis* 32 (2011) 1198–1206.
- [132] L. Zhang, Z. Zhao, S. Xu, M. Tandon, C.R. LaValle, F. Deng, Q.J. Wang, Androgen suppresses protein kinase D1 expression through fibroblast growth factor receptor substrate 2 in prostate cancer cells, *Oncotarget* 8 (2017) 12800–12811.
- [133] J. Guo, D.M. Clausen, J.H. Beumer, M.A. Parise, M.J. Egorin, K. Bravo-Altamirano, P. Wipf, E.R. Sharlow, Q.J. Wang, J.L. Eiseman, In vitro cytotoxicity, pharmacokinetics, tissue distribution, and metabolism of small-molecule protein kinase D inhibitors, kb-NB142-70 and kb-NB165-09, in mice bearing human cancer xenografts, *Cancer Chemother. Pharmacol.* 71 (2013) 331–344.
- [134] M. Tandon, J. Johnson, Z. Li, S. Xu, P. Wipf, Q.J. Wang, New pyrazolopyrimidine inhibitors of protein kinase D as potent anticancer agents for prostate cancer cells, *PLoS One* 8 (2013) e75601.
- [135] M. Tandon, J.M. Salamoun, E.J. Carder, E. Farber, S. Xu, F. Deng, H. Tang, P. Wipf, Q.J. Wang, SD-208, a novel protein kinase D inhibitor, blocks prostate cancer cell proliferation and tumor growth in vivo by inducing G2/M cell cycle arrest, *PLoS One* 10 (2015) e0119346.
- [136] M. Tandon, L. Wang, Q. Xu, X. Xie, P. Wipf, Q.J. Wang, A targeted library screen reveals a new inhibitor scaffold for protein kinase D, *PLoS One* 7 (2012) e44653.
- [137] E.T. Bowden, M. Barth, D. Thomas, R.I. Glazer, S.C. Mueller, An invasion-related complex of cortactin, paxillin and PKCmu associates with invadopodia at sites of extracellular matrix degradation, *Oncogene* 18 (1999) 4440–4449.
- [138] S. Borges, H. Doppler, E.A. Perez, C.A. Andorfer, Z. Sun, P.Z. Anastasiadis, E. Thompson, X.J. Geiger, P. Storz, Pharmacologic reversal of epigenetic silencing of the PRKD1 promoter blocks breast tumor cell invasion and metastasis, *Breast Cancer Res.* 15 (2013) R66.
- [139] C. Du, C. Zhang, S. Hassan, M.H. Biswas, K.C. Balaji, Protein kinase D1 suppresses epithelial-to-mesenchymal transition through phosphorylation of snail, *Cancer Res.* 70 (2010) 7810–7819.
- [140] L.L. Bastea, H. Doppler, B. Balogun, P. Storz, Protein kinase D1 maintains the epithelial phenotype by inducing a DNA-bound, inactive SNAIL transcriptional repressor complex, *PLoS One* 7 (2012) e30459.
- [141] T. Eiseler, M.A. Schmid, F. Topbas, K. Pfizenmaier, A. Hauser, PKD is recruited to sites of actin remodelling at the leading edge and negatively regulates cell migration, *FEBS Lett.* 581 (2007) 4279–4287.
- [142] R. Palmantier, M.D. George, S.K. Akiyama, F.M. Wolber, K. Olden, J.D. Roberts, Cis-polyunsaturated fatty acids stimulate beta1 integrin-mediated adhesion of human breast carcinoma cells to type IV collagen by activating protein kinases C-epsilon and -mu, *Cancer Res.* 61 (2001) 2445–2452.
- [143] S.B. Kennett, J.D. Roberts, K. Olden, Requirement of protein kinase C micro activation and calpain-mediated proteolysis for arachidonic acid-stimulated adhesion of MDA-MB-435 human mammary carcinoma cells to collagen type IV, *J. Biol. Chem.* 279 (2004) 3300–3307.
- [144] Q. Hao, R. McKenzie, H. Gan, H. Tang, Protein kinases D2 and D3 are novel growth regulators in HCC1806 triple-negative breast cancer cells, *Anticancer Res.* 33 (2013) 393–399.

- [145] J. Chen, L. Lu, Y. Feng, H. Wang, L. Dai, Y. Li, P. Zhang, PKD2 mediates multi-drug resistance in breast cancer cells through modulation of P-glycoprotein expression, *Cancer Lett.* 300 (2011) 48–56.
- [146] S. Borges, P. Storz, Protein kinase D isoforms: new targets for therapy in invasive breast cancers? *Expert. Rev. Anticancer Ther.* 13 (2013) 895–898.
- [147] B. Huck, S. Duss, A. Hausser, M.A. Olayioye, Elevated protein kinase D3 (PKD3) expression supports proliferation of triple-negative breast cancer cells and contributes to mTORC1-S6 K1 pathway activation, *J. Biol. Chem.* 289 (2014) 3138–3147.
- [148] S. Borges, E.A. Perez, E.A. Thompson, D.C. Radisky, X.J. Geiger, P. Storz, Effective targeting of estrogen receptor-negative breast cancers with the protein kinase D inhibitor CRT0066101, *Mol. Cancer Ther.* 14 (2015) 1306–1316.
- [149] P.A. Philip, Targeted therapies for pancreatic cancer, *Gastrointest. Cancer Res.* 2 (2008) S16–S19.
- [150] N. Bardeesy, R.A. DePinho, Pancreatic cancer biology and genetics, *Nat. Rev. Cancer* 2 (2002) 897–909.
- [151] M. Korc, Role of growth factors in pancreatic cancer, *Surg. Oncol. Clin. N. Am.* 7 (1998) 25–41.
- [152] R.H. Hruban, R.E. Wilentz, M. Goggins, G.J. Offerhaus, C.J. Yeo, S.E. Kern, Pathology of incipient pancreatic cancer, *Ann. Oncol.* 10 (Suppl. 4) (1999) 9–11.
- [153] A. Trauzold, S. Schmiedel, B. Sipos, H. Wermann, S. Westphal, C. Roder, W. Klapper, A. Arlt, L. Lehnert, H. Ungefroren, F.J. Johannes, H. Kalthoff, PKCmu prevents CD95-mediated apoptosis and enhances proliferation in pancreatic tumour cells, *Oncogene* 22 (2003) 8939–8947.
- [154] G.Y. Liou, H. Doppler, U.B. Braun, R. Panayiotou, M. Scotti Buzhardt, D.C. Radisky, H.C. Crawford, A.P. Fields, N.R. Murray, Q.J. Wang, M. Leitges, P. Storz, Protein kinase D1 drives pancreatic acinar cell reprogramming and progression to intraepithelial neoplasia, *Nat. Commun.* 6 (2015) 6200.
- [155] S. Guha, J.A. Lunn, C. Santiskulvong, E. Rozengurt, Neurotensin stimulates protein kinase C-dependent mitogenic signaling in human pancreatic carcinoma cell line PANC-1, *Cancer Res.* 63 (2003) 2379–2387.
- [156] S. Guha, O. Rey, E. Rozengurt, Neurotensin induces protein kinase C-dependent protein kinase D activation and DNA synthesis in human pancreatic carcinoma cell line PANC-1, *Cancer Res.* 62 (2002) 1632–1640.
- [157] J. Yuan, E. Rozengurt, PKD, PKD2, and p38 MAPK mediate Hsp27 serine-82 phosphorylation induced by neurotensin in pancreatic cancer PANC-1 cells, *J. Cell. Biochem.* 103 (2008) 648–662.
- [158] L. Xu, R.C. Bergan, Genistein inhibits matrix metalloproteinase type 2 activation and prostate cancer cell invasion by blocking the transforming growth factor beta-mediated activation of mitogen-activated protein kinase-activated protein kinase 2-27-kDa heat shock protein pathway, *Mol. Pharmacol.* 70 (2006) 869–877.
- [159] K.D. Shin, M.Y. Lee, D.S. Shin, S. Lee, K.H. Son, S. Koh, Y.K. Paik, B.M. Kwon, D.C. Han, Blocking tumor cell migration and invasion with biphenyl isoxazole derivative KRIBB3, a synthetic molecule that inhibits Hsp27 phosphorylation, *J. Biol. Chem.* 280 (2005) 41439–41448.
- [160] A.K. McCollum, C.J. Teneyck, B.M. Sauer, D.O. Toft, C. Erlichman, Up-regulation of heat shock protein 27 induces resistance to 17-allylamino-demethoxygeldanamycin through a glutathione-mediated mechanism, *Cancer Res.* 66 (2006) 10967–10975.
- [161] Y. Chen, A.P. Arrigo, R.W. Currie, Heat shock treatment suppresses angiotensin II-induced activation of NF-kappaB pathway and heart inflammation: a role for IKK depletion by heat shock? *Am. J. Physiol. Heart Circ. Physiol.* 287 (2004) H1104–H1114.
- [162] C. Wille, C. Kohler, M. Armacki, A. Jamali, U. Gossele, K. Pfizenmaier, T. Seufferlein, T. Eiseler, Protein kinase D2 induces invasion of pancreatic cancer cells by regulating matrix metalloproteinases, *Mol. Biol. Cell* 25 (2014) 324–336.
- [163] J. Rennecke, P.A. Rehberger, G. Furstenberger, F.J. Johannes, M. Stohr, F. Marks, K.H. Richter, Protein-kinase-Cmu expression correlates with enhanced keratinocyte proliferation in normal and neoplastic mouse epidermis and in cell culture, *Int. J. Cancer* 80 (1999) 98–103.
- [164] V.L. Ristich, P.H. Bowman, M.E. Dodd, W.B. Bollag, Protein kinase D distribution in normal human epidermis, basal cell carcinoma and psoriasis, *Br. J. Dermatol.* 154 (2006) 586–593.
- [165] P. Ivanova, G. Atanasova, Y. Poumay, V. Mitev, Knockdown of PKD1 in normal human epidermal keratinocytes increases mRNA expression of keratin 10 and involucrin: early markers of keratinocyte differentiation, *Arch. Dermatol. Res.* 300 (2008) 139–145.
- [166] A. Jadali, S. Ghazizadeh, Protein kinase D is implicated in the reversible commitment to differentiation in primary cultures of mouse keratinocytes, *J. Biol. Chem.* 285 (2010) 23387–23397.
- [167] S.N. Arun, I. Kaddour-Djebbar, B.A. Shapiro, W.B. Bollag, Ultraviolet B irradiation and activation of protein kinase D in primary mouse epidermal keratinocytes, *Oncogene* 30 (2011) 1586–1596.
- [168] M. Rashel, N. Alston, S. Ghazizadeh, Protein kinase D1 has a key role in wound healing and skin carcinogenesis, *J. Invest. Dermatol.* 134 (2014) 902–909.
- [169] V. Ryykin, M. Rashel, T. Gaddapara, S. Ghazizadeh, Opposing growth regulatory roles of protein kinase D isoforms in human keratinocytes, *J. Biol. Chem.* 290 (2015) 11199–11208.
- [170] M. Kim, H.R. Jang, J.H. Kim, S.M. Noh, K.S. Song, J.S. Cho, H.Y. Jeong, J.C. Norman, P.T. Caswell, G.H. Kang, S.Y. Kim, H.S. Yoo, Y.S. Kim, Epigenetic inactivation of protein kinase D1 in gastric cancer and its role in gastric cancer cell migration and invasion, *Carcinogenesis* 29 (2008) 629–637.
- [171] M.Y. Shabelnik, L.M. Kovalevska, M.Y. Yurchenko, L.M. Shlapatska, Y. Rzepetsky, S.P. Sidorenko, Differential expression of PKD1 and PKD2 in gastric cancer and analysis of PKD1 and PKD2 function in the model system, *Exp. Oncol.* 33 (2011) 206–211.
- [172] L. Paolucci, E. Rozengurt, Protein kinase D in small cell lung cancer cells: rapid activation through protein kinase C, *Cancer Res.* 59 (1999) 572–577.
- [173] W. Brenner, S. Beitz, E. Schneider, F. Benzing, R.E. Unger, F.C. Roos, J.W. Thuroff, C. Hampel, Adhesion of renal carcinoma cells to endothelial cells depends on PKCmu, *BMC Cancer* 10 (2010) 183.
- [174] L.M. Kovalevska, O.V. Yurchenko, L.M. Shlapatska, G.G. Berdova, S.V. Mikhailap, J. Van Lint, S.P. Sidorenko, Immunohistochemical studies of protein kinase D (PKD) 2 expression in malignant human lymphomas, *Exp. Oncol.* 28 (2006) 225–230.
- [175] E. Bernhart, S. Damm, A. Wintersperger, T. DeVaney, A. Zimmer, T. Raynham, C. Ireson, W. Sattler, Protein kinase D2 regulates migration and invasion of U87MG glioblastoma cells in vitro, *Exp. Cell Res.* 319 (2013) 2037–2048.
- [176] V. Sundram, A. Ganju, J.E. Hughes, S. Khan, S.C. Chauhan, M. Jaggi, Protein kinase D1 attenuates tumorigenesis in colon cancer by modulating beta-catenin/T cell factor activity, *Oncotarget* 5 (2014) 6867–6884.
- [177] N. Wei, E. Chu, P. Wipf, J.C. Schmitz, Protein kinase d as a potential chemotherapeutic target for colorectal cancer, *Mol. Cancer Ther.* 13 (2014) 1130–1141.
- [178] K. Bravo-Altamirano, K.M. George, M.C. Frantz, C.R. Laval, M. Tandon, S. Leimgruber, E.R. Sharlow, J.S. Lazo, Q.J. Wang, P. Wipf, Synthesis and structure-activity relationships of benzothienothiazepinone inhibitors of protein kinase D, *ACS Med. Chem. Lett.* 2 (2011) 154–159.
- [179] K.M. George, M.C. Frantz, K. Bravo-Altamirano, C.R. Laval, M. Tandon, S. Leimgruber, E.R. Sharlow, J.S. Lazo, Q.J. Wang, P. Wipf, Design, synthesis, and biological evaluation of PKD inhibitors, *Pharmaceuticals* 3 (2011) 186–228.
- [180] C.R. Laval, K. Bravo-Altamirano, K.V. Giridhar, J. Chen, E. Sharlow, J.S. Lazo, P. Wipf, Q.J. Wang, Novel protein kinase D inhibitors cause potent arrest in prostate cancer cell growth and motility, *BMC Chem. Biol.* 10 (2010) 5.
- [181] E.R. Sharlow, G. Mustata Wilson, D. Close, S. Leimgruber, M. Tandon, R.B. Reed, T.Y. Shun, Q.J. Wang, P. Wipf, J.S. Lazo, Discovery of diverse small molecule chemotypes with cell-based PKD1 inhibitory activity, *PLoS One* 6 (2011) e25134.
- [182] J. Yuan, Y. Liu, T. Tan, S. Guha, I. Gukovsky, A. Gukovskaya, S.J. Pandol, Protein kinase d regulates cell death pathways in experimental pancreatitis, *Front. Physiol.* 3 (2012) 60.
- [183] L. Monovich, R.B. Vega, E. Meredith, K. Miranda, C. Rao, M. Capparelli, D.D. Lemon, D. Phan, K.A. Koch, J.A. Chapo, D.B. Hood, T.A. McKinsey, A novel kinase inhibitor establishes a predominant role for protein kinase D as a cardiac class IIa histone deacetylase kinase, *FEBS Lett.* 584 (2010) 631–637.
- [184] E.L. Meredith, O. Ardayfio, K. Beattie, M.R. Dobler, I. Enyedy, C. Gaul, V. Hosagrahara, C. Jewell, K. Koch, W. Lee, H. Lehmann, T.A. McKinsey, K. Miranda, N. Pagratis, M. Pancost, A. Patnaik, D. Phan, C. Plato, M. Qian, V. Rajaraman, C. Rao, O. Rozhitskaya, T. Ruppen, J. Shi, S.J. Siska, C. Springer, M. van Eis, R.B. Vega, A. von Matt, L. Yang, T. Yoon, J.H. Zhang, N. Zhu, L.G. Monovich, Identification of orally available naphthyridine protein kinase D inhibitors, *J. Med. Chem.* 53 (2010) 5400–5421.
- [185] G.G. Gamber, E. Meredith, Q. Zhu, W. Yan, C. Rao, M. Capparelli, R. Burgis, I. Enyedy, J.H. Zhang, N. Soldermann, K. Beattie, O. Rozhitskaya, K.A. Koch, N. Pagratis, V. Hosagrahara, R.B. Vega, T.A. McKinsey, L. Monovich, 3,5-Diarylazoles as novel and selective inhibitors of protein kinase D, *Bioorg. Med. Chem. Lett.* 21 (2011) 1447–1451.
- [186] K.B. Harikumar, A.B. Kunnumakkara, N. Ochi, Z. Tong, A. Deorukhkar, B. Sung, L. Kelland, S. Jamieson, R. Sutherland, T. Raynham, M. Charles, A. Bagherzadeh, C. Foxton, A. Boakes, M. Farooq, D. Maru, P. Diagaradjane, Y. Matsuo, J. Sinnett-Smith, J. Gelovani, S. Krishnan, B.B. Aggarwal, E. Rozengurt, C.R. Ireson, S. Guha, A novel small-molecule inhibitor of protein kinase D blocks pancreatic cancer growth in vitro and in vivo, *Mol. Cancer Ther.* 9 (2010) 1136–1146.
- [187] I.M. Evans, A. Bagherzadeh, M. Charles, T. Raynham, C. Ireson, A. Boakes, L. Kelland, I.C. Zachary, Characterization of the biological effects of a novel protein kinase D inhibitor in endothelial cells, *Biochem. J.* 429 (2010) 565–572.
- [188] M. Golkowski, R.S. Vidadala, C.K. Lombard, H.W. Suh, D.J. Maly, S.E. Ong, Kinobead and single-shot LC-MS profiling identifies selective PKD inhibitors, *J. Proteome Res.* 16 (2017) 1216–1227.
- [189] K.B. Harikumar, A.B. Kunnumakkara, N. Ochi, Z. Tong, A. Deorukhkar, B. Sung, L. Kelland, S. Jamieson, R. Sutherland, T. Raynham, M. Charles, A. Bagherzadeh, C. Foxton, A. Boakes, M. Farooq, D. Maru, P. Diagaradjane, Y. Matsuo, J. Sinnett-Smith, J. Gelovani, S. Krishnan, B.B. Aggarwal, E. Rozengurt, C.R. Ireson, S. Guha, A novel small-molecule inhibitor of protein kinase D blocks pancreatic cancer growth in vitro and in vivo, *Mol. Cancer Ther.* 9 (2010) 1136–1146.
- [190] C.D. Chen, C.L. Sawyers, NF-kappa B activates prostate-specific antigen expression and is upregulated in androgen-independent prostate cancer, *Mol. Cell. Biol.* 22 (2002) 2862–2870.

Protein Kinase D2 Modulates Cell Cycle By Stabilizing Aurora A Kinase at Centrosomes

Adhiraj Roy¹, Maria Victoria Veroli¹, Sahdeo Prasad^{1,2}, and Qiming Jane Wang¹



Abstract

Aurora A kinase (AURKA) is a master cell-cycle regulator that is often dysregulated in human cancers. Its overexpression has been associated with genome instability and oncogenic transformation. The protein kinase D (PKD) family is an emerging therapeutic target of cancer. Aberrant PKD activation has been implicated in tumor growth and survival, yet the underlying mechanisms remain to be elucidated. This study identified, for the first time, a functional crosstalk between PKD2 and Aurora A kinase in cancer cells. The data demonstrate that PKD2 is catalytically active during the G₂–M phases of the cell cycle, and inactivation or depletion of PKD2 causes delay in mitotic entry due to downregulation of Aurora A, an effect that can be rescued by overexpression of Aurora A. Moreover, PKD2 localizes in the

centrosome with Aurora A by binding to γ -tubulin. Knockdown of PKD2 caused defects in centrosome separation, elongated G₂ phase, mitotic catastrophe, and eventually cell death via apoptosis. Mechanistically, PKD2 interferes with Fbxw7 function to protect Aurora A from ubiquitin- and proteasome-dependent degradation. Taken together, these results identify PKD as a cell-cycle checkpoint kinase that positively modulates G₂–M transition through Aurora A kinase in mammalian cells.

Implications: PKD2 is a novel cell-cycle regulator that promotes G₂–M transition by modulating Aurora A kinase stability in cancer cells and suggests the PKD2/Aurora A kinase regulatory axis as new therapeutic targets for cancer treatment. *Mol Cancer Res*; 1–13. ©2018 AACR.

Introduction

Cell cycle is propelled by well-coordinated complex signaling events by which cells grow and divide. Uncontrolled cell division, whereby cells go through the cycle unchecked, lies at the base of cancer formation and progression. Cell cycle occurs through progression of four distinct stages, namely G₀–G₁, S, G₂, and M, which are monitored by a battery of cyclin-dependent kinases (CDK) and their partner cyclins (1). Mutations in the signaling pathways, aberrant activation of CDKs, genetic lesions in the genes encoding cell-cycle-regulatory proteins result in genome instability, abnormal growth, and eventually cancer.

The protein kinase D (PKD) family of serine/threonine kinases belongs to the Ca⁺⁺/Calmodulin-dependent protein kinase (CaMK) superfamily and it consists of three isoforms in mammals, notably, PKD1, PKD2, and PKD3, which are highly conserved throughout evolution (2). Structurally, PKD possesses an N-terminal regulatory domain that contains a tandem cysteine-rich Zn-finger like motif (CRD, C1a, and C1b) and a pleckstrin homology (PH) domain, and a C-terminal catalytic domain (3, 4). In a canonical pathway, diverse physiologic factors, such as GPCR agonists, bioactive peptides (5), lipids (6), and growth factors (7) converge to the activation of PKDs through the gen-

eration of diacylglycerol (DAG) by phospholipase C (PLC) and the activation of classical or novel protein kinase C (c/nPKC; ref. 8). Activation of PKD is marked by phosphorylation of two conserved serine residues in the activation loop of PKDs and concomitant autophosphorylation of PKD1^{Ser916} or PKD2^{Ser876} are widely used as a biomarker of PKD activation (3, 9). PKD regulates a wide range of cellular processes including cell proliferation, migration/invasion, angiogenesis, protein trafficking, and gene expression. Its functional importance has been implicated in major human cancers including carcinomas of breast, skin, pancreas, and prostate (8).

Aurora kinases are master regulators of the cell cycle (10–13). Human genome encodes three isoforms of Aurora kinases; namely, Aurora kinases A, B, and C (14). Aurora A localizes on centrosomes, spindle poles, and spindles during mitosis and regulates centrosome function, spindle assembly, and mitotic progression. Expression of Aurora A is cell-cycle-regulated, that is, the levels of mRNA and protein are low in G₁ and S, increase during G₂–M, and reduce during mitotic exit. Aurora A is ubiquitinated by APC/Cdh1 and Fbxw7 ubiquitin ligases and degraded in proteasome-dependent pathways (15, 16). A number of Aurora A substrates have been reported, such as TPX2, CDC25B, and p53 (17). Aurora A is frequently overexpressed in cancer. Aberrant expression of Aurora A promotes centrosome amplification and aneuploidy, leading to genome instability and consequently oncogenic transformation (18). Thus, Aurora A represents a well-established therapeutic target for cancer. In this study, we uncovered a novel crosstalk between PKD2 and Aurora A kinase. Our study defined the functional impact of this regulatory axis on G₂–M transition and mitosis. We have shown that PKD2 is required for stabilization of Aurora A kinase, and that the abrogation of PKD2 expression or inhibition of its catalytic activity causes proteasome-mediated downregulation of Aurora A kinase displaying delayed G₂, mitotic catastrophe and cell death. Moreover, we report that

¹Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania. ²Department of Biotechnology and Immunotherapeutics, Texas Tech University, Amarillo, Texas.

Corresponding Author: Qiming Jane Wang, Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, E1354 BST, 200 Lothrop Street, Pittsburgh, PA 15261. Phone: 412-383-7754; Fax: 412-641-495; E-mail: qjw1@pitt.edu.

doi: 10.1158/1541-7786.MCR-18-0641

©2018 American Association for Cancer Research.

PKD2 localizes on the centrosomes during G₂ and that it is critical for centrosome separation in early G₂. Taken together, our study provides compelling evidence to support the role of PKD2 in G₂-M transition through modulating Aurora A stability, and aberrant PKD2 activity may contribute to oncogenesis.

Materials and Methods

Cell culture, synchronization, and treatments

Authenticated HeLa, LNCaP, and PC3 cells were obtained from ATCC. After purchase, the cell lines were expanded and frozen according to manufacturer's instructions after two to three passages. LNCaP was used for no more than 8 passages and PC3/HeLa cells were used for no more than 10 passages. All cell cultures were routinely tested for *Mycoplasma* using MycoAlert PLUS Mycoplasma Detection Kit (Lonza). HeLa, LNCaP, and PC3 cells were grown in 1× Minimum Essential Medium (MEM) supplemented with Eagle salt and L-glutamine (Invitrogen), RPMI1640 (Invitrogen), or Ham F-12 (Thermo Fisher Scientific) media, respectively, supplemented with 10% FBS and 1× penicillin/streptomycin (Thermo Fisher Scientific, MT30002CI) and maintained at 37°C in a humidified incubator containing 5% CO₂. Cells were synchronized at G₁-S or the start of M phases by double thymidine or nocodazole, respectively. Briefly, for G₁-S synchronization, cells were treated with 2 mmol/L thymidine for 16 hours, washed three times with 1× PBS, and refed into fresh culture medium without thymidine for 8 hours. The second block was performed by treating the cells with 2 mmol/L thymidine for another 16 hours. Cell arrest at the start of M phase was performed by treating the cells with nocodazole (100 ng/mL) for 16 hours. PKD inhibitors (CRT0066101 and kb-NB142-70) were used at a final concentration of 2 μmol/L for indicated times. Turbofect (Thermo Fisher Scientific) and Lipofectamine 3000 (Invitrogen) transfection reagents were used to transfect plasmids and siRNAs, respectively, according to the manufacturer's protocol.

Cell survival assay

Cell survival assay was performed by Cell Counting Kit-8 (Dojindo Laboratories) according to the manufacturer's protocol. Briefly, HeLa and PC3 cells were plated at a density of 4,000 cells per well in 96-well plate. After treating the cells with different doses of drugs (CRT0066101 and Kb-NB142-70) for 72 hours, CCK-8 solution was added to cells in each well, followed by incubation for 2 hours. Cell proliferation/viability was determined by measuring the OD at 450 nm.

Western blotting and densitometry analysis

Cells were lysed in cell lysis buffer (50 mmol/L Tris-HCl pH 7.5, 150 mmol/L NaCl, 1.5 mmol/L MgCl₂, 10% glycerol, 1% Triton X-100, 5 mmol/L EGTA, 1 mmol/L Na₃VO₄, 10 mmol/L NaF, 1 mmol/L β-glycerophosphate and protease inhibitor cocktail). Cell extracts were resolved by SDS-PAGE, transferred to nitrocellulose membrane, and probed with following antibodies: PKD2 (Cell Signaling Technology, 8188), phospho-PKD2^{Ser876} (Thermo Fisher Scientific, PA5-64538), Cyclin B1 (Cell Signaling Technology, 12231), phospho CDC25^{Thr48} (Cell Signaling Technology 12028), phospho-Histone H3^{Ser10} (Cell Signaling Technology, 3377), Aurora Kinase A (Cell Signaling Technology, 14475), Aurora Kinase B (Cell Signaling Technology, 3094), γ-tubulin (Santa Cruz Biotechnology, sc-17788), PARP (Cell

Signaling Technology, 9542), ubiquitin (P4D1, Cell Signaling Technology 3936), Lamin A (Cell Signaling Technology, 2032), Cdh1 (Santa Cruz Biotechnology, sc-56312), Fbxw7 (Abcam, EPR8069), and GAPDH (Enzo ADI-CSA-335-E). Protein half-life was measured as described previously (19). Briefly, the band intensities were quantified by ImageJ software and the values of target proteins were normalized against the band intensity of GAPDH. The measured protein intensity values were log transformed, fitted to a linear regression plot, and decay rate constant (*k*) was determined. The half-life was calculated from *k* using the formula $T_{1/2} = \ln(2)/k$.

Immunoprecipitation analysis

For immunoprecipitation assay, whole-cell extract was prepared by lysing the cells with cell lysis buffer. The lysate was centrifuged, supernatant was collected, and protein concentration was estimated using BCA protein assay kit (Pierce) according to manufacturer's protocol. Cell lysate containing equal amount of total protein was incubated with primary antibodies at 4°C overnight. Protein A/G agarose beads were added later for another 2 hours. The immunoprecipitates were collected and washed with wash buffer containing 0.5% Triton X-100. Finally, the samples were diluted with 6× SDS PAGE loading buffer, boiled for 10 minutes, and subjected to Western blotting.

Subcellular fractionation

Cells were washed with ice-cold PBS, collected by centrifugation at 1,000 rpm for 5 minutes, and resuspended in cytoplasm extraction buffer (10 mmol/L HEPES pH 7.5, 60 mmol/L KCl, 1 mmol/L EDTA, 0.1% NP-40, 1 mmol/L DTT, and protease inhibitor cocktail) for 5 minutes on ice. The cell lysate was centrifuged at 1,000 rpm for 5 minutes at 4°C and cytoplasmic fraction was separated in a new tube. The nuclei were washed in cytoplasm extraction buffer without NP-40 and resuspended in nucleus extraction buffer (20 mmol/L Tris-HCl pH 8, 400 mmol/L NaCl, 1.5 mmol/L MgCl₂, 0.2 mmol/L EDTA, 25% glycerol and protease inhibitor cocktail). The extracts were incubated on ice for 10 minutes. Both cytoplasmic and nuclear extracts were centrifuged at 14,000 rpm for 15 minutes at 4°C and transferred to new tubes.

Flow cytometry

Trypsinized cells were pelleted, washed two times with ice-cold FACS buffer (0.1% glucose in PBS), and resuspended in ice cold 70% ethanol while vortexing. Cells were fixed by incubating overnight at 4°C, washed with PBS, and incubated with 1 mL propidium iodide (PI) staining solution (50 μg/mL PI, 0.1 mg/mL RNase A in FACS buffer) for 30 minutes. Cells were analyzed in a FACSCalibur (BD Biosciences) flow cytometer and the data were analyzed by ModFit 3.0 software (BD Biosciences). A total of 10,000 events were analyzed for each sample.

Apoptosis assay

Trypsinized cells were washed twice with ice-cold PBS and stained with Annexin V-Alexa Fluor 488 antibody and propidium iodide (Molecular Probes, Thermo Fisher Scientific) according to the manufacturer's instructions. The samples were divided in two parts. One half was subjected to FACSCalibur (BD Biosciences) flow cytometer and the other part was subjected to confocal fluorescence microscopy using Alexa Fluor 488 and PI filters.

Immunofluorescence staining and microscopy

HeLa and PC3 cells were grown on poly-D-lysine-coated coverslips, washed with PBS, fixed in 4% paraformaldehyde at room temperature for 30 minutes, washed three times with 1× PBS, and blocked in 5% normal goat serum containing 0.3% Triton X-100 for 1 hour at room temperature. The cells were incubated with the primary antibodies diluted in antibody dilution buffer (1% BSA, 0.3% Triton X-100 in 1× PBS) overnight at 4°C, washed three times in 1× PBS, and incubated with secondary antibodies diluted in antibody dilution buffer for 1 hour at room temperature. The cells were counterstained with DAPI (1 µg/mL) and the coverslips were mounted in ProLong Gold antifade reagent (Invitrogen). The cells were analyzed using an Olympus Fluoview (FV1000) confocal microscope using 60 X/1.45 objectives.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Statistical analysis

Data analysis was done using the Student *t* test for comparison between two groups (two-tailed). All statistical analyses were done using GraphPad Prism IV software (GraphPad Software). All values are represented as mean ± SEM of at least three independent experiments. A *P* < 0.05 was considered statistically significant (*, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001; ****, *P* < 0.0001; ns, not significant).

Results

Elevated PKD2 activity during mitotic entry of the cell cycle


First, we examined the expression and activity of PKD (refers to PKD2 and PKD3 throughout the text) in different cell-cycle stages. PKD2 and PKD3 are major isoforms expressed in HeLa cells (Supplementary Fig. S1), similar to the prostate cancer cell line PC3, but differed from LNCaP cells (20). HeLa and PC3 cells were treated with nocodazole, a microtubule-destabilizing agent that activates the spindle assembly checkpoint and causes cell arrest in start of M-phase (prometaphase), and the activity of PKD2 was assessed after releasing from nocodazole using the phospho-PKD2^{Ser876} (p-PKD2) antibody. As shown in Fig. 1A (top) and B, p-PKD2 was detected in G₂-M-arrested HeLa cells following nocodazole synchronization and persisted till the end of mitosis. This was accompanied by increased cyclin B1, a marker for G₂ and M phases of the cell cycle, and phospho-Histone H3^{Ser10} (pHH3), which was detectable in G₂, peaks during mitosis and disappears when cells enters in G₁ (Fig. 1A; Supplementary Fig. S2). Similar results were obtained with PC3 cells where activation of PKD2 was marked by increased levels of p-PKD2 after release from nocodazole arrest (Fig. 1A, bottom). To determine whether the activation is specific to G₂-M, PKD2 activation was measured in G₁-S synchronized HeLa cells that progressed to M-phase. As shown in Fig. 1C and D, when cells were released from a double thymidine (DT) block that synchronizes cells at G₁-S border, PKD2 phosphorylation gradually elevated, starting at 4 hours when cells entered in late S-phase peaked around 10 hours in late G₂, and persisted throughout mitosis till 14 hours after thymidine release. The highest activity was detected at G₂-M border and M-phase. This pattern correlated well with increased cyclin B1 and pHH3. As controls, PKD2 remained the same under both conditions.

The intracellular distribution of PKD2 was then analyzed at different stages of the cell cycle by immunofluorescence (IF) staining. In HeLa cells, PKD2 was primarily localized in the cytoplasm and perinuclear zone in interphase. When cells progress through cell cycle, it exhibited scattered punctate staining that are minimally overlapped with mitotic bodies (spindle fibers and midbody) and excluded from condensed chromatin (Fig. 1E, top). PKD2 phosphorylation was low or undetectable during interphase and significantly increased during G₂, sustained throughout prophase, metaphase, and anaphase, before returning to baseline when the cells entered telophase. To corroborate these results in cancer, PC3 cells were stained in similar fashion. Similar results were found when different stages of the cell cycle in PC3 cells were imaged using antibodies against phospho-PKD2^{Ser876} (green) and α -tubulin (red; Fig. 1F). It is well documented that PKD, upon activation, accumulates in the nucleus (21). Indeed, we observed elevated active PKD in the nucleus during G₂ and early prophase when the nucleus was still intact, further attesting that PKD2 was activated during G₂ and mitosis. We further validated that phospho-PKD2 was increased in the nucleus during early mitosis by cell fractionation (Fig. 1G, left). HeLa and PC3 cells were arrested at the start of M phase by nocodazole and cytoplasmic (Cyto) or nuclear (Nuc) fractions were prepared. The samples were analyzed by Western blotting using phospho-PKD2 antibody and the results showed that, indeed, active PKD2 levels were elevated in early mitosis (Fig. 1G, right). Lamin A was blotted as a marker for the nuclear fraction. Taken together, these results suggested that PKD is active during G₂ and mitosis and may play an important role in cell-cycle regulation.


Inactivation or knockdown of PKD2 causes delay in G₂-M transition

The activation profile of PKD2 suggests that PKD2 may regulate G₂ and M-phase cell-cycle progression. To test this hypothesis, cells with altered PKD2 expression and activity were synchronized by DT at G₁-S border and released (Figs. 2A and 3A); cell-cycle progression was analyzed by flow cytometry. As shown in Fig. 2B and C, after thymidine release, cells reached G₂-M at 8 hours, and by 12 hours, cells resumed normal distribution. In contrast, cells treated with the PKD inhibitor CRT0066101 (CRT101) reached G₂-M at 8 hours, stayed after 12 hours, and by 14 hours there remained a large G₂-M population as compared with the control. Inactivation of PKD by another PKD inhibitor kb-NB142-70 (kb-NB) gave rise to similar results (Fig. 2B, kb-NB). Quantification analysis confirmed prolonged G₂-M phase of cell cycle in cells treated with PKC inhibitors (Fig. 2C). Thus, inactivation PKD2 prolonged G₂-M phase and resulted in delayed mitotic entry in cells.

To determine that PKD2 activity was indeed required for G₂-M transition, cells transfected with PKD2 siRNAs were synchronized to G₁-S border by DT and released; cell-cycle progression was monitored by flow cytometry (Fig. 3A). Our data showed that cells transfected with nontargeting siRNA reached G₂-M around 8 hours and returned to normal distribution at about 12–14 hours. In contrast, cells transfected with two PKD2 siRNAs (Fig. 3B) progressed slower through the cell cycle and peaked at G₂-M phase around 12 hours and remained till after 14 hours (Fig. 3C), indicating that PKD2-depleted cells progressed through G₂-M boundary at much slower rate than control cells. Knockdown of PKD2 using two siRNAs in these samples were confirmed by Western blotting (Fig. 3D). To corroborate these findings, we

**Figure 1.**

PKD is activated during mitotic entry. **A**, HeLa and PC3 cells were synchronized with nocodazole (100 ng/mL) for 16 hours and mitotic cells were released into fresh medium for indicated times. Cell extracts from each sample were subjected to Western blot analysis to analyze activation of PKD2. **B**, Densitometric quantification of phospho-PKD2^{S876} from the immunoblot of HeLa was performed. **C**, Western blot analysis of indicated proteins after the HeLa cells were released from a double thymidine (DT) block was performed. **D**, Densitometric quantification of phospho-PKD2^{S876} from the immunoblot of **C** was performed. Different phases of cell cycle are marked (G₁, S, G₂, and M). **E**, HeLa cells were subjected to immunofluorescence (IF) staining to visualize colocalization PKD2 or phospho-PKD2^{S876} with α -tubulin during different phases of cell cycle. **F**, PC3 cells were subjected to IF staining to visualize colocalization of phospho-PKD2^{S876} with α -tubulin during different phases of cell cycle. DAPI was used to stain the nucleus. Scale bar, 8 μ m. GAPDH was used as loading control in **A** and **C**. **G**, HeLa and PC3 cells were synchronized with nocodazole (100 ng/mL) for 16 hours and mitotic cells were collected by shake-off. Cytoplasmic (Cyto) and nuclear (Nuc) fractions were prepared and the samples were analyzed by Western blotting with indicated antibodies (left). Relative amount of phospho-PKD2^{S876} was quantified (right). The graphs show average of three independent experiments with error bars representing SEM (**, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; ns, not significant).

**Figure 2.**


Inactivation of PKD by CRT101 and kb-NB induces a delay in G₂-M progression. **A**, Schematic representation of experimental design is shown. **B**, Flow cytometry analysis of cells treated with DMSO, CRT101, and kb-NB were performed according to **A**. **C**, Summarizing graph of the distribution of different phases of cell cycle are shown (right). The graphs show average of three independent experiments with error bars representing SEM.

examined HeLa cells with overexpressed PKD2. As shown in Fig. 3E, overexpression of PKD2 significantly increased the number of cells entering mitosis. Taken together, our data indicate that PKD2 was activated in G₂-M and its activity was required for mitotic entry.

Abrogation of PKD2 expression sensitizes cells to mitotic catastrophe and cell death

Cells experience prolonged G₂ arrest and delayed mitotic entry may eventually enter into mitosis, however, they often suffer from

aberrant mitosis known as mitotic catastrophe, and die subsequently by apoptosis (22). Here, we examined whether mitotic catastrophe was associated with loss/inactivation of PKD2 and delayed G₂-M transition. Knockdown or inactivation of PKD2 using two siRNAs or PKD inhibitor CRT101, respectively, induced a series of mitotic defects including chromosomal misalignment at metaphase plane, abnormal chromosomal arrangement (Fig. 4A, white arrows), and formation of multiple centrosomes (Fig. 4A, red arrows). Quantification analysis indicated that depletion or inactivation of PKD2 significantly increased cell


**Figure 3.**

Abrogation of PKD2 expression delays G₂-M progression. **A**, Schematic representation of experimental design is shown. **B**, Flow cytometry analysis of cells transfected with control nontargeting siRNA (siNT) and two siRNAs against PKD2 was performed according to **A**. **C**, Summarizing graph of the distribution of different phases of cell cycle are shown. The graphs show average of three independent experiments with error bars representing SEM. **D**, Western blot analysis of the samples from **A** was performed to confirm knockdown of PKD2. GAPDH was used as loading control. **E**, PKD2 was ectopically overexpressed using a plasmid expressing Flag-PKD2 (top) and percentage of interphase and mitotic cell population was counted (bottom). The graphs show average of three independent experiments with error bars representing SEM (*, $P < 0.05$; **, $P < 0.01$; ****, $P < 0.0001$).

population with defective mitosis (Fig. 4B). It is well known that mitotic catastrophe leads to apoptosis followed by PARP and Lamin A cleavage (23, 24). Analysis of PKD2-depleted or PKD-inactivated HeLa cells showed expression of cleaved PARP^{Asp214} and Lamin A^{Asp230}, indicative of apoptotic response (Fig. 4C). To further this analysis, cells transfected with siNT and siPKD2s were stained with Annexin V-Alexa Fluor 488/propidium iodide (PI) and analyzed by flow cytometry (Fig. 4D, left). Knockdown of PKD2 caused significant cell death (PI-positive) with elevated apoptotic population (Annexin V position) as compared with the control samples (siNT; Fig. 4D, right). Treatment of both HeLa and PC3 cells using increasing concentrations of CRT101 and kb-NB reduced cell viability in a concentration-dependent manner (Fig. 4E). Taken together, our data indicated that G₂-M deficiencies caused by PKD2 depletion or inhibition could result in mitotic catastrophe and cell death followed by apoptosis.

Inactivation of PKD2 causes degradation of Aurora A, an effect that can be suppressed by overexpression of Aurora A


We showed that knockdown or inactivation of PKD2 caused prolonged G₂-M transition, delay in mitotic entry, and mitotic catastrophe (Figs. 2–4). In the order to gain insights into the role of PKD in cell cycle, we sought to identify its downstream targets. Through a whole-genome RNA-seq analysis conducted on cancer cells treated with CRT101 and kb-NB, we identified AURKA/Aurora A kinase as one of the top candidate genes altered by inhibition of PKD. Aurora A is a master regulator of G₂-M and mitosis (12, 25–27). First, using normal prostate epithelial cell line RWPE-1 and prostate adenocarcinoma (PAC) cell lines, we sought to identify whether expression level of PKD2 correlates with that of Aurora A kinase. As shown in Fig. 5A, we observed positive correlation between PKD2 and Aurora A in these cell lines. Next, we subjected HeLa cells to IF staining and checked whether PKD2 and Aurora A colocalize at different stages of

**Figure 4.**

Knockdown of PKD2 causes mitotic catastrophe followed by cell death. **A**, IF staining of cells transfected with control nontargeting siRNA (siNT) and two siRNAs against PKD2 (top) or DMSO/CRT101 (bottom) was performed to visualize γ -tubulin (green). DAPI was used to stain the nucleus. Red and white arrow heads indicate defects in spindle pole formation and misaligned chromosomes, respectively. **B**, Quantification of cell population harboring mitotic defects after the cells were treated with either si-PKD2 (top) or CRT101 (bottom). **C**, Western blot analysis was done to show cleavage of PARP/Lamin A upon depletion or inactivation of PKD2 in HeLa. GAPDH served as loading control. **D**, Microscopic analysis of cells transfected with siNT or two siPKD2s after costaining with Annexin V-Alexa Fluor 488 and propidium iodide (left). Quantification of cell population positive for Annexin V-Alexa Fluor 488 and PI (right). **E**, HeLa and PC3 cells were treated with different concentrations of PKD inhibitors (CRT101, kb-NB) and cultured for 72 hours. MTT assay was performed to assess cell viability. The graphs show average of three independent experiments with error bars representing SEM (***, $P < 0.001$; ****, $P < 0.0001$).

mitosis. Our confocal microscopy results showed that a significant portion of PKD2 (green) colocalizes with Aurora A (red) during mitosis (Fig. 5B), raising a strong possibility that PKD2 might contribute to cell cycle by regulating Aurora A. We examined the expression levels of Aurora A upon inactivation or depletion of PKD. HeLa cells were arrested at the start of M phase by nocodazole, released from the block by mitotic shake off and turnover of Aurora A protein levels was analyzed in the absence or presence

of CRT101 by Western blotting. The results showed that CRT101 caused rapid downregulation of Aurora A with < 50% remaining 2 hours after CRT101 treatment (Fig. 5C). To corroborate this finding in the context of cancer, we treated PC3 prostate adenocarcinoma and COV362 ovarian epithelial-endometroid carcinoma cell lines as described above and checked Aurora A turnover by Western blotting (Fig. 5D). As expected, Aurora A was rapidly downregulated by CRT101 in these two cell lines.

**Figure 5.**

Inactivation or knockdown of PKD2 causes downregulation of Aurora A kinase. **A**, Western blot analysis was performed to determine expression patterns of PKD2 and Aurora A in designated cell lines. **B**, HeLa cells were stained with antibodies against PKD2 (green) and Aurora A (red). IF staining was performed to visualize colocalization of PKD2 and Aurora A at different stages of cell cycle. Scale bar, 8 μ m. DAPI was used to stain the nucleus. **C**, HeLa cells were synchronized at the start of M-phase by nocodazole, released from the block and mitotic cells were cultured for indicated times with or without CRT101. Downregulation of Aurora A was analyzed by Western blotting (top). Quantification of Aurora A protein levels by densitometry scanning from immunoblots was performed (bottom). **D**, PC3 and COV362 cells were treated as described in **C** and turnover of Aurora A protein levels were analyzed by Western blotting. The expression levels of Aurora B were analyzed as control. **E**, HeLa Cells were arrested at the start of M-phase by nocodazole (noc), released in fresh medium containing DMSO or CRT101 for indicated times and cell extracts from each sample were analyzed by Western blotting using indicated antibodies (left). Aurora A was ectopically overexpressed and the cells were released from nocodazole block in fresh medium containing CRT101 for indicated times. Cell extracts from each sample were analyzed by Western blotting using indicated antibodies. Overexpression of Aurora A was confirmed by Western blotting. **F**, Cells were transfected by si-NT (control) and two different concentrations of si-PKD2. Forty-eight hours post transfection, cells were treated with cycloheximide for indicated times and cell extracts from each sample were analyzed by Western blotting using indicated antibodies (left). GAPDH served as loading control. Half-life of Aurora A protein was measured from the samples of **F** (right). The graphs show average of three independent experiments with error bars representing SEM (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ns, not significant).

Meanwhile, the downregulation of Aurora A coupled with time-dependent decline of cyclin B1, p-CDC25^{T48} and p-HH3, which are in line with a critical role of PKD activity in G₂-M transition and mitotic progression (Fig. 5E, left). In comparison, the expression levels of Aurora B did not significantly change (Fig. 5D and E, Aurora B), indicating a selective effect of CRT101 on Aurora A degradation. Importantly, when ectopically overexpressed, Aurora A completely reversed the effect of CRT101 on cell-cycle markers (Fig. 5E, right; Supplementary Fig. S3). Thus, Aurora A mediated the effect of PKD inhibition on G₂-M transition. To corroborate these findings, Aurora A protein stability

was assessed in cells transfected with siPKD2-3 (siD2) and treated with cycloheximide (CHX) that blocks protein synthesis. In the control cells (siNT), the half-life ($t_{1/2}$) of Aurora A was 49 hours. Partial knockdown of PKD2 reduced $t_{1/2}$ of Aurora A to about 28 hours, while nearly complete knockdown further reduced it to 11 hours (Fig. 5F). Similarly, knockdown of PKD3 with two siRNAs also resulted in downregulation of Aurora A (Supplementary Fig. S5). These results identify Aurora A as a key mediator of PKD2-regulated cell-cycle progression, and the activation of PKD2 at G₂-M stabilizes Aurora A kinase and promotes mitotic entry.

PKD2 and Aurora A localize at the centrosome and knockdown of PKD2 inhibits centrosome separation during G₂

Aurora A has been shown to localize at mitotic apparatus and contribute to G₂-M transition and mitotic progression (13, 22, 28, 29). Aurora A resides in the centrosome and regulates centrosome duplication and separation, inhibition of Aurora A arrests cells in G₂ and blocks mitotic entry. Here, we sought to determine whether the regulation of Aurora A by PKD also took place at the centrosome and whether it affects centrosome function. HeLa cells were costained with PKD2 and γ -tubulin, a marker of the centrosome. As shown in Fig. 6A, PKD2 (red) was found to colocalize with γ -tubulin (green) at different stages of cell cycle, and the colocalization was most prominent in the G₂ stage. Next, cells were synchronized to the start of M-phase by nocodazole and the interaction of PKD2 and γ -tubulin were examined by reciprocal immunoprecipitation. As shown in Fig. 6B, γ -tubulin was detected in immunoprecipitated PKD2 complex, and vice versa. Thus, a portion of PKD2 was present on the centrosome and PKD2

directly interacts with γ -tubulin at the centrosome during G₂-M, which was further confirmed by IF staining (Fig. 6C). Next, we examined whether PKD2 might directly interact with Aurora A. Aurora A is known to localize at the centrosome (13, 27, 30, 31). Our data confirmed that Aurora A (red) colocalized with γ -tubulin (green) during G₂ and mitosis (Supplementary Fig. S6). By costaining for PKD2 and Aurora A, we also found significant colocalization of PKD2 (green) and Aurora A (red) during G₂ and mitosis (Fig. 5A). Colocalization of PKD2 and Aurora A on the centrosome raised a possibility that they might physically interact. However, our follow-up study indicated that PKD2 failed to coimmunoprecipitate with Aurora A in G₂-M-synchronized cells (Supplementary Fig. S7). Despite the lack of direct interaction, the colocalization of PKD2 and Aurora A support the potential functional link between the two proteins in centrosome. Similar colocalization pattern was observed when the intracellular distributions of PKD3 and Aurora A were analyzed during G₂ and mitosis (Supplementary Fig. S7). Aurora A is crucial for




Figure 6.

PKD2 positively regulates centriole separation during late G₂ by colocalizing with centrosomes. **A-C**, HeLa cell was subjected to IF staining to visualize colocalization of PKD2 (red) and γ -tubulin (green). **B**, Coimmunoprecipitation assay was carried out to assess physical interaction between PKD2 and γ -tubulin. **D**, Knockdown of PKD2 by two siRNAs was confirmed by IF staining. **E**, Centrioles were visualized by IF staining of γ -tubulin in cells transfected with siNT or two siPKD2s (left). Cell population harboring unseparated centrosomes was quantified (right). At least 100 cells were counted and graph shows average number of cells with error bars representing SEM (****, $P < 0.0001$). DAPI was used to stain the nucleus.

centrosome maturation and separation during G₂ (31). Hence, we questioned whether loss of Aurora A by depletion of PKD2 caused any defect in centrosome separation. To do so, cells were transfected with two siPKD2s followed by IF staining of γ -tubulin to visualize the centrioles. Knockdown of PKD2 was confirmed by IF staining (Fig. 6D, red). Our results showed that depletion of PKD2 resulted in defects in centriole separation (Fig. 6E, left) and significantly increased cell population harboring unseparated centrosomes (Fig. 6E, right). Taken together, these results suggest that PKD positively regulates stabilization of Aurora A and centrosome separation to promote G₂-M progression.

PKD2 interferes with Fbxw7-mediated ubiquitination and downregulation of Aurora A

To assess whether PKD regulates Aurora A degradation through the proteasome degradation pathway, cells were synchronized with nocodazole, released in fresh medium containing CRT101 in the presence or absence of a proteasome inhibitor MG132, and the levels of Aurora A were examined by immunoblotting. Our data showed that Aurora A levels were dampened within 3 hours of CRT101 treatment and MG132 completely inhibited CRT101-induced Aurora A degradation (Fig. 7A). The half-life of Aurora A was measured in the presence of cycloheximide (CHX). As shown in Fig. 7B, treatment of CRT101 reduced the half-life of Aurora A protein to about 5 hours, whereas MG132 restored its half-life like control sample (nocodazole only; Fig. 7B). These results demonstrated that CRT101 directed Aurora A to the proteasomal pathway for its degradation. To further corroborate this finding, G₂-M-arrested cells were released in medium containing MG132 with or without CRT101, and cell extracts from the samples were subjected to IP using anti-Aurora A antibody, followed by immunoblotting with anti-ubiquitin antibody (Fig. 7C). We found that CRT101 increased ubiquitination of Aurora A (Fig. 7C, top) confirming that Aurora A is downregulated via ubiquitination-dependent proteasomal degradation pathway upon treatment with CRT101. Aurora A is targeted by Cdh1 and F-box protein Fbxw7 (15, 32) for degradation. Therefore, to gain insight into PKD-mediated downregulation of Aurora A, we knocked down PKD2, Cdh1, and Fbxw7 using site-specific siRNAs and analyzed the turnover of Aurora A protein levels. Knockdown of PKD2 downregulated Aurora A, whereas depletion of Cdh1 did not significantly alter Aurora A protein level (Fig. 7D). Interestingly, when Fbxw7 was depleted in PKD2 knocked down cells, Aurora A level was significantly restored as compared with siPKD2 (Fig. 7E). To further corroborate our findings, cells were transfected with nontargeting siRNA (siNT) or siFbxw7, synchronized at the start of M-phase by nocodazole, treated with CRT101 and cell extracts were analyzed by Western blotting to monitor Aurora A degradation (Fig. 7F). The immunoblot showed that Aurora A was downregulated within 3 hours of CRT101 treatment and this turnover was inhibited by Fbxw7 knockdown. These results suggested that PKD protects Aurora A from Fbxw7-mediated ubiquitination and proteasomal degradation to regulate G₂-M transition.


Discussion

PKD as a key target of the second messenger DAG regulates many important cellular functions including cell proliferation. Aberrant expression and activation of PKD have been demonstrated in many cancers and are implicated in oncogenic transformation and tumor progression. PKD2 in particular has been shown to promote tumor cell proliferation in human cancer

including carcinoma of the prostate (8). However, the mechanisms through which it regulates tumor cell proliferation have not been well defined. In this study, we identified PKD2 as a cell-cycle regulator that promotes G₂-M transition and mitotic entry. PKD2 is activated during G₂ and M phases of the cell cycle. It is localized at the centrosome and upon activation regulates centrosome duplication and separation, a key event for the cells to exit G₂ and enter mitosis. Mechanistically, we further demonstrated that PKD activity was required for Aurora A stability during G₂-M, and it acts through Fbxw7 to protect Aurora A from ubiquitination and proteasomal degradation. Our study has identified PKD2 as a potential G₂-M checkpoint kinase. Its aberrant activation may contribute to premature mitotic entry and genome instability that drive tumorigenesis.

A key finding from our study is that PKD is active during G₂ and its activity is sustained throughout mitosis (Fig. 1F). Increased PKD2 activity in G₂-M was supported by: (i) immunoblotting results showing elevated active form of PKD2 in cells entering G₂ and progressing through mitosis (Fig. 1A-D), (ii) IF staining results showing increased active PKD2 staining in cell nuclei during G₂ (Fig. 1) and prophase of mitosis, indicative of PKD activation (21). Our finding that activation of PKD is crucial for G₂-M cell-cycle progression agrees with the report by Papazyan and colleagues that demonstrated a possible connection of PKD activation during mitosis and its role in cell cycle (33). The effects of PKD inhibitors are in line with previous reports of anticancer effects of PKDis. CRT0066101 (CRT101) in particular has been shown to block cell proliferation by inducing G₂-M arrest, downregulation of mitotic regulatory proteins and induction of apoptosis in colorectal (34), bladder (35), pancreatic (36), and prostate (37) cancer cell lines. Our laboratory has also shown that kb-NB142-70, another novel PKD inhibitor causes dramatic G₂-M cell-cycle arrest in PC3 prostate cancer cells (38). Both of these two PKD inhibitors were also shown to inhibit HeLa cell proliferation (Supplementary Fig. S3). Collectively, our study has identified PKD as a crucial cell-cycle regulator that is activated during G₂-M and promotes mitotic entry. Moreover, in this study, we have provided possible mechanistic insights to PKD inhibitor-mediated cell-cycle arrest, downregulation of cell-cycle regulators and inhibition of cell proliferation, which reflected a positive role of PKD in promoting G₂-M transition by modulating Aurora A protein stability.

Aurora A modulates the function of several cell-cycle regulators critical for G₂-M transition and mitosis, such as PLK1, CDK1/cyclin B, and CDC25C (18) and is essential for key mitotic events, such as centrosome maturation and separation (31), mitotic entry and formation of mitotic spindles (27). Aurora A must be degraded to ensure proper mitotic exit. During late anaphase, it loses its interaction with TPX2 and targeted by anaphase-promoting complex/cyclosome (APC/C) ubiquitin ligase for proteasomal degradation (16). Aurora A is also downregulated by Fbxw7, a p53-dependent tumor suppressor (15). Mechanistically, Fbxw7 physically interacts with Aurora A and facilitates the ubiquitination-mediated proteasomal degradation in a Gsk3 β -dependent manner. Aberrant expression and proteostasis of Aurora A have been linked to many cancers, thus enabling it as a promising target for therapeutic interventions in cancer (18). Our study has revealed for the first time, a functional link between PKD and Aurora A kinase in G₂-M progression and mitotic entry of the cell cycle. We have shown evidence that PKD is critical for late G₂ cells to enter mitosis, and that abrogation of its activity delays G₂-M

**Figure 7.**

PKD interferes with Fbxw7-mediated ubiquitination of Aurora kinase A. **A**, Western blot analysis was performed to show that CRT101-mediated downregulation of Aurora A was proteasome-dependent. **B**, Half-life of Aurora A protein was measured from the immunoblot of **A**. **C**, HeLa cells were synchronized at the start of M phase by nocodazole and preincubated with MG132 for 30 minutes. DMSO (control) or CRT101 was added to the culture and the cells were further incubated for 3 hours. Aurora A was pulled down by IP and the protein was analyzed by Western blotting to assess its ubiquitination by CRT101. Downregulation of Aurora A by knockdown of PKD2 and Cdh1 (**D**) or PKD2 and Fbxw7 (**E**) was analyzed by Western blotting (top). Quantification of Aurora A protein levels was performed by densitometry analysis (bottom). **F**, Western blot analysis depicting that CRT101-mediated downregulation of Aurora A is inhibited by knockdown of Fbxw7. GAPDH was used as loading control. **G**, A model of regulation of Aurora A protein stability by PKD. The graph represents average of at least three independent experiments with error bars representing SEM (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ns, not significant).

transition (Figs. 2 and 3). We have shown that loss of PKD has profound effects on cell cycle, such as, (i) rapid downregulation of mitotic regulators including cyclin B1, phospho-CDC25^{T48} (Supplementary Fig. S5), (ii) induction of mitotic catastrophe

characterized by defects in spindle formation, apoptosis and cell death, (iii) defects in centriole separation in late G₂ and most importantly, (iv) degradation of Aurora A by Fbxw7-mediated proteasomal degradation. Cdh1 is a component of anaphase-

promoting complex that is responsible for substrate degradation during anaphase of mitosis (39). Whereas Fbxw7 (α , β , and γ) expression spans all cell-cycle stages (40), which may be a potential cause for siFbxw7, not siCdh1 to be effective against Aurora A turnover in an asynchronous culture where mitotic cell population is significantly low. Therefore, it is conceivable that a cell harboring inactive or no PKD2 eventually lose Aurora A at the protein level and displays somewhat an "Aurora A null" phenotype (41).

Another significant finding of our study is that PKD2 resides on the centrosome during G₂-M and controls centriole separation. We showed that Aurora A and PKD2 colocalized in centrosome in early G₂ and throughout G₂-M. It is known that the maturation and separation of centrosomes are dependent on the recruitment and activation of Aurora A kinase on centrosomes (42). It is unclear whether PKD2 plays a direct role in recruiting Aurora A to the centrosomes. We did not detect direct interaction between PKD2 and Aurora A although both proteins colocalize in the centrosomes and bind to γ -tubulin. Several possibilities may explain this result: (i) the interaction was limited to the centrosomes, which is difficult to detect using whole-cell lysates, (ii) the interaction may be low affinity and transient like many protein kinases and substrates; (iii) the regulation may be indirect through other regulators. Although Fbxw7 was found to mediate the regulation of Aurora A by PKD, it remains to be determined whether PKD directly regulates Fbxw7 activity. Functionally, it is evident that PKD activity modulates the availability/stability of Aurora A at the centrosomes. Thus, inhibition of PKD results in degradation of Aurora A, which may impede the maturation and separation of centrosomes, and thereby block the entry of cells into mitosis. Kienzle and colleagues have shown that PKD positively controls Golgi complex fragmentation at G₂ phase of the cell cycle via Raf-MEK1 pathway, thus, enabling cells to enter mitosis (43). This study is in line with our findings because centrosome duplication and separation is an early step that required for Golgi fragmentation in G₂ (44). Thus, the alteration of Aurora A stability and the resulting defects in centrosome separation may be linked to Golgi fragmentation that occurs at

G₂ phase. Beyond Aurora A, the localization of PKD2 in centrosome, the major microtubule-organizing center of the cell, raises the possibility that it might modulate other centrosomal proteins to ensure tight regulation of cell cycle.

In summary, our results provide strong evidence for a novel function of PKD2 in regulating G₂-M transition, mitotic entry and cell-cycle progression. PKD2 does so by protecting critical cell-cycle regulators including Aurora A kinase, an essential mitotic regulator from Fbxw7-mediated proteasomal degradation (Fig. 7G). The regulation of Aurora A by PKD2 provides mechanistic insights to the oncogenic effects of these kinases and may have important therapeutic implications in cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: A. Roy, Q.J. Wang

Development of methodology: A. Roy, M.V. Veroli, Q.J. Wang

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): A. Roy, S. Prasad, Q.J. Wang

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A. Roy, M.V. Veroli, Q.J. Wang

Writing, review, and/or revision of the manuscript: A. Roy, M.V. Veroli, S. Prasad, Q.J. Wang

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): Q.J. Wang

Study supervision: Q.J. Wang

Acknowledgments

This work was supported in part by the Department of Defense award PC150190 (to Q.W. Wang).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 14, 2018; revised June 18, 2018; accepted July 10, 2018; published first July 17, 2018.

References

- Otto T, Sicinski P. Cell cycle proteins as promising targets in cancer therapy. *Nat Rev Cancer* 2017;17:93–115.
- Ellwanger K, Hausser A. Physiological functions of protein kinase D in vivo. *IUBMB Life* 2013;65:98–107.
- Rozenfurt E, Rey O, Waldron RT. Protein kinase D signaling. *J Biol Chem* 2005;280:13205–8.
- Rylx A, De Kimpe L, Mikhalap S, Vantus T, Seufferlein T, Vandenhede JR, et al. Protein kinase D: a family affair. *FEBS Lett* 2003;546:81–6.
- Guo J, Gertsberg Z, Ozgen N, Sabri A, Steinberg SF. Protein kinase D isoforms are activated in an agonist-specific manner in cardiomyocytes. *J Biol Chem* 2011;286:6500–9.
- Durand N, Borges S, Storz P. Protein kinase D enzymes as regulators of EMT and cancer cell invasion. *J Clin Med* 2016;5:E20.
- Wong C, Jin ZG. Protein kinase C-dependent protein kinase D activation modulates ERK signal pathway and endothelial cell proliferation by vascular endothelial growth factor. *J Biol Chem* 2005;280:33262–9.
- Roy A, Ye J, Deng F, Wang QJ. Protein kinase D signaling in cancer: a friend or foe? *Biochim Biophys Acta* 2017;1868:283–94.
- Wang QJ. PKD at the crossroads of DAG and PKC signaling. *Trends Pharmacol Sci* 2006;27:317–23.
- Bischoff JR, Plowman GD. The Aurora/Ipl1p kinase family: regulators of chromosome segregation and cytokinesis. *Trends Cell Biol* 1999;9:454–9.
- Chan F, Sun C, Perumal M, Nguyen QD, Bavetsias V, McDonald E, et al. Mechanism of action of the Aurora kinase inhibitor CCT129202 and in vivo quantification of biological activity. *Mol Cancer Ther* 2007;6:3147–57.
- Vader G, Lens SM. The Aurora kinase family in cell division and cancer. *Biochim Biophys Acta* 2008;1786:60–72.
- Barr AR, Gergely F. Aurora-A: the maker and breaker of spindle poles. *J Cell Sci* 2007;120:2987–96.
- Nigg EA. Mitotic kinases as regulators of cell division and its checkpoints. *Nat Rev Mol Cell Biol* 2001;2:21–32.
- Kwon YW, Kim JJ, Wu D, Lu J, Stock WA Jr, Liu Y, et al. Pten regulates aurora-A and cooperates with Fbxw7 in modulating radiation-induced tumor development. *Mol Cancer Res* 2012;10:834–44.
- Lindon C, Grant R, Min M. Ubiquitin-mediated degradation of aurora kinases. *Front Oncol* 2015;5:307.
- Kollareddy M, Dzubak P, Zheleva D, Hajdud M. Aurora kinases: structure, functions and their association with cancer. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 2008;152:27–33.
- Yan M, Wang C, He B, Yang M, Tong M, Long Z, et al. Aurora-A kinase: a potent oncogene and target for cancer therapy. *Med Res Rev* 2016;36:1036–79.
- Lukasiewicz KB, Lingle WL. Aurora A. centrosome structure, and the centrosome cycle. *Environ Mol Mutagen* 2009;50:602–19.

20. Chen J, Deng F, Singh SV, Wang QJ. Protein kinase D3 (PKD3) contributes to prostate cancer cell growth and survival through a PKCepsilon/PKD3 pathway downstream of Akt and ERK 1/2. *Cancer Res* 2008;68:3844–53.
21. Rey O, Sinnett-Smith J, Zhukova E, Rozengurt E. Regulated nucleocytoplasmic transport of protein kinase D in response to G protein-coupled receptor activation. *J Biol Chem* 2001;276:49228–35.
22. Bucher N, Britten CD. G2 checkpoint abrogation and checkpoint kinase-1 targeting in the treatment of cancer. *Br J Cancer* 2008;98:523–8.
23. Kimura M, Yoshioka T, Saio M, Banno Y, Nagaoka H, Okano Y. Mitotic catastrophe and cell death induced by depletion of centrosomal proteins. *Cell Death Dis* 2013;4:e603.
24. Castedo M, Perfettini JL, Roumier T, Andreau K, Medema R, Kroemer G. Cell death by mitotic catastrophe: a molecular definition. *Oncogene* 2004;23:2825–37.
25. Hirota T, Kunitoku N, Sasayama T, Marumoto T, Zhang D, Nitta M, et al. Aurora-A and an interacting activator, the LIM protein Ajuba, are required for mitotic commitment in human cells. *Cell* 2003;114:585–98.
26. He L, Yang H, Ma Y, Pledger WJ, Cress WD, Cheng JQ. Identification of Aurora-A as a direct target of E2F3 during G₂-M cell cycle progression. *J Biol Chem* 2008;283:31012–20.
27. Fu J, Bian M, Jiang Q, Zhang C. Roles of Aurora kinases in mitosis and tumorigenesis. *Mol Cancer Res* 2007;5:1–10.
28. Dutertre S, Cazales M, Quaranta M, Froment C, Trabut V, Dozier C, et al. Phosphorylation of CDC25B by aurora-A at the centrosome contributes to the G₂-M transition. *J Cell Sci* 2004;117:2523–31.
29. Jang CY, Coppinger JA, Seki A, Yates JR 3rd, Fang G. Plk1 and aurora A regulate the depolymerase activity and the cellular localization of Kif2a. *J Cell Sci* 2009;122:1334–41.
31. Lukasiewicz KB, Lingle WL, Aurora A. centrosome structure, and the centrosome cycle. *Environ Mol Mutagen* 2009;50:602–19.
31. Dutertre S, Descamps S, Prigent C. On the role of aurora-A in centrosome function. *Oncogene* 2002;21:6175–83.
32. Taguchi S, Honda K, Sugiura K, Yamaguchi A, Furukawa K, Urano T. Degradation of human Aurora-A protein kinase is mediated by hCdh1. *FEBS Lett* 2002;519:59–65.
33. Papazyan R, Doche M, Waldron RT, Rozengurt E, Moyer MP, Rey O. Protein kinase D isozymes activation and localization during mitosis. *Exp Cell Res* 2008;314:3057–68.
34. Wei N, Chu E, Wipf P, Schmitz JC. Protein kinase d as a potential chemotherapeutic target for colorectal cancer. *Mol Cancer Ther* 2014;13:1130–41.
35. Li QQ, Hsu I, Sanford T, Railkar R, Balaji N, Sourbier C, et al. Protein kinase D inhibitor CRT0066101 suppresses bladder cancer growth in vitro and xenografts via blockade of the cell cycle at G₂-M. *Cell Mol Life Sci* 2017;75:939–63.
36. Harikumar KB, Kunnumakkara AB, Ochi N, Tong Z, Deorukhkar A, Sung B, et al. A novel small-molecule inhibitor of protein kinase D blocks pancreatic cancer growth in vitro and in vivo. *Mol Cancer Ther* 2010;9:1136–46.
37. Tandon M, Johnson J, Li Z, Xu S, Wipf P, Wang QJ. New pyrazolopyrimidine inhibitors of protein kinase d as potent anticancer agents for prostate cancer cells. *PLoS One* 2013;8:e75601.
38. Lavallo CR, Bravo-Altamirano K, Giridhar KV, Chen J, Sharlow E, Lazo JS, et al. Novel protein kinase D inhibitors cause potent arrest in prostate cancer cell growth and motility. *BMC Chem Biol* 2010;10:5.
39. Pesin JA, Orr-Weaver TL. Regulation of APC/C activators in mitosis and meiosis. *Annu Rev Cell Dev Biol* 2008;24:475–99.
40. Sionov RV, Netzer E, Shaulian E. Differential regulation of FBXW7 isoforms by various stress stimuli. *Cell Cycle* 2013;12:3547–54.
41. Sasai K, Parant JM, Brandt ME, Carter J, Adams HP, Stass SA, et al. Targeted disruption of Aurora A causes abnormal mitotic spindle assembly, chromosome misalignment and embryonic lethality. *Oncogene* 2008;27:4122–7.
42. Cervigni RI, Barretta ML, Persico A, Corda D, Colanzi A. The role of Aurora-A kinase in the Golgi-dependent control of mitotic entry. *Bioarchitecture* 2011;1:61–5.
43. Kienzle C, Eisler SA, Villeneuve J, Brummer T, Olayioye MA, Hausser A. PKD controls mitotic Golgi complex fragmentation through a Raf-MEK1 pathway. *Mol Biol Cell* 2013;24:222–33.
44. Corda D, Barretta ML, Cervigni RI, Colanzi A. Golgi complex fragmentation in G₂-M transition: an organelle-based cell-cycle checkpoint. *IUBMB Life* 2012;64:661–70.

Molecular Cancer Research

Protein Kinase D2 Modulates Cell Cycle By Stabilizing Aurora A Kinase at Centrosomes

Adhiraj Roy, Maria Victoria Veroli, Sahdeo Prasad, et al.

Mol Cancer Res Published OnlineFirst July 17, 2018.

Updated version	Access the most recent version of this article at: doi: 10.1158/1541-7786.MCR-18-0641
------------------------	--

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://mcr.aacrjournals.org/content/early/2018/08/31/1541-7786.MCR-18-0641 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.