

AWARD NUMBER: W81XWH-16-1-0278

TITLE: Synthetic Lethal Metabolic Targeting of Senescent Cells after Androgen Deprivation Therapy

PRINCIPAL INVESTIGATOR: Jarrard, David F.

CONTRACTING ORGANIZATION: University of Wisconsin System

REPORT DATE: SEPTEMBER 2020

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE SEPTEMBER 2020		2. REPORT TYPE Final		3. DATES COVERED 1 July 2016 - 30 June 2020	
4. TITLE AND SUBTITLE Synthetic Lethal Metabolic Targeting of Senescent Cells after Androgen Deprivation Therapy				5a. CONTRACT NUMBER W81XWH-16-1-0278	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) David Jarrard E-Mail: jarrard@urology.wisc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Wisconsin System 21 N Park St., Ste. 6401 Madison, WI 53715-1218				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Progression to castrate resistant prostate cancer for men with advanced prostate cancer (PC) results after the initiation of androgen deprivation therapy (ADT). One underutilized strategy that has the potential to dramatically improve outcomes is eradicating persistent senescent cancer cells that remain after ADT and likely play a key role in castration-resistant PC. We have demonstrated that ADT induces senescence in androgen-dependent PC cells and acts synergistically with the diabetes medication metformin to induce apoptosis in PC cells. ADT activates Akt, AMPK, mTORC1 and XIAP at various time points and silencing XIAP augments apoptosis induced by ADT, pointing to XIAP as a promising drug target to increase the activity of ADT in advanced prostate cancer. We have also shown that metformin enhances the antitumor activity of ADT in two PDX PC models. Analysis of the VA database revealed that metformin use in patients with advanced prostate cancer receiving ADT is associated with improved OS and cancer-specific survival. Our results point to metformin as a novel therapeutic strategy for castration-resistant PC.					
15. SUBJECT TERMS prostate cancer, androgen deprivation therapy, senescence, proteotoxic stress, xenograft models, metformin, synthetic lethality					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
Unclassified	Unclassified	Unclassified	Unclassified	96	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Page

1. Introduction	4
2. Keywords	5
3. Accomplishments.....	5
4. Impact	15
5. Changes/Problems	16
6. Products.....	16
7. Participants & Other Collaborating Organizations	17
8. Special Reporting Requirements	18
8. Appendices	18

Title: **Synthetic lethal metabolic targeting of senescent cells after androgen deprivation therapy**

1. INTRODUCTION

Progression to castrate resistant prostate cancer for men with advanced prostate cancer (PC) results after the initiation of androgen deprivation therapy (ADT). One underutilized therapeutic strategy that has the potential to dramatically improve outcomes is eradicating the persistent cancer cells that remain after ADT and likely play a key role in the development of castration-resistant PC. The initiation of ADT induces susceptibilities in PC cells that make them amenable to synergistic treatment and improved cell killing.

Androgen withdrawal in murine xenografts and human PC tissues is associated with a decrease in the proliferative index, but surprisingly low levels of apoptosis. We and others have demonstrated that a substantial portion of these persistent cells express markers of cellular senescence, a terminal growth arrest characterized by exit from the cell cycle and senescence-associated β -galactosidase expression. These senescent cells, although not proliferating, generate a protumor response, the senescent secretory phenotype, that may be detrimental to the patient and must be removed. However, the unique metabolic phenotype expressed by these persistent senescent cells is characterized by increased protein synthesis and notably an amplified proteotoxic stress response (PSR), a conserved survival pathway characterized by induction of multiple heat shock protein (Hsp) families coordinated by the master transcriptional regulator Hsf1. It is our overall hypothesis that the new senescent phenotype induced in prostate cancer cells by ADT may result in unique vulnerabilities to drugs targeting pathways such as the PSR that are critical for survival in the senescent state.

In preliminary data activation of the PSR in these residual cancer cells may represent a pathway critical for the survival of senescent PC cells. Further experiments have identified one agent, metformin, a widely-used, nontoxic oral antidiabetic drug that we propose to repurpose as synthetic lethal therapy in combination with ADT. We postulate that that metformin is synthetic lethal with ADT because it disables the principal PSR pathway mediated by Hsf1 in senescent PC cells that are already experience high levels of proteotoxic stress.

In Aim 1 we will examine the activity of metformin in eradicating senescent PCs following ADT in cellular models. In addition, we will determine whether metformin's actions are specifically mediated by inactivation of Hsf1 and resultant disruption of the PSR cell survival pathway mediated by Hsp27, Hsp70 and Hsp 90. In vitro and xenograft PC models utilizing overexpression of a phosphorylation-resistant Hsf1 mutant will be used to interrogate the specific role of the Hsf1-mediated PSR in the synthetic lethal response. PC has a variable response to ADT. The ability of metformin to clear senescent cancer cells after ADT will be examined in Aim 2 in a series of human PC xenografts that exhibit variable responses to ADT. We will utilize a xenograft system consisting of human prostate cancer tumors that can be exposed to drug combinations in a physiologically relevant setting, the growth easily tracked, and the tumor readily harvested for detailed examination. Experiments will test whether synchronous ADT-metformin or their stepwise use leads to better tumor regression and longer survival. In addition, markers of response will be investigated in the tumors focusing on the Hsps examined in Aim 1. Finally, in Aim 3 we will employ a health sciences research approach using the National Department of Veterans Affairs Corporate Data Warehouse to investigate a retrospective cohort of patients on ADT (~260,000 men), 8% of whom are on metformin (~21,000), to determine PC-specific mortality, biochemical recurrence-free survival and skeletal related events. This will provide further evidence for the implementation of this novel synthetic lethal therapeutic strategy.

Our studies have the potential to lead to a new treatment paradigm for PC by specifically targeting a unique vulnerability of senescent PCs (the PSR) that persist following ADT and likely contribute to androgen-resistance. The proposed study directly addresses mechanisms of resistance for men with high-risk cancer and furthermore, since metformin may mitigate the metabolic side effects of ADT, may improve the physical health of men with PC. When completed our new synthetic lethal approach to PC can be readily translated into the clinic since both ADT and metformin are safe and currently in use.

2. KEYWORDS

prostate cancer, androgen deprivation therapy, senescence, proteotoxic stress, xenograft models, metformin, synthetic lethality

3. ACCOMPLISHMENTS

SPECIFIC AIM 1:

Major Task 1: Determine whether the ADT-metformin synergistic response is mediated via disruption of the Hsf1-mediated proteolytic stress response (PSR).

Subtask 1: Characterize the effects of metformin on the viability of senescent PC cells following ADT (Jarrard/Cryns)

- PC cells will be treated with vehicle or bicalutamide for 4 days followed by metformin or vehicle for 2-4 days
- Score senescent PC cells using SA- β -gal activity, GLB1 immunostaining and flow cytometry
- Evaluate apoptosis using co-immunofluorescence with GLB1 and active caspase-3 Ab and by annexin V labeling of GLB1-flourescent PC cells
- Cell lines: LNCaP, CWR22Rv1, VCaP in subtasks 1-3

-COMPLETED

We have optimized the dosing and time to achieve a maximal coefficient index for synthetic lethality for 2 cell lines, LNCaP and LAPC4. VCaP was not utilized because of variable androgen responsiveness, and CWR22RV1 was removed from the study due to the inconsistent response to androgen deprivation. LAPC4 was used as an alternate cell line. In addition, we adopted another approach for inducing senescence: charcoal-stripped serum (CSS) instead of FBS.

We have demonstrated that the combination of androgen deprivation therapy (ADT, bicalutamide or CSS) and metformin inhibit prostate cancer cell growth more robustly than either agent alone (**Figure 1**).




Figure 1. Combination of ADT with metformin decreases the growth of androgen-dependent prostate cancer cells compared to single-agent use. A&C, Cells were cultured in medium containing different percentages of FBS/CSS for 6 days and then followed by addition of metformin for 24 or 48 hours. Cell proliferation was measured by DNA assay. Con: 10% FBS; CSSL: 8% CSS + 2% FBS; CSSH: 10% CSS; ML, metformin low dose (mM): LNCaP, 0.1; LAPC4, 2.5. MH, metformin high dose (mM): LNCaP, 1; LAPC4, 5. **B&D**, Cells were treated with bicalutamide for 6 days, followed by addition of metformin for 24 or 48 hours. Con: 10% FBS treated with vehicles. BL, bicalutamide low dose (μ M): LNCaP, 1; LAPC4, 20. BH, bicalutamide high dose (μ M): LNCaP, 5; LAPC4, 30. Metformin doses are the same as A&C. Data are shown as percentage of the control group, Mean $\pm$ SD. Synergistic effect was calculated by Calcsyn, moderate synergy (combination index CI: 0.7-0.85) +; synergy (CI: 0.3-0.7) ++, and strong synergy (CI: 0.1-0.3) +++.

Senescence was quantitated using SA- β -gal activity, flow cytometry and western blotting for p16 and p27. We have demonstrated that ADT induces senescence in androgen-dependent prostate cancer cells (**Figure 2**).




Figure 2. ADT induces phenotypic characteristics of cellular senescence in androgen-dependent prostate cancer cells. **A**, LNCaP or LAPC4 cells were cultured in CSS medium or treated with bicalutamide for 8 days, and then stained for SA- β -gal activity (magnification $\times 100$). Con: 10% FBS in RPMI media; CSS: 10% Charcoal-stripped serum in RPMI media ; Bic, bicalutamide in complete RPMI media, LNCaP, 5 μ M; LAPC 30 μ M. **B**, Flow cytometry was used to examine the cellular complexity side scatter. The percentage of cells with side-scatter is shown in graph (Mean $\pm$ SD) * t-test, $p < 0.05$. ADT increases the fraction of cells with elevated cellular complexity and size. **C**, LNCaP and LAPC4 cells were cultured in CSS medium or treated with bicalutamide for days indicated, then whole cell lysates were harvested and analyzed for p16 and p27 expression by western blot. 30 μ g of protein was loaded, and α -tubulin was used as a loading control.

Apoptosis occurs maximally 48 hr after exposure to ADT in LNCaP and LAPC4 prostate cancer cells (**Table 1**). Apoptosis increases with the addition of metformin to the senescent cells.

Table 1. Addition of metformin increases apoptosis in senescence-induced cancer cells

	LNCaP	LAPC4
CSS	0.3	1.29
Bic	0.87	0.95
Met	1.21	1.13
CSS + Met	2.51	1.72
Bic + Met	2.00	2.97

Table 1: Apoptosis was assayed using Caspase-Glo and shown as fold change relative to control, mean value. CSS: 10% CSS; Bic, bicalutamide (μ M), LNCaP, 5; LAPC4 30. Met: metformin (mM): LNCaP, 1; LAPC4, 5.

Subtask 2: Characterize the effects of metformin on the PSR of senescent PC cells following ADT (Cryns/Jarrard).


- PC cells will be treated with ADT and/or metformin as in subtask 1
- Collect GLB1-fluorescent and non-fluorescent cells by FACs and determine Hsf1, Hsp27, Hsp70 and Hsp90 mRNA and protein levels.
- Perform co-immunofluorescence with GLB1 and Hsf1, Hsp27, Hsp70, Hsp90, p-AMPK and AMPK
- Determine whether AMPK binds directly to Hsf1 by co-immunoprecipitation.

Milestones: We predict that metformin will increase binding of AMPK to Hsf1 and inhibit the PSR.

-COMPLETED


We have examined the proteotoxic stress response (PSR) in multiple prostate cancer cell lines in response to androgen deprivation therapy with and without metformin treatment. Analyses of the entire population of treated prostate cancer cells has yielded results that depend on the prostate cancer cell line and individual heat shock proteins with regard to the effects of both androgen deprivation and metformin treatment. For example, we observed increased HSP90 in LNCaP cells, but not LAPC4 cells, with exposure to ADT, and decreased HSP27 in LAPC4 cells in response to ADT (**Figure 3**). Other HSPs were also not consistently altered.

Figure 3: Expression of Hsps in PC cells exposed to ADT. LNCaP and LAPC4 Cells were treated with bicalutamide for 6 days, and then followed by treatment of metformin for 1 or 2 days. Bic, bicalutamide (μ M), LNCaP, 5; LAPC4 30. Met: metformin (mM): LNCaP, 1; LAPC4, 5.



Since we did not observe consistent PSR induction in prostate cancer cells following ADT, we have explored alternative signaling pathway(s) that may be associated with cell survival after ADT. AKT plays a critical role in cell survival, senescence and apoptosis-resistance (1). We observed that activated p-AKT (T308) increased in prostate cancer cells in response to ADT (**Figure 4**). AMPK is a cellular sensor that plays a key role in regulating energy homeostasis (2). We also observed that p-AMPK increased in prostate cancer cells 2-4 days after ADT treatment when senescence markers were minimally induced, consistent with AMPK activation. Subsequently, p-AMPK levels decreased in late terminal senescence (8-10 days).

Figure 4: Time-dependent AKT and AMPK signaling alterations induced by ADT. LNCaP and LAPC cells were cultured with media containing 10% CSS or treated with complete culture media containing bicalutamide (LNCaP 5 μ M, LAPC4 30 μ M) as for the indicated number of days. 30 μ g of protein was used for each sample.



AMPK is a known suppressor of mTORC1 signaling through phosphorylation of TSC2 (3). Persistent activation of AKT and inhibition of AMPK after long-term exposure to ADT was associated with mTORC1 activation as determined by increased p-p70S6K, a downstream target of mTORC1 (**Figure 5**). Activation of mTORC1 may serve as one of the pathways by which senescent prostate cancer cells persist. Consistent with this idea, addition of metformin, an activator of AMPK, to ADT, resulted in decreased p-p70S6K (**Figure 5**) and a reduction in the senescence marker p16 (data not shown), suggesting clearance of senescent cells through metformin's effect on AMPK activation and subsequent inhibition of mTORC1.




Figure 5. Late ADT induces mTORC1 signaling through activation of AKT and inhibition of AMPK. Metformin addition reactivates AMPK and inhibits mTORC1 signaling. PC cells were cultured in 10% CSS medium (**A**) or treated with bicalutamide (**B**) for 6 days followed by addition of metformin. Whole cell lysates were harvested and analyzed by western blotting. 30 µg of protein was used for each sample. Bic: bicalutamide (mM), LNCaP, 5; LAPC4, 30. Met: metformin (mM): LNCaP, 1; LAPC4, 5.

expression were decreased, and apoptosis increased to a greater extent in ADT-treated cells with XIAP knockdown than in control media, indicating that a significant part of the survival response that ADT induces is XIAP-dependent (**Figure 6B and C**). We then examined whether overexpression of XIAP alters the response of PC cells to ADT and metformin. XIAP was overexpressed in LNCaP cells using a pcDNA3-XIAP vector, while the empty vector was used as a control. After stable cell lines were established, cell growth, apoptosis and protein expression were analyzed (**Figure 6D-F**). Overexpression of XIAP abrogated the synergistic decrease in growth seen with metformin following senescence induction by ADT. In addition,

XIAP overexpression reversed the increased apoptosis after metformin exposure or the combination of ADT and metformin (**Figure 6F**). Moreover, the senescence marker p16 was reproducibly increased compared to the cells transfected with control vector, suggesting increased cell survival (**Figure 6E**). Akt signaling increased in cells overexpressing XIAP compared to controls, consistent with a known feedback loop. These data support a role for the anti-apoptotic protein XIAP in the response of PC cells to ADT and metformin.




Figure 6: XIAP inhibits apoptosis in senescent PC cells. **A-C:** LNCaP cells were transfected with non-silencing control siRNA (si-Control) or siRNA targeting XIAP (si-XIAP), 24hr later were replaced with fresh media containing either 10% FBS or CSS. **A.** Cell proliferation was measured with DNA assay at 2 and 3 days after switched to CSS media, data shown as Mean $\pm$ SD, * $p < 0.05$. Eight replicates in each group, three independent experiments. **B.** Whole protein lysates were harvested for checking the denoted proteins' expression by western blot. **C.** Apoptosis was measured with caspase-Glo, normalized with cell number. The fold change relative to cells transfected with si-Control and cultured in media containing 10% FBS, data shown as Mean $\pm$ SD, * $p < 0.05$, comparison between CSS si-Control vs CSS si-XIAP. Eight replicates in each group, three independent experiments. **D-E:** XIAP overexpression abrogated metformin induced apoptosis in senescent cells. LNCaP cells were transfected with either pcDNA3-XIAP or pcDNA3 plasmids as control. After stable cell lines have been established, cells were cultured with either Media containing FBS or CSS followed by metformin treatment for 2 days and then cell proliferation (**D**), protein expression (**E**) and apoptosis (**F**) were analyzed as above. Apoptosis was assayed using Caspase-Glo, shown as fold change relative to control, Mean $\pm$ SD value. CSS: 10% CSS; Met: metformin (1 mM). Statistical analysis using one-way ANOVA demonstrates p values between different groups CSS/Met compared to CSS or Met alone (pcDNA control: 0.008, 0.0004, pcDNA-XIAP: 0.1015, 0.006, respectively).

Subtask 3: Determine the functional role of site-specific phosphorylation of Hsf1 on the effects of metformin on senescent PCs following ADT (Cryns/Jarrard).

- Stably transduce PC cells with vector, WT Hsf1 or mutant S121A Hsf1 by lentiviral transduction.

- Treat PC cells stably expressing vector, WT or S121A mutant Hsf1 with ADT and/or metformin as in subtask 1.
- Perform cell viability (subtask 1) and molecular characterization (subtask 2) assays on senescent PCs and the entire population of PCs.

Milestones: We predict that the S121A mutant Hsf1 will abrogate the effects of metformin on cell death and PSR following ADT.

Due to the inconsistent alterations of HSPs by ADT in the investigated androgen-dependent prostate cancer cell lines, we altered our focus to AKT, AMPK, mTORC1 and XIAP as described in the previous section and published manuscript.

Subtask 4: Determine the function of site-specific phosphorylation of Hsf1 on the antitumor effects of ADT and metformin in vivo (Jarrard/Cryns).

- Male nude mice with LnCaP and CWR22rv1 flank tumors stably expressing WT or S121A mutant Hsf1 will be randomized to 4 groups (8 mice per group): (1) vehicle + sham operation; (2) vehicle + castration; (3) metformin + sham operation; and (4) metformin + castration. Metformin will be tried simultaneously (groups 3 and 4) or sequentially (groups 5 and 6). Tumor size will be assessed weekly and serum PSA recorded. To assess senescent cell clearing, a parallel experiment will be performed using the same 4 treatment groups (8 mice per group) except that mice will be euthanized 4 weeks after castration and tumors harvested for analysis. (16X(6X3)) for 2 experiments (total 576)
- Mouse tissues will be analyzed for SA- β -gal activity, HP1 γ , GLB1, p27, Hsp27, Hsp70, Hsp90, Ki67, active caspase-3 and TUNEL staining

We did not investigate the function of site-specific phosphorylation of Hsf1 on the antitumor effects of ADT and metformin *in vivo* due to the inconsistent data regarding the PSR (Aim 1, Subtask 3).

Major Task 2: Examine the synthetic lethal response involving ADT-metformin in vivo in cancers of variable androgen sensitivity and test markers of response.

Subtask 1: To determine the optimal schedule for combining ADT and metformin and assess whether metformin eradicates senescent PC cells following ADT (Jarrard).

-COMPLETED

- We obtained approval by the USAMRMC ORP Animal Care and Use Review Office (ACURO), in addition to the local Institutional Animal Care and Use Committee (IACUC).
- Four groups were randomized using 2 xenografts (77 and 147) when flank tumors are 100mm³ to examine the i) castrated and treated with metformin (CM), ii) castrated and treated with vehicle control (CC), iii) sham surgery and treated with metformin (SM), and iv) sham surgery and treated with vehicle control (SC). Metformin was administered one week after surgery in drinking water at a dose of 2g / liter. Post-ADT Tumor growth and body weight were assessed weekly.




Figure 7: Metformin and ADT suppress tumor growth in patient-derived xenografts (PDXs) more robustly than ADT alone. A. LuCaP 77, B. LuCaP 147. The growth rates are shown as percentage of tumor size compared to the time point of sham surgery/castration (median $\pm$ SE). One-way ANOVA was used to analyze the differences between any 2 groups. ** $p < 0.01$ is considered as statistically significant between castration plus metformin treated group compared to castration alone (N=8 mice per group) Sham: sham surgery control, Cast: castration, Met: sham surgery with metformin treated, Cast+Met: castration and metformin treated.

The growth of both PC PDXs was initially inhibited by castration; however, tumors began regrowing after 4-6 weeks (**Figure 7**). Metformin also modestly inhibited tumor growth compared to controls in both models. With the combination treatment, LuCaP 77 tumors at 7 weeks were significantly smaller than those in the castration ($p < 0.01$) or metformin ($p < 0.01$) alone groups. In the LuCaP 147 PDX model, tumors were significantly smaller at 6-8 weeks post-surgery in the combination therapy compared to those in the castration ($p < 0.01$) or metformin groups ($p < 0.01$).

We conclude from these animal studies that combined treatment with ADT and simultaneous metformin leads to an improved tumor response in prostate cancer PDX models.

A pilot study we performed using a smaller number of animals (8 vs 8) that underwent ADT and metformin treatment simultaneously versus delayed metformin treatment (10 days) suggested that simultaneous use of these 2 agents versus 10 d delay led to no appreciable differences in outcomes. This experiment was not further pursued.

Subtask 2: To determine whether PSR markers predict improved response to ADT-Metformin (Jarrard/Cryns).

- Xenograft tumors from the 2A will be sectioned and immunofluorescence will be used and quantitated using the automated Vectra™ system for Hsp27, Hsp70 and Hsp90. The proteolytic stress response (PSR) represented by these 3 genes in castrated animals harvested at 4wk (group i) will be statistically compared to tumor response, survival, PSA, and other markers including GLB1 in ADT-Metformin groups (iii and iv).

-COMPLETED

Tissue Microarrays (TMA) were constructed from 58 tissue specimens (29 in duplicate) from the Xenograft 77 model and 54 tissue specimens (27 in duplicate) from the Xenograft 147 model. Antibodies for Ki67 and active cleaved caspase-3 (CC3) were used to quantify proliferation and apoptosis. For image analysis and quantification of the staining intensity, the VECTRA system was used. Cores with $< 5\%$ epithelial component or loss of tissue were excluded from the analysis. Nuance system and Inform 1.2™ software (Caliper Life Sciences, Hopkinton, MA)

were used to for building spectral libraries on the basis of target signals of the two stained parameters. Expression of Ki67 (proliferation) and cleaved caspase 3 (apoptosis) were evaluated by immunohistochemistry. H&E staining was used to differentiate nuclear and cytoplasmic tissue compartments. Two cores from the same tumor were averaged to give a more precise estimate.

Ki67 expression decreased in both the castration and castration plus metformin groups (**Figure 8**). A difference in CC3-positivity in tumors from different treatment groups was not detected ($p=0.1905$) in contrast to our *in vitro* data showing induction of apoptosis (**Table 1 and Figure 6**). We did not further pursue the staining of HSPs in this TMA due to the inconsistent activation of HSPs *in vitro*.




Figure 8. Castration, metformin and the combination reduce proliferation in a prostate cancer PDX model. Ki67 expression in LuCaP 147 tumors. Expression of Ki67 protein was detected in this TMA constructed from LuCaP 17 xenograft tumors, with duplicate cores per tissue. The VECTRA system was used for image capture and quantification of the staining intensity. **A.** Images of Ki67 staining. The brown color is positive staining. H&E was used as counter-staining. **B.** Data is shown as the percentage of Ki67-positive tumor cells (mean $\pm$ SE). A one-way ANOVA was used to analyze the differences between any 2 groups. * $p < 0.05$, ** $p < 0.01$.

Major Task 3: Determine whether metformin combined with ADT results in improved cancer-specific survival and longer time to secondary interventions in patients on these agents.

Subtask 1: We propose to utilize a robust observational cohort from the national Veterans Affairs (VA) database to specifically evaluate our hypothesis that metformin improves PC response to ADT, thereby directly examining the patient relevance of our preclinical data in validated patient population. Approvals (Jarrard/Richards).

-COMPLETED

Subtask 2: Data collection, organization with exclusion and inclusion from 2000-2008 (Jarrard/Richards).

-COMPLETED

Subtask 3: Analysis of primary and secondary predictive variables (Jarrard/Richards). Evaluate and control for other covariates including other diabetes medication administration history, age, race, Charlson-comorbidity score, agent orange exposure, family history of prostate cancer, tobacco use, blood type, local therapy (surgery or

radiation), date of prostate cancer diagnosis, stage at diagnosis, Gleason score, and other medication administration history (finasteride, aspirin, and docetaxol).

-COMPLETED

Milestone(s) Achieved:

Using national Veterans Affairs databases, we identified all men diagnosed with PCa between 2000-2008 that were treated with ADT with follow-up through October of 2015. We excluded patients that were treated with ADT for ≤ 6 months or were receiving ADT concurrently with localized radiation therapy. We split these patients into three cohorts: 1. Patients without diabetes 2. Diabetics on metformin 3. Diabetics not treated with metformin. Our primary outcome was overall survival (OS) and secondary outcomes included skeletal related events (SRE) and prostate-cancer specific survival. Cox proportional hazards ratios were calculated for overall and disease specific survival.

The total cohort after exclusions consisted of 87,344 patients of which 53,893 (61%) were non-diabetics, 14,517 (17%) were diabetics on metformin, and 18,934 (22%) were diabetics not receiving metformin. The mean age was 75 ± 11 years in the non-diabetics, 71 ± 12 in the diabetics on metformin, and 75 ± 10 in the diabetics no metformin ($p < 0.001$). The median OS was 7.1 years in the non-diabetics, 9.1 years in the diabetics on metformin, and 7.4 years in the diabetics not treated with metformin.




Figure 9. Kapan-Meier analysis of prostate cancer specific survival improves for men on metformin at the time of ADT initiation. Performed through the VA system.

0.78) and to a lesser effect diabetics not treated with metformin (HR 0.87, 95% CI 0.81- 0.93) with non-diabetics as referent group.

We conclude that metformin use in Veterans with advanced prostate cancer receiving ADT is associated with improved OS and cancer-specific survival. The impact of metformin in prostate cancer patients should be evaluated in a prospective clinical trial.

Multivariable Cox proportional hazards analysis assessing for predictors of overall survival showed improved survival in diabetics on metformin (HR 0.77, 95% CI 0.74-0.81) vs. diabetics not treated with metformin (HR 0.99, 95% CI 0.95-1.03) with non-diabetics as referent group. Multivariable Cox proportional hazards analysis assessing for predictors of SRE revealed no association between metformin use (HR 0.99, 95% CI 0.92-1.07) and SRE. Lastly, multivariable Cox proportional hazards analysis assessing for predictors of prostate-cancer specific survival showed improved survival in diabetics on metformin (HR 0.72, 95% CI 0.67-

Completed and abstract presented at the American Urological Association Meeting May 2017 and the GU ASCO meeting Feb 2017. Paper published J Urol 2018 (included in the appendix). Paper in press in Molecular cancer therapeutics (included in the appendix).

References

1. Manning, B. D. and A. Toker (2017). AKT/PKB Signaling: Navigating the Network. Cell 169(3): 381-405.
2. Hardie, D. G., et al. (2012). AMPK: a nutrient and energy sensor that maintains energy homeostasis. Nat Rev Mol Cell Biol 13(4): 251-262.
3. Hindupur, S. K., et al. (2015). The opposing actions of target of rapamycin and AMP activated protein kinase in cell growth control. Cold Spring Harb Perspect Biol 7(8): a019141.

Opportunities for training and professional development?

These include an oncology fellow Dr Shiva Damoradan who has recently taken a clinical position at the University of Toledo. Additional trainees include Nathan Damaschke a graduate student who performed work with the tumor analysis. He is now doing a postdoctoral training at Northwestern University.

How were the results disseminated to communities of interest?

Abstract presentations at the American Urological Association Meeting (5/2017 in Boston MA), GU ASCO Meeting (Feb 2017 in Orlando, FL), the Prostate Cancer Foundation meeting (10/2017 in Washington DC) and multiple publications (see Products).

What do you plan to do during the next reporting period to accomplish the goals and objectives?

This is the final report. Future goals will focus on a trial involving metformin and ADT.

4. IMPACT

A growing body of evidence indicates that ADT induces a susceptible therapeutic niche that may be exploited to enhance therapeutic response. In the current study, we demonstrate for the first time a synergistic inhibitory effect of metformin combined with ADT on growth in human PC cell lines *in vitro* and PDX tumors *in vivo*. We find a novel role for XIAP, an anti-apoptotic protein driven by increased AKT signaling in PC cells after ADT. These observations mechanistically support our recent study demonstrating improved survival in patients taking metformin at the time of initiating ADT for advanced PC. The fate of PC cells after ADT includes apoptosis, quiescence and other phenotypes. In the current work, we demonstrate that subpopulations of androgen-sensitive PC cells after ADT exhibit molecular markers characteristic of senescence, such as increased SA-beta-gal-staining, cellular complexity and expression of cyclin dependent kinase inhibitor 2A p16INK4a. Apoptosis occurs early after ADT and is limited. In a series of xenograft tumors *in vivo*, surgical castration shows a similar induction of senescence with decreased proliferation as determined by Ki-67 expression. We also demonstrate that persistent activation of AKT and inhibition of AMPK following long-term ADT activating mTOR1 signaling may serve as one of the pathways by which senescent cells persist and represent a potential for therapeutic exploitation. A synergistic reduction in cell number was noted with the addition of metformin to ADT *in vitro*, as well as a reduction in xenograft tumor growth with the combination *in vivo*, which is through the activation of AMPK by metformin leading to downregulation of the anti-apoptosis protein XIAP and induction of cell death. These studies provide a molecular rationale for the observed clinical observation that metformin at the time of ADT improves both overall and cancer-specific survival in patients with advanced PCa. We would anticipate that other agents that activate AMPK efficiently may also

act synergistically with ADT. Trials are indicated to further test this inexpensive minimally toxic drug in combination with ADT.

5. CHANGES/PROBLEMS

- We are using a modified approach to identify senescent prostate cancer cells after treatment as noted under Major Task 1, Subtask 1. We have also utilized a different approach employing charcoal-stripped serum to remove androgens.
- We observed prostate cell line-specific and Hsp-specific alterations in response to ADT (Major Task 1, Subtask 2) Given inconsistent activation of the proteotoxic stress response, we did not investigate the functional role of Hsf1 (the major transcription factor for HSPs), on the effects of metformin on senescent PCs following ADT under Major Task 1, Subtasks 3 & 4.
- Alternatively we have investigated alternative signaling pathway(s) that may be associated with cell survival after ADT. We found that Akt, AMPK, mTORC1 and the antiapoptotic protein XIAP are induced at various time points during ADT and that silencing XIAP augmented apoptosis by ADT.

6. PRODUCTS

Abstracts:

Metformin Use Associated with Improved Survival in Veterans with Advanced Prostate Cancer: A Large Observational Study. Kyle A. Richards, Jinn-ing Liou, Vincent Cryns, Tracy M. Downs, E. Jason Abel, and David F. Jarrard (American Urological Association Meeting. Boston MA 2017).

THE IMPACT OF METFORMIN USE ON ONCOLOGIC OUTCOMES IN PROSTATE CANCER PATIENTS RECEIVING DOCETAXEL CHEMOTHERAPY Andrew Truong, Gray Roberge, Jinn-ing Liou, E. Jason Abel, Tracy Downs, David Jarrard, Kyle Richards, Madison, WI, American Urological Association Meeting, San Francisco CA, 2018).

Androgen Deprivation Therapy Induces Unique Anti-Warburg Metabolomic Alteration in Hormone Sensitive Prostate Cancer Cells. Tariq A. Khemees, Mikolaj Filon, Bing Yang, Amani Gillette, Joseph Gawdzik, Megan Showalter, Oliver Fiehn, Melissa Skala and David F. Jarrard. (American Urological Association Meeting, Chicago IL, 2020).

Publications:

1. Strelakova E, Malin D, Rajanala H, **Cryns VL**. Metformin sensitizes triple-negative breast cancer to proapoptotic TRAIL receptor agonists by suppressing XIAP expression. *Breast Cancer Res Treat* 2017 163:435-447.
2. **Richards KA**, Liou JI, **Cryns VL**, Downs TM, Abel EJ, **Jarrard DF**. Metformin Use Is Associated with Improved Survival in Patients with Advanced Prostate Cancer on Androgen Deprivation Therapy. *J Urol*. 2018 Dec;200(6):1256-1263.
3. Etheridge T, Damodaran S, Schultz A, Richards KA, Gawdzik J, Yang B, **Cryns V**, **Jarrard DF**. Combination therapy with androgen deprivation for hormone sensitive prostate cancer: A new frontier. *Asian J Urol*. 2019 Jan;6(1):57-64.
4. Damodaran S, Lang JM, **Jarrard DF**. Targeting Metastatic Hormone Sensitive Prostate Cancer: Chemohormonal Therapy and New Combinatorial Approaches. *J Urol*. 2019 May;201(5):876-885.
5. Yang B, Damodaran S, Khemees TA, Filon MJ, Schultz A, Gawdzik J, Etheridge T, Malin D, Richards KA, **Cryns V**, **Jarrard DF**. Synthetic Lethal Metabolic Targeting of Androgen Deprived Prostate Cancer Cells with Metformin, *Molecular Cancer Therapeutics*, in press.

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

Senior key personnel have been working on the project since the initiation of the project with no changes.

The following individuals have worked on the project:

Name: David F. Jarrard, MD

Project Role: Principal Investigator

Researcher Identifier (e.g., ORCID ID): [0000-0001-8444-7165](#)

Nearest person month worked: 1.2

Contribution to Project: David Jarrard has conceived and designed the study, reviewed all of the data and the analysis of all of the results on the project, wrote and revised the manuscript.

Name: Vince Cryns, MD

Project Role: Co-Principal Investigator

Researcher Identifier (e.g., ORCID ID): 0000-0003-0355-2268

Nearest person month worked: 2.4

Contribution to Project: Vince Cryns has conceived and designed the study, reviewed all of the data and the analysis of all of the results on the project, wrote and revised the manuscript.

Name: Kyle Richards, MD

Project Role: Co- Investigator

Researcher Identifier (e.g., ORCID ID):

Nearest person month worked: 0.3

Contribution to Project: Dr Richards has reviewed the data and the analysis of all of Aim 3 on the project, co-wrote and revised the manuscript.

Name: Bing Yang, MD, PhD

Project Role: Researcher

Researcher Identifier (e.g., ORCID ID):

Nearest person month worked: 3.6

Contribution to Project: Bing Yang has prepared all the PCa cell lines used in this study and performed the analysis on the cell lines, organized the data and assisting with the mouse studies.

Name: Shiva DAMODARAN

Project Role: Post-Doctoral Fellow

Researcher Identifier (e.g., ORCID ID):

Nearest person month worked: 4

Contribution to Project: Shiva has optimized the animal xenografts and metformin treatment and is currently performing the animal studies.

Name: Joe Gawdzik

Project Role: Post-Doctoral Fellow

Researcher Identifier (e.g., ORCID ID):

Nearest person month worked: 1

Contribution to Project: Joe has assisted with the cell culture experiments for Task 1 and data analysis and collection.

Name: Jinn-ing Liou
Project Role: Researcher
Researcher Identifier (e.g., ORCID ID):
Nearest person month worked: 0.3
Contribution to Project: Jinn-ing has generated the data and the analysis of all of Aim 3 on the project, co-wrote and revised the manuscript.

Name: Dmitry Malin, PhD
Project Role: Associate Scientist
Researcher Identifier (e.g., ORCID ID): 0000-0002-5728-7511
Nearest person month worked: 6
Contribution to Project: Dmitry Malin has analyzed human PCa cell lines for markers of proteotoxic stress in response to androgen deprivation therapy with or without metformin treatment and assisted with the design of these experiments.

Name: Zhanhai Li, PhD
Project Role: Biostatistician
Researcher Identifier (e.g., ORCID ID):
Nearest person month worked: 0.36
Contribution to Project: Dr. Sampene has provided biostatistical support for these studies.

Name: Elena Strekalova, PhD
Project Role: Assistant Scientist
Researcher Identifier (e.g., ORCID ID): 0000-0001-9271-5465
Nearest person month worked: 2.5
Contribution to Project: Elena Strekalova has worked together with Dr. Malin to characterize the therapeutic response of PCa cell lines to androgen deprivation therapy and metformin.

8. SPECIAL REPORTING REQUIREMENTS

Duplicative reports have been submitted for both the Initiating PI and the Partnering PI

9. APPENDICES

Manuscripts



HHS Public Access

Author manuscript

Breast Cancer Res Treat. Author manuscript; available in PMC 2019 May 31.

Published in final edited form as:

Breast Cancer Res Treat. 2017 June ; 163(3): 435–447. doi:10.1007/s10549-017-4201-0.

Metformin Sensitizes Triple-Negative Breast Cancer to Proapoptotic TRAIL Receptor Agonists by Suppressing XIAP Expression

Elena Strekalova, Dmitry Malin, Harisha Rajanala, and Vincent L. Cryns

Department of Medicine, University of Wisconsin Carbone Cancer Center, University of Wisconsin School of Medicine and Public Health, Madison, WI USA

Abstract

Purpose: Despite robust antitumor activity in diverse preclinical models, TNF-related apoptosis-inducing ligand (TRAIL) receptor agonists have not demonstrated efficacy in clinical trials, underscoring the need to identify agents that enhance their activity. We postulated that the metabolic stress induced by the diabetes drug metformin would sensitize breast cancer cells to TRAIL receptor agonists.

Methods: Human triple (estrogen receptor, progesterone receptor and HER2)-negative breast cancer (TNBC) cell lines were treated with TRAIL receptor agonists (monoclonal antibodies or TRAIL peptide), metformin or the combination. The effects on cell survival, caspase activation, and expression of TRAIL receptors and the antiapoptotic protein XIAP were determined. In addition, XIAP was silenced by RNAi in TNBC cells and the effects on sensitivity to TRAIL were determined. The antitumor effects of metformin, TRAIL or the combination were evaluated in an orthotopic model of metastatic TNBC.

Results: Metformin sensitized diverse TNBC cells to TRAIL receptor agonists. Metformin selectively enhanced the sensitivity of transformed breast epithelial cells to TRAIL receptor agonist-induced caspase activation and apoptosis with little effect on untransformed breast epithelial cells. These effects of metformin were accompanied by robust reductions in the protein levels of XIAP, a negative regulator of TRAIL-induced apoptosis. Silencing XIAP in TNBC cells mimicked the TRAIL-sensitizing effects of metformin. Metformin also enhanced the antitumor effects of TRAIL in a metastatic murine TNBC model.

Conclusions: Our findings indicate that metformin enhances the activity of TRAIL receptor agonists, thereby supporting the rationale for additional translational studies combining these agents.

Correspondence to: Vincent L. Cryns, Department of Medicine, MFCB 4144, 1685 Highland Avenue, Madison, WI 53705 USA. Phone: 608-262-4786; Fax: 608-263-9983. vlcryns@medicine.wisc.edu.

Compliance with Ethical Standards

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

Ethical approval: This article does not contain any studies with human participants performed by any of the authors.

Keywords

metformin; breast cancer; TRAIL; metastasis; apoptosis; therapeutics

Introduction

Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL/Apo2L) and agonistic mAbs targeting its proapoptotic receptors (TRAIL-R1/DR4 and TRAIL-R2/DR5) selectively activate the caspase-8/10-mediated extrinsic apoptotic pathway in transformed cells and exhibit robust antitumor effects in diverse murine models of cancer [1, 2]. Despite promising preclinical results, TRAIL receptor agonists have failed to demonstrate significant efficacy, either alone or in combination with chemotherapy, in multiple clinical trials in advanced malignancies [3–8]. These disappointing results in clinical trials have been attributed to a number of factors, including *de novo* and/or acquired resistance to TRAIL receptor agonists, inadequate receptor oligomerization on ligand binding, limiting procaspase-8/10 activation, and lack of biomarkers to predict treatment response [1, 2].

In an effort to enhance the activity of TRAIL receptor agonists, these proapoptotic agents have been combined with TRAIL-sensitizing agents (*e.g.*, histone deacetylase inhibitors, PPAR γ agonists, aspirin, mTOR and other kinase inhibitors) to augment its antitumor effects [9–13]. More recently, we have identified a novel nutritional intervention that selectively sensitizes breast cancer cells to TRAIL-R2 agonists [14]. Specifically, we demonstrated that depletion of the essential amino acid methionine metabolically primes breast cancer cells to respond to the agonistic TRAIL-R2 mAb lexatimumab by increasing cell surface expression of TRAIL-R2. Moreover, dietary methionine restriction enhanced the antitumor activity of lexatimumab against mammary tumors and lung metastases in an orthotopic model of metastatic breast cancer [14]. Hence, methionine restriction metabolically primes breast cancer cells to targeted agents that activate cell death by exposing a targetable vulnerability, namely, enhanced cell surface expression of TRAIL-R2.

Intriguingly, the diabetes medication metformin mimics many of the effects of methionine restriction, including disruption of methionine metabolism via inhibition of the functionally linked folate cycle in the one-carbon metabolic pathway, inhibition of the mechanistic target of rapamycin (mTOR), broad antitumor activity, improved insulin sensitivity, and prolonged lifespan [15–19]. Metformin use has been associated with reduced incidence of a broad range of tumor-types and reduced cancer mortality in many epidemiologic studies [19–22]. However, the antitumor mechanisms of metformin are not well understood. Both direct tumor effects (activation of AMPK with resultant inactivation of mTORC1, inhibition of mitochondrial complex I, and suppression of nuclear translocation of NF κ B and Stat3 phosphorylation) and indirect systemic effects (reduction in insulin and IGF-1 levels) have been reported and implicated in its antitumor effects [18, 19]. Consistent with preclinical findings, metformin treatment of newly diagnosed breast cancer patients prior to surgery improves metabolic indices, increases tumor apoptosis and decreases tumor proliferation [23].

Given the similarities noted between methionine restriction and metformin, we postulated that metformin may also metabolically prime breast cancer cells to respond to proapoptotic TRAIL receptor agonists. Here we report that metformin sensitizes TNBC cells to TRAIL receptor agonists *in vitro* and *in vivo*. Metformin selectively enhances the sensitivity of transformed breast epithelial cells to TRAIL receptor agonist-induced caspase activation and apoptosis with little effect on untransformed breast epithelial cells. These effects of metformin are accompanied by a robust reduction in the expression level of the antiapoptotic protein XIAP. Metformin also enhances the antitumor effects of TRAIL in a murine model of TNBC. Collectively, our findings indicate that metformin enhances the clinical activity of TRAIL receptor agonists and suggests that additional translational studies combining these agents are warranted.

Methods and materials

Cell culture and reagents

Human MDA-MB-231 and GILM2 TNBC cells stably expressing mCherry and parental GILM2 TNBC cells were maintained as described [24]. MDA-MB-468 TNBC cells were cultured in DMEM media supplemented with 10% FBS and 100 IU/mL penicillin/streptomycin (ThermoFisher Scientific/Gibco). Human MCF-10A breast epithelial cells stably expressing oncogenic H-RasV12 or vector were maintained as described [25]. Mapatumumab and lexatumumab were kindly provided by Dr. Robin Humphreys (Human Genome Sciences). Recombinant TRAIL peptide (amino acids 95–281) was purified as described [11].

Crystal violet cell survival assay

A crystal violet cell survival assay was performed as described [10]. Cells were seeded on 6-well plates (3×10^5 cells/well) overnight. Cells were then preincubated with vehicle or metformin (1 mM for MDA-MB-468 cells or 5 mM for MDA-MB-231-mCherry and GILM2 cells) for 48 hours, and treated with vehicle or TRAIL, mapatumumab or lexatumumab (each 1.5 μ g/ml for MDA-MB-231-mCherry and GILM2 cells or 4 μ g/ml for MDA-MB-468 cells) for an additional 24 hours (MDA-MB-231-mCherry and GILM2 cells) or 48 hours (MDA-MB-468 cells) before staining with crystal violet. The percentage cell confluence of crystal violet-positive cells was determined in 3 fields per treatment condition using NIH ImageJ software. Cells in each field were colored using “color threshold”. Percentage confluence of colored cells was quantified using “analyze particles” which reports the percent area in each field occupied by colored cells.

Immunoblotting

Whole-cell lysates were immunoblotted as described [25] using primary Abs against XIAP, β -actin (Sigma-Aldrich), PARP (BD Biosciences) and procaspase-3 (Cell Signaling Technology).

Caspase-3/7 activity assay

The Caspase-Glo 3/7 Assay System (Promega) was used to measure caspase-3/7 activity in cell lysates according to the manufacturer’s instructions. Briefly, cells were seeded in 96-

well plates (2.5×10^3 cells/well) overnight. The next day, cells were preincubated with vehicle or metformin (5 mM) for 48 hours and then treated with vehicle or TRAIL (1.5 $\mu\text{g/ml}$ for MDA-MB-231-mCherry, GILM2, MCF-10A and MCF-10A-RAS cells or 4 $\mu\text{g/ml}$ for MDA-MB-468 cells) for an additional 24 hours. Caspase-3/7 activity was normalized to cell number and expressed as fold activity compared to vehicle-treated cells.

TRAIL receptor cell surface expression

Cell surface expression of TRAIL receptors (TRAIL-R1 and TRAIL-R2) was determined by flow cytometry using TRAIL-R1, TRAIL-R2 or control IgG1 mAb conjugated with allophycocyanin (BioLegend) as described previously [14].

XIAP siRNA experiments

siRNAs targeting the sequences GAAGCUAGAUUAAAGUCCU (sil-XIAP) or CAGUGAAGACCCUUGGGAA (si2-XIAP) in human XIAP and non-silencing control siRNA were purchased from Sigma-Aldrich. Cells were transfected with siRNAs using Lipofectamine RNAiMAX Reagent (ThermoFisher Scientific) according to the manufacturer's protocol.

Real-time PCR

Real-time PCR for TRAIL-R1, TRAIL-R2 and GAPDH was performed as described previously [14]. Primers for XIAP (forward 5-AGTGCCACGCAGTCTACAAA, reverse 5-GCATGTGTCTCAGATGGCCT) were purchased from Integrated DNA Technologies and real-time PCR was performed using the same methods. A comparative C_t method was used to normalize RNA expression in samples to the controls in each experiment.

Orthotopic Model of Metastatic TNBC

GILM2-mCherry TNBC cells (2×10^6) were resuspended in Matrigel (BD Biosciences) and injected bilaterally into the 4th mammary gland ducts of 4-to 5-week-old female NOD *scid* IL2 receptor γ chain knockout (NSG) mice (Jackson Laboratory). Mice were randomized into four treatment groups (10 mice per group) three weeks after tumor inoculation: 1) PBS vehicle *i.p.* daily; 2) metformin 2 mg/ml in the drinking water; 3) TRAIL (10 mg/kg *i.p.* daily), or (4) metformin (2 mg/ml in the drinking water) plus TRAIL (10 mg/kg *i.p.* daily). Mammary tumor volume was calculated as described [26]. Lung metastases were visualized by fluorescence microscopy in isolated whole lungs and scored using NIH ImageJ analysis as described [26]. All animal experiments were carried out as part of an IACUC-approved protocol at the University of Wisconsin-Madison.

Tumor apoptosis assay

Formalin-fixed, paraffin-embedded tumor tissue sections were analyzed for active caspase-3 expression by immunohistochemistry using an Ab against cleaved caspase-3 (Cell Signaling) as described [26].

Statistics

The statistical significance of differences between groups was determined by ANOVA with Bonferroni posttests using GraphPad Prism 4 software.

Results

Metformin sensitizes TNBC cells to TRAIL receptor agonists

To determine whether metformin sensitizes TNBC to TRAIL receptor agonists, three human TNBC cell lines (MDA-MB-231-mCherry, GILM2 and MDA-MB-468) were preincubated with vehicle or metformin and then treated with TRAIL, mapatumumab or lexatumumab. Metformin sensitized all three TNBC cell lines to TRAIL receptor agonists, with the most robust effects observed in MDA-MB-231-mCherry and GILM2 cells. (Fig. 1). A dose-response experiment in MDA-MB-468 cells identified 0.5 mM metformin as the minimal concentration needed to sensitize these TNBC cells to TRAIL receptor agonists (Fig. S1). In addition, preincubation with metformin sensitized human HT29 colon adenocarcinoma and DU145 prostate cancer cell lines to TRAIL receptor agonists (Fig. S2). Collectively, these results demonstrate that metformin augments the cytotoxicity of TRAIL receptor agonists against a broad range of tumor cell types, including TNBC cells.

Metformin sensitizes transformed breast epithelial cells to TRAIL receptor agonists

To determine whether metformin enhanced the cytotoxicity of TRAIL receptor agonists preferentially in transformed cells, we utilized human MCF-10A breast epithelial cells transformed by oncogenic H-RasV12 and untransformed MCF-10A cells stably expressing empty vector. Strikingly, metformin robustly sensitized MCF-10A-Ras cells to TRAIL and lexatumumab, but had a minimal effect on the sensitivity of untransformed MCF-10A-Vector cells to these agents (Fig. 2). In contrast, metformin had only a modest effect on the cytotoxicity of mapatumumab against MCF-10A-Ras cells. Collectively, these results indicate that metformin preferentially sensitizes transformed cells to TRAIL receptor agonists and support the potential tumor-selectivity of this therapeutic combination.

Metformin enhances TRAIL-induced caspase activation in TNBC cells

To investigate whether metformin promotes TRAIL-induced caspase activation, TNBC cells were preincubated with metformin, treated with vehicle or TRAIL and then analyzed by immunoblotting. Metformin promoted TRAIL-induced proteolysis of the caspase substrate PARP as detected by diminished full-length PARP and/or increased cleavage product compared to vehicle-treated cells (Fig. 3A). In addition, metformin enhanced TRAIL-induced procaspase-3 proteolytic processing as detected by decreased procaspase-3 levels and/or increased cleaved subunit compared to vehicle-treated cells. Similarly, pretreatment of HT29 and DU145 carcinoma cells with metformin augmented TRAIL-induced caspase activation (Fig. S3). To quantitate more precisely the effects of metformin and TRAIL on caspase activation, we utilized a caspase-3/-7 activity assay. Metformin robustly enhanced TRAIL-induced caspase-3/-7 activity in all three TNBC cells compared to cells treated with either agent alone (Fig. 3B). Metformin also enhanced TRAIL-induced caspase activation in MCF-10A-Ras cells, while untransformed MCF-10-Vector cells were not sensitive to this

combination (Fig. 3C and 3D). Consistent with its reported mechanism of action [18, 19], metformin increased p-AMPK levels and selectively inhibited mTORC1 as determined by reduced phosphorylation of the mTORC1 substrate p-S6 and no effect on the mTORC2 substrate p-Akt (Fig. S4). Taken together, these findings indicate that metformin potentially augments caspase activation and apoptosis by TRAIL in a broad range of tumor cell types and provide additional evidence for the tumor-selectivity of this combination.

Metformin does not alter cell surface expression of TRAIL receptors in TNBC cells

To determine whether metformin sensitizes cancer cells to TRAIL by increasing the expression of its proapoptotic receptors (TRAIL-R1 and TRAIL-R2), TNBC cells were treated with metformin and TRAIL receptor mRNA levels were measured by real-time PCR. Metformin treatment resulted in a modest increase in TRAIL-R2 mRNA levels in MDA-MB-468 TNBC cells, but had little effect on TRAIL-R2 levels in the other TNBC cell lines or on TRAIL-R1 mRNA levels in any of the TNBC cell lines (Fig. 4A). Furthermore, metformin did not significantly affect cell surface expression of either proapoptotic TRAIL-R1 or TRAIL-R2 receptors as determined by flow cytometry (Fig. 4B). Metformin also did not alter protein levels of MAGED2 (Fig. S5), which we previously demonstrated to be downregulated by methionine restriction [14]. These findings indicate that the TRAIL-sensitizing effects of metformin are not due to enhanced TRAIL receptor expression or cell surface localization in TNBC cells.

Metformin reduces XIAP expression in TNBC cells and silencing XIAP sensitizes TNBC cells to TRAIL

The antiapoptotic X-linked inhibitor of apoptosis protein (XIAP) has been demonstrated to confer resistance to TRAIL-induced apoptosis by suppressing caspase activation [27, 28]. Therefore, we postulated that metformin might sensitize cancer cells to TRAIL by downregulating XIAP. Consistent with this hypothesis, treatment of MDA-MB-231-mCherry, GILM2 and MDA-MB-468 TNBC cells with metformin resulted in a reduction in XIAP protein levels (Fig. 5A, left panel). In contrast, treatment with metformin did not affect XIAP mRNA levels (Fig. 5A, right panel), indicating that metformin regulates XIAP via a posttranscriptional mechanism. To examine the functional role of XIAP downregulation in the TRAIL-sensitizing effects of metformin, MDA-MB-231 cells were transfected with a scrambled siRNA (si-Control) or one of two different siRNAs targeting XIAP (si-1 XIAP and si-2 XIAP). Both siRNAs targeting XIAP reduced XIAP protein levels compared to the scrambled control siRNA (Fig. 5B). Notably, silencing XIAP in MDA-MB-231-mCherry cells had a modest effect on cell viability and robustly sensitized these cells to TRAIL treatment compared to a scrambled control siRNA (Fig. 5C). To examine whether silencing XIAP enhanced TRAIL-induced caspase activation, MDA-MB-231-mCherry cells were transfected with siRNAs targeting XIAP or a scrambled control and then treated overnight with vehicle or TRAIL. Silencing XIAP enhanced TRAIL-induced cleavage of the caspase substrate PARP (reduction of full-length PARP and/or increased cleavage product), while levels of procaspase-3 proteolysis (reduction of procaspase-3 levels) were comparable in TRAIL-treated cells transfected with scrambled siRNAs or siRNAs targeting XIAP (Fig. 5D). Collectively, these results indicate that metformin sensitizes TNBC cells to TRAIL-induced apoptosis by downregulating expression of XIAP.

Metformin enhances the antitumor effects of TRAIL in an orthotopic model of metastatic TNBC

To determine whether metformin augments the antitumor effects of TRAIL in an orthotopic model of metastatic TNBC, we treated female NSG mice bearing established GILM2-mCherry mammary tumors with vehicle, metformin alone, TRAIL alone or the combination of metformin and TRAIL. Under the conditions tested, metformin had no significant effect on mammary tumor growth or lung metastases. In contrast, TRAIL inhibited mammary tumor growth, but the combination of TRAIL and metformin was more effective than TRAIL alone (Fig. 6A). Both therapies, TRAIL alone or in combination with metformin inhibited lung metastases to a comparable degree (Fig. 6B). Moreover, both TRAIL alone or in combination with metformin induced apoptosis in mammary tumors and lung metastatic lesions as determined by active, cleaved caspase-3 immunostaining (Fig. 6C). Consistent with our *in vitro* findings, the combination of TRAIL and metformin resulted in more apoptosis induction in mammary tumors and lung metastases than either TRAIL or metformin alone. Importantly, metformin treatment reduced XIAP protein levels in mammary tumors (Fig. 6D), consistent with our *in vitro* findings. None of the treatments affected the body weight of the mice at the end of the study compared to control vehicle-treated mice (Fig. 6E). Collectively, these findings indicate that metformin enhances the antitumor activity of TRAIL against mammary tumors and lung metastases *in vivo* and provide preclinical evidence supporting additional translational studies investigating the combination of metformin and TRAIL receptor agonists in metastatic TNBC.

Discussion

Based largely on epidemiologic data and promising preclinical studies, the oral diabetes medication metformin has been incorporated into a multitude of clinical trials in diverse early-stage and advanced malignancies, either as monotherapy or in combination with cytotoxic agents or radiation [19]. We have demonstrated that metformin sensitizes TNBC cells to proapoptotic TRAIL receptor agonists, while untransformed breast epithelial cells are largely resistant to this combination. Although other groups have recently demonstrated that metformin enhances TRAIL-induced apoptosis in cultured cancer cells [29–32], we have provided the first *in vivo* evidence for the therapeutic utility of this combination in an orthotopic model of metastatic TNBC that recapitulates many features of the human disease [24]. Specifically, we showed that metformin augments the antitumor effects of TRAIL against mammary tumors and increases apoptosis in mammary tumors and lung metastases compared to treatment with either agent alone. Importantly, these antitumor effects *in vivo* were observed at metformin doses that were well tolerated by the mice, a critical point because the TRAIL-sensitizing effects of metformin *in vitro* typically require millimolar concentrations of metformin in standard cell culture media due to their supraphysiologic glucose concentration [33, 34]. Given the lack of efficacy of TRAIL receptor agonists in clinical trials in advanced solid tumors [3–8], our findings are particularly significant from a translational perspective because they suggest that metformin may enhance the efficacy of TRAIL receptor agonists in clinical trials. As such, our results provide the first murine model evidence for the clinical utility of the combination of metformin and TRAIL receptor

agonists, thereby providing critical preclinical evidence to support additional translational studies in poor-prognosis TNBC, which currently lack targeted therapies [35].

Indeed, metformin is a particularly attractive TRAIL-sensitizing agent based on its well documented safety in both diabetic and non-diabetic patients, its low cost and its beneficial impact on metabolic health by improving insulin sensitivity [18, 19]. In addition, our results indicate that the combination of metformin and TRAIL receptor agonists activate apoptosis in p53 mutant TNBC cells and preferentially induce apoptosis in transformed breast epithelial cells with little effect on untransformed breast epithelial cells. Although the mechanism of this tumor-selectivity remains poorly understood, our findings suggest that the toxicity of the combination therapy may be modest. Moreover, our studies may provide insights into the reduced cancer incidence in individuals treated with metformin reported in many epidemiologic studies [19–22]: metformin may enhance the sensitivity of nascent tumors to TRAIL-dependent immune surveillance pathways.

Mechanistically, we have demonstrated that metformin sensitizes TNBC cells to TRAIL receptor agonists by downregulating the antiapoptotic protein XIAP. XIAP is a direct inhibitor of caspases that confers resistance to TRAIL-induced caspase-3 activation and apoptosis, while induction of TRAIL-induced apoptosis requires XIAP to be displaced from caspases by mitochondrial release of Smac/DIABLO [27, 28]. We observed that metformin reduces protein expression of XIAP but does not alter XIAP mRNA levels, thereby pointing to a post-translational mechanism. Intriguingly, rapamycin was previously demonstrated to negatively regulate protein translation of XIAP [36], suggesting that metformin may reduce XIAP protein levels via inhibition of mTORC1. Moreover, metformin treatment reduced XIAP protein levels in mammary tumors in our murine model of TNBC, indicating that the dosing used in our study was sufficient to target this pathway *in vivo*. To determine the functional relevance of the downregulation of XIAP protein by metformin, we used RNAi and observed that XIAP silencing robustly sensitizes TNBC cells to TRAIL-induced caspase activation and cell death, underscoring the importance of this molecular event in sensitizing TNBC cells to TRAIL. Our results are also consistent with a prior report demonstrating that a small molecule inhibitor of XIAP (embelin) sensitizes inflammatory breast cancer cells lines to TRAIL [37].

Notably, we did not observe any significant effects of metformin on the cell surface expression of TRAIL receptors in contrast to other reports that metformin increases TRAIL-R2 protein levels, although the cellular localization of the receptor was not delineated in these prior studies [29, 31]. Additionally, metformin did not affect protein levels of MAGED2, indicating that metformin sensitizes TNBC cells to TRAIL receptor agonists by a different mechanism than methionine restriction, which increases cell surface expression of TRAIL-R2 by downregulating expression of MAGED2, an inhibitor of TRAIL receptor expression [14, 38]. Furthermore, other groups have implicated downregulation of the antiapoptotic protein FLIP as an important molecular event in the TRAIL-sensitizing effects of metformin [12, 39]. Although our results do not exclude the potential contribution of FLIP or other molecular targets, our studies point to the downregulation of XIAP by metformin as a functionally relevant event for its TRAIL-sensitizing effects.

In conclusion, we have demonstrated that metformin downregulates XIAP and sensitizes clinically aggressive TNBC cells to TRAIL receptor agonists. We have also shown that the combination of metformin and TRAIL has robust antitumor activity in a murine model of metastatic TNBC which recapitulates features of the human disease. Given the lack of targeted therapies for TNBC and the largely disappointing activity of TRAIL receptor agonists in clinical trials to date, our results point to the combination of metformin and TRAIL receptor agonists as a promising approach to enhance activity of these proapoptotic agents. Finally, our studies suggest that inhibition of XIAP expression by metformin may be a useful biomarker to predict response to this combined therapy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are indebted to Dr. Robin Humphreys for providing mapatumumab and lexatumumab.

Funding: This study was funded by grants from the Breast Cancer Research Foundation (VLC), the Department of Defense PC150221P1 (VLC), and P30CA14520 University of Wisconsin Comprehensive Cancer Center core facility support.

References

1. Ashkenazi A (2015) Targeting the extrinsic apoptotic pathway in cancer: lessons learned and future directions. *J Clin Invest* 125(2):487–9 [PubMed: 25642709]
2. de Miguel D, Lemke J, Anel A, Walczak H, Martinez-Lostao L (2016) Onto better TRAILs for cancer treatment. *Cell Death Differ* 23(5):733–47 [PubMed: 26943322]
3. Plummer R, Attard G, Pacey S, Li L, Razak A, Perrett R, Barrett M, Judson I, Kaye S, Fox NL et al. (2007) Phase I and pharmacokinetic study of lexatumumab in patients with advanced cancers. *Clin Cancer Res* 13(20):6187–94 [PubMed: 17947486]
4. Soria JC, Smit E, Khayat D, Besse B, Yang X, Hsu CP, Reese D, Wiezorek J, Blackhall F (2010) Phase Ib study of dulanermin (recombinant human Apo2L/TRAIL) in combination with paclitaxel, carboplatin, and bevacizumab in patients with advanced non-squamous non-small-cell lung cancer. *J Clin Oncol* 28(9):1527–33 [PubMed: 20159815]
5. Herbst RS, Eckhardt SG, Kurzrock R, Ebbinghaus S, O'Dwyer PJ, Gordon MS, Novotny W, Goldwasser MA, Tohny TM, Lum BL et al. (2010) Phase I dose-escalation study of recombinant human Apo2L/TRAIL, a dual proapoptotic receptor agonist, in patients with advanced cancer. *J Clin Oncol* 28(17):2839–46 [PubMed: 20458040]
6. Leong S, Cohen RB, Gustafson DL, Langer CJ, Camidge DR, Padavic K, Gore L, Smith M, Chow LQ, von Mehren M et al. (2009) Mapatumumab, an antibody targeting TRAIL-R1, in combination with paclitaxel and carboplatin in patients with advanced solid malignancies: results of a phase I and pharmacokinetic study. *J Clin Oncol* 27(26):4413–21 [PubMed: 19652058]
7. Merchant MS, Geller JI, Baird K, Chou AJ, Galli S, Charles A, Amaoko M, Rhee EH, Price A, Wexler LH et al. (2012) Phase I trial and pharmacokinetic study of lexatumumab in pediatric patients with solid tumors. *J Clin Oncol* 30(33):4141–7 [PubMed: 23071222]
8. Fuchs CS, Fakih M, Schwartzberg L, Cohn AL, Yee L, Dreisbach L, Kozloff MF, Hei YJ, Galimi F, Pan Y et al. (2013) TRAIL receptor agonist conatumumab with modified FOLFOX6 plus bevacizumab for first-line treatment of metastatic colorectal cancer: A randomized phase 1b/2 trial. *Cancer* 119(24):4290–8 [PubMed: 24122767]
9. Inoue S, Mai A, Dyer MJ, Cohen GM (2006) Inhibition of histone deacetylase class I but not class II is critical for the sensitization of leukemic cells to tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis. *Cancer Res* 66(13):6785–92 [PubMed: 16818655]

10. Lu M, Strohecker A, Chen F, Kwan T, Bosman J, Jordan VC, Cryns VL (2008) Aspirin sensitizes cancer cells to TRAIL-induced apoptosis by reducing survivin levels. *Clin Cancer Res* 14(10): 3168–76 [PubMed: 18483385]
11. Lu M, Kwan T, Yu C, Chen F, Freedman B, Schafer JM, Lee EJ, Jameson JL, Jordan VC, Cryns VL (2005) Peroxisome proliferator-activated receptor γ agonists promote TRAIL-induced apoptosis by reducing survivin levels via cyclin D3 repression and cell cycle arrest. *J Biol Chem* 280(8):6742–51 [PubMed: 15569667]
12. Zhao L, Yue P, Khuri FR, Sun SY (2013) mTOR complex 2 is involved in regulation of Cbl-dependent c-FLIP degradation and sensitivity of TRAIL-induced apoptosis. *Cancer Res* 73(6): 1946–57 [PubMed: 23319802]
13. Abdulghani J, Allen JE, Dicker DT, Liu YY, Goldenberg D, Smith CD, Humphreys R, El-Deiry WS (2013) Sorafenib sensitizes solid tumors to Apo2L/TRAIL and Apo2L/TRAIL receptor agonist antibodies by the Jak2-Stat3-Mcl1 axis. *PloS One*. 8(9):e75414 [PubMed: 24086526]
14. Strekalova E, Malin D, Good DM, Cryns VL (2015) Methionine deprivation induces a targetable vulnerability in triple-negative breast cancer cells by enhancing TRAIL receptor-2 expression. *Clin Cancer Res* 21(12):2780–91 [PubMed: 25724522]
15. Corominas-Faja B, Quirantes-Pine R, Oliveras-Ferraro C, Vazquez-Martin A, Cufi S, Martin-Castillo B, Micol V, Joven J, Segura-Carretero A, Menendez JA (2012) Metabolomic fingerprint reveals that metformin impairs one-carbon metabolism in a manner similar to the antifolate class of chemotherapy drugs. *Aging (Albany NY)* 4(7):480–98 [PubMed: 22837425]
16. Cabreiro F, Au C, Leung KY, Vergara-Irigaray N, Cocheme HM, Noori T, Weinkove D, Schuster E, Greene ND, Gems D (2013) Metformin retards aging in *C. elegans* by altering microbial folate and methionine metabolism. *Cell* 153(1):228–39 [PubMed: 23540700]
17. Martin-Montalvo A, Mercken EM, Mitchell SJ, Palacios HH, Mote PL, Scheibye-Knudsen M, Gomes AP, Ward TM, Minor RK, Blouin MJ, et al. (2013) Metformin improves healthspan and lifespan in mice. *Nat Commun* 4:2192 [PubMed: 23900241]
18. Foretz M, Guigas B, Bertrand L, Pollak M, Viollet B (2014) Metformin: from mechanisms of action to therapies. *Cell Metab* 20(6):953–66 [PubMed: 25456737]
19. Chae YK, Arya A, Malecek MK, Shin DS, Carneiro B, Chandra S, Kaplan J, Kalyan A, Altman JK, Platanias L et al. (2016) Repurposing metformin for cancer treatment: current clinical studies. *Oncotarget* 7(26):40767–80 [PubMed: 27004404]
20. Libby G, Donnelly LA, Donnan PT, Alessi DR, Morris AD, Evans JM (2009) New users of metformin are at low risk of incident cancer: a cohort study among people with type 2 diabetes. *Diabetes Care* 32(9):1620–5 [PubMed: 19564453]
21. Currie CJ, Poole CD, Jenkins-Jones S, Gale EA, Johnson JA, Morgan CL (2012) Mortality after incident cancer in people with and without type 2 diabetes: impact of metformin on survival. *Diabetes Care* 35(2):299–304 [PubMed: 22266734]
22. Zhang P, Li H, Tan X, Chen L, Wang S (2013) Association of metformin use with cancer incidence and mortality: a meta-analysis. *Cancer Epidemiol* 37(3):207–18 [PubMed: 23352629]
23. Niraula S, Dowling RJ, Ennis M, Chang MC, Done SJ, Hood N, Escallon J, Leong WL, McCready DR, Reedijk M et al. (2012) Metformin in early breast cancer: a prospective window of opportunity neoadjuvant study. *Breast Cancer Res Treat* 135(3):821–30 [PubMed: 22933030]
24. Malin D, Strekalova E, Petrovic V, Deal AM, Al Ahmad A, Adamo B, Miller CR, Ugolkov A, Livasy C, Fritchie K et al. (2014) aB-crystallin: a novel regulator of breast cancer metastasis to the brain. *Clin Cancer Res* 20(1):56–67 [PubMed: 24132917]
25. Moyano JV, Evans JR, Chen F, Lu M, Werner ME, Yehiely F, Diaz LK, Turbin D, Karaca G, Wiley E et al. (2006) aB-crystallin is a novel oncoprotein that predicts poor clinical outcome in breast cancer. *J Clin Invest* 116(1):261–70 [PubMed: 16395408]
26. Malin D, Chen F, Schiller C, Koblinski J, Cryns VL (2011) Enhanced metastasis suppression by targeting TRAIL receptor 2 in a murine model of triple-negative breast cancer. *Clin Cancer Res* 17(15):5005–15 [PubMed: 21653692]
27. Cummins JM, Kohli M, Rago C, Kinzler KW, Vogelstein B, Bunz F (2004) X-linked inhibitor of apoptosis protein (XIAP) is a nonredundant modulator of tumor necrosis factor-related apoptosis-

- inducing ligand (TRAIL)-mediated apoptosis in human cancer cells. *Cancer Res* 64(9):3006–8 [PubMed: 15126334]
28. Deng Y, Lin Y, Wu X. (2002) TRAIL-induced apoptosis requires Bax-dependent mitochondrial release of Smac/DIABLO. *Genes Dev* 16(1):33–45 [PubMed: 11782443]
 29. Do MT, Kim HG, Choi JH, Jeong HG (2014) Metformin induces microRNA-34a to downregulate the Sirt1/Pgc-1alpha/Nrf2 pathway, leading to increased susceptibility of wild-type p53 cancer cells to oxidative stress and therapeutic agents. *Free Radic Biol Med* 74:21–34 [PubMed: 24970682]
 30. Nazim UM, Moon JH, Lee JH, Lee YJ, Seol JW, Eo SK, Lee JH, Park SY (2016) Activation of autophagy flux by metformin downregulates cellular FLICE-like inhibitory protein and enhances TRAIL-induced apoptosis. *Oncotarget* 7(17):23468–81 [PubMed: 26992204]
 31. Tanaka R, Tomosugi M, Horinaka M, Sowa Y, Sakai T (2015) Metformin Causes G1-Phase Arrest via Down-Regulation of MiR-221 and Enhances TRAIL Sensitivity through DR5 Up-Regulation in Pancreatic Cancer Cells. *PloS One* 10(5):e0125779 [PubMed: 25955843]
 32. Zhang T, Wang X, He D, Jin X, Guo P (2014) Metformin sensitizes human bladder cancer cells to TRAIL-induced apoptosis through mTOR/S6K1-mediated downregulation of c-FLIP. *Anticancer Drugs* 25(8):887–97 [PubMed: 24714080]
 33. Chandel NS, Avizonis D, Reczek CR, Weinberg SE, Menz S, Neuhaus R, Christian S, Haegbarth A, Algire C, Pollak M (2016) Are metformin doses used in murine cancer models clinically relevant? *Cell Metab* 23(4):569–70 [PubMed: 27076070]
 34. Wahdan-Alaswad R, Fan Z, Edgerton SM, Liu B, Deng XS, Arnadottir SS, Richer JK, Anderson SM, Thor AD (2013) Glucose promotes breast cancer aggression and reduces metformin efficacy. *Cell Cycle* 12(24):3759–69 [PubMed: 24107633]
 35. Herold CI, Anders CK (2013) New targets for triple-negative breast cancer. *Oncology (Williston Park)* 27(9):846–54 [PubMed: 24282978]
 36. Yan H, Frost P, Shi Y, Hoang B, Sharma S, Fisher M, Gera J, Lichtenstein A (2006) Mechanism by which mammalian target of rapamycin inhibitors sensitize multiple myeloma cells to dexamethasone-induced apoptosis. *Cancer Res* 66(4):2305–13 [PubMed: 16489035]
 37. Allensworth JL, Aird KM, Aldrich AJ, Batinic-Haberle I, Devi GR (2012) XIAP inhibition and generation of reactive oxygen species enhances TRAIL sensitivity in inflammatory breast cancer cells. *Mol Cancer Ther* 11(7): 1518–27 [PubMed: 22508521]
 38. Tseng HY, Chen LH, Ye Y, Tay KH, Jiang CC, Guo ST, Jin L, Hersey P, Zhang XD (2012) The melanoma-associated antigen MAGE-D2 suppresses TRAIL receptor 2 and protects against TRAIL-induced apoptosis in human melanoma cells. *Carcinogenesis* 33(10):1871–81 [PubMed: 22791814]
 39. Panner A, James CD, Berger MS, Pieper RO (2005) mTOR controls FLIPS translation and TRAIL sensitivity in glioblastoma multiforme cells. *Mol Cell Biol* 25(20):8809–23 [PubMed: 16199861]




Figure 1. Metformin sensitizes TNBC cells to TRAIL receptor agonists.

Crystal violet cell survival assay of TNBC cells preincubated with vehicle or metformin (1 mM for MDA-MB-468 cells or 5 mM for MDA-MB-231-mCherry and GILM2 cells) for 48 hours, and treated with vehicle or TRAIL, mapatumumab or lexatumumab (each 1.5 μ g/ml for MDA-MB-231-mCherry and GILM2 cells or 4 μ g/ml for MDA-MB-468 cells) for an additional 24 hours (MDA-MB-231-mCherry and GILM2 cells) or 48 hours (MDA-MB-468 cells). Left panels: representative images. Right panels: the percentage confluence of crystal

violet-positive cells was scored (mean $\pm$ SEM, n = 3). In all panels, *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. Metformin sensitizes transformed cells breast epithelial cells to TRAIL receptor agonists.

Crystal violet cell survival assay of MCF-10A-Vector or MCF-10A-Ras cells preincubated with vehicle or metformin (5 mM) for 48 hours, and then treated with vehicle or TRAIL, mapatumumab or lexatumumab (each 1.5 μ g/ml) for an additional 24 hours. Left panels: representative images. Right panels: the percentage confluence of crystal violet-positive cells was scored (mean $\pm$ SEM, n = 3). In all panels, *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$.




Figure 3. Metformin enhances TRAIL-induced caspase activation in TNBC cells.

A, Immunoblots of TNBC cells preincubated with vehicle or metformin (5 mM) for 48 hours and then treated with vehicle or TRAIL (1.5 μ g/ml for MDA-MB-231-mCherry and GILM2 cells or 4 μ g/ml for MDA-MB-468 cells) for 16 hours. **B**, TNBC cells were preincubated with vehicle or metformin (5 mM) for 48 hours, treated with vehicle or TRAIL (1.5 μ g/ml for MDA-MB-231-mCherry and GILM2 cells or 4 μ g/ml for MDA-MB-468 cells) for 24 hours, and caspase-3/7 activity was measured (mean $\pm$ SEM, n=3). **C**, Immunoblots of MCF-10A breast epithelial cells stably expressing vector or H-RasV12

treated preincubated with vehicle or metformin (5 mM) for 48 hours and then treated with vehicle or TRAIL (1.5 µg/ml) for 16 hours. **D**, MCF-10A-Vector or MCF-10A-RAS cells were preincubated with vehicle or metformin (5 mM) for 48 hours, treated with vehicle or TRAIL (1.5 µg/ml) for 24 hours, and caspase-3/7 activity was measured (mean ± SEM, n=3). In **(B)** and **(D)**, **, $P < 0.01$ and ***, $P < 0.001$.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 4. Metformin does not alter cell surface expression of TRAIL receptors in TNBC cells. **A**, TNBC cells were treated with vehicle or metformin (5 mM) for 72 hours. TRAIL-R2 (left panel) and TRAIL-R1 (right panel) mRNA levels were determined by real-time PCR and were normalized to expression in vehicle-treated TNBC cells. **B**, TNBC cells were treated with vehicle or MF (5 mM) for 72 hours and then incubated with IgG, TRAIL-R1 or TRAIL-R2 Ab. Cell surface expression of TRAIL receptors was determined by flow cytometry. Grey bar: TNBC cells incubated with IgG. Blue line: TNBC cells treated with vehicle and incubated with TRAIL-R1 (bottom panels) or TRAIL-R2 (top panels) Ab. Red

line: TNBC cells treated with metformin and incubated with TRAIL-R1 (bottom panels) or TRAIL-R2 (top panels) Ab.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 5. Metformin reduces XIAP protein expression in TNBC cells and silencing XIAP sensitizes TNBC cells to TRAIL.

A, Immunoblot of XIAP protein levels (left panel) and real-time PCR analysis of XIAP mRNA levels (right panel) in TNBC cells treated with vehicle or metformin (5 mM) for 72 hours (immunoblots) or 48 hours (real-time PCR). **B**, MDA-MB-231-mCherry cells were transfected with a non-silencing control siRNA (si-Control) or one of two different siRNAs targeting XIAP (si1-XIAP or si2-XIAP). Immunoblot of XIAP expression 48 hours after transfection. **C**, Crystal violet cell survival assay of MDA-MB-231-mCherry cells

transfected with control or XIAP siRNAs, and 48 hours later treated with vehicle or TRAIL (0.3 $\mu\text{g/ml}$) for 24 hours. Left panel: representative images. Right panel: quantification of cell confluence in 3 fields for each treatment (mean $\pm$ SEM, $n = 3$) *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$. **D**, Silencing XIAP enhances TRAIL-induced caspase activation in TNBC cells. MDA-MB-231-mCherry cells were transfected with control or XIAP siRNAs, and 48 hours later were treated with vehicle or TRAIL (0.3 $\mu\text{g/ml}$) for 16 hours. PARP (full length and caspase-cleaved) and procaspase-3 levels were determined by immunoblotting.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 6. Metformin enhances the antitumor effects of TRAIL in an orthotopic model of metastatic TNBC.

Three weeks after intraductal injection of tumor cells, female NSG mice with GILM2-mCherry mammary tumors were randomized into 4 groups (10 mice per group): control group (PBS *i.p.* daily for three weeks), metformin (2 mg per ml in drinking water for three weeks), TRAIL (10 mg/kg *i.p.*, daily for three weeks) or metformin plus TRAIL (same dosing as in single agent arms). **A**, the percentage of the original mammary tumor volume (at 3 weeks) in each group (mean $\pm$ SEM, $n = 10$ mice per group). **B**, the percentage of the

surface area occupied by lung metastases (mean $\pm$ SEM, n = 10 mice per group). **C**, the percentage active caspase-3-positive tumor cells in mammary tumors (left panel) or metastatic lung tumors (right panel) after treatment (mean $\pm$ SEM, n = 3 tumors per group). **D**, immunoblot of XIAP expression in mammary tumors from control vehicle-treated or metformin-treated mice (n=3 mammary tumors per group). XIAP expression normalized to β -actin was measured by Image J analysis. **E**, body weight of the mice in each treatment group at the end of the study (mean $\pm$ SEM, n = 10 mice per group). In all panels, **, $P < 0.01$, ***, $P < 0.001$ versus vehicle-treated mice or the indicated comparison.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Metformin Use is Associated with Improved Survival for Patients with Advanced Prostate Cancer on Androgen Deprivation Therapy



Kyle A. Richards,* Jinn-ing Liou, Vincent L. Cryns, Tracy M. Downs, E. Jason Abel and David F. Jarrard*

From the Section of Urology, Department of Surgery, William S. Middleton Memorial Veterans Hospital (KAR) and Departments of Urology (KAR, TMD, EJA, DFJ) and Medicine (JL, VLC), University of Wisconsin-Madison, Madison, Wisconsin

Abbreviations and Acronyms

ADT = androgen deprivation therapy

CDW = Corporate Data Warehouse

CSS = cancer specific survival

DM = diabetes mellitus

IPSW = inverse propensity score weighted

OS = overall survival

PCa = prostate cancer

PSA = prostate specific antigen

SRE = skeletal related event

VA = Veterans Administration

Purpose: Metformin is commonly prescribed for patients with type 2 diabetes mellitus. We hypothesized that metformin plus androgen deprivation therapy may be beneficial in combination. Our objective was to assess this combination in a retrospective cohort of patients with advanced prostate cancer.

Materials and Methods: Using national Veterans Affairs databases we identified all men diagnosed with prostate cancer between 2000 and 2008 who were treated with androgen deprivation therapy with followup through May 2016. Study exclusions included treatment with androgen deprivation therapy for 6 months or longer, or receipt of androgen deprivation therapy concurrently with localized radiation. Three patient cohorts were developed, including no diabetes mellitus, diabetes mellitus with no metformin and diabetes mellitus with metformin. Cox proportional HRs were calculated for overall survival, skeletal related events and cancer specific survival.

Results: After exclusions the cohort consisted of 87,344 patients, including 61% with no diabetes mellitus, 22% with diabetes mellitus and no metformin, and 17% with diabetes mellitus on metformin. Cox proportional hazard analysis of overall survival showed improved survival in men with diabetes mellitus on metformin (HR 0.82, 95% CI 0.78–0.86) compared to those with diabetes mellitus who were not on metformin (HR 1.03, 95% CI 0.99–1.08). The reference group was men with no diabetes mellitus. Cox proportional hazard analysis of predictors of skeletal related events revealed a HR of 0.82 (95% CI 0.72–0.93) in men with diabetes mellitus on metformin. Cox proportional hazard analysis of cancer specific survival showed improved survival in men with diabetes mellitus

Accepted for publication June 13, 2018.

No direct or indirect commercial incentive associated with publishing this article.

The corresponding author certifies that, when applicable, a statement(s) has been included in the manuscript documenting institutional review board, ethics committee or ethical review board study approval; principles of Helsinki Declaration were followed in lieu of formal ethics committee approval; institutional animal care and use committee approval; all human subjects provided written informed consent with guarantees of confidentiality; IRB approved protocol number; animal approved project number.

Supported by Department of Defense Grant PC150221 via the Prostate Cancer Research Program, and William S. Memorial Veterans Hospital resource and facility use.

The contents do not represent the views of the United States Department of Veterans Affairs or the United States Government.

* Correspondence: Department of Urology, Urology Veterans Affairs, University of Wisconsin-Madison, 1685 Highland Ave., Madison, Wisconsin 53705 (telephone: 608-262-0759; FAX: 608-262-6453; e-mail: richardsk@urology.wisc.edu) and Department of Urology, University of Wisconsin Carbone Cancer Center, University of Wisconsin-Madison, 1685 Highland Ave., Madison, Wisconsin 53705 (telephone: 608-262-0759; FAX: 608-262-6453).

Editor's Note: This article is the first of 5 published in this issue for which category 1 CME credits can be earned. Instructions for obtaining credits are given with the questions on pages 1382 and 1383.

on metformin (HR 0.70, 95% CI 0.64–0.77) vs those with diabetes mellitus without metformin (HR 0.93, 95% CI 0.85–1.00). The reference group was men with no diabetes mellitus.

Conclusions: Metformin use in veterans with prostate cancer who receive androgen deprivation therapy is associated with improved oncologic outcomes. This association should be evaluated in a prospective clinical trial.

Key Words: prostatic neoplasms, metformin, gonadotropin-releasing hormone, analogs, derivatives, and diabetes mellitus

THE past decade has witnessed remarkable advances with 6 new therapies approved by the United States FDA (Food and Drug Administration) for the treatment of men with advanced PCa.¹ Despite these advances nearly 27,000 men died of PCa in 2017, highlighting the ongoing need for additional therapeutic options in men in whom conventional treatments fail.²

ADT remains the standard first line approach for metastatic PCa. It leads to regression but rarely to cure as hormone insensitive disease invariably develops from resistant clones. These cells that remain after the initiation of ADT represent an underexplored therapeutic niche which may improve therapy. In support a recent randomized clinical trial demonstrated that up-front chemotherapy with ADT improved survival by 10.5 months vs ADT alone in hormone naïve patients, suggesting that initiating ADT induces susceptibilities in PCa cells that make them amenable to synergistic treatments.³

Metformin, a commonly used insulin sensitizer, is a first line agent for patients with type 2 DM. There is scientific evidence for the antineoplastic effects that metformin may have for various cancers but its impact in men with advanced PCa and its usefulness in combination with other treatments remain poorly studied.^{4,5} Metformin activates AMPK (AMP-activated protein kinase), which inhibits mTOR (mammalian target of rapamycin), a central regulator of cell growth.^{6,7} ADT has been shown to induce senescence in androgen sensitive cells, a phenotype with high glycolysis and proteolytic turnover.^{8–10}

Given these data, we hypothesized that metformin may be beneficial in combination with ADT to target PCa cells that persist after ADT, leading to improved survival. To test this approach we performed a large observational study evaluating the impact of metformin use on cancer outcomes in men with PCa who were being treated with ADT.

MATERIALS AND METHODS

Data Source

The study was approved by local institutional review boards. The VA provides care to more than 20 million

veterans at a total of more than 1,400 centers. All care processes are captured via the VistA (Veterans Information System Technology Architecture) electronic health record, which provides a longitudinal view of patients receiving care nationwide, including diagnoses, procedures, medications, laboratory findings, physiological measurements, text notes and reports.¹¹ Data are aggregated from individual VistA systems to the VA CDW, where the data are prepared for use.

Study Population

To develop a cohort of men with PCa on ADT we identified all 558,252 men diagnosed with PCa (ICD-9 code 185) in the VA CDW from 2000 to 2008. In this cohort we included only the 129,672 men receiving ADT by querying the pharmacy domain for VA formulary approved ADT medications, including leuprolide, goserelin, bicalutamide, flutamide and nilutamide, from 2000 through May 31, 2016. These were the only approved ADT medications on formulary during the study period.

We excluded from study 33,312 patients with no information on the ADT medication supply days, quantity or dose, those on ADT for 6 months or less and/or 10,960 receiving ADT concurrent with primary radiation therapy of the prostate, leaving a final cohort of 87,344 patients for our analytical file. ADT was entered as a time dependent variable in the models. Longitudinal data on patients were compiled until death or until the study end of May 31, 2016, at which point they were censored.

We divided the study population into 3 cohorts and defined DM in the VA using a previously published algorithm with ICD-9 codes 250.00 or 250.02.¹² Comparator groups included 1) no DM, 2) DM and no prescription of metformin for 180 days or longer during the study period and 3) DM with a prescription of metformin for 180 days or longer during the study period.

Outcomes of Interest

The primary outcome of interest in this study was OS. Secondary outcomes of interest included SRE and death from PCa (CSS). The dependent variable used in our analyses was the interval from the ADT starting date to death from any cause, SRE and/or death from PCa. SRE served as a surrogate for progression using a previously described claims based model to identify SRE.¹³

Predictors and Measures

The metformin group consisted of patients for whom metformin was prescribed for 180 days or longer. We did not exclude patients with exposure to insulin or other glucose lowering medications because the impact on

cancer outcomes is conflicting.^{14,15} Prior clinical trials on metformin consisted of at least 24 weeks of exposure. Therefore, we chose to define drug use as at least 180 days based on this and other studies.^{12,16} There were no metformin users in the no DM group. Metformin use was entered as a time dependent variable in the models, allowing for patients to move from a period of exposure to a period of nonexposure.

Covariates adjusted for in the analyses included the demographic and clinical characteristics of each patient, including age at ADT initiation, race, the Charlson comorbidity index, Agent Orange exposure, PSA at ADT initiation, diagnosis year, Gleason score, local therapy receipt,¹⁷ docetaxel receipt and insulin use.

Statistical Analysis

Medians were compared by the Mann-Whitney U test. The Fisher exact and chi-square tests were used to compare categorical variables. We performed multivariable Cox proportional hazard analyses to assess for independent predictors of OS, SRE and CSS. We then calculated a propensity score by multinomial logistic regression and used it to adjust the IPSW in the final models.¹⁸ We constructed IPSW Kaplan-Meier curves of OS, SRE and CSS, and performed the log rank test. We also performed sensitivity analysis of CSS to account for competing risks as a result of death from other causes using a subdistribution hazard model adapted for time dependent covariates.^{19,20} Finally, we performed subset IPSW multivariable Cox proportional hazard analyses to assess for independent predictors of OS, SRE and CSS in patients with PSA greater than 20 ng/ml at ADT initiation. Statistical significance was considered at 2-sided $p < 0.05$ and statistical analysis was performed with Stata® 14.

RESULTS

The total cohort available for analysis after exclusions consisted of 87,344 patients, including 53,893 (61%) in the no DM group, 18,934 (22%) in the DM without metformin group and 14,517 (17%) in the DM plus metformin group. The metformin group was younger with a median age of 71.0 years (IQR 64–76) compared to the no DM group (75.0, IQR 69–80) and the DM without metformin group (75.0, IQR 69–79, $p < 0.001$, see table).

The OS was longest in the metformin group as represented by the IPSW Kaplan-Meier curve ($p = 0.005$, fig. 1). The adjusted Cox proportional hazard multivariable analysis identified that the metformin group was associated with improved OS (HR 0.82, 95% CI 0.78–0.86, $p < 0.001$) vs the DM without metformin group (HR 1.03, 95% CI 0.99–1.08, $p = 0.18$) with the no DM group as the reference group. A dose-response relationship was observed in the cumulative duration of metformin use before and after IPSW with 36 months or more found to be most protective (HR 0.69, 95% CI

0.65–0.74, $p < 0.001$, supplementary table 1, <http://jurology.com/>).

The proportion of patients with SREs was highest in the metformin group at 11.1% but time to SRE was also longest in the metformin group as represented by the IPSW Kaplan-Meier curve ($p = 0.005$, fig. 2). The adjusted Cox proportional hazard multivariable analysis identified that the metformin group was associated with a decreased risk of SRE (HR 0.84, 95% CI 0.74–0.96, $p = 0.009$) vs the DM without metformin group (HR 1.08, 95% CI 0.96–1.23, $p = 0.20$) with the no DM group as the reference group. A dose-response relationship was observed in the cumulative duration of metformin use before and after IPSW with 36 months or more found to be most protective (HR 0.70, 95% CI 0.59–0.83, $p < 0.001$, supplementary table 2, <http://jurology.com/>).

The proportion of patients documented to have died of PCa was lowest in the metformin group at 9.3% as shown by the IPSW Kaplan-Meier curve ($p < 0.001$, fig. 3). The adjusted Cox proportional hazard multivariable analysis identified that the metformin group was associated with improved CSS (HR 0.70, 95% CI 0.64–0.77, $p < 0.001$) vs the DM without metformin group (HR 0.93, 95% CI 0.85–1.00, $p = 0.0054$) with the no DM group as the reference group. A dose-response relationship was observed in the cumulative duration of metformin use before and after IPSW with 36 months or more found to be most protective (HR 0.58, 95% CI 0.51–0.66, $p < 0.001$, supplementary table 3, <http://jurology.com/>). After accounting for competing risks as a result of death from other causes the decreased risk observed between metformin for 36 months or greater and prostate cancer mortality remained statistically significant (HR 0.66, 95% CI 0.58–0.75, $p < 0.001$).

The subset Cox proportional hazard multivariable analyses to assess for independent predictors of OS, SRE and CSS in patients with PSA greater than 20 ng/ml at the time of ADT initiation revealed no change in the noted associations (supplementary tables 1, 3 and 4, <http://jurology.com/>). However, the association with SRE was no longer statistically significant (supplementary tables 2, 5 and 6, <http://jurology.com/>).

DISCUSSION

This large observational study revealed that metformin use was associated with improved oncologic outcomes in men with PCa on ADT. Prior studies evaluating the impact of metformin in men with PCa focused on disease at diagnosis or early treatment. To our knowledge the current study is unique in evaluating the impact of metformin in men on ADT as these drugs may have an additive effect.

Characteristics of 87,344 patients with prostate cancer on androgen deprivation therapy

	No Diabetes		Diabetes		p Value
			No Metformin	Metformin	
No. pts	53,893		18,934	14,517	
Median age (IQR)	75.0 (69–80)		75.0 (69–79)	71.0 (64–76)	<0.001
No. race (%):					<0.001
White	35,416 (65.7)		11,141 (58.8)	8,760 (60.3)	
Black	8,791 (16.3)		4,707 (24.9)	3,337 (23.0)	
Other	9,686 (18.0)		3,086 (16.3)	2,420 (16.7)	
No. Charlson comorbidity score (%):					<0.001
0–1	42,490 (78.8)		14,065 (74.3)	10,960 (75.5)	
2–3	10,477 (19.4)		3,937 (20.8)	2,936 (20.2)	
Greater than 3	926 (1.7)		932 (4.9)	621 (4.3)	
No. Agent Orange exposure (%)	1,804 (3.4)		696 (3.7)	975 (6.7)	<0.001
No. mos ADT (%):					<0.001
Less than 12	16,744 (31.1)		5,360 (28.3)	3,865 (26.6)	
12–Less than 24	14,509 (26.9)		4,903 (25.9)	3,629 (25.0)	
24–Less than 36	7,925 (14.7)		2,894 (15.3)	2,230 (15.4)	
36 or Greater	14,715 (27.3)		5,777 (30.5)	4,793 (33.0)	
Median ng/dl PSA IQR):*					<0.001
Less than 4	14,191 (26.3)		5,332 (28.2)	4,591 (31.6)	
4–10	7,738 (14.4)		2,970 (15.7)	2,809 (19.4)	
Greater than 10	16,768 (31.1)		5,921 (31.3)	4,253 (29.3)	
Missing	15,196 (28.2)		4,711 (24.9)	2,864 (19.7)	
No. diagnosis yr (%):					<0.001
2000–2004	41,496 (77.0)		15,225 (80.4)	10,453 (72.0)	
2005–2008	12,397 (23.0)		3,709 (19.6)	4,064 (28.0)	
No. Gleason score (%):					<0.001
6	3,487 (6.5)		1,438 (7.6)	1,402 (9.7)	
7	4,542 (8.4)		1,630 (8.6)	1,719 (11.8)	
8–10	6,094 (11.3)		2,126 (11.2)	1,985 (13.7)	
Missing	39,770 (73.8)		13,740 (72.6)	9,411 (64.8)	
No. local therapy (%)	3,964 (7.4)		1,387 (7.3)	1,788 (12.3)	<0.001
No. docetaxel (%)	1,803 (3.4)		508 (2.7)	584 (4.0)	<0.001
No. insulin (%)			8,755 (46.2)	9,297 (64.0)	<0.001
No. vital status (% deceased)	42,133 (78.2)		15,215 (80.4)	9,512 (65.5)	<0.001
Median yrs overall survival (IQR)	5.1 (2.5–8.8)		5.4 (2.7–9.0)	6.8 (3.5–10.1)	<0.001
No. prostate cancer death (%)	5,522 (10.3)		1,959 (10.4)	1,337 (9.2)	<0.001
Skeletal related event:					
No. pts (%)	4,863 (9.0)		1,833 (9.7)	1,609 (11.1)	<0.001
Median yrs to event (IQR)	4.7 (2.2–8.3)		4.9 (2.3–8.5)	6.1 (2.8–9.5)	<0.001

* At androgen deprivation therapy initiation.

Residual cancer cells after ADT are characterized by metabolic abnormalities which may be targeted preferentially by metformin.¹⁰ Capturing prescription medication use is vital for this type of analysis and VA databases provided an ideal platform to perform this study since approximately 83% of VA enrollees who use VA pharmacy benefits fill prescriptions through a VA pharmacy.²¹ Additionally, the VA provides continuous and equal access care for the majority of these veterans as monitored through 1 health care record, making outcomes easier to determine.

Our analysis, which controlled for multiple variables, identified that metformin use was associated with improved OS (HR 0.82) in dose dependent fashion. CSS also improved (HR 0.70), specifically in men with DM receiving metformin compared to the other groups. It was difficult to clearly define the patients in whom ADT was initiated for metastatic hormone sensitive PCa in this data set. However, controlling for PSA and performing subset analysis

of patients with PSA greater than 20 ng/ml at ADT initiation confirmed the overall and cancer specific survival advantage to being on metformin. In this higher PSA subset there were improved outcomes in patients at higher risk for metastatic disease, the group in which ADT is typically initiated for modern, hormone sensitive PCa. The recognition of increased cardiac, bone density and other side effects has led to delaying ADT in many patients with micrometastatic disease.²²

To our knowledge studies to date have not focused on a potential additive role of metformin at the time of ADT initiation. In a meta-analysis of 21 eligible studies metformin receipt was associated with decreased PCa risk (OR 0.91) and biochemical recurrence following treatment (HR 0.81) but not with improved OS in patients with PCa (HR 0.86, 95% CI 0.64–1.14).²³ Our data do not discount a role for metformin in improving disease in the castrate resistant state. In a phase 2 clinical trial of metformin in 44 men with progressive castrate




Figure 1. Kaplan-Meier curve of overall survival stratified by nondiabetes, DM plus metformin and DM without metformin after IPSW adjustment (log rank test $p = 0.005$).

resistant PCa, Metformin Hydrochloride as First-Line Therapy in Treating Patients With Locally Advanced or Metastatic Prostate Cancer (ClinicalTrials.gov NCT01243385), 36% of the men were free of progression at the 12-week followup with no grade 3 or 4 toxicity, suggesting some activity in this space.²⁴ The multi-arm, multistage, randomized STAMPEDE (Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy, ClinicalTrials.gov NCT00268476) clinical trial is currently recruiting patients in a metformin plus ADT arm to assess the safety and

efficacy of this approach. In addition, the randomized, prospective, phase 3 PRIME (Metformin in Patients Initiating ADT as Prevention and Intervention of Metabolic Syndrome, ClinicalTrials.gov NCT03031821) clinical trial is under way to assess the proportion of patients in whom metabolic syndrome develops.

The duration of metformin receipt may influence outcomes as suggested by our data and those of others. Margel et al performed a retrospective cohort study to evaluate associations of the cumulative duration of antidiabetic drug use after PCa




Figure 2. Kaplan-Meier curve of skeletal related events stratified by nondiabetes, DM plus metformin and DM without metformin after IPSW adjustment (log rank test $p = 0.005$).




Figure 3. Kaplan-Meier curve of prostate cancer specific mortality stratified by nondiabetes, DM plus metformin and DM without metformin after IPSW adjustment (log rank test $p = 0.001$).

diagnosis with CSS and OS in patients with type 2 DM.²⁵ Each additional 6 months of metformin resulted in an adjusted CSS HR of 0.76 (95% CI 0.64–0.89). However, no relationship was seen between the cumulative use of other antidiabetic drugs and CSS or OS. Furthermore, we found similar adjusted HRs of OS and CSS in our cohort, noting that our study included patients without DM as a functional control group and all study patients were on ADT. This highlighted the difference in our study design.

In addition, we found that metformin was associated with a reduced risk of SRE, which we used as a measure of progression. Notably progression was not assessed in the study by Margel et al.²⁵ There was an increased incidence of SRE in the metformin group but when controlling for time and other covariates, the risk of SRE was attenuated in the metformin group. We chose the SRE algorithm as a measure of progression since we thought that it was a more sensitive measure in this patient population, given the low rate of chemotherapy or novel anti-androgen therapies.

In our study we aimed to specifically assess the effects of metformin in patients on ADT based on the potential for an additive benefit of these 2 agents in preclinical studies.^{6–10} In vitro and in vivo studies suggested that combining metformin with bicalutamide would result in reduced proliferation of androgen receptor positive cells and apoptosis of androgen receptor negative cells.²⁶ ADT induces senescence in a population of PCa cells,²⁷ which generates inherent susceptibilities that may be used. These cells have high levels of protein

turnover and gluconeogenesis, rendering them susceptible to proteolytic inhibitors and agents that alter sugar metabolism.¹⁰ Metformin activates AMPK, a sensor of cellular energy change, and switches on energy producing pathways as well as inhibiting mTOR.^{6,7} This leads to apoptosis of these residual cells, providing a molecular rationale for this response.

Other studies showed that long-term ADT use may also induce metabolic syndrome and in turn increase the risk of cardiovascular morbidity.²⁸ Metformin may have benefits in reducing these effects, in addition to the direct antineoplastic activity.

There are several limitations to our study. 1) This was a retrospective observational study with potentially unmeasured confounding variables and/or missing variables. 2) Because national VA data are developed as an administrative data set via the CDW, we could not account for drug discontinuation reasons, key variable mis-coding, complete laboratory data on the entire cohort, socioeconomic status, body mass index, exercise, smoking, local therapies received outside the VA or stage. In addition, we could not account for other potential health benefits of metformin which may have impacted our results, including an improvement in DM and cardiovascular health. However, our large sample size and our propensity score matching enabled us to control for other important confounding factors. 3) Finally, our population of aging veterans may lack external validity. Additional studies are warranted in other populations.

CONCLUSIONS

Metformin use was associated with improved OS, SRE and CSS in men with PCa who were also receiving ADT. We believe that these findings may

be related to an additive antineoplastic effect between metformin and ADT. Additional studies are warranted to further validate these findings and establish causation via well designed clinical trials.

REFERENCES

1. United States Food and Drug Administration: Hematology/Oncology (Cancer) Approvals and Safety Notifications. Available at <https://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm279174.htm>. Accessed August 15, 2017.
2. Siegel RL, Miller KM and Jemal A: Cancer statistics, 2017. *Ca Cancer J Clin* 2017; **67**: 7.
3. Sweeney CJ, Chen YH, Carducci M et al: Chemohormonal therapy in metastatic hormone-sensitive prostate cancer. *N Engl J Med* 2015; **373**: 737.
4. Jiralerspong S, Palla SL, Giordano SH et al: Metformin and pathologic complete responses to neoadjuvant chemotherapy in diabetic patients with breast cancer. *J Clin Oncol* 2009; **27**: 3297.
5. Pollak M: Metformin and other biguanides in oncology: advancing the research agenda. *Cancer Prev Res (Phila)* 2010; **3**: 1060.
6. Rocha GZ, Dias MM, Rapelle ER et al: Metformin amplifies chemotherapy-induced AMPK activation and antitumoral growth. *Clin Cancer Res* 2011; **17**: 3993.
7. Dowling RJ, Zakikhani M, Fantus IG et al: Metformin inhibits mammalian target of rapamycin-dependent translation initiation in breast cancer cells. *Cancer Res* 2007; **67**: 10804.
8. Ewald JA, Desotelle JA, Church DR et al: Androgen deprivation induces senescence characteristics in prostate cancer cells in vitro and in vivo. *Prostate* 2013; **73**: 337.
9. Pernicova Z, Slabakova E, Kharaschvili G et al: Androgen depletion induces senescence in prostate cancer cells through down-regulation of Skp2. *Neoplasia* 2011; **13**: 526.
10. Dorr JR, Yu Y, Milanovic M et al: Synthetic lethal metabolic targeting of cellular senescence in cancer therapy. *Nature* 2013; **501**: 421.
11. United States Department of Veterans Affairs: Health Services Research and Development—VA Informatics and Computing Infrastructure (VINCI). Available at https://www.hsrd.research.va.gov/for_researchers/vinci/. Accessed June 12, 2018.
12. Chen-Pin W, Javier H, Lorenzo C et al: Statins and finasteride use differentially modify the impact of metformin on prostate cancer incidence in men with type 2 diabetes. *Ann Transl Med Epidemiol* 2014; **1**: pii:1004.
13. Aly A, Onukwughu E, Woods C et al: Measurement of skeletal related events in SEER-Medicare: a comparison of claims-based methods. *BMC Med Res Methodol* 2015; **15**: 65.
14. Hou G, Zhang S, Zhang X et al: Clinical pathologic characteristics and prognostic analysis of 1013 breast cancer patients with diabetes. *Breast Cancer Res Treat* 2013; **137**: 807.
15. Curie CJ, Poole CD, Jenkins-Jones S et al: Mortality after incident cancer in people with and without type 2 diabetes: impact of metformin on survival. *Diabetes Care* 2012; **35**: 299.
16. Curie CJ, Poole CD and Gale EA: The influence of glucose-lowering therapies on cancer risk in type 2 diabetes. *Diabetologia* 2009; **52**: 1766.
17. Ritch CR, Graves AJ, Keegan KA et al: Increasing use of observation among men at low risk for prostate cancer mortality. *J Urol* 2015; **193**: 801.
18. Austin PC: An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate Behav Res* 2011; **46**: 399.
19. Fine JP and Gray RJ: A proportional hazards model for subdistribution of a competing risk. *J Am Stat Assoc* 1999; **94**: 496.
20. Ruan PK and Gray RJ: Analyses of cumulative incidence functions via non-parametric multiple imputation. *Stat Med* 2008; **27**: 5709.
21. PBM Drug Utilization Annual Report. Washington, D. C.: Veterans Affairs Pharmacy Benefits Management Services 2016.
22. Chen DY, See LC, Liu JR et al: Risk of cardiovascular ischemic events after surgical castration and gonadotropin-releasing hormone agonist therapy for prostate cancer: a nationwide cohort study. *J Clin Oncol* 2017; **35**: 3697.
23. Yu H, Yin L, Jiang X et al: Effect of metformin on cancer risk and treatment outcome of prostate cancer: a meta-analysis of epidemiological observational studies. *PLoS One* 2014; **9**: e116327.
24. Rothermundt C, Hayoz S, Templeton AJ et al: Metformin in chemotherapy-naïve castration-resistant prostate cancer: a multicenter phase 2 trial (SAKK 08/09). *Eur Urol* 2014; **66**: 468.
25. Margel D, Urbach DR, Lipscombe LL et al: Metformin use and all-cause and prostate cancer-specific mortality among men with diabetes. *J Clin Oncol* 2013; **31**: 3069.
26. Colquhoun AJ, Venier NA, Vandersluis AD et al: Metformin enhances the antiproliferative and apoptotic effect of bicalutamide in prostate cancer. *Prostate Cancer Prostatic Dis* 2012; **15**: 346.
27. Blute ML Jr, Damaschke N, Wagner J et al: Persistence of senescent prostate cancer cells following prolonged neoadjuvant androgen deprivation therapy. *PLoS One* 2017; **12**: e0172048.
28. Bosco C, Crawley D, Adolfsson J et al: Quantifying the evidence for the risk of metabolic syndrome and its components following androgen deprivation therapy for prostate cancer: a meta-analysis. *PLoS One* 2015; **10**: e0117344.

EDITORIAL COMMENT

Metformin has been proposed to have efficacy in prostate cancer through several putative mechanisms. They include effects on insulin responsive prostate cancers via attenuation of hyperinsulinemia, inhibition of oxidative phosphorylation causing energetic stress in cancer cells and potentially delaying the

development of castrate resistant prostate cancer, which hyperinsulinemia can potentiate.¹ Retrospective studies have supported the hypothesis that metformin can improve outcomes in patients with prostate cancer (reference 25 in article).² However, this report is by far the largest observational study to demonstrate a strong

association between metformin use and improved oncologic outcomes.

The time has come to confirm these findings as well as any benefit of decreasing the metabolic morbidities of androgen deprivation in prospective clinical trials. Two phase III studies are presently under way in this arena. The ongoing multi-arm comparative STAMPEDE study recently added a metformin arm to evaluate overall survival in patients with advancing or metastatic prostate cancer. The PRIME study compares metformin to placebo in patients in whom intermittent androgen deprivation therapy is initiated for metabolic morbidity and

efficacy outcomes. These studies could confirm the benefit of one of the simplest and cost-effective therapies for prostate cancer in a long time.

Bernhard J. Eigel


*BC Cancer
Vancouver, British Columbia
and*

Nawaid Usmani

*Cross Cancer Institute
Edmonton, Alberta
Canada*

REFERENCES

1. Foretz M, Guigas B, Bertrand L et al: Metformin: from mechanisms of action to therapies. *Cell Metab* 2014; **20**: 953.
2. Spratt DE, Zhang C, Zumsteg ZS et al: Metformin and prostate cancer: reduced development of castration-resistant disease and prostate cancer mortality. *Eur Urol* 2013; **63**: 709.



Review

Combination therapy with androgen deprivation for hormone sensitive prostate cancer: A new frontier



Tyler Etheridge^a, Shivashankar Damodaran^a, Adam Schultz^a,
Kyle A. Richards^a, Joseph Gawdzik^a, Bing Yang^a,
Vincent Cryns^{b,c}, David F. Jarrard^{a,c,*}

^a Department of Urology, University of Wisconsin-Madison, Madison, WI, USA

^b Department of Medicine, University of Wisconsin-Madison, Madison, WI, USA

^c University of Wisconsin-Madison, Carbone Cancer Center, University of Wisconsin-Madison, Madison, WI, USA

Received 24 April 2018; received in revised form 1 June 2018; accepted 22 June 2018
Available online 15 September 2018

KEYWORDS

Prostate cancer;
Cellular senescence;
Androgen deprivation
therapy;
Combination therapy;
Synthetic lethal
targeting;
Metformin;
Statins

Abstract Androgen deprivation therapy (ADT) has been the standard of care for the last 75 years in metastatic hormone sensitive prostate cancer (PCa). However, this approach is rarely curative. Recent clinical trials have demonstrated that ADT combined with other agents, notably docetaxel and abiraterone, lead to improved survival. The mechanisms surrounding this improved cancer outcomes are incompletely defined. The response of cancer cells to ADT includes apoptosis and cell death, but a significant fraction remains viable. Our laboratory has demonstrated both *in vitro* and *in vivo* that cellular senescence occurs in a subset of these cells. Cellular senescence is a phenotype characterized by cell cycle arrest, senescence-associated β -galactosidase (SA- β -gal), and a hypermetabolic state. Positive features of cellular senescence include growth arrest and immune stimulation, although persistence may release cytokines and growth factors that are detrimental. Senescent tumor cells generate a catabolic state with increased glycolysis, protein turnover and other metabolic changes that represent targets for drugs, like metformin, to be applied in a synthetic lethal approach. This review examines the response to ADT and the putative role of cellular senescence as a biomarker and therapeutic target in this context.

© 2019 Editorial Office of Asian Journal of Urology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Department of Urology, University of Wisconsin-Madison, Madison, WI, USA.

E-mail address: jarrard@urology.wisc.edu (D.F. Jarrard).

Peer review under responsibility of Second Military Medical University.

1. Introduction

Prostate cancer (PCa) is the most common male malignancy in the western world and is a leading cause of cancer death [1]. For the last 75 years, advanced disease has been managed with androgen deprivation therapy (ADT), an approach that leads to disease regression, but rarely cure. Castration-resistant PCa generally results in death over 30–40 months depending on disease extent at diagnosis [2]. Multiple phase III clinical trials (ChemoHormonal Therapy Versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer [CHAARTED], A Randomized, Double-blind, Comparative Study of Abiraterone Acetate Plus Low-Dose Prednisone Plus Androgen Deprivation Therapy [ADT] Versus ADT Alone in Newly Diagnosed Subjects with High-Risk, Metastatic Hormone-naïve Prostate Cancer [mHNPc] [LATITUDE], Randomized Phase III Trial Comparing an Association of Hormonal Treatment and Docetaxel Versus the Hormonal Treatment Alone in Metastatic Prostate Cancers [GETUG-AFU 15], and Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy [STAMPEDE]) have recently demonstrated that in metastatic hormone-sensitive PCa, combining either docetaxel chemotherapy or abiraterone (an androgen signaling inhibitor) at the time of initiating ADT markedly improves overall survival in men [3–6]. These clinical observations suggest that ADT induces susceptibilities in PCa cells that make them amenable to synergistic treatment and improve cell killing. This underexplored area has the potential to have a major impact on PCa survival by improving the cancer response to ADT.

One phenotype that arises with extended replication or after cell stress is cellular senescence, in which cells are arrested yet viable [7]. Cellular senescence can also be induced after exposure to oxidative stress, DNA damaging agents, and ADT [8]. In breast cancer, another hormonally dependent cancer, senescence results after tamoxifen exposure in estrogen receptor positive cells [9]. Beneficial aspects of cellular senescence include a therapeutic arrest and immune stimulation. However, negative features include reentry of a subset of these persistent cells into the cell cycle and a secretory phenotype that may induce the growth of surrounding tumor cells. Thus, removal of these cells may lead to improved cancer outcomes.

The late Dr. Donald Coffey was always a strong supporter of “thinking outside the box”. His work led to remarkable advances in their own right, but placed in the context of his mentees he had an even greater impact of the field. Dr. Coffey often marveled at nature and would often state “You are going to be surprised at the simplicity and beauty of the real answer”. This review, dedicated to him, focuses on a surprising and major shift in our approach to advanced hormone sensitive PCa, that of the role of ADT and combined therapy. It is an area that requires significant research effort to understand and refine recent clinical discoveries. We will review cancer cell responses to ADT, seek to understand the clinical responses and mechanisms underlying these new combinatorial therapies with ADT, and discuss putative targeting of senescent cells after ADT as a therapeutic niche to improve outcomes.

2. Cancer cell responses to ADT

One of the best studied responses to ADT is apoptosis or programmed cell death. Kyprianou and colleague [10,11] performed some of the early work that androgen deprivation leads to apoptosis in pre-clinical PCa models. Androgen withdrawal in murine xenografts and human PCa tissues is associated with a decrease in the proliferative index, but surprisingly low levels of apoptosis encompassing only 2%–3% of cells [12–14]. Apoptosis typically occurs early after ADT within the first 72 h [15]. In addition to apoptosis, other cell death mechanisms induced with ADT include autophagy, necrosis, and necroptosis [16].

Autophagy is an evolutionarily conserved catabolic pathway that targets cellular organelles and cytoplasmic constituents to the lysosomes for degradation. Autophagy, although a cell death pathway, can also function as a survival mechanism exploited by cancer cells under various physiological stresses [17,18]. Androgen deprivation and tissue hypoxia, conditions which occur in prostate tissue after surgical or medical castration, lead to an increased adenosine monophosphate (AMP)-activated protein kinase (AMPK) activity in a threshold dependent manner in murine xenografts [19]. Increased AMPK activity is associated with greater cell survival and the induction of autophagy. *In vitro* in hormone-sensitive PCa cells, autophagy has been reported to occur after bicalutamide, but this occurs at much higher toxic doses than those seen with the induction of cellular senescence [20].

Some tumor cells become quiescent after ADT and have the potential to reactivate. PCa stem-like cells are typically androgen receptor (AR) negative and may represent part of the normally quiescent cancer stem cell population that emerges and expands after ADT [21]. These cells are rare, but express a specific surface antigen profile (CD44⁺/α₂β₁^{hi}/CD133⁺) when isolated from primary PCa tissues and show high levels of clonogenic ability [22].

3. Cellular senescence as a response to ADT in PCa

The phenotype of these residual cells after ADT is complex, but cellular senescence represents an intriguing response that has potential for therapeutic exploitation. Replicative senescence was first described as a phenotype in primary cells after extensive culture and replicative exhaustion *in vitro*, which was linked to telomere shortening [23]. More recently, DNA damage, increased oncogenic signaling, and oxidative stress have been found to result in induced or accelerated senescence [24]. The propensity of tumor cells to undergo senescence induction in response to multiple anti-cancer therapies has been demonstrated [25]. Notably, multiple solid tumor cell lines have shown dose-dependent cellular senescence induction in response to DNA damaging agents such as doxorubicin, diaziquone, cisplatin, and to a lesser degree ionizing radiation. Recently reported data suggest enzalutamide, an antiandrogen, potentiates DNA damage caused by radiation and significantly increases radiation-induced cellular senescence in LNCaP cells [26].

Senescent cells remain viable and metabolically active, but are permanently growth-arrested (Table 1) [24]. They are persistent, in contrast to cells undergoing programmed responses including apoptosis, autophagy, and/or mitotic catastrophe using conventional cytotoxic agents. Growth arrest is achieved and maintained in either G₁ or G₂/M phase, in part, by the increased expression of specific cyclin-dependent kinase inhibitors (CDKIs), including p16^{Ink4a}, p21^{Waf1/Cip1} and p27^{Kip1} [27]. Interestingly, transformed neoplastic cells that lack cellular senescence-associated tumor suppressor genes present in non-transformed cells (e.g., p53 and Rb) retain the capacity to become senescent with exposure to doxorubicin, docetaxel, and other chemotherapy agents [28,29].

Cultured *in vitro*, senescent cells develop a distinct and recognizable flattened and enlarged morphology with a prominent nucleus and increased cytoplasmic granularity. Most notably, these cells can be visualized using a staining technique based on senescence-associated β -galactosidase (SA- β -gal) activity [30]. This technique, which stains lysosomes in the perinuclear compartment blue at pH 6.0, is a widely accepted and utilized marker of cellular senescence, but has limitations *in vivo*, as it is not applicable to formalin-fixed archival tissues.

To resolve this issue, we validated one of the first antibodies to SA- β -gal protein (GLB1) *in vitro*, subsequently allowing the biology of cellular senescence in PCa *in vivo* to be interrogated [31]. In hormonally intact prostate tissues, quantitative imaging detects increased GLB1 expression in high-grade prostatic intraepithelial neoplasia (HGPIN) known to contain senescent cells compared to benign prostate tissues [31]. This work also demonstrated that in intermediate grade PCa increased GLB1 predicts prostate-specific antigen (PSA)-free survival. Furthermore, senescent cells are found less commonly in high grade (Gleason score 8–10) versus intermediate grade (Gleason score 6–7) cancers. These findings support a tumor suppression aspect of cellular senescence seen in skin and many other aging organs.

These studies in our laboratory, and subsequently others, have demonstrated that cellular senescence is induced in androgen sensitive cells after ADT [32,33]. Increased expression of the senescence-related proteins

GLB1, the CDKI p27^{Kip1}, and chromatin-regulating heterochromatin protein 1 γ (HP1 γ) are detected in 50%–80% of androgen sensitive LNCaP cells after being cultured in androgen-free media [8]. In mice bearing LuCaP xenograft tumors *in vivo*, surgical castration similarly increases senescent markers [8]. In another study, immunohistochemistry of human prostate tumors removed after ADT induced with goserelin acetate (Zoladex) showed a similar induction of GLB1, HP1 γ , and decreased Ki-67 [34]. ADT induces a cellular growth arrest consistent with cellular senescence, including hypophosphorylation of Rb, reduction of cyclin-dependent kinase activity, and a G₁/S block [27]. More recently, we have found that in patients undergoing ADT prior to prostate removal for cancer, elevated cellular GLB1 levels measured by VECTRA automated immunoquantitation are noted in as many as 50% of cancer cells [35]. GLB1 levels increase in tumors with longer ADT duration.

Despite permanent growth arrest, senescent cells are metabolically active and have increased energy demand, protein turnover, and glycolysis [36]. The increased energy demand is marked by activation of AMPK, a cellular energy sensor [37]. Increased glucose transport and amplified glycolytic enzyme and pyruvate kinase expression promote the utilization of glucose through non-oxidative glycolysis, similar to the Warburg phenomenon described in tumor cells [36]. Additionally, higher amino acid transport and increased protein synthesis generate cytokines that characterize the proinflammatory secretory-associated senescent phenotype (SASP). These metabolic changes present an opportunity for improved therapeutic approaches when combined with ADT.

Interestingly, AR signaling may play an important role in the cellular senescence response. Blocking the AR, or paradoxically applying supraphysiologic levels of androgens, such as those used in bipolar androgen therapy, may induce cellular senescence [38]. This may represent part of the biphasic growth response seen in AR expressing PCa cells, and induction of senescence in this context may function through the Src-P13K-Akt signaling pathway. Pharmacologic induction of senescence has also been demonstrated in murine xenografts and human PCa tissues using the naturally occurring AR antagonist atraric acid [39].

Table 1 Summary of clinical trial results using ADT in combination with other therapies for the treatment of metastatic hormone sensitive PCa.

Clinical trial	ADT+	Number of patients	Follow-up (month)	HR 95% CI	Reference
GETUG-AFU 15 ^a	Docetaxel	385	50	1.01 (0.75–1.36)	Gravis et al., 2013 [40]
CHAARTED ^b	Docetaxel	790	28.9	0.61 (0.47–0.80)	Sweeney et al., 2015 [3]
LATITUDE ^c	Abiraterone	1199	30.4	0.62 (0.51–0.76)	Fizazi et al., 2017 [4]
STAMPEDE-Doc ^d	Docetaxel	2962	43	0.78 (0.66–0.93)	James et al., 2016 [5]
STAMPEDE-Abi ^e	Abiraterone	1917	40	0.63 (0.52–0.76)	James et al., 2017 [6]

PCa, prostate cancer; HR, hazard ratio; CI, confidential interval.

^a Randomized Phase III Trial Comparing an Association of Hormonal Treatment and Docetaxel Versus the Hormonal Treatment Alone in Metastatic Prostate Cancers (GETUG-AFU 15).

^b ChemoHormonal Therapy Versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer (CHAARTED).

^c A Randomized, Double-blind, Comparative Study of Abiraterone Acetate Plus Low-Dose Prednisone Plus Androgen Deprivation Therapy (ADT) Versus ADT Alone in Newly Diagnosed Subjects with High-Risk, Metastatic Hormone-naïve Prostate Cancer (mHNPc) (LATITUDE).

^d Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy (STAMPEDE) with Docetaxel (Doc).

^e Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy (STAMPEDE) with Abi (Abiraterone).

4. Clinical trials suggest unique PCa susceptibilities after ADT

The application of ADT has always been approached as the sole systemic therapy for metastatic hormone sensitive PCa, traditionally using agents that target the substrate (testosterone) or block the AR. In a major paradigm shift, several recent trials using either docetaxel or androgen signaling inhibitors in combination with ADT for metastatic hormone sensitive PCa have demonstrated remarkable improvements in survival. This suggests ADT induces susceptibilities in cancer cells that may be exploited. The phase III CHAARTED trial (ChemoHormonal Therapy Versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer), demonstrated upfront chemotherapy with concurrent docetaxel and ADT improves survival by 13.6 months versus ADT alone in hormone naïve patients (Table 1) [3]. The docetaxel group also experienced a significant delay in biochemical, symptomatic, or radiographic progression compared to controls (20.1 vs. 11.7 months; hazard ratio [HR] = 0.61; 95% confidence interval [CI] = 0.51 to 0.72; $p < 0.001$). These results were confirmed in the STAMPEDE (Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy) [5] and GETUG-AFU 15 trials (Randomized Phase III Trial Comparing an Association of Hormonal Treatment and Docetaxel Versus the Hormonal Treatment Alone in Metastatic Prostate Cancers) [40]. Docetaxel functions to inhibit the transport of the AR, potentiating ADT action [41]. It is also a microtubule inhibitor having direct toxicity to cells. We anticipate this combination results in fewer persistent senescent cells than ADT alone post-treatment, but further mechanistic study is required regarding the synergistic activity of this combination.

The phase III LATITUDE trial (A Randomized, Double-blind, Comparative Study of Abiraterone Acetate Plus Low-Dose Prednisone Plus Androgen Deprivation Therapy [ADT] Versus ADT Alone in Newly Diagnosed Subjects with High-Risk, Metastatic Hormone-naïve Prostate Cancer [mHNPC]) demonstrated upfront therapy with concurrent abiraterone and luteinizing hormone-releasing hormone agonists improves overall survival versus ADT alone (median not reached versus 34.7 months, HR = 0.62; 95% CI: 0.51–0.76; $p < 0.001$) in hormone naïve patients [4]. The abiraterone group also experienced a significant delay in radiographic progression-free survival compared to control (33.0 months vs. 14.8 months; HR = 0.47; 95% CI: 0.39–0.55; $p < 0.001$). The abiraterone and ADT arm of the STAMPEDE trial also showed improved outcomes [6]. Abiraterone irreversibly inhibits the Cytochrome P450 17 α -hydroxylase/17,20-lyase (CYP17) enzyme expressed in adrenal tissues, reducing androgen synthesis from all sources [42]. Although the combination of CYP17 inhibition and ADT demonstrates more effective androgen depletion than either agent alone [43], resistance to abiraterone develops, likely through increase CYP17 expression [44], enzymatic alterations [45], and/or gain of function 3 β -hydroxysteroid dehydrogenase type 1 mutation [46]. These clinical observations suggest that the initiation of ADT induces susceptibilities in PCa cells that make them amenable to synergistic treatment and improved cell killing. Further

study is required to delineate the synergistic activity of abiraterone and ADT, along with clearly defining the mechanisms of resistance in castration-resistant PCa.

5. Exploiting cellular senescence as synthetic therapy

Cellular senescence may have detrimental features. Oncogene-induced cellular senescence imposes selective pressures that promote the outgrowth of senescence-resistant aggressive tumor cell subpopulations [47]. Although cellular senescence is cytostatic and offers a potential survival advantage in some models, long-term exposure of surrounding cells to the SASP, and resulting proinflammatory cytokines and growth factors, may have deleterious effects on surrounding cells [7]. In a recent *Nature* paper, it was reported that lymphoma cells released from chemotherapy-induced senescence results in a population of cells exhibiting a stem cell phenotype that exhibits highly aggressive growth potential upon escape from cell-cycle blockade [48]. This population is enriched in relapsing hematologic tumors. It has been proposed that ADT-induced cellular senescence might play a role in the chemoresistance that arises with intermittent ADT [32].

The cellular senescence phenotype presents unique opportunities that have the potential to be exploited for therapeutic cure (Fig. 1) [49]. Dörr and colleagues [36] have induced senescence in lymphoma cells and used compounds that target the inhibition of glucose transporters, glycolytic enzymes, and adenosine triphosphate (ATP) depletion to generate synthetic lethality in cancer cells (Table 2). These combinations lead to improved survival and elimination of cancer cells through caspase-12- and caspase-3-mediated endoplasmic-reticulum-related apoptosis [36]. Additionally, higher amino acid transport and increased protein synthesis generate cytokines that characterize the proinflammatory SASP. These findings highlight the hypercatabolic nature of senescent cells after induction with ADT and other agents that is therapeutically exploitable by synthetic lethal metabolic targeting. These approaches have been largely unexplored to date in PCa, but as outlined below, interesting supportive data exist.

The therapeutic induction of cellular senescence from chemotherapeutics, such as cyclophosphamide, results in augmented protein translation with the resultant accumulation of misfolded proteins, which activates the conserved proteotoxic stress response (PSR) [36]. The substantial proteotoxic stress in cellular senescence evokes energy-consuming countermeasures for cell survival. The PSR is orchestrated by the transcription factor heat shock factor 1 (Hsf1), which regulates the gene expression of several families of molecular chaperones (heat shock protein (Hsp) 27, Hsp70, Hsp90 and others) that attenuate proteotoxic stress and promote cell survival by preventing protein aggregation, refolding non-native proteins, or delivering them to the proteasome [54]. Anabolic tumor cells are more dependent on the Hsf1-mediated PSR than normal cells for cell survival [55]. The PSR may represent an additional novel targetable vulnerability of senescent transformed cells that can be exploited therapeutically, a hypothesis that has yet to be tested.




Figure 1 Synthetic lethal targeting of ADT induced cellular senescence for improved prostate cancer cell killing. ADT, androgen deprivation therapy.

Table 2 Presence of the cellular senescence phenotype offers therapeutic opportunities.	
Characteristic	Synthetic lethal targeting of senescent cells
Cell morphology	
SA-β-gal positive	
Enlarged cell with prominent nuclei and cytoplasmic granularity	
Metabolic alterations	
Hypermetabolic	Glucose transport inhibitors phloretin, cytochalasin B, 2-deoxy-D-glucose [36]
Glycolysis	AMPK inhibitor compound C [36]/AMPK activator and mTOR inhibitor metformin [50,51]
High protein turnover	Lysosomal V-ATPase inhibitors bafilomycin A1 and concanamycin A [36]
Secretory phenotype	
Pro-inflammatory cytokines	Tumor microenvironment cancer-based immunotherapy [52]
Growth factors	Growth factor and growth factor receptor inhibitors (e.g. VEGF, IGF-1 inhibitors) [53]

AMPK, adenosine monophosphate-activated protein kinase; ATPase, adenosine triphosphatase; IGF-1, insulin-like growth factor 1; mTOR, mammalian target of rapamycin; SA-β-gal, senescence-associated β-galactosidase; VEGF, vascular endothelial growth factor.

Metformin is an oral anti-diabetic agent in the biguanide class that has generated interest as a cancer therapy. Metformin is an intriguing drug that may not only enhance chemotherapy response for established tumors, but also demonstrates anticancer activity as a single agent [56]. Metformin inhibits the mammalian target of rapamycin (mTOR), a central regulator of cell growth and survival, in part by activating AMPK [51]. Additionally, metformin inhibits Hsf1 by activating AMPK, which in turn phosphorylates Hsf1 on Ser 121 to inhibit its activity [57]. This disables the protective effects of Hsf1 against proteotoxic stress. Synergistic activity after neoadjuvant chemotherapy was suggested in an analysis showing improved clinical response in diabetic patients with breast cancer receiving metformin and neoadjuvant chemotherapy [58]. These patients received doxorubicin, which is well known for its senescence-inducing properties in prostate and breast cancer [28,29].

Given the metabolic susceptibilities that ADT induces, metformin leads to increased cell kill when combined with ADT. Metformin has been combined with the antiandrogen bicalutamide *in vitro* and in animal models [59]. This combination significantly reduced clonogenicity ($p < 0.005$) and tumor growth with greater effects in AR-positive cells. In unpublished data, we have found LNCaP, LaPC4, and CWR22 PCa lines all demonstrate synthetic lethal responses as calculated by CalcuSyn (Biosoft, Cambridge, UK) to low dose bicalutamide (1–5 $\mu\text{mol/L}$) to induce senescence initially, followed by metformin at low dose (0.1–1 mmol/L) [60]. Increased apoptosis peaks 2 days after metformin application.

In a large observational study of 87 344 veterans, we demonstrated that metformin combined with ADT improved overall survival (HR = 0.82, 95% CI: 0.78–0.86), reduced the risk of skeletal related events (HR = 0.84, 95% CI: 0.74–0.96), and improved cancer specific survival

(HR = 0.70, 95% CI: 0.64–0.77) [61]. Skeletal related events, defined as pathologic fracture, spinal cord compression, and/or necessity for bone radiation or surgery due to pain or impending fracture, was used as one surrogate for disease progression. Although retrospective observational studies do not prove causality, these data merit further clinical trial investigation. Notably, metformin is an inexpensive, widely used drug associated with minimal side effects even when used in the non-diabetic population. Therefore, few barriers exist to prevent its clinical translation in PCa.

Statins are another widely used and inexpensive drug that may provide synergistic lethality with ADT given the metabolic profile of senescent cells. Statins are a class of oral anti-hypercholesterolemia agents that inhibit 3-hydroxy-3-methylglutaryl-CoA (HMG CoA) reductase. Although they are associated with myopathy and elevated transaminases, the side effect profile is favorable [62]. The use of statins reduces PSA *in vivo* [63], through downregulation of AR expression and activity due to enhanced proteolysis of the AR protein [64]. In LNCaP cells, the combination of simvastatin and ADT demonstrates increased growth inhibition in AR positive lines [65]. Statins also compete with the dihydrotestosterone precursor for Solute Carrier Organic Anion Transporter Family Member 2B1 mediated transport, thus decreasing androgen availability in PCa cells [66]. Men with hormone sensitive PCa undergoing ADT and taking statin drugs experienced a delayed time to disease progression compared to men not taking statins (median 27.5 months; 95% CI: 21.1–37.7 vs. 17.4 months; 95% CI: 14.9–21.1). Simvastatin has also been shown to decrease the secretory phenotype of senescent human fibroblasts, including interleukin-6, by inhibiting protein prenylation [67]. Whether statins directly induce cell death in senescent cells is unknown as of yet.

Other characteristics of senescent cells include a hypermetabolic phenotype comprised of enhanced glycolysis and protein turnover, providing a critical cellular senescence associated metabolic liability that may be targeted therapeutically. Senescent cells are selectively susceptible to inhibition of glucose transporters (e.g. cytochalasin B), or to the pharmacological competitor (2-deoxy-D-glucose) [36]. Senescent cells also rely on an intact lysosomal protein degradation machinery to buffer proteotoxic stress. Exposure of senescent cells to bafilomycin A1 or concanamycin A, specific inhibitors of lysosomal V-ATPases, or to a cocktail of lysosomal protease inhibitors all generate increased death in therapy induced senescent cells [36]. With use of these inhibitor in cell culture, senescent cell death occurred at significantly higher rates. They provide an avenue for therapeutic exploitation that should be examined further.

6. Conclusion and future directions

ADT has been used as the primary approach to advanced hormone sensitive PCa for over 75 years. Recent clinical trials have demonstrated combining ADT with other agents improves survival. ADT and docetaxel (or abiraterone) should be considered standard of care for patients with

metastatic hormone sensitive PCa given recent trial results. Research is needed to define and understand these observations to further improve our progress in this area. Cellular senescence is a distinctive phenotype characterized by metabolic alterations and growth arrest. ADT induced cellular senescence in PCa may be deleterious; however, it may offer a unique opportunity for synthetic lethal targeting of residual PCa cells. Several recent retrospective hypothesis-generating studies suggest combining ADT with agents that target metabolism, including metformin, may improve patient survival. Future work should focus on further delineating the response of tumors to ADT and cellular senescence biology in this context.

Conflicts of interest

The study was supported by DOD Prostate Cancer Research Program PC150221, R. Stephenson Family Fund.

Author contributions

Study concept and design: Kyle A. Richards, Bing Yang, Vincent Cryns, David F. Jarrard.

Drafting of manuscript: Tyler Etheridge, Shivashankar Damodaran, Adam Schultz, Kyle A. Richards, Joseph Gawdzik, Bing Yang, Vincent Cryns, David F. Jarrard.

References

- [1] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2017. *CA Cancer J Clin* 2017;67:7–30.
- [2] Moul JW. Hormone naïve prostate cancer: predicting and maximizing response intervals. *Asian J Androl* 2015;17: 929–35.
- [3] Sweeney CJ, Chen YH, Carducci M, Liu G, Jarrard DF, Eisenberger M, et al. Chemohormonal therapy in metastatic hormone-sensitive prostate cancer. *N Engl J Med* 2015;373: 737–46.
- [4] Fizazi K, Tran N, Fein L, Matsubara N, Rodriguez-Antolin A, Alekseev BY, et al. Abiraterone plus prednisone in metastatic, castration-sensitive prostate cancer. *N Engl J Med* 2017;377: 352–60.
- [5] James ND, Sydes MR, Clarke NW, Mason MD, Dearnaley DP, Spears MR, et al. Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multi-arm, multistage, platform randomised controlled trial. *Lancet* 2016;387:1163–77.
- [6] James ND, de Bono JS, Spears MR, Clarke NW, Mason MD, Dearnaley DP, et al. Abiraterone for prostate cancer not previously treated with hormone therapy. *N Engl J Med* 2017; 377:338–51.
- [7] Ewald JA, Desotelle JA, Wilding G, Jarrard DF. Therapy-induced senescence in cancer. *J Natl Cancer Inst* 2010;102: 1536–46.
- [8] Ewald JA, Desotelle JA, Church DR, Yang B, Huang W, Laurila TA, et al. Androgen deprivation induces senescence characteristics in prostate cancer cells *in vitro* and *in vivo*. *Prostate* 2013;73:337–45.
- [9] Tuttle R, Miller KR, Maiorano JN, Termuhlen PM, Gao Y, Berberich SJ. Novel senescence associated gene, YPEL3, is repressed by estrogen in ER+ mammary tumor cells and

- required for tamoxifen-induced cellular senescence. *Int J Cancer* 2012;130:2291–9.
- [10] Kyprianou N, Isaacs JT. Activation of programmed cell death in the rat ventral prostate after castration. *Endocrinology* 1988;122:552–62.
 - [11] Kyprianou N, English HF, Isaacs JT. Programmed cell death during regression of PC-82 human prostate cancer following androgen ablation. *Cancer Res* 1990;50:3748–53.
 - [12] Agus DB, Cordon-Cardo C, Fox W, Drobnjak M, Koff A, Golde DW, et al. Prostate cancer cell cycle regulators: response to androgen withdrawal and development of androgen independence. *J Natl Cancer Inst* 1999;91:1869–76.
 - [13] Ohlson N, Wikström P, Stattin P, Bergh A. Cell proliferation and apoptosis in prostate tumors and adjacent non-malignant prostate tissue in patients at different time-points after castration treatment. *Prostate* 2005;62:307–15.
 - [14] Westin P, Stattin P, Damber JE, Bergh A. Castration therapy rapidly induces apoptosis in a minority and decreases cell proliferation in a majority of human prostatic tumors. *Am J Pathol* 1995;146:1368–75.
 - [15] Smitherman AB, Gregory CW, Mohler JL. Apoptosis levels increase after castration in the CWR22 human prostate cancer xenograft. *Prostate* 2003;57:24–31.
 - [16] Zhang KX, Firus J, Prieur B, Jia W, Rennie PS. To die or to survive, a fatal question for the destiny of prostate cancer cells after androgen deprivation therapy. *Cancers (Basel)* 2011;3:1498–512.
 - [17] Mathew R, Karantza-Wadsworth V, White E. Role of autophagy in cancer. *Nat Rev Cancer* 2007;7:961–7.
 - [18] Gerland LM, Peyrol S, Lallemand C, Branche R, Magaud JP, Ffrench M. Association of increased autophagic inclusions labeled for beta-galactosidase with fibroblastic aging. *Exp Gerontol* 2003;38:887–95.
 - [19] Chhipa RR, Wu Y, Ip C. AMPK-mediated autophagy is a survival mechanism in androgen-dependent prostate cancer cells subjected to androgen deprivation and hypoxia. *Cell Signal* 2011;23:1466–72.
 - [20] Boutin B, Tajeddine N, Vandersmissen P, Zanou N, Van Schoor M, Mondin L, et al. Androgen deprivation and androgen receptor competition by bicalutamide induce autophagy of hormone-resistant prostate cancer cells and confer resistance to apoptosis. *Prostate* 2013;73:1090–102.
 - [21] Ojo D, Lin X, Wong N, Gu Y, Tang D. Prostate cancer stem-like cells contribute to the development of castration-resistant prostate cancer. *Cancers (Basel)* 2015;7:2290–308.
 - [22] Collins AT, Berry PA, Hyde C, Stower MJ, Maitland NJ. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res* 2005;65:10946–51.
 - [23] Hayflick L. The limited in vitro lifespan of human diploid cell strains. *Exp Cell Res* 1965;25:585–621.
 - [24] Campisi J. Cancer, aging and cellular senescence. *In Vivo* 2000;14:183–8.
 - [25] Gewirtz DA, Holt SE, Elmore LW. Accelerated senescence: an emerging role in tumor cell response to chemotherapy and radiation. *Biochem Pharmacol* 2008;76:947–57.
 - [26] Ghashghaei M, Paliouras M, Heravi M, Bekerat H, Trifiro M, Niazi TM, et al. Enhanced radiosensitization of enzalutamide via schedule dependent administration to androgen-sensitive prostate cancer cells. *Prostate* 2018;78:64–75.
 - [27] Bringold F, Serrano M. Tumor suppressors and oncogenes in cellular senescence. *Exp Gerontol* 2000;35:317–29.
 - [28] Chang BD, Broude EV, Dokmanovic M, Zhu H, Ruth A, Xuan Y, et al. A senescence-like phenotype distinguishes tumor cells that undergo terminal proliferation arrest after exposure to anticancer agents. *Cancer Res* 1999;59:3761–7.
 - [29] Schwarze SR, Fu VX, Desotelle JA, Kenowski ML, Jarrard DF. The identification of senescence-specific genes during the induction of senescence in prostate cancer cells. *Neoplasia* 2005;7:816–23.
 - [30] Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that identifies senescent human cells in culture and in aging skin *in vivo*. *Proc Natl Acad Sci USA* 1995;92:9363–7.
 - [31] Wagner J, Damaschke N, Yang B, Truong M, Guenther C, McCormick J, et al. Overexpression of the novel senescence marker β -galactosidase (GLB1) in prostate cancer predicts reduced PSA recurrence. *PLoS One* 2015;10:e0124366.
 - [32] Burton DG, Giribaldi MG, Munoz A, Halvorsen K, Patel A, Jorda M, et al. Androgen deprivation-induced senescence promotes outgrowth of androgen-refractory prostate cancer cells. *PLoS One* 2013;8:e68003.
 - [33] Pernicová Z, Slabáková E, Kharaishvili G, Bouchal J, Král M, Kunická Z, et al. Androgen depletion induces senescence in prostate cancer cells through down-regulation of Skp2. *Neoplasia* 2011;13:526–36.
 - [34] Bartkova J, Rezaei N, Liontos M, Karakaidos P, Kletsas D, Issaeva N, et al. Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature* 2006;444:633–7.
 - [35] Blute ML, Damaschke N, Wagner J, Yang B, Gleave M, Fazli L, et al. Persistence of senescent prostate cancer cells following prolonged neoadjuvant androgen deprivation therapy. *PLoS One* 2017;12:e0172048.
 - [36] Dörr JR, Yu Y, Milanovic M, Beuster G, Zasada C, Däbritz JH, et al. Synthetic lethal metabolic targeting of cellular senescence in cancer therapy. *Nature* 2013;501:421–5.
 - [37] Jones RG, Plas DR, Kubek S, Buzzai M, Mu J, Xu Y, et al. AMP-activated protein kinase induces a p53-dependent metabolic checkpoint. *Mol Cell* 2005;18:283–93.
 - [38] Roediger J, Hessenkemper W, Bartsch S, Manvelyan M, Huettner SS, Liehr T, et al. Supraphysiological androgen levels induce cellular senescence in human prostate cancer cells through the Src-Akt pathway. *Mol Cancer* 2014;13:214.
 - [39] Hessenkemper W, Roediger J, Bartsch S, Houtsmuller AB, van Royen ME, Petersen I, et al. A natural androgen receptor antagonist induces cellular senescence in prostate cancer cells. *Mol Endocrinol* 2014;28:1831–40.
 - [40] Gravis G, Fizazi K, Joly F, Oudard S, Priou F, Esterni B, et al. Androgen-deprivation therapy alone or with docetaxel in non-castrate metastatic prostate cancer (GETUG-AFU 15): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2013;14:149–58.
 - [41] Zhu ML, Horbinski CM, Garzotto M, Qian DZ, Beer TM, Kyprianou N. Tubulin-targeting chemotherapy impairs androgen receptor activity in prostate cancer. *Cancer Res* 2010;70:7992–8002.
 - [42] Thakur A, Roy A, Ghosh A, Chhabra M, Banerjee S. Abiraterone acetate in the treatment of prostate cancer. *Biomed Pharmacother* 2018;101:211–8.
 - [43] Attard G, Reid AH, Yap TA, Raynaud F, Dowsett M, Settatree S, et al. Phase I clinical trial of a selective inhibitor of CYP17, abiraterone acetate, confirms that castration-resistant prostate cancer commonly remains hormone driven. *J Clin Oncol* 2008;26:4563–71.
 - [44] Mostaghel EA, Marck BT, Plymate SR, Vessella RL, Balk S, Matsumoto AM, et al. Resistance to CYP17A1 inhibition with abiraterone in castration-resistant prostate cancer: induction of steroidogenesis and androgen receptor splice variants. *Clin Cancer Res* 2011;17:5913–25.
 - [45] Cai C, Chen S, Ng P, Bubley GJ, Nelson PS, Mostaghel EA, et al. Intratumoral *de novo* steroid synthesis activates androgen receptor in castration-resistant prostate cancer and is upregulated by treatment with CYP17A1 inhibitors. *Cancer Res* 2011;71:6503–13.

- [46] Chang KH, Li R, Kuri B, Lotan Y, Roehrborn CG, Liu J, et al. A gain-of-function mutation in DHT synthesis in castration-resistant prostate cancer. *Cell* 2013;154:1074–84.
- [47] Halazonetis TD, Gorgoulis VG, Bartek J. An oncogene-induced DNA damage model for cancer development. *Science* 2008;319:1352–5.
- [48] Milanovic M, Fan DNY, Belenki D, Däbritz JHM, Zhao Z, Yu Y, et al. Senescence-associated reprogramming promotes cancer stemness. *Nature* 2018;553:96–100.
- [49] Cairney CJ, Bilsland AE, Evans TR, Roffey J, Bennett DC, Narita M, et al. Cancer cell senescence: a new frontier in drug development. *Drug Discov Today* 2012;17:269–76.
- [50] Rocha GZ, Dias MM, Ropelle ER, Osório-Costa F, Rossato FA, Vercesi AE, et al. Metformin amplifies chemotherapy-induced AMPK activation and antitumoral growth. *Clin Cancer Res* 2011;17:3993–4005.
- [51] Dowling RJ, Zakikhani M, Fantus IG, Pollak M, Sonenberg N. Metformin inhibits mammalian target of rapamycin-dependent translation initiation in breast cancer cells. *Cancer Res* 2007;67:10804–12.
- [52] Nakamura K, Smyth MJ. Targeting cancer-related inflammation in the era of immunotherapy. *Immunol Cell Biol* 2017;95:325–32.
- [53] Macfarlane RJ, Chi KN. Novel targeted therapies for prostate cancer. *Urol Clin North Am* 2010;37:105–19, Table of Contents.
- [54] Westerheide SD, Morimoto RI. Heat shock response modulators as therapeutic tools for diseases of protein conformation. *J Biol Chem* 2005;280:33097–100.
- [55] Dai C, Whitesell L, Rogers AB, Lindquist S. Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. *Cell* 2007;130:1005–18.
- [56] Foretz M, Guigas B, Bertrand L, Pollak M, Viollet B. Metformin: from mechanisms of action to therapies. *Cell Metab* 2014;20:953–66.
- [57] Dai S, Tang Z, Cao J, Zhou W, Li H, Sampson S, et al. Suppression of the HSF1-mediated proteotoxic stress response by the metabolic stress sensor AMPK. *EMBO J* 2015;34:275–93.
- [58] Jiralerspong S, Palla SL, Giordano SH, Meric-Bernstam F, Liedtke C, Barnett CM, et al. Metformin and pathologic complete responses to neoadjuvant chemotherapy in diabetic patients with breast cancer. *J Clin Oncol* 2009;27:3297–302.
- [59] Colquhoun AJ, Venier NA, Vandersluis AD, Besla R, Sugar LM, Kiss A, et al. Metformin enhances the antiproliferative and apoptotic effect of bicalutamide in prostate cancer. *Prostate Cancer Prostatic Dis* 2012;15:346–52.
- [60] McCormick JR, Blute ML, Yang B, Damaschke N, Jarrard DF. Synthetic lethal metabolic targeting of cellular senescence in prostate cancer with the repurposed drug metformin. *J Urol* 2016;195:e673–4.
- [61] Richards KA, Liou JI, Cryns VL, Downs TM, Abel EJ, Jarrard DF. Metformin use is associated with improved survival in patients with advanced prostate cancer on androgen deprivation therapy. *J Urol* 2018;200:1256–63.
- [62] Armitage J. The safety of statins in clinical practice. *Lancet* 2007;370:1781–90.
- [63] Hamilton RJ, Goldberg KC, Platz EA, Freedland SJ. The influence of statin medications on prostate-specific antigen levels. *J Natl Cancer Inst* 2008;100:1511–8.
- [64] Yang L, Egger M, Plattner R, Klocker H, Eder IE. Lovastatin causes diminished PSA secretion by inhibiting AR expression and function in LNCaP prostate cancer cells. *Urology* 2011;77:1508. e1501–7.
- [65] Syväälä H, Pennanen P, Bläuer M, Tammela TL, Murtola TJ. Additive inhibitory effects of simvastatin and enzalutamide on androgen-sensitive LNCaP and VCaP prostate cancer cells. *Biochem Biophys Res Commun* 2016;481:46–50.
- [66] Harshman LC, Wang X, Nakabayashi M, Xie W, Valenca L, Werner L, et al. Statin use at the time of initiation of androgen deprivation therapy and time to progression in patients with hormone-sensitive prostate cancer. *JAMA Oncol* 2015;1:495–504.
- [67] Liu S, Uppal H, Demaria M, Desprez PY, Campisi J, Kapahi P. Simvastatin suppresses breast cancer cell proliferation induced by senescent cells. *Sci Rep* 2015;5:17895.

Targeting Metastatic Hormone Sensitive Prostate Cancer: Chemohormonal Therapy and New Combinatorial Approaches



Shivashankar Damodaran, Joshua M. Lang and David F. Jarrard*

From the Department of Urology, School of Medicine and Public Health (SD, DFJ) and Carbone Cancer Center (JML, DFJ), University of Wisconsin-Madison, Madison, Wisconsin

Abbreviations and Acronyms

ADT = androgen deprivation therapy
AR = androgen receptor
CHAARTED = Chemohormonal Therapy versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer
CRPC = castration resistant PC
GETUG-AFU = Groupe d'Étude des Tumeurs Uro-Genital and Association Française d'Urologie trial
HSPC = hormone sensitive PC
mHSPC = metastatic HSPC
PC = prostate cancer
PSA = prostate specific antigen
QOL = quality of life
STAMPEDE = Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy

Purpose: Androgen deprivation therapy alone has been the standard of care for metastatic hormone sensitive prostate cancer for the last 75 years. This review focuses on recent trials and mechanisms which highlight the new paradigm of combining androgen deprivation therapy with other agents, changing the treatment of patients with prostate cancer who have advanced disease.

Materials and Methods: We searched the peer reviewed literature on the PubMed® and Web of Science® databases through January 2018 using the key words, "metastatic hormone sensitive prostate cancer," "metastatic castration sensitive prostate cancer," "docetaxel," "abiraterone" and "senescence in cancer." ClinicalTrials.gov was queried for ongoing studies. Relevant data recently presented at major urology and medical oncology meetings were also evaluated.

Results: Recently published, phase III trials using androgen deprivation therapy combinations for metastatic hormone sensitive prostate cancer can be broadly grouped into chemohormonal studies (docetaxel) or trials of androgen signaling inhibitors. The CHAARTED (Chemohormonal Therapy versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer) and STAMPEDE (Systemic Therapy in Advancing or Metastatic Prostate Cancer: Evaluation of Drug Efficacy) studies showed a survival advantage when combining androgen deprivation therapy with chemotherapy, as well as increased time to progression to castration resistant status. The abiraterone arm of the STAMPEDE and LATITUDE trials, which analyzed combining androgen deprivation therapy with abiraterone, revealed improved overall and progression-free survival. Androgen deprivation therapy generates a number of phenotypes in resistant cancer cells, including quiescence, autophagy and cellular senescence. Senescent cells represent a metabolic target for synergistic lethality with drugs such as metformin. Ongoing trials are under way to examine the effect of combining newer antiandrogens and novel drugs with androgen deprivation therapy in patients with metastatic hormone sensitive prostate cancer.

Accepted for publication August 13, 2018.

The corresponding author certifies that, when applicable, a statement(s) has been included in the manuscript documenting institutional review board, ethics committee or ethical review board study approval; principles of Helsinki Declaration were followed in lieu of formal ethics committee approval; institutional animal care and use committee approval; all human subjects provided written informed consent with guarantees of confidentiality; IRB approved protocol number; animal approved project number.

Supported in part by Department of Defense Prostate Cancer Research Program funding (PC150221).

No direct or indirect commercial, personal, academic, political, religious or ethical incentive is associated with publishing this article.

*Correspondence: Department of Urology, Translational Research, Carbone Cancer Center, 7037, Wisconsin Institutes of Medical Research, 1111 Highland Ave., Madison, Wisconsin 53792 (telephone: 608-263-9534; e-mail: jarrard@urology.wisc.edu).

Conclusions: Combination therapy has evolved as the standard of care for metastatic hormone sensitive prostate cancer. The ideal combination is tailored to patients after individualized counseling taking into account general health and comorbid illness status.

Key Words: prostatic neoplasms, castration-resistant; neoplasm metastasis; drug therapy, combination; antineoplastic agents; cellular senescence

PROSTATE cancer is the most common cancer in males with an estimated new case incidence of 164,690 and an estimated mortality of 29,430 expected in 2018. Despite an overall 5-year survival rate of 98.2%, mHSPC has a dismal 30% 5-year survival rate.¹ Conventional treatment of mHSPC has been ADT since the landmark discovery by Huggins and Hodges in 1941 demonstrating the hormonal sensitivity of PC.² Metastatic HSPC treated with ADT transitions to the CRPC stage with a median survival of approximately 3 years.² Progression to CRPC is associated with a deterioration in QOL. In the recently concluded STAMPEDE trial patients were found to spend three-quarters of overall survival after the mHSPC diagnosis in the CRPC state, highlighting the significant alteration in the natural history of the disease with newer treatments.³ Treatment continues to evolve with a focus on disease management at earlier time points.

Approaches to improve the response rate to ADT or decrease side effects have included intermittent hormone therapy, use of an antiandrogen with medical or surgical castration, or antiandrogens alone.⁴ An analysis of these trials detailed the minimal benefits of these approaches, including only 2% to 3% improvement in 5-year survival with a wide range of uncertainty.⁴ For a number of years researchers had considered earlier application of cytotoxic therapy in an effort to delay progression to castrate disease. The addition of chemotherapy to ADT was tested in the last 30 years and this approach was used in a number of published randomized trials, as summarized by Millikan et al.⁵ However, the lack of cytotoxic therapy, which improved survival in CRPC cases, led to minimal advances. In CRPC more recently docetaxel chemotherapy resulted in an incremental 2.5-month improvement in median survival, leading to its approval in 2004.⁶ These findings lead to the initiation of trials earlier in the disease in patients with a larger disease burden at ADT initiation.

ADT induces a number of unique responses in prostate cancer cells, of which some lead to cellular persistence and the development of castration resistance. Recent phase III trials have demonstrated striking improvements in patient survival with combined ADT and docetaxel as well as ADT and androgen synthesis inhibitors. The synergistic targeting of hormonally sensitive prostate cancer with ADT combined with other novel agents is a

new chapter in the evolution of prostate cancer treatment. These trials and new approaches are reviewed in this study, emphasizing that combination therapy is now the standard of care in patients with mHSPC.

METHODS

We performed PubMed and Web of Science database searches of the peer reviewed mHSPC literature on the mechanisms of cellular persistence after ADT as well as combination therapies that use ADT with another therapeutic agent (fig. 1). Original studies of this subject as well as a small number of reviews were analyzed for strengths and weaknesses. We provide a comprehensive review of prospective, phase III trials of combination therapy with ADT in the setting of mHSPC, the synergy mechanism, side effects and QOL. The mechanisms of cellular persistence after ADT are also discussed with special emphasis on senescence. Combination therapies ongoing and to be considered in the near future are examined.

RESULTS

Metastatic Hormone Sensitive Prostate Cancer

Chemohormonal Therapy Trials. Early studies of chemotherapy combined with ADT were less optimistic, in part due to toxicity and the lack of active agents. Millikan et al performed a phase III trial in 286 patients with mHSPC who received 3, 8-week cycles of ketoconazole and doxorubicin alternating with vinblastine and estramustine, given in addition to standard ADT vs ADT alone.⁵ No difference in time to progression or overall survival was noted. Furthermore, 51% of patients experienced grade 3 or worse adverse events, including thromboembolic events. Another trial featured mitomycin, cyclophosphamide, epirubicin and fluorouracil with no improvement in survival.⁷ In 2004 improved survival in CRPC with docetaxel chemotherapy subsequently led to consideration of its use at the time of ADT initiation in patients with a higher tumor burden.

The CHARTED study was the first trial to show that adding 6 cycles of docetaxel 75 mg/m² every 3 weeks to standard ADT significantly improved outcomes in men with mHSPC (table 1).⁸ The 13.6-month improved overall survival in the chemohormonal group compared to the ADT group (median overall survival 57.6 vs 44 months, HR 0.61, 95% CI 0.47–0.80, *p* <0.001) represented one of the




Figure 1. Review criteria used for study selection. *CSPC*, castration sensitive prostate cancer.

largest improvements in survival in patients with advanced PC in the modern era. Time to the CRPC transition was also prolonged in the chemohormonal therapy arm (20.2 vs 11.2 months, HR 0.61, 0.51–0.72). On stratified analysis of patients with high volume disease, defined as visceral metastases, or 4 or more bone lesions with at least 1 lesion beyond the vertebral bodies and pelvis, there was almost 17-month improved overall survival, which was not seen for low volume disease (51.2 vs 34.4 months, HR 0.63, 0.50–0.79, $p < 0.0001$). This was attributable to the higher proportion of hormone resistant cells in the high volume disease subpopulation, which contributed to resistance to hormonal manipulation.

The STAMPEDE study uses a unique phase II/III trial design to investigate new agents under the umbrella of a single trial. Additional arms are added to the study as new approaches are designed.⁹ Patients were initially stratified into 4 groups which received ADT alone, ADT plus docetaxel, ADT plus zoledronate and ADT plus docetaxel plus zoledronate, respectively. In addition to metastatic disease, the study also included lymph node involvement, high risk locally advanced disease (T3/4 plus Gleason score 8–10, T3/4 plus prostate specific antigen 40 ng/ml or greater or Gleason score 8–10 plus prostate specific antigen 40 ng/ml or greater) and recurrent disease previously treated with definitive local therapy. There was a 10-month improvement

Table 1. Phase III combination trials with ADT in metastatic hormone sensitive prostate cancer

	ADT + Docetaxel/ADT			ADT + Abiraterone/ADT	
	GETUG-AFU 15	CHAARTED	STAMPEDE-Docetaxel	LATITUDE	STAMPEDE-Abiraterone
No. pts	192/193	397/393	592/1,184	597/602	960/957
Median age	63/64	64/63	65/65	68/67	67/67
Median survival:					
Overall*	58.9/54.2 Mos	57.6/44 Mos	81/71 Mos	66%/49%	83%/76%
HR (95% CI)	0.88 (0.68–1.14)	0.61 (0.47–0.80)	0.78 (0.66–0.93)	0.62 (0.51–0.76)	0.63 (0.52–0.76)
Progression-free†	22.9/12.9 Mos	20.2/11.7 Mos	37/20 Mos	33/14.8 Mos	75%/45%
HR (95% CI)	0.72 (0.57–0.91)	0.61 (0.51–0.72)	0.61 (0.53–0.70)	0.47 (0.39–0.55)	0.29 (0.25–0.34)

* CHAARTED and STAMPEDE 5-year survival, and LATITUDE and STAMPEDE-Abiraterone 3-year survivor rate with survival in STAMPEDE including M0 (nonmetastatic) and M1 (metastatic) subsets at 60 vs 45 months for STAMPEDE-docetaxel in M1 subset combination and ADT arms, respectively.

† CHAARTED time to CRPC, STAMPEDE clinical, radiographic and biochemical failure-free survival and LATITUDE radiographic survival.

in overall survival in the chemohormonal therapy arm compared to the ADT arm (81 vs 71 months, HR 0.78, 95% CI 0.66–0.93, $p = 0.006$). The secondary end points of failure-free survival and time to the first skeletal related event were also significantly better in the chemohormonal therapy arm.

A third study using this approach, the GETUG-AFU 15 trial, which also compared chemohormonal therapy to ADT, did not report improved outcomes. However, post hoc analysis of data on the high volume disease subset showed a nonsignificant 20% reduction in death.^{10,11} A recent meta-analysis of aggregate data of patients with high volume disease from the CHAARTED and GETUG-AFU 15 trials showed a survival advantage with a pooled HR of 0.68 (95% CI 0.56–0.82, $p < 0.00$).¹² There was also significant heterogeneity in the treatment effect in the combination treatment arm between the high and low volume subgroups ($p = 0.017$). This suggests that other approaches, notably a biomarker based on tumor genotype, might better inform patient selection for treatment.

In GETUG-AFU and CHAARTED side effects were more common in the chemohormonal therapy arm. The most common grade 3 or greater adverse events in GETUG-AFU and CHAARTED were neutropenia at 32% and 12%, febrile neutropenia at 7% and 6%, and fatigue at 7% and 4%, respectively. In each study there were negligible grade 3 or greater adverse events in the ADT arm. Diarrhea, stomatitis, and motor and sensory neuropathy were the less common adverse effects, which developed in less than 1% of the population in the CHAARTED study. STAMPEDE reported additional toxicity in the chemohormonal therapy arm compared with that of ADT alone (grade 3 or greater adverse events in 52% vs 32% of patients). This was mostly due to toxicity during the first 6 months on trial, when grade 3 or greater adverse events were reported in 36% of the chemohormonal therapy arm vs 17% in the ADT arm. A 1-year analysis of 1,998 patients with available profiles revealed a balanced rate of grade 3 or greater adverse events of 10% in each arm. There were 2 deaths in the chemohormonal therapy arm and 72 patients (13%) discontinued treatment. QOL assessment was done 3, 6, 9 and 12 months after randomization in CHAARTED using the FACT (Functional Assessment of Cancer Therapy)-Prostate score. Although QOL scores with docetaxel decreased at 3 months, it was better at 12 months in patients who received docetaxel vs ADT alone.¹³

Chemohormonal therapy may improve survival because of the existence and emergence of hormonally resistant cellular clones during the CRPC transition. Interestingly, each trial that demonstrated a survival advantage had a time delay between the start of ADT and chemotherapy,

including 120 days in CHAARTED and 90 days in STAMPEDE. In GETUG-AFU 15 the patients were required to enroll within 2 months of starting ADT and almost half of them enrolled within 15 days of starting ADT. Some have suggested that this might explain the decreased survival benefit with chemohormonal therapy in this trial. A timed sequence of chemohormonal therapy contributes to maximum synergy by targeting hormone resistant cells when they are most vulnerable.¹⁴ Microtubule targeting chemotherapy inhibited nuclear translocation of androgen receptor in preclinical studies, which also potentially contributes to synergy with ADT.¹⁵

Combined Androgen Deprivation Therapy and Androgen Synthesis Inhibitors. Abiraterone inhibits cytochrome P-450 CYP17, a critical enzyme in androgen biosynthesis in the testes, adrenal gland and prostate. Its active D4A metabolite contributes to its antitumor effects through blockade of multiple steroidogenic enzymes and antagonism of the androgen receptor.¹⁶ Approval of this drug in the prechemotherapy and post-chemotherapy era led to its application to earlier disease. Resistance to ADT is driven in part by up-regulation of androgen receptor signaling through adrenal androgen production, intratumor testosterone production and modification of androgen receptors.¹⁷ The neoadjuvant combination of abiraterone plus prednisone and ADT markedly reduced the tumor burden in men with newly diagnosed, high risk, localized prostate cancer, suggesting a potential role for inhibiting extragonadal androgen biosynthesis before the emergence of resistant clones.¹⁸ These findings led to 2 randomized, phase III trials testing the efficacy of abiraterone and ADT in mHSPC.

In STAMPEDE-Abiraterone the investigators reported outcomes of the combination of abiraterone and prednisone at the time of ADT initiation.¹⁹ They used a multistage, multi-arm setting similar to that of previous trials enrolling patients with newly diagnosed metastatic, node positive or high risk locally advanced disease. Patients who relapsed with high risk features after previous treatment with radical surgery or radiotherapy were also included. The group consisted of 1,917 patients, of whom 52% had metastatic disease, 20% had node positive or node indeterminate nonmetastatic disease and 28% had node negative, high risk nonmetastatic disease. Patients were randomized to receive ADT alone or a combination of ADT plus abiraterone 1,000 mg plus prednisolone 5 mg. This trial also mandated radiotherapy in patients with node negative nonmetastatic disease and provided the option of radiotherapy for patients with node positive nonmetastatic disease. Treatment continued until PSA, radiological or clinical

progression. In patients in whom radiotherapy was planned treatment was administered for 2 years or until any type of progression, whichever came first. Results demonstrated improved overall survival at 3 years in the combination therapy group with a 37% reduction in the relative risk compared to ADT (death HR 0.63, 95% CI 0.52–0.76, $p < 0.001$). Failure-free survival showed a 71% relative risk reduction (treatment failure HR 0.29, 95% CI 0.25–0.34, $p < 0.001$).

In the multicenter, phase III LATITUDE trial 1,199 patients with newly diagnosed mHSPC were enrolled within 3 months of diagnosis and randomized to receive a combination of ADT plus abiraterone 1,000 mg plus prednisolone 5 mg or ADT alone plus placebo.²⁰ The trial was done at a total of 235 sites in 34 countries in Europe, the Asia-Pacific region, Latin America and Canada. Patients needed at least 2 of 3 high risk features, including Gleason score 8 or greater, visceral metastasis and 3 or more bone lesions on imaging. At 3 years two-thirds of the patients in the combination group survived compared to only half of those in the placebo (ADT) group with a relative risk reduction of 38% (HR 0.62, 95% CI 0.51–0.76, $p < 0.001$). The risk reduction was similar to that in STAMPEDE at the end of 3 years of followup (38% vs 37%).⁹ Radiographic progression-free survival was improved in the combination treatment group with a 53% risk reduction (HR 0.47, 95% CI 0.39–0.55, $p < 0.001$).

Abiraterone was well tolerated with adverse events primarily related to elevated mineralocorticoids. Grade 3 hypertension was reported in 20% of patients in LATITUDE compared to 10% on placebo. Hypokalemia was also higher in the abiraterone arm, requiring treatment discontinuation in 2 patients. STAMPEDE reported grade 3 or greater events in 15% of the combined therapy arm compared to 11% in the ADT arm.⁹ Hypertension, elevated transaminases and respiratory events were reported more often in the combination therapy arm. Both studies mentioned that these side effects were medically manageable and seldom caused life threatening adverse events.

Combination therapy with abiraterone is based on the hypothesis that incomplete blockade of androgen production by ADT leads to tumor cell adaptation. By blocking CYP17 almost total suppression of extragonadal androgen production, especially from within the tumor cell, contributes to improved tumor clearance when used synergistically with ADT.²¹ Median time to start abiraterone was 8 weeks after initiating ADT in the STAMPEDE trial and at the same time in the LATITUDE trial. Thus, synchronous targeting with combination therapy does not appear to alter outcomes, unlike chemohormonal therapy, which requires sequential

targeting. In this context the accessory sources of androgen production are blocked, leading to almost total androgen suppression, in contrast to chemohormonal therapy, in which the hormone resistant population is targeted.

Mechanisms of Cellular Persistence after Androgen Deprivation Therapy

Two distinct and not mutually exclusive mechanisms have been proposed in the transition from hormone sensitive cells to CRPC. Tumor cells may acquire new alterations which enable them to survive in the castrated state (adaptation) or preexisting cells capable of surviving hormonal therapy may be selected after a course of ADT (selection).²² Increased levels of intratumor androgens due to incomplete blockade by ADT, amplification of the AR gene, splice variations and a gain of function mutations, changes in the expression of coregulatory molecules to stimulate transcription after antiandrogen binding and bypass of the AR signaling pathway are examples of adaptive mechanisms.^{23,24} The selection model is based on the hypothesis that clones of cells with inherent resistance to ADT are selected after ADT application. Therefore, a heterogeneous population of androgen dependent and androgen independent cells exists before ADT initiation.²⁵ The finding of quiescent stem cells²⁶ and AR deficient neuroendocrine cells, which are inherently resistant to ADT, supports this hypothesis.²⁷ In this context chemohormonal therapy targets selection mechanisms and abiraterone targets adaptive mechanisms.

ADT causes cellular changes in prostate cancer tissue. Apoptosis, autophagy, necrosis and necroptosis are the cell death mechanisms activated after initiating ADT.²⁸

Senescence is a less studied adaptive mechanism in androgen sensitive cells.²⁹ Replicative senescence was first described as a phenotype in primary cells after extensive culture and replicative exhaustion in vitro that was linked to telomere shortening.³⁰ Induced or accelerated senescence in cancer cells results from DNA damage, increased oncogenic signaling and oxidative stress. Senescent cells remain viable and metabolically active, in contrast to apoptosis or autophagy, in which growth is arrested.³¹ Markers of cellular senescence include senescence associated GLB1 (β -galactosidase) expression. Tissues from patients who underwent neoadjuvant ADT before radical prostatectomy demonstrated viable cells that frequently expressed GLB1 and accumulated after ADT.³² Although senescence is cytostatic, cells express a SASP (secretory associated senescent phenotype). The resulting proinflammatory cytokines and growth factors may have permissive effects on surrounding

cancer cells. The unique metabolic phenotype expressed by these persistent senescent cells is characterized by a high metabolic rate, increased glycolysis, high rates of protein synthesis and down-regulation of the AMPK (5' adenosine monophosphate activated protein kinase) pathway.³³ A strategy to remove these persistent senescent cells may maximize responses to ADT and improve long-term patient outcomes.

Given the unique phenotype of these residual cancer cells, it is interesting to speculate that other less toxic agents that alter metabolism might impact ADT outcomes. One agent is the well-known oral glyburide drug metformin. Metformin directly acts on tumor cells by inhibiting the respiratory mitochondrial electron transport chain inhibiting gluconeogenesis and decreasing glucose uptake, as well as activating AMPK.³⁴ It also inhibits fatty acid synthesis, lipid peroxidation and the Krebs cycle, which are crucial for PC cell survival.³⁵ Metformin represses AR mediated signaling in hormone sensitive cell lines³⁶ and enhances the antiproliferative and apoptotic effect of the antiandrogen bicalutamide.³⁷ A putative role for metformin acting synergistically with chemotherapy has been noted in therapy resistant stem cells in breast and pancreatic tumors.³⁸ Interest has also arisen in examining concurrent use of statins and ADT, given inhibition of the HMG-CoA (3-hydroxy-3-methylglutaryl-coenzyme A) enzyme at the rate limiting step in the mevalonate pathway of cholesterol synthesis and the potential importance in cancer.² Previous combination trials using celecoxib and zoledronic acid combined with ADT in patients starting long-term hormonal therapy did not show a survival advantage.^{9,39}

In a recently published retrospective study of more than 87,000 patients who were placed on ADT for advancing PC, metformin improved overall and cancer specific survival, and reduced skeletal metastases compared to diabetic men on insulin or nondiabetic men.⁴⁰ These data suggest that combining metformin with ADT could synergistically eliminate persistent cancer cells after ADT and delay the onset of CRPC. This approach ultimately requires a prospective trial. STAMPEDE is currently evaluating the combination of metformin and ADT (arm K) against the standard of care ADT (arm A) and this ongoing analysis is expected to shed light on this synergistic approach.⁴¹

Patient Selection for Combination Therapy of Metastatic Hormone Sensitive Prostate Cancer

Combined therapy now represents a standard of care in men with mHSPC.

ADT plus docetaxel can be offered to patients with mHSPC who are eligible for chemotherapy,

particularly those with a high metastatic burden or a rapid pace of disease (fig. 2).⁸ Barriers to docetaxel use include advanced patient age, poor performance status, coexisting illness and patient preference. Hematological side effects, neuropathy and fatigue are more common for chemohormonal therapy than for ADT and chemotherapy related deaths, although rare at 1% to 3%, were documented in all 3 randomized trials in which docetaxel was added. The CHAARTED and STAMPEDE docetaxel phase III trials used an 18-week course of therapy (6 cycles, each consisting of 3 weeks), and 26% and 23% of patients, respectively, in the chemohormonal therapy arm did not complete the full course of therapy. This becomes even more important in community practice, where patients with mHSPC are commonly older than those enrolled in the CHAARTED and STAMPEDE clinical trials.⁴²

Abiraterone has a better side effect profile than docetaxel and, being an oral agent, it is easier to administer in the urologist office. Of patients 12% discontinued therapy due to toxic effects. In LATITUDE 88% of patients completed therapy without a dose modification.²⁰ In the abiraterone arm of STAMPEDE only a few patients discontinued therapy due to toxicity.¹⁹ In LATITUDE 63% of patients reported grade 3 or 4 adverse events in the abiraterone group compared to 48% in the placebo group.²⁰ In STAMPEDE 47% of patients in the combination group reported grade 3 or greater adverse events compared to 33% in the placebo group.¹⁹ Most of the latter adverse events were related to mineralocorticoid side effects, including hypertension, fluid retention and hypokalemia. Altered liver transaminases also occur more frequently with abiraterone and must be monitored. A meta-analysis of these trials revealed a threefold increase in grade 3 or greater cardiac and hepatic events, and a twofold increase in grade 3 or greater vascular events in the abiraterone combination group.⁴³

Quality of life and side effects are important to consider in most men who were otherwise asymptomatic. In patients with low volume disease or significant comorbid conditions ADT alone remains an appropriate treatment option, which should be discussed in individualized counseling. The duration of treatment with abiraterone is longer at 2 years or more. This raises concerns about safety, especially in patients with preexisting risk factors for cardiovascular disorders and stroke. STAMPEDE excluded men with a significant cardiac history, limiting generalizations of benefit or toxicity in those patients. A short course of docetaxel might be preferred in patients with good performance status to avoid the long-term effects of steroids and the toxicity associated with abiraterone, including hyperglycemia, cardiovascular risks, osteopenia




Figure 2. Optimizing treatment of newly diagnosed mHSPC. 1, patients with poor performance status or desire to avoid therapy side effects may elect ADT alone. 2, CHAARTED showed 37% relative risk reduction in high volume metastatic subgroup (more than 4 metastases) compared to 27% overall. 3, LATITUDE included patients with de novo metastatic disease and 2 of 3 high risk features, namely Gleason score 8 or greater, visceral metastasis and 3 or more bone lesions on imaging. 4, treatment depends on life expectancy and performance status.

and/or osteoporosis. The requirement for concurrent prednisone with abiraterone can limit its use in patients with brittle diabetes, patients with chronic gastric ulcers and patients with infection.

This raises the question of the ideal therapeutic agent in the setting of mHSPC. To our knowledge a direct head-to-head comparison of ADT plus abiraterone or docetaxel has not been performed, limiting conclusions. A recent analysis of STAMPEDE indicated that contemporaneously randomized patients showed no evidence of a difference in overall or PC specific survival, or in symptomatic skeletal events.⁴⁴ Interestingly, failure-free survival favored abiraterone, likely reflecting the PSA response and the mechanism of action. The docetaxel cohort had more durable survival after failure. Toxicity was similar between the arms with an 11% prevalence of grade 3 or 4 toxicity at 1 year. The question has been raised whether a combination of abiraterone plus docetaxel (and ADT) might lead to an additive benefit in survival. Data on this will emerge from the ongoing PEACE1 (Phase III Study for Patients with Metastatic Hormone-naïve Prostate Cancer) trial.

The cost of long-term abiraterone treatment is also a factor that physicians and patients should consider before starting therapy. While the cost of docetaxel for a 6-cycle course is estimated to be about \$20,000, the cost of abiraterone for a 2-year

course can exceed \$120,000 per patient.^{45,46} While cost analyses have been completed for abiraterone in men with CRPC, to our knowledge they have not been reported in the HSPC setting. Given the extended treatment duration of abiraterone in HSPC cases, often exceeding 2 years, the potential costs can be significant. The fluid nature of prescription drug coverage across different insurers, especially for oral agents, has made it difficult to predict year-to-year costs of these agents. This represents a new world for many prescribers in which monitoring patient costs for these agents is a critical issue requiring close collaboration with oncology pharmacists. The emergence of new assistance programs for these expensive therapies requires dedicated staff to guide patients through these applications.

Ongoing and Future Trials

Recent positive phase III trials have led to a wealth of new trials in the mHSPC space. Combining ADT with the androgen axis inhibitors enzalutamide, apalutamide and orteronel are ongoing (table 2). Enzalutamide is an androgen signaling inhibitor with multiple actions, including blocking AR translocation to the nucleus, AR binding to DNA and receptor mediated DNA transcription.⁴⁷ It has been approved in men with advancing CRPC before and after docetaxel chemotherapy, and recently for

Table 2. Ongoing phase III trials of combination therapy with ADT in metastatic hormone sensitive prostate cancer

Trial Type/Name	ClinicalTrials.gov Identifier	Phase/No. Enrolled	Intervention	Primary		
				End Point	Estimated Completion	
Hormonal: ENZAMET	NCT02446405	III/1,125	ADT + enzalutamide vs ADT + nonsteroidal anti-inflammatory drug	Overall survival	9/2020	
ARCHES	NCT02677896	III/1,100	ADT + enzalutamide vs ADT + placebo	Radiographic progression-free survival	4/2020	
STAMPEDE	NCT00268476	II/III/—	ADT vs ADT + abiraterone + enzalutamide (Arm J)	Overall survival	2020	
TITAN	NCT02488318	III/1,000	ADT + apalutamide vs ADT + placebo	Overall + radiographic progression-free survival	11/2020	
SWOG S1216	NCT01809691	III/1,304	ADT + orteronel vs ADT + bicalutamide	Overall survival	3/2022	
Metabolic (STAMPEDE)	NCT00268476	II/III/—	ADT vs ADT + metformin (Arm K)	Overall survival	2020	
Chemotherapy (PEACE1)	NCT01957436	III/1,168	ADT +/- docetaxel +/- abiraterone +/- radiation	Overall + radiographic progression-free survival	5/2019	

nonmetastatic CRPC with rapidly rising PSA, as shown in PROSPER (Safety and Efficacy Study of Enzalutamide in Patients With Nonmetastatic Castration-Resistant Prostate Cancer). It is being evaluated as combination therapy with ADT for mHSPC separately as well as an adjunct with another androgen axis inhibitor, abiraterone. Apalutamide, which is similar to enzalutamide in action, was recently approved for nonmetastatic CRPC with rapidly rising PSA while on ADT (SPARTAN, Study of Apalutamide [ARN-509] in Men With Non-Metastatic Castration-Resistant Prostate Cancer).⁴⁸ Orteronel is a selective nonsteroidal inhibitor of 17,20 lyase, a key enzyme in androgen synthesis. It has shown significant activity in the setting of CRPC.⁴⁹ Combination therapy with these additional androgen axis inhibitors is expected to contribute to the evolving landscape of mHSPC management.

The synergistic combination of chemotherapy with androgen axis inhibition was also evaluated in combination with radiation therapy (table 2). Metformin is also being assessed in combination with ADT in arm K of STAMPEDE. The side effect profile and low cost would make this an attractive adjunct in the management of prostate cancer if the results show a synergistic benefit. Finally, ADT induces an AR specific T-cell response, suggesting that ADT combined with AR directed immunotherapy might be an alternate approach to prevent the development of AR over expressing CRPC clones.⁵⁰ This approach represents an intriguing and potentially less toxic strategy for future trials.

CONCLUSIONS

It is remarkable that for more than half a century after its introduction androgen suppression remained the preferred front line approach to the treatment of hormonally sensitive, metastatic prostate cancer. With the development of cytotoxic regimens with clinically relevant activity in CRPC earlier combination trials in men with mHSPC have demonstrated significant improvements in survival and QOL. ADT and docetaxel or abiraterone should be considered the standard of care in these patients.

ADT induces adaptive changes in PC cells and selects resistant clones, leading to castration resistance. The susceptibilities generated by ADT can be synergistically targeted to improve survival outcomes and delay the onset of CRPC, as shown in several recent combination trials. Novel approaches targeting metabolic pathways have the potential to be synthetically lethal with ADT by targeting the multiple susceptibilities induced by ADT. This area needs to be explored in further clinical trials.

REFERENCES

- National Cancer Institute Surveillance, Epidemiology, and End Results Program: Cancer Stat Facts: Prostate Cancer. Available at <https://seer.cancer.gov/statfacts/html/prost.html>. Accessed March 15, 2018.
- Tangen CM, Hussain MH, Higano CS et al: Improved overall survival trends of men with newly diagnosed M1 prostate cancer: a SWOG phase III trial experience (S8494, S8894 and S9346). *J Urol* 2012; **188**: 1164.
- James ND, Spears MR, Clarke NW et al: Survival with newly diagnosed metastatic prostate cancer in the "docetaxel era": data from 917 patients in the control arm of the STAMPEDE trial (MRC PR08, CRUK/06/019). *Eur Urol* 2015; **67**: 1028.
- Maximum androgen blockade in advanced prostate cancer: an overview of the randomised trials. Prostate Cancer Trialists' Collaborative Group. *Lancet* 2000; **355**: 1491.
- Millikan RE, Wen S, Pagliaro LC et al: Phase III trial of androgen ablation with or without three cycles of systemic chemotherapy for advanced prostate cancer. *J Clin Oncol* 2008; **26**: 5936.
- Tannock IF, de Wit R, Berry WR et al: Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer. *N Engl J Med* 2004; **351**: 1502.
- Schweizer MT and Antonarakis ES: Chemotherapy and its evolving role in the management of advanced prostate cancer. *Asian J Androl* 2014; **16**: 334.
- Sweeney CJ, Chen YH, Carducci M et al: Chemohormonal therapy in metastatic hormone-sensitive prostate cancer. *N Engl J Med* 2015; **373**: 737.
- James ND, Sydes MR, Clarke NW et al: Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet* 2016; **387**: 1163.
- Gravis G, Fizazi K, Joly F et al: Androgen-deprivation therapy alone or with docetaxel in non-castrate metastatic prostate cancer (GETUG-AFU 15): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2013; **14**: 149.
- Gravis G, Boher JM, Joly F et al: Androgen deprivation therapy (ADT) plus docetaxel versus ADT alone in metastatic non castrate prostate cancer: impact of metastatic burden and long-term survival analysis of the randomized phase 3 GETUG-AFU15 trial. *Eur Urol* 2016; **70**: 256.
- Gravis G, Boher JM, Chen YH et al: Burden of metastatic castrate naive prostate cancer patients, to identify men more likely to benefit from early docetaxel: further analyses of CHAARTED and GETUG-AFU15 studies. *Eur Urol* 2018; **73**: 847.
- Morgans AK, Chen YH, Sweeney CJ et al: Quality of life during treatment with chemohormonal therapy: analysis of e3805 chemohormonal androgen ablation randomized trial in prostate cancer. *J Clin Oncol* 2018; **36**: 1088.
- Eigl BJ, Eggener SE, Baybik J et al: Timing is everything: preclinical evidence supporting simultaneous rather than sequential chemohormonal therapy for prostate cancer. *Clin Cancer Res* 2005; **11**: 4905.
- Zhu ML, Horbinski CM, Garzotto M et al: Tubulin-targeting chemotherapy impairs androgen receptor activity in prostate cancer. *Cancer Res* 2010; **70**: 7992.
- Li Z, Bishop AC, Alyamani M et al: Conversion of abiraterone to D4A drives anti-tumour activity in prostate cancer. *Nature* 2015; **523**: 347.
- Yamaoka M, Hara T and Kusaka M: Overcoming persistent dependency on androgen signaling after progression to castration-resistant prostate cancer. *Clin Cancer Res* 2010; **16**: 4319.
- Taplin ME, Montgomery B, Logothetis CJ et al: Intense androgen-deprivation therapy with abiraterone acetate plus leuprolide acetate in patients with localized high-risk prostate cancer: results of a randomized phase II neoadjuvant study. *J Clin Oncol* 2014; **32**: 3705.
- James ND, de Bono JS, Spears MR et al: Abiraterone for prostate cancer not previously treated with hormone therapy. *N Engl J Med* 2017; **377**: 338.
- Fizazi K, Tran N, Fein L et al: Abiraterone plus prednisone in metastatic, castration-sensitive prostate cancer. *N Engl J Med* 2017; **377**: 352.
- Danila DC, Morris MJ, de Bono JS et al: Phase II multicenter study of abiraterone acetate plus prednisone therapy in patients with docetaxel-treated castration-resistant prostate cancer. *J Clin Oncol* 2010; **28**: 1496.
- Zong Y and Goldstein AS: Adaptation or selection—mechanisms of castration-resistant prostate cancer. *Nat Rev Urol* 2013; **10**: 90.
- Feldman BJ and Feldman D: The development of androgen-independent prostate cancer. *Nat Rev Cancer* 2001; **1**: 34.
- Scher HI and Sawyers CL: Biology of progressive, castration-resistant prostate cancer: directed therapies targeting the androgen-receptor signaling axis. *J Clin Oncol* 2005; **23**: 8253.
- Isaacs JT and Coffey DS: Adaptation versus selection as the mechanism responsible for the relapse of prostatic cancer to androgen ablation therapy as studied in the Dunning R-3327-H adenocarcinoma. *Cancer Res* 1981; **41**: 5070.
- Clevers H: The cancer stem cell: premises, promises and challenges. *Nat Med* 2011; **17**: 313.
- Bonkhoff H: Neuroendocrine cells in benign and malignant prostate tissue: morphogenesis, proliferation, and androgen receptor status. *Prostate Suppl* 1998; **8**: 18.
- Zhang KX, Firus J, Prieur B et al: To die or to survive, a fatal question for the destiny of prostate cancer cells after androgen deprivation therapy. *Cancers (Basel)* 2011; **3**: 1498.
- Ewald JA, Desotelle JA, Church DR et al: Androgen deprivation induces senescence characteristics in prostate cancer cells in vitro and in vivo. *Prostate* 2013; **73**: 337.
- Hayflick L: The limited in vitro lifetime of human diploid cell strains. *Exp Cell Res* 1965; **37**: 614.
- Campisi J: Cancer, aging and cellular senescence. *In Vivo* 2000; **14**: 183.
- Blute ML Jr, Damaschke N, Wagner J et al: Persistence of senescent prostate cancer cells following prolonged neoadjuvant androgen deprivation therapy. *PLoS One* 2017; **12**: e0172048.
- Guo YN, Wang JC, Cai GY et al: AMPK-mediated downregulation of connexin43 and premature senescence of mesangial cells under high-glucose conditions. *Exp Gerontol* 2014; **51**: 71.
- Owen MR, Doran E and Halestrap AP: Evidence that metformin exerts its anti-diabetic effects through inhibition of complex 1 of the mitochondrial respiratory chain. *Biochem J* 2000; **348**: 607.
- Zadra G, Photopoulos C and Loda M: The fat side of prostate cancer. *Biochim Biophys Acta* 2013; **1831**: 1518.
- Wang Y, Liu G, Tong D et al: Metformin represses androgen-dependent and androgen-independent prostate cancers by targeting androgen receptor. *Prostate* 2015; **75**: 1187.
- Colquhoun AJ, Venier NA, Vandersluis AD et al: Metformin enhances the antiproliferative and apoptotic effect of bicalutamide in prostate cancer. *Prostate Cancer Prostatic Dis* 2012; **15**: 346.
- Hirsch HA, Iliopoulos D, Tschlis PN et al: Metformin selectively targets cancer stem cells, and acts together with chemotherapy to block tumor growth and prolong remission. *Cancer Res* 2009; **69**: 7507.
- James ND, Sydes MR, Mason MD et al: Celecoxib plus hormone therapy versus hormone therapy alone for hormone-sensitive prostate cancer: first results from the STAMPEDE multiarm, multistage, randomised controlled trial. *Lancet Oncol* 2012; **13**: 549.
- Richards KA, Liou JI, Cryns VL et al: Metformin use is associated with improved survival in patients with advanced prostate cancer on

- androgen deprivation therapy. *J Urol* 2018; **200**: 1256.
41. Gillessen S, Gilson C, James N et al: Repurposing metformin as therapy for prostate cancer within the STAMPEDE trial platform. *Eur Urol* 2016; **70**: 906.
42. Gerritse FL, Meulenbeld HJ, Roodhart JM et al: Analysis of docetaxel therapy in elderly (≥ 70 years) castration resistant prostate cancer patients enrolled in the Netherlands Prostate Study. *Eur J Cancer* 2013; **49**: 3176.
43. Rydzewska LHM, Burdett S, Vale CL et al: Adding abiraterone to androgen deprivation therapy in men with metastatic hormone-sensitive prostate cancer: a systematic review and meta-analysis. *Eur J Cancer* 2017; **84**: 88.
44. Sydes MR, Spears MR, Mason MD et al: Adding abiraterone or docetaxel to long-term hormone therapy for prostate cancer: directly randomised data from the STAMPEDE multi-arm, multi-stage platform protocol. *Ann Oncol* 2018; **29**: 1235.
45. Ramaekers BLT, Riemsma R, Tomini F et al: Abiraterone acetate for the treatment of chemotherapy-naïve metastatic castration-resistant prostate cancer: an evidence review group perspective of an NICE single technology appraisal. *Pharmacoeconomics* 2017; **35**: 191.
46. Pilon D, Queener M, Lefebvre P et al: Cost per median overall survival month associated with abiraterone acetate and enzalutamide for treatment of patients with metastatic castration-resistant prostate cancer. *J Med Econ* 2016; **19**: 777.
47. Hoffman-Censits J and Kelly WK: Enzalutamide: a novel antiandrogen for patients with castration-resistant prostate cancer. *Clin Cancer Res* 2013; **19**: 1335.
48. Smith MR, Saad F, Chowdhury S et al: Apalutamide treatment and metastasis-free survival in prostate cancer. *N Engl J Med* 2018; **378**: 1408.
49. Fizazi K, Jones R, Oudard S et al: Phase III, randomized, double-blind, multicenter trial comparing orteronel (TAK-700) plus prednisone with placebo plus prednisone in patients with metastatic castration-resistant prostate cancer that has progressed during or after docetaxel-based therapy: ELM-PC 5. *J Clin Oncol* 2015; **33**: 723.
50. Olson BM, Gamat M, Seliski J et al: Prostate cancer cells express more androgen receptor (AR) following androgen deprivation, improving recognition by AR-specific T cells. *Cancer Immunol Res* 2017; **5**: 1074.

Synthetic Lethal Metabolic Targeting of Androgen Deprived Prostate Cancer Cells with Metformin

Bing Yang¹, Shivashankar Damodaran¹, Tariq A. Khemees¹, Mikolaj J. Filon¹, Adam Schultz¹, Joseph Gawdzik¹, Tyler Etheridge¹, Dmitry Malin², Kyle A. Richards¹, Vincent L. Cryns^{2,3} and David F. Jarrard^{1,3,4} *

1. Department of Urology, School of Medicine and Public Health, University of Wisconsin, Madison, WI 53705
2. Department of Medicine, School of Medicine and Public Health, University of Wisconsin, Madison, WI 53705
3. Carbone Comprehensive Cancer Center, University of Wisconsin, Madison, WI 53705
4. Molecular and Environmental Toxicology Program, University of Wisconsin, Madison, WI 53706

Running title: Metformin and ADT

Keywords: Prostate cancer, androgen deprivation therapy, senescence, metformin

Corresponding author: David Jarrard, University of Wisconsin-Madison, 7037 WIMR, 1111 Highland Avenue, Madison WI 53705. Phone: (608) 262-0759. Email: jarrard@urology.wisc.edu.

Conflict of interest: There is no conflict interest for all authors.

Word count: 4,677 total words of text (exclusive of references),

5 Figures and 1 table.

Abstract

The initiation of androgen deprivation therapy (ADT) induces susceptibilities in prostate cancer (PC) cells that make them vulnerable to synergistic treatment and enhanced cell death. Senescence results in cell cycle arrest, but cells remain viable. In this study, we investigated the mechanisms by which PC cells undergo senescence in response to ADT, and determined whether an FDA approved antidiabetic drug metformin has a synergistic effect with ADT in PC both *in vitro* and *in vivo*. Our results show that longer term exposure to ADT induced senescence associated with p16^{INK4a} and/or p27^{kip2} induction. The activation of PI3K/Akt and inactivation of AMPK in senescent cells resulted in mTORC1 activation. In addition, the anti-apoptotic protein XIAP expression was increased in response to ADT. Addition of metformin following ADT induced apoptosis, attenuated mTOR activation by ADT, reduced senescent cell number *in vitro* and inhibited tumor growth in PC PDX models. This study suggests that combining ADT and metformin may be a feasible therapeutic approach to remove persistent PC cells after ADT.

Introduction

The conventional treatment of metastatic hormone-sensitive prostate cancer (mHSPC) has been Androgen Deprivation Therapy (ADT) since its discovery by Huggins and Hodges in 1941 (1). ADT leads to tumor regression, but is rarely curative, as castrate resistant PC (CRPC) develops over 36-40 months. The initiation of ADT may induce susceptibilities in PC cells that make them vulnerable to synergistic treatments that enhance cell death. The recently published Phase III trials combining chemotherapy with ADT (2,3) and abiraterone with ADT (4,5) delayed the onset of CRPC. Thus, one underutilized therapeutic strategy that has the potential to improve outcomes is eradicating the persistent cancer cells that remain after ADT and likely play a key role in the development of CRPC.

Studies have demonstrated *in vitro* and *in vivo* that senescence is induced in androgen sensitive cells after ADT (6-8). Cellular senescence can also be caused by oxidative stress and DNA damaging agents. These senescent cells, although not proliferating, generate a protumor response, the senescent secretory phenotype, characterized by proinflammatory cytokines and growth factors that contribute to PC progression (9,10). Senescent cells are metabolically active and have increased energy demand, protein turnover, and glycolysis (11). Thus, a strategy to eradicate these persistent senescent cells is required to maximize response to ADT and improve long-term patient outcomes.

To target the metabolic therapeutic opportunity that senescence induction presents after ADT, consideration was given to metformin, a widely-used, nontoxic oral antidiabetic agent in the biguanide class. Metformin has multifaceted antitumor activity, including modulating the systemic effects of ADT (12-14). We recently retrospectively analyzed whether advanced PC patients on metformin and ADT had improved survival compared to patients on ADT alone (15).

In a cohort of 87,344 patients with advanced PC, veterans on metformin had improved cancer-specific and overall survival to control patients. In the current study, we examine whether ADT-metformin has a synthetic lethal effect directly on PC cells utilizing *in vitro* and *in vivo* models of PC with variable androgen sensitivity.

Materials and Methods

Cell lines and cell culture

Androgen-dependent LNCaP (ATCC) were routinely cultured in in RPMI-1640 medium (ATCC) supplemented with 10% fetal bovine serum (FBS; Gibco, Invitrogen) and 100 units/ml penicillin/streptomycin (PS, Cell grow), and LAPC4 PC cells were cultured in DMEM medium with 10% FBS and 1% PS. Both were grown at 37°C in 5% CO₂. LNCaP cells were authenticated and tested mycoplasma-free and purchased from the ATCC and used for this project at passage 2. LAPC4 cells were authenticated in March 2014 with STR matching analysis, and only passed once before being used in the described experiments. Mycoplasma testing is performed monthly during cell cultures using MycoProbe Mycoplasma Detection Kit (R&D Systems, #CUL001B). To generate LNCaP cells stably expressing human XIAP, pcDNA3-XIAP plasmid (Addgene, #11833) and empty pcDNA3 vector were transfected into cells using Lipofectamine 3000 (ThermoFisher Scientific). Cells were then cultured in the presence of 700 µg/ml gentamicin (ThermoFisher Scientific) for two weeks to select for XIAP-overexpressing cells, which was confirmed by immunoblotting.

Senescence induction by androgen-deprivation therapy (ADT) followed by metformin treatment

Two ADT approaches were utilized. The first one was to culture the cells in 10% charcoal-stripped serum (CSS) instead of FBS. The second method was to treat the cells with the antiandrogen medication bicalutamide (Sigma, St. Louis, MO). A time course for CSS exposure as well as a time course and dose-response study for bicalutamide and metformin were performed. The optimal duration for induction of senescence by CSS and bicalutamide was 6 days. The doses of bicalutamide and metformin that resulted in 30% and 50% growth inhibition were determined: bicalutamide, 1 and 5 μ M for LNCaP cells and 20 and 30 μ M for LAPC4 cells; metformin, 0.1 and 1 mM for LNCaP cells and 2.5 and 5 mM for LAPC4 cells. Metformin was administered 1 to 3 days after 6-day senescence induction. All controls were treated with vehicles; there were at least three replicates per condition and at least three independent experiments.

Cell growth assay

DNA levels of control and treated cells were quantitated with a fluorescent DNA assay, as an indicator of cell growth, the details of which have been published previously (16). Briefly, 2×10^3 cells were plated in 96-well culture plates containing regular growth medium with FBS. The following day, the media was switched to either CSS medium or regular media containing two concentrations of bicalutamide. After 6 days with ADT, the cells were treated with metformin. When the treatments were completed, plates were washed with 25% PBS once, and then added 100 μ l of sterilized nuclease-free water and frozen at -80° C. Subsequently plates were thawed, incubated with Hoechst 33258 dye at final concentration of 6.7 μ g/mL in high salt TNE buffer, and scanned on a plate reader using 360 nm excitation / 460 nm emission.

Western blotting and antibodies used

Cells were seeded on 6-well plates, the duration and doses for CSS, bicalutamide and metformin were exactly same as described in the growth assay. After harvesting, the cells were lysed in a Tris buffer (pH 8.0) NaCl, EDTA, NP-40 and protease and phosphatase inhibitor cocktail (Invitrogen), incubated on ice for a minimum 30 minutes. Protein concentration was measured using BCA, 20 µg of total protein was run on a 4–12% Bis-Tris pre-cast NuPAGE gel (Invitrogen) and subsequently transferred onto a PVDF membrane (Biorad). Blots were then probed with antibodies overnight at 4° C against the following proteins: AR (SC-7305, Santa Cruz), p16^{INK4} (ab108349, Abcam), p27 (M7203, Dako), phospho-Akt (2065, Cell Signaling), Phospho-S6K1 (P01475, Boster), phospho-AMPK (2535, Cell Signaling), phospho-ACC(D7D11, Cell Signaling), XIAP (2045S, Cell Signaling), cleaved-PARP (9546, Cell Signaling) and α -tubulin (CP06, Calbiochem). Following incubation with the appropriate secondary horseradish peroxidase-conjugated antibodies (Jackson ImmunoResearch), blots were developed using Pierce® ECL solution (Thermo Fisher) and then imaged with Syngene (Cambridge). The blots were probed with each antibody individually; the stripping buffer (Tris pH 6.8 containing SDS and 2-mecapthoethanol) was used to remove the binding of the primary antibody between changes of different primary antibodies.

Detection of cellular senescence

Senescence-associated beta-galactosidase activity (SA- β -gal) was performed by standard protocol (17). Briefly, cells were fixed in 2% formaldehyde and 1% glutaraldehyde in PBS for 5 min at room temperature, washed twice in PBS and incubated overnight at 37°C in freshly-made staining solution containing 1mg/ml of 5-bromo-4-chloro-3-indolyl-b-galactoside (Sigma), 150

mM NaCl, 2 mM MgCl₂, 5 mM K₄[Fe(CN)₆]·3H₂O, 5 mM K₃[Fe(CN)₆] and 40 mM NaPi, pH 6.0. The cells were washed the following day with PBS and images were captured using an Evos microscope (ThermoFisher Scientific). For each analysis, five photographs were taken from each of five independent replicates per condition. To confirm our visual assessment of cellular enlargement and complexity, flow cytometry was utilized to evaluate side scatter (SSC), a parameter commonly increased in senescence (10). After treatment, cells were harvested using trypsin and then live cells were analyzed immediately using flow cytometry (MACSQuant 10). The protein expression of the senescence markers *p16* and *p27* was also detected by western blotting.

Measurement of apoptosis

Apoptosis was measured by caspase 3 activity using Caspase-Glo® 3/7 Assay (Promega) following the manufacturer's instructions. Briefly, control and treated cells were plated in triplicate in a white-walled 96-well plate. A blank well with vehicle and cell culture medium but without cells was utilized as a control. The same number of cells and treatment groups were plated into a black-walled 96-well plate at the same time to define the cell number by DNA assay. One hundred µl of Caspase-Glo reagent (reagent to cell culture media 1:1 ratio) was added to each well of plates containing blank which has vehicle and cell culture medium without cells, the plate was covered and contents mixed gently using a plate shaker at 300–500 rpm for 30 seconds. After incubation at room temperature for 1 hour, the luminescence was measured using a plate-reading illuminometer (PE EnSpire). Apoptosis was calculated as luminescence of each sample subtracted from the blank, and then normalized to cell number. Apoptosis was presented as fold change normalized to control.

XIAP siRNA experiments

Cells were transfected with siRNAs targeting the sequence CAGUGAAGACCCUUGGGAA (si-XIAP) in human XIAP or non-silencing control siRNAs (Sigma-Aldrich) using Lipofectamine RNAiMAX Reagent (ThermoFisher Scientific) as per the manufacturer's instructions.

Mouse Xenograft studies

Animal work performed in this study was approved by the IACUC at University of Wisconsin-Madison. Patient-derived xenografts LuCaP 77 and 147 were kindly provided by Dr. Eva Corey at the University of Washington. The clinicopathological and patient characteristics were well defined and reflect the molecular heterogeneity of PC (18). Xenograft (LuCaP) 77 was derived from a PC patient with a femur metastasis and disease for 15 years prior to harvest, and LuCaP 147 from a patient with liver metastases and more aggressive disease for 2 years prior to harvest. LuCaP 77 shows an intermediate response to castration and LuCaP 147 demonstrates less response by tumor volume, serum PSA level and survival after ADT. Six-week old severe combined immunocompromised mice (CB-17 strain SCID, Code #236) were procured from Charles River. LuCaP 77 or LuCaP 147 was implanted into the mouse flank region bilaterally. When flank tumors were 100 mm³, half of the mice were castrated and the other half treated with sham surgery.

The animals were then divided into 4 groups, 16 mice per group: (1) castrated and treated with metformin (CM), (2) castrated and treated with vehicle control (CC), (3) sham surgery and treated with metformin (SM), and (4) sham surgery and treated with vehicle control (SC).

Metformin was administered one week after surgery in drinking water at a dose of 2g / liter (19). Metformin undergoes renal excretion and has a mean plasma elimination half-life after oral administration of between 4.0 and 8.7 hours (20). Tumor growth and body weight were measured weekly and half of the animals in each group were harvested 4 weeks after metformin administrated to permit an assessment of senescence, apoptosis and molecular characteristics of the tumors. The remaining animals were monitored and sacrificed when tumor size exceeded 17 mm in accordance with CACC guidelines.

Tissue Microarrays (TMA) were constructed from the above mouse tumors, with duplicate cores per tissue. The VECTRA system (Perkin Elmer) was used for image analysis and quantification of the staining intensity (21). Cores with <5% epithelial component or loss of tissue were excluded from the analysis. Nuance system and Inform 1.2™ software (Caliper Life Sciences) were utilized for building spectral libraries on the basis of target signals of the stained parameters. This system allows automated quantitation of colorimetric/ fluorescent staining on a per-cell basis and selection of cellular subsets (nucleus versus cytoplasmic) for analysis of target signals. Expression of *Ki67* (proliferation) and cleaved caspase 3 (apoptosis) were evaluated by immunohistochemistry. H&E staining was used to differentiate nuclear and cytoplasmic tissue compartments. Two cores from the same tumor were averaged to give a more precise estimate. We also detected the evaluated protein expression identified *in vitro* in the xenografts of all above groups.

Statistical Analysis

Data were shown as a mean $\pm$ s.d., with a minimum of three replicates in all *in vitro* experiments. For *in vivo* studies, the data are shown as mean $\pm$ s.e. The significance of the

differences between two groups was analyzed with one-way ANOVA, $p < 0.05$ was considered significant. CalcuSyn Version 2.0 was used to quantify the synergism of combined drug effects.

Results

Combining metformin with ADT induces synergistic growth inhibition in PC cell lines compared to monotherapy.

To understand the phenotype of hormone-sensitive PC cells after ADT, we characterized the LNCaP and LAPC4 cell lines for senescence and other characteristics. Cells exposed to either charcoal stripped serum (CSS) or the androgen receptor antagonist bicalutamide for 6 days demonstrate an enlarged and flattened morphology, increased SA- β -gal activity representing an expanded lysosomal compartment, and increased side scatter (SSC), a flow cytometry parameter increased in senescence (6,10) (Fig. 1A and B). Time-dependent induction of senescence was confirmed by the increased expression of the senescence markers p16 and p27 on western blot analysis (Fig. 1C). Expression of p16 increased in terminal senescence after treatment with either CSS or bicalutamide, while p27 was increased in early senescence. Senescent markers and morphology appear maximally induced at 6-10 days, confirming previous data (6).

Given data showing rapid gluconeogenesis and increased proteolytic stress in therapy-induced senescence (11), we investigated whether metformin, an oral antidiabetic agent in the biguanide class, acts in a synthetic lethal manner to cause apoptosis in cells first treated with ADT. To test for synergy we utilized a standard approach described by Chang, et al (22) that utilizes CalcuSyn™, a software program that quantifies synergy based on cell counts with high

and low dose combinations (Supplementary Table S1). Dosing of bicalutamide, CSS and metformin were chosen based on 20-30% (low) or 50-60% (high) cell growth inhibition using a DNA assay at 6 days in LNCaP and LAPC4 cells (Supplementary Fig. S1). Exposure to either ADT or metformin alone caused a significant reduction in growth in both cell lines ($p < 0.05$) compared to the controls (Fig. 2A-D and Supplementary Fig. S2). Combining bicalutamide or CSS with metformin resulted in further growth inhibition compared with treatment with either monotherapy. Combination of bicalutamide followed by metformin showed strong synergy ($CI = 0.1-0.3$) in both cell lines, and CSS with metformin displayed a moderate calculated synergistic effect ($CI = 0.5-0.8$) (Supplementary Table S1).

To identify the extent of cell death in PC cells treated with the combination of ADT and metformin, we evaluated apoptosis using the Caspase-Glo Assay measuring multiple caspases. DNA levels in control and treated cell cultures were simultaneously measured to reflect cell number. PC cells exposed to ADT for 6 days followed by 2 days treatment with metformin demonstrated a 2-3 fold increase in Caspase-Glo activity with the combination therapy compared to ADT or metformin alone (Table 1). The clearance of these senescent cells is also reflected by decreases in the expression of the senescence markers *p16* after metformin treatment (Fig. 2E and F) as well as a decrease in SA-B-gal stained cells in ADT/Met treated cultures (24% decreased for LNCaP). At both 24 and 48 hours after metformin addition, a comparison of ADT plus Metformin versus the ADT shows a decrease in senescence marker expression in both LNCaP and LAPC4 cells.

Akt and AMPK signaling pathways are associated with cell survival after ADT through upregulation of the anti-apoptotic protein XIAP.

Akt plays a critical role in cell survival, senescence and apoptosis (23,24). Investigating responses to ADT and metformin using western blot at multiple timepoints, we observed that activated p-Akt levels increase with the duration of ADT (Supplementary Fig. S3). The effect of Akt on the key downstream mediator mTORC1 was measured using phospho-p70S6K (*p-PS6K*). Increased p-PS6K confirms the activation of PI3K/Akt/mTORC1 signaling. Both elevated *p-Akt* and *p-PS6K* are decreased with the subsequent addition of metformin to ADT-treated LNCaP and LAPC4 cells consistent with removal of senescent cells (Fig. 3A and B).

AMPK is a cellular sensor that plays a key role in maintaining energy homeostasis. We observed that phospho-AMPK increases in PC cells 2-4 days after ADT treatment when senescence markers are minimally induced (Supplementary Fig. S3). Subsequently, AMPK activity is downregulated in late terminal senescence (8-10 days). In late ADT treated cells, the addition of metformin activates AMPK, notably reversing the inhibition of AMPK in senescent cells with longer-term treatment ADT (Fig. 3C). The alteration of AMPK activity by ADT or metformin was confirmed by phosphorylation its downstream protein ACC at Ser79. Less phosphorylation of ACC on Ser79 was observed when AMPK activity is downregulated in late ADT treated cells, and increased when AMPK activity is resumed by the addition of metformin. AMPK is a known suppressor of mTORC1 signaling through phosphorylation of TSC2 (25) and we find increasing expression of phosphorylated mTOR1 (pAKT-T308) with time of ADT exposure (Supplementary Fig. S3). AMPK also activates autophagy (25), but an assessment of ADT did not activate autophagy during early senescence in either LNCaP or LAPC4 cells (Supplementary Fig. S4) when AMPK was induced.

Cell senescence is associated with a reduced susceptibility to cell death (26). To determine the basis for metformin to induce increased cell death in ADT-treated cells we

investigated the role of the anti-apoptotic protein XIAP, a gene known to be positively regulated by AKT signaling (27,28). ADT in both LNCaP and LAPC4 cultures induced XIAP protein expression (Fig. 3A and B). The addition of metformin to cell cultures after ADT resulted in reduced XIAP expression compared to ADT alone shown schematically in Fig. 3A and B.

Given these findings, we further investigated the role XIAP plays in the cell response to ADT and metformin. First, LNCaP cells were transfected with control or XIAP-targeted siRNAs. XIAP knockdown reduced cell growth in both ADT exposed and control media compared to si-Control (Fig. 4A). pAkt and p-p70S6K expression were decreased, and apoptosis increased to a greater extent in ADT-treated cells with XIAP knockdown than in control media, indicating that a significant part of the survival response that ADT induces is XIAP-dependent (Fig. 4B and C).

We then examined whether overexpression of XIAP alters the response of PC cells to ADT and metformin. XIAP was overexpressed in LNCaP cells using a pcDNA3-XIAP vector, while the empty vector was used as a control. After stable cell lines were established, cell growth, apoptosis and protein expression were analyzed (Figs 4D-F). Overexpression of XIAP abrogated the synergistic decrease in growth seen with metformin following senescence induction by ADT. In addition, XIAP overexpression reversed the increased apoptosis after metformin exposure or the combination of ADT and metformin (Fig. 4F). Moreover, the senescence marker p16 was reproducibly increased compared to the cells transfected with control vector, suggesting increased cell survival (Fig. 4E). Akt signaling increased in cells overexpressing XIAP compared to controls, consistent with a known feedback loop (29). These data support a role for the anti-apoptotic protein XIAP in the response of PC cells to ADT and metformin.

Metformin and ADT suppresses tumor growth in patient-derived xenografts (PDXs) more robustly than ADT alone.

The role of metformin in ADT-treated cancer was then tested on human tumors passaged *in vivo*. Patient-derived PC xenografts LuCaP 77 and LuCaP 147 were implanted and passaged in immune-deficient mice as previously described (18). Both PDX models exhibit some growth inhibition in response to castration with LuCaP 77 showing an intermediate and LuCaP 147 a lesser response to castration (18). When flank tumors reached 100 mm³ mice were castrated or treated with sham surgery and metformin or placebo administered one week later. Tumor size was calculated weekly as volume and presented as a percentage of the tumor size at castration.

The growth of both PDXs was initially inhibited by castration but began regrowth after 3-4 weeks (Fig. 5). Metformin alone modestly inhibited tumor growth compared to controls in both models. With the combination treatment, LuCaP 77 tumors at 7 weeks were significantly smaller than those in the castration ($p<0.01$) or metformin ($p<0.005$) alone groups (Fig. 5A). In the LuCaP 147 PDX model, tumors were significantly smaller at 6-8 weeks post-surgery in the combination therapy compared to those in the castration ($p<0.005$) or metformin alone groups ($p<0.01$, Fig. 5B).

Protein expression of the proliferation marker Ki67 and the apoptosis marker cleaved caspase 3 (CC3) was analyzed using quantitative VECTRA immunostaining. Ki67 expression decreased in the castration and castration plus metformin groups (Fig. 5C and D). A difference in CC3-positivity in tumors from different treatment groups was not detected ($p=0.1905$). In addition, western blot analysis of remaining samples was performed which demonstrated decreased pAMPK and increased XIAP expression from the mice post-castration at one month, while metformin attenuated these effects in the majority of tumors analyzed (Fig. 5E).

Discussion

A growing body of evidence indicates that ADT induces a susceptible therapeutic niche that may be exploited to enhance therapeutic response (30). As a sole treatment for PC, metformin demonstrates little benefit. This has led investigators to examine its role as an additive or synergistic treatment in combination with other drugs (31-33). In the current study, we demonstrate for the first time a synergistic inhibitory effect of metformin combined with ADT on growth in human PC cell lines *in vitro* and PDX tumors *in vivo*. We find a novel role for XIAP, an anti-apoptotic protein driven by increased AKT signaling in PC cells after ADT. These observations mechanistically support our recent study demonstrating improved survival in patients taking metformin at the time of initiating ADT for advanced PC (15).

The fate of PC cells after ADT includes apoptosis, quiescence and other phenotypes. In the current work, we demonstrate that subpopulations of androgen-sensitive PC cells after ADT exhibit molecular markers characteristic of senescence, such as increased SA- β -gal-staining, cellular complexity measured by flow cytometry and expression of cyclin dependent kinase inhibitor 2A p16^{INK4a}. Senescence is mediated by p16^{INK4a}, p53/p21 or p27^{kip} depending on the cell type, stressor and duration of stress (34). We found that CSS as well as bicalutamide induces a senescent phenotype in the observed cells predominantly associated with an accumulation of p16, but p27 is an earlier transient marker consistent with previous senescence studies (6,8). Apoptosis occurs early after ADT and is limited. In a series of xenograft tumors *in vivo*, surgical castration shows a similar induction of senescence with decreased proliferation as determined by

Ki-67 expression (6). Autophagy is induced by AMPK and had been reported to occur in LNCaP cells after high dose bicalutamide (35). However, autophagy was not strongly induced in our model (Supplementary Fig. S4), potentially because of the lower, more physiologically relevant doses of bicalutamide employed in our studies. In addition, CSS reduced androgen receptor (AR) expression in both LNCaP and LAPC4 cells, while bicalutamide effects were more variable. AR expression increased at 48hr when metformin was added to ADT, compared to metformin alone. These effects may reflect the removal of senescent cells by metformin.

In order to define the events contributing to this response we exposed cells sequentially first to ADT and characterized the activated signaling pathways before adding metformin. We found that both Akt and AMPK signaling pathways are activated initially in multiple cell lines with shorter-term (4 day) exposure to ADT. Continued expression of the PI3K/Akt/mTOR signaling pathway that regulates cellular survival, differentiation and stem cell-like properties is noted at extended 8 day timepoints (36-39). However, AMPK signaling is repressed later after ADT exposure (Supplementary Fig. S3 and Fig. 3) when the full phenotype of senescence is most marked. AMPK acts as a cellular energy sensor that stimulates glycolysis by activating PFKFB2/3 and negatively regulates the mTOR signaling pathway through the phosphorylation of the transcription initiation factor TSC2 (25). The balance between activation of mTORC1 via AKT activation and inhibition of mTORC1 via AMPK activation may restrict cell growth in early ADT. However, with extended ADT when senescence is robustly induced, AMPK phosphorylation is almost completely inhibited. Persistent activation of AKT and inhibition of AMPK activating mTORC1 signaling (Fig. 3C) may serve as one of the pathways by which senescent cells persist and represents a potential for therapeutic exploitation.

A synergistic reduction in cell number was noted with the addition of metformin to ADT *in vitro*, as well as a reduction in xenograft tumor growth with the combination *in vivo*. In addition to a decrease in the number of cells with the combination, we observed increased apoptosis and a reduction in the senescence marker p16, suggesting clearance of senescent cells. In these cultures, the addition of metformin induces AMPK signaling and represses pAkt (Fig. 3). The finding of Akt induction after ADT, prompted an examination of XIAP (X-linked inhibitor of apoptosis protein), a gene previously found to be upregulated in castration-resistant PC (8). We observed XIAP levels increase in response to ADT with a similar time course as PI3K/Akt activation. Akt phosphorylates XIAP at Ser 87, leading to its stabilization and promoting cell survival (40). In addition, metformin has been shown to downregulate XIAP in breast cancer through the inhibition of mTORC1 (19). A seminal observation of the current study is that the addition of metformin downregulates XIAP in ADT-induced senescent PC cells, a process mimicked by siRNA knockdown of XIAP (Fig. 4) making them more susceptible to apoptosis.

We have also demonstrated that knockdown of the apoptosis-inhibitor XIAP mimics the effects of metformin in sensitizing PC cells to ADT. Our findings are consistent with a previous report that metformin sensitizes breast cancer cells to TRAIL-induced apoptosis by downregulating expression of XIAP (19). To determine the functional role of XIAP downregulation by metformin on the response to ADT, we stably overexpressed XIAP in LNCaP cells and observed that XIAP overexpression abrogated the synergistic effect of ADT and metformin on cell growth and increased apoptosis (Fig 4). Metformin inhibits senescent cell growth by inhibiting AKT and activating AMPK signaling, and these changes were reversed by XIAP overexpression. There is a feedback loop between XIAP and AKT whereby XIAP

regulates the activation of AKT, and this may explain the increased pAKT levels with XIAP overexpression (29). AMPK signaling induction by metformin was not affected by overexpression of XIAP. These experiments point to a key role of XIAP downregulation by metformin in augmenting cell death by ADT, but we do not discount the potential involvement of other cell survival pathways.

Conclusions

The exposure of PC cells to ADT generates dichotomous responses inducing both survival and cell death pathways. We demonstrate that AMPK signaling is inactivated over time, while AKT and the mTOR1 signaling pathways are induced. The addition of metformin activates AMPK, leading to downregulation of the anti-apoptotic protein XIAP and induction of cell death. These studies provide a molecular rationale for the observed clinical observation that metformin treatment at the time of ADT improves both overall and cancer-specific survival in patients with advanced PCa (41). We would anticipate that other agents that activate AMPK efficiently may also act synergistically with ADT. AMPK activator-AICAR has been reported to decrease growth of prostate cancer cells (LNCaP cells and Du145) *in vitro* (42,43). A number of these agents such as MT 63-78 and Salicylic acid are being developed clinically (44-46).

Acknowledgments

This research is supported by the U.S. Department of Defense Prostate Cancer Research Program (grant no. W81XWH-16-1-0278 to D.F. Jarrard and W81XWH-16-1-0279 to V. Cryns).

References

1. Denmeade SR, Isaacs JT. A history of prostate cancer treatment. *Nature reviews Cancer* 2002;2:389-96.
2. Sweeney CJ, Chen YH, Carducci M, Liu G, Jarrard DF, Eisenberger M, *et al.* Chemohormonal Therapy in Metastatic Hormone-Sensitive Prostate Cancer. *The New England journal of medicine* 2015;373:737-46.
3. James ND, Sydes MR, Clarke NW, Mason MD, Dearnaley DP, Spears MR, *et al.* Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet* 2016;387:1163-77.
4. James ND, de Bono JS, Spears MR, Clarke NW, Mason MD, Dearnaley DP, *et al.* Abiraterone for Prostate Cancer Not Previously Treated with Hormone Therapy. *N Engl J Med.* 2017;377:338-351.
5. Fizazi K, Tran N, Fein L, Matsubara N, Rodriguez-Antolin A, Alekseev BY, *et al.* Abiraterone plus Prednisone in Metastatic, Castration-Sensitive Prostate Cancer. *N Engl J Med.* 2017;377(4):352-360.
6. Ewald JA, Desotelle JA, Church DR, Yang B, Huang W, Laurila TA, *et al.* Androgen deprivation induces senescence characteristics in prostate cancer cells in vitro and in vivo. *The Prostate* 2013;73:337-45.
7. Pernicová Z, Slabáková E, Kharaishvili G, Bouchal J, Král M, Kunická Z, *et al.* Androgen depletion induces senescence in prostate cancer cells through down-regulation of Skp2. *Neoplasia* 2011;13:526-36.
8. Burton DG, Giribaldi MG, Munoz A, Halvorsen K, Patel A, Jorda M, *et al.* Androgen deprivation-induced senescence promotes outgrowth of androgen-refractory prostate cancer cells. *PLoS One* 2013;8:e68003.
9. Taddei ML, Cavallini L, Comito G, Giannoni E, Folini M, Marini A, *et al.* Senescent stroma promotes prostate cancer progression: the role of miR-210. *Mol Oncol* 2014;8:1729-46.
10. Ewald JA, Desotelle JA, Wilding G, Jarrard DF. Therapy-Induced Senescence in Cancer. *Journal of the National Cancer Institute* 2010;102:1536-46.
11. Dörr JR, Yu Y, Milanovic M, Beuster G, Zasada C, Däbritz JH, *et al.* Synthetic lethal metabolic targeting of cellular senescence in cancer therapy. *Nature* 2013;501(7467):421-5.
12. Whitburn J, Edwards CM, Sooriakumaran P. Metformin and Prostate Cancer: a New Role for an Old Drug. *Curr Urol Rep* 2017;18:46.
13. Saraei P, Asadi I, Kakar MA, Moradi-Kor N. The beneficial effects of metformin on cancer prevention and therapy: a comprehensive review of recent advances. *Cancer Manag Res* 2019;11:3295-313.
14. Nobes JP, Langley SE, Klopper T, Russell-Jones D, Laing RW. A prospective, randomized pilot study evaluating the effects of metformin and lifestyle intervention on patients with prostate cancer receiving androgen deprivation therapy. *BJU Int* 2012;109(10):1495-502.

15. Richards KA, Liou JI, Cryns VL, Downs TM, Abel EJ, Jarrard DF. Metformin Use is Associated with Improved Survival for Patients with Advanced Prostate Cancer on Androgen Deprivation Therapy. *J Urol* 2018;200(6):1256-63.
16. Mehraein-Ghomi F, Kegel SJ, Church DR, Schmidt JS, Reuter QR, Saphner EL, *et al.* Targeting androgen receptor and JunD interaction for prevention of prostate cancer progression. *The Prostate* 2014;74:792-803.
17. Dimri GP, Lee XH, Basile G, Acosta M, Scott C, Roskelley C, *et al.* A Biomarker That Identifies Senescent Human-Cells in Culture and in Aging Skin in-Vivo. *Proc Natl Acad Sci U S A* 1995;92:9363-7.
18. Nguyen HM, Vessella RL, Morrissey C, Brown LG, Coleman IM, Higano CS, *et al.* LuCaP Prostate Cancer Patient-Derived Xenografts Reflect the Molecular Heterogeneity of Advanced Disease and Serve as Models for Evaluating Cancer Therapeutics. *The Prostate* 2017;77(6):654-71.
19. Strekalova E, Malin D, Rajanala H, Cryns VL. Metformin sensitizes triple-negative breast cancer to proapoptotic TRAIL receptor agonists by suppressing XIAP expression. *Breast Cancer Res Treat* 2017;163(3):435-47.
20. Scheen AJ. Clinical pharmacokinetics of metformin. *Clin Pharmacokinet* 1996;30(5):359-71.
21. Huang W, Hennrick K, Drew S. A colorful future of quantitative pathology: validation of Vectra technology using chromogenic multiplexed immunohistochemistry and prostate tissue microarrays. *Hum Pathol* 2013;44(1):29-38.
22. Chang TT, Chou TC. Rational approach to the clinical protocol design for drug combinations: a review. *Acta Paediatr Taiwan* 2000;41(6):294-302.
23. Edlind MP, Hsieh AC. PI3K-AKT-mTOR signaling in prostate cancer progression and androgen deprivation therapy resistance. *Asian J Androl* 2014;16(3):378-86.
24. Bitting RL, Armstrong AJ. Targeting the PI3K/Akt/mTOR pathway in castration-resistant prostate cancer. *Endocr Relat Cancer* 2013;20(3):R83-99.
25. Jeon SM. Regulation and function of AMPK in physiology and diseases. *Exp Mol Med* 2016;48:e245.
26. Hampel B, Malisan F, Niederegger H, Testi R, Jansen-Dürr P. Differential regulation of apoptotic cell death in senescent human cells. *Exp Gerontol* 2004;39:1713-21.
27. Asselin E, Wang Y, Tsang BK. X-linked inhibitor of apoptosis protein activates the phosphatidylinositol 3-kinase/Akt pathway in rat granulosa cells during follicular development. *Endocrinology* 2001;142:2451-7.
28. Phillipps HR, Hurst PR. XIAP: a potential determinant of ovarian follicular fate. *Reproduction* 2012;144:165-76.
29. Hussain AR, Uddin S, Ahmed M, Bu R, Ahmed SO, Abubaker J, *et al.* Prognostic significance of XIAP expression in DLBCL and effect of its inhibition on AKT signalling. *J Pathol* 2010;222:180-90.
30. Bao BY, Pao JB, Huang CN, Pu YS, Chang TY, Lan YH, *et al.* Significant associations of prostate cancer susceptibility variants with survival in patients treated with androgen-deprivation therapy. *Int J Cancer* 2012;130:876-84.
31. Sun S, Gong F, Liu P, Miao Q. Metformin combined with quercetin synergistically repressed prostate cancer cells via inhibition of VEGF/PI3K/Akt signaling pathway. *Gene* 2018;664:50-7.

32. Delma MI. Three May Be Better Than Two: A Proposal for Metformin Addition to PI3K/Akt Inhibitor-antiandrogen Combination in Castration-resistant Prostate Cancer. *Cureus* 2018;10:e3403.
33. Babcook MA, Shukla S, Fu P, Vazquez EJ, Puchowicz MA, Molter JP, *et al.* Synergistic simvastatin and metformin combination chemotherapy for osseous metastatic castration-resistant prostate cancer. *Mol Cancer Ther* 2014;13:2288-302.
34. Schwarze SR, Shi Y, Fu VX, Watson PA, Jarrard DF. Role of cyclin-dependent kinase inhibitors in the growth arrest at senescence in human prostate epithelial and uroepithelial cells. *Oncogene* 2001;20:8184-92.
35. Boutin B, Tajeddine N, Vandersmissen P, Zanou N, Van Schoor M, Mondin L, *et al.* Androgen deprivation and androgen receptor competition by bicalutamide induce autophagy of hormone-resistant prostate cancer cells and confer resistance to apoptosis. *Prostate* 2013;73:1090-102.
36. Dubrovskaya A, Kim S, Salamone RJ, Walker JR, Maira SM, García-Echeverría C, *et al.* The role of PTEN/Akt/PI3K signaling in the maintenance and viability of prostate cancer stem-like cell populations. *Proc Natl Acad Sci U S A* 2009;106:268-73.
37. Courtney KD, Corcoran RB, Engelman JA. The PI3K pathway as drug target in human cancer. *J Clin Oncol* 2010;28:1075-83.
38. Liu P, Cheng H, Roberts TM, Zhao JJ. Targeting the phosphoinositide 3-kinase pathway in cancer. *Nat Rev Drug Discov* 2009;8:627-44.
39. Lim M, Chuong CM, Roy-Burman P. PI3K, Erk signaling in BMP7-induced epithelial-mesenchymal transition (EMT) of PC-3 prostate cancer cells in 2- and 3-dimensional cultures. *Horm Cancer* 2011;2:298-309.
40. Nakhaei P, Sun Q, Solis M, Mesplede T, Bonneil E, Paz S, *et al.* I κ B kinase ϵ -dependent phosphorylation and degradation of X-linked inhibitor of apoptosis sensitizes cells to virus-induced apoptosis. *J Virol* 2012;86:726-37.
41. Etheridge T, Damodaran S, Schultz A, Richards KA, Gawdzik J, Yang B, *et al.* Combination therapy with androgen deprivation for hormone sensitive prostate cancer: A new frontier. *Asian J Urol* 2019;6:57-64.
42. Sauer H, Engel S, Milosevic N, Sharifpanah F, Wartenberg M. Activation of AMP-kinase by AICAR induces apoptosis of DU-145 prostate cancer cells through generation of reactive oxygen species and activation of c-Jun N-terminal kinase. *Int J Oncol* 2012;40:501-8.
43. Xiang X, Saha AK, Wen R, Ruderman NB, Luo Z. AMP-activated protein kinase activators can inhibit the growth of prostate cancer cells by multiple mechanisms. *Biochem Biophys Res Commun* 2004;321:161-7.
44. Hung LVM, Moon JY, Ryu JY, Cho SK. Nootkatone, an AMPK activator derived from grapefruit, inhibits KRAS downstream pathway and sensitizes non-small-cell lung cancer A549 cells to adriamycin. *Phytomedicine* 2019;63:153000.
45. Schiewer MJ, Knudsen KE. AMPed up to treat prostate cancer: novel AMPK activators emerge for cancer therapy. *EMBO Mol Med* 2014;6(4):439-41.
46. Zadra G, Photopoulos C, Tyekucheva S, Heidari P, Weng QP, Fedele G, *et al.* A novel direct activator of AMPK inhibits prostate cancer growth by blocking lipogenesis. *EMBO Mol Med* 2014;6:519-38.

Table 1. Addition of metformin increases apoptosis in cancer cells treated with androgen deprivation.

	LNCaP	LAPC4
FBS	0.99 ± 0.09	1.00 ± 0.11
CSS	0.92 ± 0.13	1.29 ± 0.27
Met	1.21 ± 0.17	1.13 ± 0.12
CSS + Met	2.51 ± 0.04	1.72 ± 0.18
Bic	0.87 ± 0.05	0.95 ± 0.32
Bic + Met	2.00 ± 0.18	2.97 ± 0.04

Apoptosis was assayed using Caspase-Glo, shown as fold change relative to control, Mean ± SD value. CSS: 10% CSS; Bic, bicalutamide (μM), LNCaP, 5; LAPC4 30. Met: metformin (mM): LNCaP, 1; LAPC4, 5. Statistical analysis using one-way ANOVA demonstrates increased apoptosis with CSS/Met compared to CSS or Met alone (P-values: LNCaP 0.0039, 0.0067 and LAPC4 0.242, 0.005, respectively). Bic/Met demonstrates increased apoptosis compared to Bic or Met alone (P-values LNCaP 0.0056, 0.063, LAPC4 0.0083, 0.0016, respectively). Data shown as Mean ± SD.

Figure legends

Figure 1. ADT induces phenotypic characteristics of cellular senescence in androgen-dependent prostate cancer cells. **A**, LNCaP or LAPC4 cells were cultured in CSS medium or treated with bicalutamide for 6 days, and then stained for SA- β -gal activity, magnification $\times 100$. Con: 10% FBS in RPMI media; CSS: 10% Charcoal-stripped serum in RPMI media ; Bic, bicalutamide in complete RPMI media, LNCaP, 5 μ M; LAPC 30 μ M. **B**, Flow cytometry was used to examine the cellular complexity side scatter, percentage of cells with side-scatter is shown in graph , Mean $\pm$ SD, * t-test, $p < 0.05$. Triplicate in each group, three independent experiments. ADT increases the fraction of cells with elevated cellular complexity and size. **C**, LNCaP and LAPC4 cells were cultured in CSS medium or treated with bicalutamide for days indicated, then whole protein lysates were harvested for analysis of senescent proteins p16 and p27 expression.

Figure 2. Combination of ADT with metformin decreases the growth of androgen-dependent prostate cancer cells compared to single-agent use. LNCaP or LAPC4 cells were cultured in medium containing different percentage of FBS/CSS (**A**, **C**) or bicalutamide (**B**, **D**) for 6 days followed by addition of high or low dose metformin for 48 hours. Cell proliferation was measured with DNA assay. Synergistic effects were calculated by Calcsyn, moderate synergy (combination index CI: 0.7-0.85) +; synergy (CI: 0.3-0.7) ++, strong synergy (CI: 0.1-0.3) +++. Conditions for CSS included: Con: 10% FBS; CSSL: 8% CSS + 2% FBS; CSSH: 10% CSS; ML, metformin low dose (mM): LNCaP, 0.1; LAPC4, 2.5. MH, metformin high dose (mM): LNCaP, 1; LAPC4 5. Conditions for bicalutamide: Con: 10% FBS treated with vehicles. BL, bicalutamide low dose (μ M): LNCaP, 1; LAPC4, 20. BH, bicalutamide high dose (μ M): LNCaP, 5; LAPC 30. Data shown as percentage to the control group, Mean $\pm$ SD. Eight replicates in each group, three independent experiments. Cells were cultured in 10% CSS medium (**E**) or treated with bicalutamide (**F**), metformin added and lysates blotted for senescence-associated proteins p16. Clearance of senescent cells is represented by decreased p16 expression.

Figure 3. Late ADT induces mTOR1 signaling through activation of AKT and inhibition of AMPK. Metformin addition reactivates AMPK and inhibits mTOR1 signaling. PC cells were cultured in 10% CSS medium (**A**) or bicalutamide (**B**) for 6 days to become senescent followed by addition of metformin, then whole protein lysates harvested for western blot. Bic: bicalutamide (μ M), LNCaP, 5; LAPC4, 30. Met: metformin (mM): LNCaP, 1; LAPC4, 5. **C**. Schema of the underlying mechanism of ADT and metformin in androgen-sensitive prostate cells. Androgen-sensitive PC cells senesce with longer-term exposure of ADT. PI3K/Akt signaling was activated and AMPK signaling was inactivated in senescent cells, phosphorylation of Ser79 on ACC-a downstream protein reflecting AMPK activity was also detected. Both cascades activate the downstream mediator mTOR1 signaling pathway. In addition, anti-apoptosis protein XIAP was upregulated in senescence. All these together potentiate outgrowth of androgen-refractory PC cell subpopulation. Metformin was able to induce cell death in senescence by inhibition of mTOR1 signaling through AMPK activation and Akt inhibition. Charcoal-stripped serum reduced androgen receptor (AR) expression in both LNCaP and LAPC4

cells while bicalutamide effects were more variable. AR expression increased at 48hr when Met is added to ADT compared to Met alone.

Figure 4. XIAP inhibits apoptosis in senescent PC cells. A-C: LNCaP cells were transfected with non-silencing control siRNA (si-Control) or siRNA targeting XIAP (si-XIAP), 24hr later were replaced with fresh media containing either 10% FBS or CSS. **A.** Cell proliferation was measured with DNA assay at 2 and 3 days after switched to CSS media, data shown as Mean $\pm$ SD, * $p < 0.05$. Eight replicates in each group, three independent experiments. **B.** Whole protein lysates were harvested for checking the denoted proteins' expression by western blot. **C.** Apoptosis was measured with caspase-Glo, normalized with cell number. The fold change relative to cells transfected with si-Control and cultured in media containing 10% FBS, data shown as Mean $\pm$ SD, * $p < 0.05$, comparison between CSS si-Control vs CSS si-XIAP. Eight replicates in each group, three independent experiments. **D-E:** XIAP overexpression abrogated metformin induced apoptosis in senescent cells. LNCaP cells were transfected with either pcDNA3-XIAP or pcDNA3 plasmids as control. After stable cell lines have been established, cells were cultured with either Media containing FBS or CSS followed by metformin treatment for 2 days and then cell proliferation (**D**), protein expression (**E**) and apoptosis (**F**) were analyzed as above. Apoptosis was assayed using Caspase-Glo, shown as fold change relative to control, Mean $\pm$ SD value. CSS: 10% CSS; Met: metformin (1 mM). Statistical analysis using one-way ANOVA demonstrates p values between different groups CSS/Met compared to CSS or Met alone (pcDNA control: 0.008, 0.0004, pcDNA-XIAP: 0.1015, 0.006, respectively).

Figure 5. Metformin and ADT suppresses tumor growth in patient-derived xenografts (PDXs) more robustly than ADT alone. A. LuCaP 77, B. LuCaP 147. The growth rates are shown as percentage of tumor size compared to the time point of sham surgery/castration, Median $\pm$ SE. One way Anova was used to analyze the differences between any 2 groups. * $p < 0.01$ is considered as statistically significant between castration plus metformin treated group compared to castration alone, N=8 mice per group. Sham: sham surgery control, Cast: castration, Met: sham surgery with metformin treated, Cast +Met: castration and metformin treated. **C:** Ki67 expression in LuCaP 147 tumors. Expression of Ki67 protein was detected in mouse tissue array constructed from above mouse tumors, with duplicate cores per tissue. VECTRA system was used for image analysis and quantification of the staining intensity. Nuance system and inform 1.2™ software. Data is shown as percentage of Ki67 positive staining cells, Mean $\pm$ SE, one way Anova was used to analyze the differences between any 2 groups. * $p < 0.05$, ** $p < 0.01$. **D.** Images of ki67 staining, the brown color is positive staining, H&E was used as counter-staining. Magnification (20X). The protein expression of key molecules in the LuCaP xenograft tumors are shown for XG147 (**E**) and XG77 (**F**).

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

