

AWARD NUMBER: W81XWH-19-1-0462

TITLE: Targeting the Endotheliopathy of Trauma in Hemorrhagic Shock and Traumatic Brain Injury with Freeze-Dried Platelets

PRINCIPAL INVESTIGATOR: Shibani Pati MD PhD

CONTRACTING ORGANIZATION: University of California, San Francisco, CA

REPORT DATE: September 2021

TYPE OF REPORT: Annual Report

PREPARED FOR: U.S. Army Medical Research and Development Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE					<i>Form Approved OMB No. 0704-0188</i>	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.						
1. REPORT DATE (DD-MM-YYYY) September 2021		2. REPORT TYPE Annual Report			3. DATES COVERED (From - To) 15Aug2020-14Aug2021	
4. TITLE AND SUBTITLE Targeting the Endotheliopathy of Trauma in Hemorrhagic Shock and Traumatic Brain Injury with Freeze-Dried Platelets					5a. CONTRACT NUMBER W81XWH-19-1-0462	
					5b. GRANT NUMBER BA180248	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Shibani Pati MD PhD PI Professor UCSF Alpa Mahuvakar PhD Researcher UCSF					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California San Francisco 513 Parnassus Avenue HSE760 San Francisco, CA 94113					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Development Command Fort Detrick, Maryland 21702-5012					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The goal of this project is to test the therapeutic potential of freeze dried platelets-FDPlts (Thrombosomes)- in disease conditions characterized by 1) inflammation, 2) vascular instability and 3) coagulation disturbances, which are all components of the endotheliopathy of trauma (EOT) (refs). Aside from their hemostatic properties, this proposal aims to also determine the mechanisms of action of the freeze dried platelets (FDPlts (Thrombosomes)) on the EOT in traumatic brain injury (TBI) and shock induced acute lung injury (ALI); all conditions with few if any effective treatment options. We hypothesize that FDPlts (Thrombosomes) will have potent hemostatic properties comparable to fresh platelets and that they will attenuate and mitigate the endotheliopathy of trauma (EOT) in TBI and HS induced ALI. We hypothesize that FDPlts (Thrombosomes) can be used as a stand-alone early therapy to mitigate outcomes in trauma.						
15. SUBJECT TERMS TBI- Traumatic brain Injury HS- Hemorrhagic Shock, ALI- Acute Lung Injury						
16. SECURITY CLASSIFICATION OF: a. REPORT U			17. LIMITATION OF ABSTRACT UU		18. NUMBER OF PAGES 22	
b. ABSTRACT U			c. THIS PAGE U		19a. NAME OF RESPONSIBLE PERSON USAMRMC	
19b. TELEPHONE NUMBER (Include area code)						

TABLE OF CONTENTS

	<u>Page</u>
1. Introduction	4
2. Keywords	4
3. Accomplishments	5-15
4. Impact	15-16
5. Changes/Problems	16-17
6. Products	17-19
7. Participants & Other Collaborating Organizations	19-21
8. Special Reporting Requirements	22
9. Appendices	22

1. INTRODUCTION:

Currently in blood-banking practice in the US, platelets are stored in incubators at 22°C, with gentle agitation for up to 5 days. The main reason for this practice of storage at 22°C is to allow for adequate circulating numbers of platelets post transfusion and to avoid the risk of bacterial contamination. It has been shown that storage of platelets at 22°C for 5 days is associated with a decline in function of the platelets, also known as a storage lesion. One option is for blood banks to store platelets at 4°C, which is currently approved for 3 days of storage; however, diminished function of 4°C platelets has also been reported. Alternatively, a freeze-dried platelet (FDPlts) product can circumvent these challenges by providing hemostasis, prolonging the shelf life of platelets without cold storage and significantly enhancing the utilization and safety of transfused platelet units. FDPlts (Thrombosomes), made by Cellphire Inc., are an infusible freeze-dried platelet-derived hemostatic agent, stabilized with trehalose and polysucrose prior to and during freeze-drying. They can be stored at room temperature with prolonged shelf life (>1 yr), eliminating the need for bacterial testing, and logistically allow for platelet availability in remote and austere conditions.

Characterization studies demonstrate that FDPlts (Thrombosomes) express markers such as P-selectin and phosphatidylserine, hence indicating that they are activated. FDPlts (Thrombosomes) have demonstrated to have potent hemostatic properties. Canines undergoing coronary artery bypass grafting (CABG) treated with fresh platelets or FDPlts (Thrombosomes) showed a dose dependent decrease in blood loss. FDPlts (Thrombosomes) also deliver hemostatic efficacy in uncontrolled arterial bleeding in rats and New Zealand white rabbits (NZWR) with busulfan induced thrombocytopenia. Thus, FDPlts (Thrombosomes) are primed hemostatic agents that can be used towards the treatment of acute uncontrolled hemorrhage in bleeding patients. Safety studies with FDPlts (Thrombosomes) have been performed in several species including non-human primates and humans. No evidence of systemic thrombosis or non-specific thrombosis has been noted, which is a concern when utilizing an activated platelet product. **The goal of this project is to test the therapeutic potential of freeze dried platelets-FDPlts (Thrombosomes)- in disease conditions characterized by 1) inflammation, 2) vascular instability and 3) coagulation disturbances, which are all components of the endotheliopathy of trauma (EOT) (refs).** Aside from their hemostatic properties, this proposal aims to also determine the mechanisms of action of the freeze dried platelets (FDPlts (Thrombosomes)) on the EOT in traumatic brain injury (TBI) and shock induced acute lung injury (ALI); all conditions with few if any effective treatment options.

We hypothesize that FDPlts (Thrombosomes) will have potent hemostatic properties comparable to fresh platelets and that they will attenuate and mitigate the endotheliopathy of trauma (EOT) in TBI and HS induced ALI. We hypothesize that **FDPlts (Thrombosomes) can be used as a stand-alone early therapy** to mitigate outcomes in trauma.

2. KEYWORDS:

Hemorrhagic shock, Freeze-dried platelets, Thrombosomes, Traumatic brain injury, Inflammation, Vascular dysfunction, endotheliopathy of trauma

3. ACCOMPLISHMENTS:

What were the major goals of the project?

Specific Aim 1: Characterize <i>in vitro</i> effects of Thrombosomes	Months	UCSF (Pati)
Major Task 1: Months 1-6: Obtain approval from institutional IACUC and ACURO for HS model.	100% complete	Dr. Pati
<i>Milestone(s) Achieved: ACURO approval for HS and TBI model completed</i>	6	
Major Task 2: FDP-Thrombosomes effects on endothelial permeability and signaling (Aim 1)		Dr. Pati
Subtask 1: Grow PECs (pulmonary endothelial cells) and brain endothelial cells (BECS) to sufficient quantities for <i>in vitro</i> assays Start ECIS and endothelial functional assays of platelet groups	3-6	100% completed
Subtask 2: Complete ECIS assays of platelet groups on all endothelial cells. Conduct Western Blots of endothelial signaling pathways and staining of ECs for junctional and cytoskeletal markers	7-9	20% completed
Subtask 3: Continue and complete Western Blots of endothelial signaling pathways and staining of endothelial cells for junctional markers.	10-12	
<i>Milestone(s) Achieved: Comparison of Platelet groups on EC permeability and PEC and BEC signaling</i>	12	
Major Task 3: Effect of FDP (Thrombosomes) on HS induced ALI- 3 hour model (Aim 2)-		Dr. Pati
Subtask 1: HS induced ALI Model acute three hour study (51 animals)	6-9	100% completed
Subtask 2: HS induced ALI Model acute three hour study (51 animals)	10-12	
Subtask 3: Sectioning of HS induced ALI 3 hour Model (102 animals)	13-15	100% completed
Subtask 4: Tissue analysis 3 hour HS model (102 animals)	16-18	100% completed
<i>Milestone(s) Achieved: Determine efficacy and optimal dose of</i>	18	

<i>FDP in vivo to modulate EOT/ALI in 3 hour model of HS</i>		
Major Task 4 Effect of FDP (Thrombosomes) on HS induced ALI-24 hour model (Aim 2)		Dr. Pati
Subtask 1: HS induced ALI Model 24 hour study (84 animals)	19-24	50% completed
Subtask 2: Sectioning of HS induced ALI 24 hour Model (84 animals)	25-27	
Subtask 3: Tissue analysis 24 hour HS model (84 animals)	28-30	
<i>Milestone(s) Achieved: Determine efficacy and optimal dose of FDP in vivo to modulate EOT/ALI in 24 hour model of HS</i>	27	
Major Task 5 Effect of FDP (Thrombosomes) in TBI (Aim 3)		Dr. Pati
Subtask 1 TBI – optimizing dose of FDP (perform surgeries and collect tissue at 3 day time point) – Total of 80 mice	13-16	100%
Subtask 2 - Tissue analysis (barrier permeability – 80 mice)	16-18	100%
Subtask 3 TBI – optimizing timing of delivery of FDP (perform surgeries and collect tissue at 1 day and 3 day time point) – Total of 70 mice	18-21	
Subtask 4 - Tissue analysis (sectioning and staining – 70 mice)	19-24	100%
Subtask 5 – Setting up behavior tests with control age and strain matched mice	18-24	50% completed
Subtask 6 –Behavior test optimization – data analysis	24-30	
Subtask 7 - TBI – Acute time point surgeries and tissue collection (168 mice)	25-31	
Subtask 8 TBI – Acute time point tissue sectioning, staining, and analysis (168 mice)	31-36	
Subtask 9 TBI – chronic time point surgeries and tissue collection, sectioning, staining (115 mice)	35-42	
Subtask 10 TBI – tissue analysis and behavior analysis for chronic time point mice (115 mice) and overall data analysis	40-48	
<i>Milestone(s) Achieved: Complete studies of FDP effects in TBI.</i>	48	
Major Task 6: Measure Thrombosome effects on endothelial glycocalyx and clot formation		Dr. Pati
Intravital Microscopy of mice (64 animals)	31-36	20% completed
Intravital Microscopy of mice (64 animals)	37-42	
<i>Milestone(s) Achieved: Completion of testing for FDP effects on endothelial glycocalyx and clot formation</i>	36	
<i>Milestone(s) Achieved: Thrombosome production</i>		
Major Task 7: Submit abstracts to meetings and manuscripts. Submit final report to DOD	48	Dr. Pati

What was accomplished under these goals?

The major work completed during this time was as follows:

1. Even under limited working capacity due to COVID19 restrictions, we have completed the acute time point studies with FDPlts in mouse model of hemorrhagic shock. This work was presented at the AAST conference (September 2020) and published in Journal of Trauma and Acute Care (see Appendix).
2. We have started in vitro endothelial signaling pathway analysis using FDPlts (Thrombosomes) and fresh platelets as controls.
3. We have completed traumatic brain injury (TBI) experiments with FDPlts (Thrombosomes) and fresh platelets as controls for barrier permeability analysis.
4. Due to space constraints, the behavior room was reestablished at UCSF. We have all the equipment now set up in the behavior room in the barrier facility.

Methods:

Cell Culture

Human umbilical cord derived endothelial cells (HUVECs) and brain derived endothelial cells (BECs) were grown under standard tissue culture conditions in 5% CO₂.

Human platelets and freeze-dried platelets (FDPlts)

Units of leukoreduced apheresis platelets stored in plasma were obtained from Bonfils Blood Center, Denver, CO on Day 1. All platelets were tested for bacterial infection by the Bonfils Blood Bank and Blood Centers of the Pacific) and found to be negative. Freeze-dried platelets (FDPlts) were obtained from Cellphire, Inc.

Transendothelial electric resistance (TEER) The integrity of HUVEC and BEC monolayers was measured using an electric cell-substrate impedance sensing system (ECIS 1600, Applied BioPhysics, Troy, NY). An increase or decline in TEER across the cell monolayers indicated accordingly, decreased or increased endothelial paracellular permeability. Cells were grown to confluence on L-cysteine reduced, Fibronectin-pre-coated 96-well plates containing electrodes in each well. Cells were treated with platelets or FDPlts ($50 \times 10^3/\mu\text{l}$ in 200 μL EBM-2 medium) and then challenged with thrombin. Monolayer resistance at 4/16/64 kHz was analyzed in 5-min intervals. Data were normalized to the mean resistance of cell monolayers before the treatments and plotted as area under the curve. 4- wells were tested/group. Statistical analysis conducted was by standard Student t test and one way ANOVA for comparison between groups.

Immunostaining of endothelial tight and adherens junction proteins

HUVECs were grown to confluence on collagen coated coverslips prior to treatment. FDPlts were reconstituted in pyrogen free, sterile water and added to the HUVEC monolayer at a concentration of 50×10^6 particles/ml for 30 minutes, followed by a 0.2U/ml thrombin challenge for 5 minutes at 37°C. Plts were added in the same concentrations. Junction proteins VE-cadherin (Cell Signaling, Danvers, MA), and zonula occludens-1 (ZO-1) (Invitrogen, Carlsbad, CA) were stained. F-actin was detected with Texas Red Phalloidin (Cell Signaling). Three coverslips per treatment group were prepared, and four images were captured per coverslip.

Images were captured at 20x magnification using the Nikon Eclipse 80i microscope (Nikon, Melville, NY) with SPOT RT-sCMOS camera (SPOT Imaging, Sterling Heights, MI). Gap junction measurements were performed using ImageJ, an open source platform for biological image analysis.

2-Dimensional Difference in Gel Electrophoresis (2D-DIGE) and Mass Spectrometry

Thrombosomes and fresh platelets (Day 1 plts) were sent to Applied Biomics (Hayward, California) for 2-Dimensional Difference in Gel Electrophoresis (2D-DIGE) and mass spectrometry analysis. Samples were matched as pairs, and labeled with CyDye2 or CyDye5, to enable simultaneous co-separation and analysis of samples on a single multiplexed gel. 2D gel electrophoresis was used to separate the labeled samples on a single 2D gel using isoelectric focusing and SDS polyacrylamide gel electrophoresis (SDS-PAGE). ImageQuant and DeCyder computer software was used to quantify the difference in protein ratios between samples in order to identify protein differences between the matched pair. The EttanTM spot picker (GE Healthcare Life Systems, >99.9% accuracy) was then used to identify spots that represent proteins for which there existed significant differences. A p-value of <0.05 was used to identify proteins of interest. Selected spots were then identified using spectrometry matrix-assisted laser desorption/ionization time-of-flight/time-of-flight (MALDI-TOF/TOF). Extracted peptides were spotted into wells of a MALDI plate, and mass spectrometry was performed using Applied Biomics Proteomics Analyzer. The 10-20 most abundant peptides were then further fragmented and tandem mass spectrometry was performed to identify the proteins. Specific protein identification was performed by matching peptide mass mapping and peptide fragmentation mapping in primary sequence databases.

Scanning Electron Microscopy

FDPlts were reconstituted then washed and fixed in 3% glutaraldehyde, followed by serial dehydration in 50%, 70%, 90%, and 100% ethanol, and mounted onto ITO CTD coverslips (SPI, NY), then coated with gold-palladium (3nm). FDPlts were examined and imaged using an SEM (S4300, Hitachi, Tokyo, Japan) with an electron energy of 1.0–1.5 keV, at 1000x and 10,000x magnification.

Western Blot Analysis

Western blot experiments were carried out as previously described (Gibb, et al, 2015). Briefly, an equal number of platelets and FDPlts were collected, and lysed with RIPA lysis buffer. Resulting lysates were denatured with NuPage LDS sample buffer (Invitrogen, NY), and separated by SDS-PAGE, transferred to polyvinylidene difluoride membranes, and blotted with anti-tubulin (T6199) and anti-acetylated tubulin (T7451), (Sigma, St. Louis, MO) and IR dye-conjugated secondary antibodies (Licor, NY). Protein was detected and intensity quantitated using the Odyssey imaging system (Licor, NY).

Flow Cytometry

FDPlts were characterized by size and expression of human integrin- α 2b (CD41) (Biolegend, San Diego, CA) and glycoprotein1b (CD42b) (Biolegend, San Diego, CA). Surface levels of P-selectin (CD62P) (Biolegend, San Diego, CA) were compared using the geometric mean fluorescence intensity (gMFI) of expression. All events were analyzed using computer software (FlowJo, Version 9.7, Tree Star, Inc., Ashland, OR).

Protein Analysis - Luminex Assay

A Luminex panel targeting the proteins angiogenin, angiopoietin-1, BDNF, CD40 ligand, FGF, IFN- γ , IL-1 β , IL-6, IL-8, IL-10, PDGF-AA, PDGF-BB, thrombomodulin, thrombospondin-2, Tie-2, TNF- α , VEGF, and vWF-A2 was created (R&D Systems, Minneapolis, MN). FDPlts and day one platelets reconstituted and pelleted and lysed with RIPA buffer containing HALT, and diluted using diluents supplied by the manufacturer. This multiple analyte assay was performed using a MAGPIX instrument, and the resulting data were analyzed using xPONENT software.

Animal Protocols

The animal studies were performed with approval of the Institutional Animal Care and Use Committee at UCSF. The experiments were conducted in compliance with the National Institutes of Health guidelines on the use of laboratory animals and the Department of Defense Animal Care and Use

Review Office. All animals were housed in a room with access to food and water ad libitum, controlled temperature and 12:12-hour light-dark cycles.

Traumatic Brain Injury (TBI) model

The controlled cortical impact model (CCI) of traumatic brain injury (TBI) was performed on 10-12 week old C57Bl/6J mice (Jackson Laboratories, Sacramento, CA). Briefly, isoflurane-anesthetized mice received a 5mm craniotomy and an impact was made using Leica Impactor 1 on the right parietal cortex, with the following setting to generate a moderate-severe injury level. A 3mm piston impacted the cortex at a velocity of 4.5 mm/s, to a depth of 1.2 mm, with a dwell time of 300 ms. Animals were injected with saline or FDPIts (1.5×10^6 particles/g body weight) at two hours and twenty four hours post-injury. Blood brain barrier permeability was evaluated at 3 days post-injury. An hour before sacrifice, animals were retro-orbitally injected with IR dye conjugated-10 kD dextran and allowed to circulate. At the end of the hour circulation, mice were transcardially perfused with ice cold PBS and brains were then removed and sliced at 2mm thickness. Dye intensity was quantitated using the Odyssey imaging system (Licor, NY). Brain slices were then fixed with 4% PFA overnight and dehydrated for 3-5 days in 30% sucrose solution then embedded in OCT. Embedded brains were further sectioned to 20 μ m thickness using a Leica 1950 cryostat (Leica Biosystems, Wetzlar, Germany).

Statistical analyses and power calculation

Statistical analyses were performed using Prism 8.3 software (Graphpad Inc. San Diego, CA), multiple group comparisons were determined by One way ANOVA with post-hoc Tukey tests for all in vitro experiments. In vivo experiments were analyzed by RM two way aANOVA. An α of 0.05 was preset as the cutoff for statistical significance. All data are represented as mean $\pm$ SD.

Results:




Figure 1: In vitro characterization of FDPIts:

A) SEM imaging shows that FDPIts have activated morphology and presence of exposed extracellular vesicles as compared to day 1 platelets. B) Flow cytometry analysis demonstrates an increased expression of CD62p on FDPIts as compared to day 1 fresh platelets. C) Deacetylation of tubulin is a marker of platelet activation. Here we see increased deacetylation on freeze dried platelets (across multiple lots) as compared to day 1 platelets.

	Plt	FDPlt			
		CT1	CT2	CT3	NT
Ang1	88038.4	98771.6	104446.2	103060.5	142533.9
Ang2	N/A	667.0	724.5	731.7	968.5
BDNF	37233.7	94509.9	98948.5	96882.0	104522.4
vWF-A2	418.9	567.5	682.4	657.8	927.6
Thrombo Modulin	3647.5	6679.0	7407.0	7004.8	8306.7
Thrombo spondin2	1179.4	4065.0	4421.1	4131.8	4020.5
CD40	14745.6	16458.2	17447.9	16880.1	19359.9
PDGF-AA	6185.1	14500.5	16488.8	15859.1	15265.3
PDGF-BB	30314.3	83742.4	99735.0	99660.0	99831.4
VEGF	315.7	352.2	399.1	370.3	627.9
EGF	36.0	236.4	328.5	251.6	351.3
FGF	250.5	75.7	95.3	93.3	102.6
TNF α	23.1	24.4	28.5	25.8	29.9
IFN γ	619.4	329.7	395.0	385.7	610.1
IL-1 β	53.0	38.9	40.7	42.5	55.7
IL-6	34.5	23.5	25.3	26.2	36.4
IL-10	5.5	7.8	12.3	10.9	12.0
CXCL8	33.8	52.3	65.1	62.1	63.6
Angiogenin	7778.2	18194.0	19958.8	19014.2	19429.2
Tie2	660.7	832.5	919.8	882.2	907.2

Table 1: Factor content in the Thrombosomes is preserved irrespective of the freeze drying process and change in morphology. More importantly, they are enriched in potent vascular regulators and growth factors, as seen in the graphical representation below.





Figure 2: FDPIts show dose dependent mitigation of thrombin induced endothelial cell permeability. A) Three doses ($10 \times 10^6/\text{ml}$, $50 \times 10^6/\text{ml}$, and $100 \times 10^6/\text{ml}$) of FDPIts were tested in the presence and absence of thrombin challenge on human brain microvascular endothelial cells in ECIS. Top panels show ECIS tracings and bottom panels show quantitation. FDPIts show a dose dependent protection on endothelial cell permeability. B) Fresh apheresis platelets ($50 \times 10^6/\text{ml}$) from 3 different donors and pooled donors were tested in the presence and absence of thrombin challenge on human brain microvascular endothelial cells in ECIS. Donor dependent variability in EC protection is observed, furthermore, FDPIts are more protective than either single donor or pooled donor fresh Plts. One-way ANOVA with Bonferroni post-hoc analysis. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ vs thrombin or control.






Figure 3: Consistent mitigation of thrombin induced endothelial cell permeability across multiple FDPIs lots. Four different lots of FDPIs were tested in the presence and absence of thrombin challenge on human brain microvascular endothelial cells in ECIS. Top panels show ECIS tracings and bottom panels show quantitation. FDPIs show protection on endothelial cell permeability across multiple lots. One-way ANOVA with Bonferroni post-hoc analysis, **** $p < 0.0001$.





Figure 4: Traumatic brain injury induced vascular permeability at 3 days post-injury. Mice were subjected to controlled cortical impact injury or sham surgery (anesthesia and placement on stereotactic device with incision). Permeability was measured by using a 10kD IR-680 tagged dextran. ID- Ipsilateral Dorsal, IV – Ipsilateral Ventral, CD- Contralateral Dorsal, CV- Contralateral Ventral. Repeated measures two-way ANOVA showing a significant effect of region (ID>all other regions) and sham vs TBI in ID region. **** $p < 0.0001$. $n = 5-6$ per group

Figure 5: Treatment with FDPlts Attenuates Barrier Disruption at 3 days Post-injury

Mice were subjected to TBI and treated with either saline or FDPlts at 2h and 24 h post-injury and barrier permeability was measured at 3 days post-injury using a 10kD IR-680 tagged dextran. Both the saline and FDPlts treated groups show focal increased barrier permeability in the ipsilateral dorsal quadrant as compared to the other three quadrants and the FDPlts treated mice have significantly less disruption as compared to the saline treated mice. Black bars = saline treatment; grey bars = FDPlts treatment; ID- Ipsilateral Dorsal, IV – Ipsilateral Ventral, CD- Contralateral Dorsal, CV- Contralateral Ventral. Repeated measures two-way ANOVA showing a significant effect of region (ID>all other regions) and saline vs. FDPlts in ID region, $p<0.0001$.



What opportunities for training and professional development has the project provided?

Nothing to Report

How were the results disseminated to communities of interest?

This work on role of FDPIts in mouse model of hemorrhagic shock was presented at the AAST conference (September 2020) and was published in Journal of Trauma and Acute Care (see Appendix).

What do you plan to do during the next reporting period to accomplish the goals?

- We will continue the cell culture experiments to examine the signaling cascade of endothelial cell and platelet and freeze dried platelet cross talk (Major task 2, subtasks 2 and 3).
- Start behavior study set up run of mice (Major task 5, subtask 6)
- Start long-term studies in mouse hemorrhagic shock model (Major task 4)
- Continue studies in traumatic brain injury (Major task 5)
- Continue intravital microscopy studies (Major task 6)

4. IMPACT:

What was the impact on the development of the principal discipline(s) of the project?

The protein content of FDPPlts is similar to fresh platelets, suggesting that the freeze drying process does not affect the protein content. FDPPlts attenuated vascular permeability in vitro in brain microvascular endothelial cells and showed lot to lot consistency unlike fresh platelets that had donor-to-donor variability. FDPPlts also attenuated blood brain barrier permeability in a mouse model of TBI. These data suggest that the effect of FDPPlts may be a suitable alternative to fresh Plts in modulating endotheliopathy associated with injury.

What was the impact on other disciplines?

This finding could be of great utility for storage and availability of platelets in remote and austere environments and also be a safer alternative from an infectious standpoint of bacterial contamination. Eventually this could change Blood Banking practice in the US and military settings.

What was the impact on technology transfer?

Nothing to report.

What was the impact on society beyond science and technology?

Nothing to Report

5. CHANGES/PROBLEMS:

Nothing to Report

Actual or anticipated problems or delays and actions or plans to resolve them

1. Due to space constraints, the behavior room was reestablished at UCSF. We have set up behavior room in the barrier facility with new equipment and will now set up all tests.

Changes that had a significant impact on expenditures

Nothing to Report

Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents

Significant changes in use or care of human subjects

Not applicable

Significant changes in use or care of vertebrate animals

None

Significant changes in use of biohazards and/or select agents

None

6. PRODUCTS:

- **Publications, conference papers, and presentations**

Journal publications.

Our first manuscript was published in Journal of Trauma and Acute Care
(Published article is attached in the appendix).

Books or other non-periodical, one-time publications.

Nothing to Report

Other publications, conference papers and presentations.

Nothing to report

- **Website(s) or other Internet site(s)**

Nothing to Report

- **Technologies or techniques**

Nothing to Report

- **Inventions, patent applications, and/or licenses**

Nothing to Report

- **Other Products**

Nothing to Report

7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

What individuals have worked on the project?

PATI, SHIBANI - 2.88

ZHANG,HAOQIAN - 3.92

MIYAZAWA,BYRON YUUJI – 4.39

MAHUVAKAR,ALPA – 4.46

Name: Shibani Pati MD PhD

Project Role: Principal Investigator

Nearest person month worked: 2.88 cal months

Contribution to Project: Dr.Pati is the principal investigator on this grant and has been overseeing the planning, data analysis and execution of the entire grant.

Name: Alpa Mahuvakar PhD.

Project Role: Research Scientist

Nearest person month worked: 4.46 cal months

Contribution to Project: Dr. Mahuvakar is a co-investigator and is involved in planning of studies, analysis of data, running/execution of all research experiments, including in vitro endothelial cell signaling and traumatic brain injury.

Name: Byron Miyazawa

Project Role: Research Associate

Nearest person month worked: 4.39 cal months

Contribution to Project: Running/execution of platelet assays, endothelial assays, staining of endothelium and signaling analysis

Name: Haoqian Zhang PhD.

Project Role: Research Scientist

Nearest person month worked: 3.92 cal months

Contribution to Project: In vitro blood brain barrier organoid assays and intravital imaging

Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?

None

What other organizations were involved as partners?

Cellphire Inc. Rockville MD
Cellphire provided the Thrombosome product needed for these studies.

8. SPECIAL REPORTING REQUIREMENTS

COLLABORATIVE AWARDS:

QUAD CHARTS:

9. APPENDICES:



Freeze_dried_platelet
s_promote_clot_forma