



**US Army Corps  
of Engineers®**  
Engineer Research and  
Development Center



*Dredging Operations Technical Support Program*

## **Birds of the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020**

Michael P. Guilfoyle, Ruth A. Beck, Bill Williams,  
Shannon J. Reinheimer, Lyle D. Burgoon, Sam S. Jackson,  
Sherwin M. Beck, Burton C. Suedel, and Richard A. Fischer

September 2022



**The U.S. Army Engineer Research and Development Center (ERDC)** solves the nation's toughest engineering and environmental challenges. ERDC develops innovative solutions in civil and military engineering, geospatial sciences, water resources, and environmental sciences for the Army, the Department of Defense, civilian agencies, and our nation's public good. Find out more at [www.erdclibrary.on.worldcat.org/discovery](http://www.erdclibrary.on.worldcat.org/discovery).

To search for other technical reports published by ERDC, visit the ERDC online library at [www.erdclibrary.on.worldcat.org/discovery](http://www.erdclibrary.on.worldcat.org/discovery).

The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.

The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

# **Birds of the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020**

Michael P. Guilfoyle, Lyle D. Burgoon, Sam S. Jackson,  
Burton C. Suedel, and Richard A. Fischer

*Environmental Laboratory  
US Army Engineer Research and Development Center  
3909 Halls Ferry Road  
Vicksburg, MS 39180-6199*

Ruth A. Beck

*Department of Biology  
College of William and Mary  
Williamsburg, Virginia 23185*

Bill Williams

*Coastal Virginia Wildlife Observatory, Waterbird Team  
154 Lakewood Drive  
Williamsburg, Virginia 23185*

Shannon J. Reinheimer

*US Army Corps of Engineers  
Norfolk District  
803 Front Street  
Norfolk, Virginia 23510*

Sherwin M. Beck

*P.O. Box 270  
Barhamsville, Virginia 23011*

Final report

Approved for public release; distribution is unlimited.

Prepared for US Army Corps of Engineers  
Washington, DC 20314-1000

Under Funding Account Code U4372823  
AMSCO Code 086000

## Abstract

This report presents the results of a long-term trend analyses of seasonal bird community data from a monitoring effort conducted on the Craney Island Dredged Material Management Area (CIDMMA) from 2008 to 2020, Portsmouth, VA. The USACE Richmond District collaborated with the College of William and Mary and the Coastal Virginia Wildlife Observatory, Waterbird Team, to conduct year-round semimonthly area counts of the CIDMMA to examine species presence and population changes overtime. This effort provides information on the importance of the area to numerous bird species and bird species' groups and provides an index to those species and group showing significant changes in populations during the monitoring period. We identified those species regionally identified as Highest, High, and Moderate Priority Species based on their status as rare, sensitive, or in need of conservation attention as identified by the Atlantic Coast Joint Venture (ACJV), Bird Conservation Region (BCR), New England/Mid-Atlantic Bird Conservation Area (BCR 30). Of 134 ranked priority species in the region, the CIDMMA supported 102 of 134 (76%) recognized in the BCR, including 16 of 19 (84%) of Highest priority ranked species, 47 of 60 (78.3%) of High priority species, and 39 of 55 (71%) of Moderate priority species for BCR 30. All bird count and species richness data collected were fitted to a negative binomial (mean abundance) or Poisson distribution (mean species richness) and a total of 271 species and over 1.5 million birds were detected during the monitoring period. Most all bird species and species groups showed stable or increasing trends during the monitoring period. These results indicate that the CIDMMA is an important site that supports numerous avian species of local and regional conservation concern throughout the year.

**DISCLAIMER:** The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products. All product names and trademarks cited are the property of their respective owners. The findings of this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

**DESTROY THIS REPOR WHEN NO LONGER NEEDED. DO NOT RETURN IT TO THE ORIGINATOR.**

# Contents

|                                                                                                                                                                                                           |             |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Abstract.....</b>                                                                                                                                                                                      | <b>ii</b>   |
| <b>Figures and Tables.....</b>                                                                                                                                                                            | <b>iv</b>   |
| <b>Preface .....</b>                                                                                                                                                                                      | <b>vii</b>  |
| <b>Executive Summary .....</b>                                                                                                                                                                            | <b>viii</b> |
| <b>1 Introduction .....</b>                                                                                                                                                                               | <b>1</b>    |
| 1.1 Background .....                                                                                                                                                                                      | 1           |
| 1.2 National initiatives.....                                                                                                                                                                             | 3           |
| 1.2.1 <i>The North American Bird Conservation Initiative</i> .....                                                                                                                                        | 3           |
| 1.2.2 <i>Craney Island Long-term Bird Management Plan</i> .....                                                                                                                                           | 4           |
| 1.3 Objectives.....                                                                                                                                                                                       | 10          |
| 1.4 Approach .....                                                                                                                                                                                        | 11          |
| 1.5 Scope.....                                                                                                                                                                                            | 11          |
| <b>2 Study Area and Methods.....</b>                                                                                                                                                                      | <b>13</b>   |
| 2.1 Study area .....                                                                                                                                                                                      | 13          |
| 2.2 Seasonal bird community monitoring .....                                                                                                                                                              | 15          |
| 2.3 Initial mean abundance and species richness trends .....                                                                                                                                              | 16          |
| 2.3.1 <i>Mean abundance trends</i> .....                                                                                                                                                                  | 16          |
| 2.3.2 <i>Mean species richness trends</i> .....                                                                                                                                                           | 18          |
| <b>3 Results.....</b>                                                                                                                                                                                     | <b>19</b>   |
| 3.1 Seasonal survey monitoring results .....                                                                                                                                                              | 19          |
| 3.2 Examples of priority species .....                                                                                                                                                                    | 22          |
| 3.2.1 <i>Highest priority species</i> .....                                                                                                                                                               | 22          |
| 3.2.2 <i>High priority species</i> .....                                                                                                                                                                  | 23          |
| 3.2.3 <i>Moderate priority species</i> .....                                                                                                                                                              | 25          |
| 3.3 Annual trends for species and species groups.....                                                                                                                                                     | 26          |
| 3.4 Species richness.....                                                                                                                                                                                 | 44          |
| <b>4 Discussion.....</b>                                                                                                                                                                                  | <b>46</b>   |
| <b>References.....</b>                                                                                                                                                                                    | <b>49</b>   |
| <b>Acronyms and Abbreviations.....</b>                                                                                                                                                                    | <b>53</b>   |
| <b>Unit Conversion Factors.....</b>                                                                                                                                                                       | <b>54</b>   |
| <b>Appendix A: Table of Counts of Birds Detected During Seasonal Surveys on<br/>the Craney Island Dredged Material Management Area, USACE Norfolk<br/>District, Portsmouth, Virginia, 2008-2020 .....</b> | <b>55</b>   |
| <b>Report Documentation Page</b>                                                                                                                                                                          |             |

# Figures and Tables

## Figures

|                                                                                                                                                                                                                                                                                                                                                                 |    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 1-1. A Black-necked Stilt chick on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. This is one of many nesting species monitored and protected on the facility. (Photo credit: USACE Norfolk District). ....                                                                                                                   | 5  |
| Figure 1-2. Signage on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, indicating an area closed to protect nesting birds. (Photo credit: S. J. Reinheimer, USACE Norfolk District). ....                                                                                                                                             | 6  |
| Figure 1-3. On-site construction on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. Efforts are made to ensure construction does not disturb nesting birds. (Photo credit: S. J. Reinheimer, USACE Norfolk District). ....                                                                                                            | 7  |
| Figure 1-4. Pond and mudflat habitats can be created, and water levels managed to provide habitat for migratory shorebirds and wintering waterfowl on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. (Photo credit: S. J. Reinheimer, USACE Norfolk District). ....                                                                  | 8  |
| Figure 1-5. Figure showing the North, Center, and South Cells on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. Portsmouth, Virginia. (Photo credit: USACE Norfolk District). ....                                                                                                                                                   | 10 |
| Figure 2-1. Overview of the Craney Island Dredged Material Management Area, Portsmouth, Virginia. ....                                                                                                                                                                                                                                                          | 14 |
| Figure 3-1. The American Oystercatcher is a species documented on the Craney Island Dredged Material Management Area that has been ranked as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservations Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Rhododentries, Wikimedia Commons). ....                                    | 22 |
| Figure 3-2. The Ruddy Turnstone is a wintering species documented on the Craney Island Dredged Material Management Area that has been ranked as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservations Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Kate Perez, Wikimedia Commons). ....                                    | 22 |
| Figure 3-3. The Piping Plover is a federally listed endangered species that has been documented on the Craney Island Dredged Material Management Area and has been listed as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008). (Photo Credit: Shutterstock.com, Wikimedia Commons). .... | 23 |
| Figure 3-4. The Black-bellied Plover is a migratory species documented on the Craney Island Dredged Material Management Area that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture) (Photo Credit: Public Domain, Wikimedia Commons). ....                                     | 23 |
| Figure 3-5. The Northern Bobwhite is a species documented on the Craney Island Dredged Material Management Area that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Uatrok77, Wikimedia Commons). ....                                                  | 24 |

|                                                                                                                                                                                                                                                                                                                                       |    |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 3-6. The Semipalmated Sandpiper is a migratory species documented on the Craney Island Dredged Material Management Areas that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Venture 2008) (Photo Credit: USFWS Northeast Region, Wikimedia Commons)..... | 24 |
| Figure 3-7. The Royal Tern is a species documented on the Craney Island Dredged Material Management Areas that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Region Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: defaulder, Wikimedia Commons).....                   | 25 |
| Figure 3-8. The Red-necked Phalarope is a migratory species documented on the Craney Island Dredged Material Management Areas that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: USFWS, Wikimedia Commons).....          | 25 |
| Figure 3-9. The Ruddy Duck is a wintering species documented on the Craney Island Dredged Material Management Area that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Public Domain, Wikimedia Commons).....             | 26 |
| Figure 3-10. Seasonal mean abundance trends for Blackbirds, Crows and Jays on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....                                                                                                                                                               | 27 |
| Figure 3-11. Seasonal mean abundance trends for Dabbling Ducks on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                                            | 28 |
| Figure 3-12. Seasonal mean abundance trends for Diving and Sea Ducks on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                                      | 29 |
| Figure 3-13. Seasonal mean abundance trends for Flycatchers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....                                                                                                                                                                              | 30 |
| Figure 3-14. Seasonal mean abundance trends for Geese and Swans on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                                           | 31 |
| Figure 3-15. Seasonal mean abundance trends for Grassland Birds and Sparrows on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                              | 32 |
| Figure 3-16. Seasonal mean abundance trends for Gulls, Terns and Skimmers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....                                                                                                                                                                | 33 |
| Figure 3-17. Seasonal mean abundance trends for Herons and Egrets on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                                         | 34 |
| Figure 3-18. Seasonal mean abundance trends for Other Nonpasserines on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.....                                                                                                                                                                       | 35 |

|                                                                                                                                                                      |    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 3-19. Seasonal mean abundance trends for Other Passerines on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....        | 36 |
| Figure 3-20. Seasonal mean abundance trends for Pelicans and Cormorants on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. .... | 37 |
| Figure 3-21. Seasonal mean abundance trends for Raptors and Vultures on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....    | 38 |
| Figure 3-22. Seasonal mean abundance trends for Sandpipers and Plovers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....  | 39 |
| Figure 3-23. Seasonal mean abundance trends for Shorebirds on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....              | 40 |
| Figure 3-24. Seasonal mean abundance trends for Swallows and Swifts on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....     | 41 |
| Figure 3-25. Seasonal mean abundance trends for Thrushes on the Craney Island Dredged Material Management Areas, Portsmouth, Virginia, 2008-2020. ....               | 42 |
| Figure 3-26. Seasonal mean abundance trends for Warblers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....                | 43 |
| Figure 3-27. Seasonal mean abundance trends for Woodpeckers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....             | 44 |
| Figure 3-28. Seasonal mean species richness for avian species groups on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020. ....    | 45 |

## Tables

|                                                                                                                                                                                                                                                                                                                                                                     |    |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Table 2-1. Total number of survey days, and total survey hours for all birds detected on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008 to 2020. ....                                                                                                                                                                               | 16 |
| Table 3-1. Total cumulative detections of the New England/Mid-Atlantic Coast Bird Conservation Region (BCR 30) Highest Priority, High Priority, and Moderate Priority Species of regional or local conservation concern (from Atlantic Coast Venture 2008) detected on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008 to 2020. .... | 20 |

## Preface

This technical report (TR) was funded by the Dredging Operations Technical Support (DOTS) program, Dr. Burton Suedel, Program Manager; Funding Account Code U4372823, AMSCO Code 086000. Portions of the fieldwork were funded by the US Army Corps of Engineers, Norfolk District. This TR was completed by the Ecological Resources Branch (ERB) of the Ecosystem Evaluation and Engineering Division of the US Army Engineer Research and Development Center - Environmental Laboratory (ERDC-EL-EEE). At the time of publication, Mr. Joey Minter was Chief, ERB; and Dr. Mark Farr was Division Chief, EEE. The Deputy Director of ERDC-EL was Dr. Brandon Lafferty, and the Director was Dr. Edmond J. Russo, Jr.

COL Christian Patterson was Commander of ERDC, and Dr. David W. Pittman was the Director of ERDC.

This work represents more than 2,600 hours of observations by Ruth A. Beck, Bill Williams and the Waterbird Team of the Coastal Virginia Wildlife Observatory and others on 403 days in the field from May 2008 to August 2020. This report is dedicated to the life and career of Ruth A. Beck, who passed in 2015. During her tenure as a professor at the College of William and Mary, Williamsburg, VA, she, and her graduate students dedicated much of their time and effort to monitor and study the bird communities on the Craney Island Dredged Material Area (CIDMMA). It is due to her efforts that these data have been collected and made available for this report. Since Ruth's May 2015 passing, CIDMMA has been monitored weekly (weather permitting) by members of the Coastal Virginia Wildlife Observatory's Waterbird Team at their own expense. Those team members include Bill Williams (Team Leader), Robert Ake, Edward S. Brinkley, Andrew Hawkins, Alex Minarik, Lee Schuster, Brian Taber, and David Youker.

## Executive Summary

This study compiled and analyzed data from regularly conducted bird community surveys performed largely by the Coastal Virginia Wildlife Observatory's Waterbird Team from May 2008 to August 2020 on the Craney Island Dredged Material Management Area (CIDMMA), Portsmouth, Virginia. Preliminary statistical analyses of population trend results are provided from seasonal monitoring of the bird communities using the CIDMMA. The Craney Island facility is a confined disposal facility (CDF), located at the confluence of the James and Elizabeth Rivers, near Hampton Roads, VA. This facility is approximately 2,500 acres (1,011 ha) and receives dredged material from US Army Corps of Engineers (USACE) maintenance dredging operations and new work in the Norfolk Harbor and adjacent waters. These maintenance operations stem from the construction and maintenance actions of private interests, municipalities, and government agencies on the federal navigation channels and overall maintenance and improvement for the US Navy and the Virginia Port Authority facilities. The Craney Island facility possesses approximately 8 miles (12.8 km) of primary containment dikes and 6 miles (9.7 km) of shoreline consisting of beaches, marshes, and hardened sections. The area is accessible through a perimeter road where the containment dikes meet the mainland.

To extend the life of the facility, the area was divided into three large subcontainment cells, where each alternatively receives dredged material on an annual basis. The dredged material is disposed into the cells from the eastside of the facility by use of hydraulic pipelines from a cutterhead dredge. The dredged material is in a slurry form as deposited and moves from the east to the west side of a cell, where the sediment separates from fine-grained to coarse-grained material. As a result, large sandy berms form on sections of the cell, and lower portions of the cell maintain a shallow water depth with associated mud and sandy flats. In cells not receiving dredged material, heavy equipment is used to repurpose the dredged material for building and enhancing the perimeter containment dikes to increase capacity of a cell to receive continued dredged material for subsequent years.

It was soon recognized that the various open sandy, wetland, and mudflat habitats in the facility were attracting large numbers and species of coastal birds for use as foraging, roosting, and nesting sites. Spring and summer

bird communities were surveyed, and nesting birds were monitored during the nesting season from 1975 to the present. In 2012, a Long-Term Bird Management Plan (LTBMP) was drafted for implementation; however, adoption of the plan has not been formalized by the USACE Norfolk District. Many of the specific elements in the LTBMP have been ongoing since before the plan was written and continue to be implemented to this day. Specific elements of the plan include (1) continued regularly performed monitoring of the four seasons' (winter, spring, summer, and fall) bird communities occurring at least semimonthly, (2) monitoring of dredged material deposition operations and construction actions to ensure that nesting birds are not disturbed nor that eggs or nests are lost during these actions, (3) monitoring and management of water levels to ensure proper water depths for migrating shorebirds and overwintering waterfowl, (4) when necessary, predator control and management actions may be employed to promote successful nesting by birds, and (5) management and monitoring of road maintenance, and onsite mowing actions to ensure minimal or no birds or nests are disturbed or lost.

In this report, monitoring efforts are documented to have detected 271 species, including species with documented counts suitable for initial trend analyses. Total counts of these 271 species were greater than 1.5 million birds over the monitoring period. We focused results on those species regionally identified as Highest, High, and Moderate Priority Species based on their status as rare, sensitive, or in need of conservation attention as identified by the Atlantic Coast Joint Venture (ACJV), Bird Conservation Region (BCR), New England/Mid-Atlantic Bird Conservation Area (BCR 30). Of 134 ranked priority species in the region, the CIDMMA supported 102 of 134 (76%) recognized in the BCR, including 16 of 19 (84%) of Highest priority ranked species, 47 of 60 (78.3%) of High priority species, and 39 of 55 (71%) of Moderate priority species for BCR 30. These results indicate that the CIDMMA is an important site that supports numerous avian species of local and regional conservation concern throughout the year.

In addition, we provide preliminary results of seasonal mean abundance and species richness trend analyses. A more formal analyses for a future report or manuscript is planned. From this analyses, most all individual species showed stable non-significantly increasing or decreasing mean abundance trends. Increases in all seasons for the American Avocet, and for the Ring-billed Gull during the winter and spring are noticeable

exceptions. Similarly, significant decreases in mean abundance trends documented for the Black Tern and Least Sandpiper during the fall, and *Calidris* spp. during the winter are also exceptions. Mean species richness increased significantly during the monitoring period for most groups, except decreases for shorebirds during the winter and thrush species during the fall; no changes occurred for woodpeckers during any season. In general, these results show that the Craney Island facility is important for attracting and maintaining stable seasonal populations for a wide diversity of local and regional avian species, including many Highest, High, and Moderate Priority Species in the New England/Mid-Atlantic Bird Conservation Region.

*The contents of this report are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such commercial products.*

# 1 Introduction

## 1.1 Background

The US Army Corps of Engineers (USACE) is responsible for maintaining our coastal infrastructure, including ports, harbors, and shoreline stabilization features; and for maintaining the Intracoastal Waterway System (ICWW) (Environmental Protection Agency [EPA]/USACE 2007, Guilfoyle et al. 2019). Dredging operations are used regularly to maintain depths of navigation channel, ports and harbors sufficient to support international and national commerce. Natural coastal sedimentation processes are altered by maintenance dredging and shoreline armoring, both of which may reduce natural formation of coastal habitats, including beaches, sand spits, mud flats, marshes, and other wetland habitats (Guilfoyle et al. 2019). These changes can reduce or degrade important coastal habitat for local and regional floral and faunal species. In addition, maintenance dredging operations can yield significant amounts of sediment material that must be disposed locally, or at nearshore or offshore deposition sites. Dredged material from ports, harbors, and navigation channels can be deposited in confined disposal facilities (CDFs) that can also be managed to support seasonal bird communities (Guilfoyle et al. 2020a, b). Moreover, dredged material from ports and harbors in large urban areas may also be contaminated by urban and industrial pollutants, including heavy metals and other toxic chemicals used in industrial processes. Deposition of contaminated dredged material in CDFs is often required to reduce exposure to the human and wildlife populations and are designed to minimize contaminated discharge into the local environment (USACE and the EPA 2003).

Beneficial uses of dredged material for habitat and wetland creation often target improved habitat conditions for coastal and inland waterbirds and comprise an established feature of the USACE dredging operations program (Soots and Landin 1978; Fischer et al. 2004, 2010; Guilfoyle et al. 2006, 2007). Engineering operations providing environmental and other benefits are consistent with the USACE Engineering With Nature (EWN®) initiative, which seeks to integrate infrastructure development and environmental enhancements to advance science and engineering practices within the USACE (Banks et al. 2013; Bridges et al. 2018; Guilfoyle et al. 2020a).

Engineering operations and the use of dredged material to create habitat for native floral and faunal species have often been combined to mitigate the loss of habitat resulting from coastal engineering projects. However, the USACE can be proactive, as with the Craney Island facility, to provide environmental and other benefits through regular infrastructure maintenance activities (Bridges et al. 2018; Guilfoyle et al. 2020a). On the CIDMMA, a specific Long-term Bird Management Plan (Beck 2012) was developed to promote habitat for seasonal avian species, including many breeding species in the region. On the USACE Savannah District operations on the Savannah Harbor Navigation Project (SHNP), Dredged Material Containment Areas (DMCAs), the effort to ensure sufficient habitat for seasonal and nesting birds, provides a similar framework for the consistent availability of critical open water, sandy berms, wetland, and mudflat habitats for seasonal waterbirds (Calver et al. 2016; Guilfoyle et al. 2020a, b).

Urban and industrial ports and harbors often have significant impacts on the availability of coastal habitats, including open sandy beach, fresh and saltwater marshes, nondegraded estuaries, sand spits and extend mudflats. Such habitats are usually lost or degraded by large-scale changes from urban and industrial expansion and development, accumulated waste and pollution, sea-level rise, global climate change, and the modification of natural sedimentation processes that serve to shift sediment away from navigation channels. However, ecosystem restoration and mitigation actions are often limited in their application since results of these actions are not reported or evaluated outside of government bureaucracy and literature. Future efforts should work to include long-term monitoring of restoration and management plans and document success and limitations of these efforts (Calver et al. 2016; Guilfoyle 2019, 2020b).

In this effort, the circumstances of the SHNP DMCAs and the CIDMMA have numerous similarities. Both facilities have initiated long-term management plans to improve habitats for seasonal coastal waterbirds (and other birds) and both have made efforts to share their data and document the year-round benefits their actions have on local and regional birds of conservation concern (e.g., Calver et al. 2016; Guilfoyle et al. 2020a, b). This report serves to introduce issues, management actions, and current knowledge on the relative abundance, richness, and seasonal trends of bird communities on the CIDMMA in much the same way that

Calver et al. (2016) introduces management and monitoring of bird communities on the SHNP DMCA's.

## **1.2 National initiatives**

### **1.2.1 The North American Bird Conservation Initiative**

The North American Bird Conservation Initiative (NABCI) was established in 2000 and integrated efforts to manage multiple groups of birds (e.g., shorebirds, waterfowl, and landbirds) into a more effective and efficient approach to bird conservation (NABCI Committee 2000). This integration includes efforts for specific taxa within the NABCI including the North American Waterfowl Management Plan (US Fish and Wildlife Service [USFWS] 1998), Partner's in Flight (PIF) (Pashley et al. 2000), United States Shorebird Conservation Plan (Brown et al. 2001), North American Waterbird Conservation Plan (Fitzgerald and Pashley 2000), and Waterbird Conservation for the Americas (Kuslan et al. 2002), into an overall approach to conserve all bird species (US NABCI Committee 2000). Currently, the NABCI functions as a voluntary, international coalition of government agencies, conservation groups, academic institutions, private businesses, and citizens dedicated to the conservation of birds and their habitats through cooperative efforts in North America and the Neotropics.

The NABCI has built upon physiographic regions developed by PIF, into Bird Conservation Regions (BCRs) developed by the Atlantic Coast Joint Venture (ACJV) (2008). As with PIF, bird conservation plans have been developed for most BCRs. The purpose of these plans is to highlight those species in the region in most need of conservation. Individual species and associated habitats are described, along with recommendations and goals to build population sustainability within these regions. Using this approach, it is hoped that federal and state listed threatened and endangered species can eventually be delisted, or to minimize the probability that rare and sensitive species will be listed in the future. The plans identify the general habitat requirements of priority species at the site-level, and then identify the quantity and quality of habitat required by birds at the landscape scale. Most conservation plans for each BCR have been drafted and are available at the NABCI website <https://nabci-us.org/resources/bird-conservation-regions-map/>. The CIDMMA lies in the New England/Mid-Atlantic Region (BCR #30) and a full conservation plan for BCR 30 can be found at <https://acjv.org/bcr-planning/>.

The NABCI uses a species prioritization scheme developed by PIF to determine which species in each region are in most need of conservation attention (Carter et al. 2000; Pashley et al. 2000). The scheme ranks each species of North American breeding bird by BCR based upon seven measures of conservation “vulnerability.” These factors include (1) relative abundance (interspecific); (2) size of breeding range; (3) size of non-breeding range; (4) threats to the species on the breeding grounds; (5) threats to the species on the wintering grounds; (6) current known population trends; and (7) relative density (intraspecific) in a given planning unit compared to the maximum reached within its range. While focus is generally on breeding birds, information for all birds, including Neotropical migrants, Nearctic migrants, and resident species are included in the measure of conservation vulnerability. For the New England/Mid-Atlantic BCR (Atlantic Coast Joint Venture 2008), birds have been grouped into three categories of priority levels, including, Highest, High, and Moderate. Species categorized as ‘Highest Priority’ are those continental or regional species in need of immediate management attention; ‘High Priority’ are continental or regional species in need of management attention; and ‘Moderate Priority’ are those continental or regional species where monitoring is recommended to ensure long-term persistence of extant populations (Atlantic Coast Joint Venture 2008).

### **1.2.2 Craney Island Long-term Bird Management Plan**

A long-term Bird Management Plan (LTBMP) was written for the Craney Island facility by Beck (2012). Although not formally adopted by the USACE Norfolk District, this plan has been followed since its development. Moreover, through consultation with researchers at the College of William and Mary, and through contractors, much of the efforts to monitor and conserve coastal birds have been in operation since 1975. Primary actions are performed, in consultation and coordination with USACE Norfolk District, to conserve and protect avian populations (from Beck 2012):

1. **Seasonal monitoring.** Monitoring actions are used to document presence and use by avian populations, particularly federally and state listed species, plus local or regional species of concern (Figure 1-1). Monitoring includes regularly conducted surveys of the CIDMMA and monitoring of active nesting sites by coastal birds, particularly federal or state listed species. The effort provides information to ensure active

use areas or nesting sites will remain undisturbed by USACE facility operations.

**Figure 1-1.** A Black-necked Stilt chick on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. This is one of many nesting species monitored and protected on the facility. (Photo credit: USACE Norfolk District).



2. **Management actions for dredged material placement operations.** This effort serves to ensure ongoing dredged material deposition does not flood or disturb nesting birds. The facility will be monitored to ensure that all back-ponded nesting sites will avoid flooding during large-scale inflow actions. Areas with high use by nesting birds, especially high sandy berms used by Least Terns, will be located and signs posted to maintain appropriate buffers from ongoing actions (Figure 1-2).

Figure 1-2. Signage on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, indicating an area closed to protect nesting birds. (Photo credit: S. J. Reinheimer, USACE Norfolk District).



3. **Management during borrow and dike construction activities.**  
Borrow and dike construction measures are actively employed to retain capacity of the facility to receive future dredged material (Figure 1-3). Generally, this action occurs on the eastern portion of a containment cell not currently receiving dredge material. Least Terns commonly use borrow sites for nesting; therefore, multiple borrow sites are constructed such that borrow sites without nesting terns can be used for source material to increase level of containment dikes. Heavy construction equipment is used during dike construction, and truck-haul operations are monitored to ensure nesting birds are not disturbed. Routes can be changed, or access restricted to protect nesting birds; when needed, all traffic along specific roads can be closed until after 1 August of that year.

Figure 1-3. On-site construction on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. Efforts are made to ensure construction does not disturb nesting birds. (Photo credit: S. J. Reinheimer, USACE Norfolk District).



4. **Management of containment cell dewatering and pond water levels.** Water resources on the facility provide foraging ponds and mudflats for many migratory shorebirds and breeding wading birds, and deeper ponds provide important habitat for wintering waterfowl (Figure 1-4). The size, depth, and duration of these habitats often depend upon the hydraulic actions to deposit dredged material into the facility. Planned deposition can be performed to actively provide water levels to meet water quality requirements; however, many of these areas also meet the needs of shorebirds, waterfowl, and other species. A cell must be periodically dewatered to permit settling of the sediments and extend the life of the containment cell and the capacity of the facility to receive future deposition. Ponds, mudflats, and other water-based habitats can be created and/or maintained through significant precipitation, providing important foraging habitat for some species. Borrow sites often hold water for longer periods due to regular precipitation events. These deeper water holding sites can provide important habitat for over-wintering waterfowl and other coastal species.

Figure 1-4. Pond and mudflat habitats can be created, and water levels managed to provide habitat for migratory shorebirds and wintering waterfowl on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. (Photo credit: S. J. Reinheimer, USACE Norfolk District).



5. **Management actions for mowing, road, and site maintenance.** Roads are regularly maintained on the Craney Island facility. Damage from precipitation and storms must be repaired to maintain normal operations. Areas on the roads can be weak and susceptible to damage or washboard conditions due to use by heavy construction vehicles. Mowing also occurs regularly to maintain shoulders and to reduce invasive phragmites and other invasive plants. To protect nesting grassland birds, mowing is minimized from May 1 to July 15 every year.
6. **Management actions during military landing operations.** The US Navy has previously used the CIDMMA for specific training activities including helicopter touch-and-go and night-time training operations. These training operations were limited to facility sections that avoid nesting birds. Approaching flight paths and landing zones were also directed away from active nesting sites. These actions have not been permitted since 2013 and are not anticipated to be conducted in the future.
7. **Unavoidable impacts.** Occasionally, accidents, or specific planned actions occur that may disturb active nesting sites, or areas used by foraging or roosting birds. In such cases, the USACE will initiate formal consultations with the USFWS under the Endangered Species Act

(ESA) as needed. Collaborative partnerships with other federal agencies and contractors can help coordinate actions and minimize impacts to nesting, foraging, and roosting birds.

**8. Other Conservation Recommendations:**

- a. **Substrate augmentation:** The LTBMP recommends construction of a permanent nesting site for terns and American Oystercatchers on an area exterior to the containment cells. For comparison, a suite of nesting birds on the Hampton Roads Bridge Tunnel South Island is provided as an example of birds that could be attracted to such a site. Note that an American Oystercatcher pair produced at least one young at CIDMMA in 2020: this is the first recorded successful breeding of this species on the facility.
- b. **Pond management:** The LTBMP recommends the monitoring and synchronizing of water levels within the three existing cells (Figure 1-5) and a proposed fourth cell. Multiple species have specific water depth requirements that could be targeted. This effort would require implementation in a manner consistent with ongoing objectives for dredged material placement and future facility capacity.
- c. **Public awareness:** The LTBMP recommends regular updates to the CIDMMA website to inform the general public on ongoing plans for eastward expansion, updated satellite imagery, links to pages documenting nesting waterbirds, results of seasonal surveys, visits by local bird clubs (e.g., Audubon Society), links to nearby parks and refuges (e.g., Hoffer Creek Wildlife Preserve and the Great Dismal Swamp), and links to the VA Birding and Wildlife Trail sponsored by the Virginia Department of Wildlife Resources. This can become a challenging enterprise requiring thoughtful consideration by the USACE and CIDMMA staff. Currently, the facility is closed to all non-designated personnel.

Figure 1-5. Figure showing the North, Center, and South Cells on the Craney Island Dredged Material Management Area, Portsmouth, Virginia. Portsmouth, Virginia. (Photo credit: USACE Norfolk District).



### 1.3 Objectives

The objective of this study was to use long-term mean abundance and species richness trends of individual bird species and species groups using seasonal habitats to assess the value of habitats provided by the Craney Island facility for a variety of coastal bird species in the New England/Mid-Atlantic BCR. Ideally, monitoring and assessment of mitigation or ecosystem restoration efforts should be compared to regional reference sites, where the goal of the effort is to achieve comparable measures of biological metrics between the reference sites and newly created or restored areas. As with the SHNP DMCA, the CIDMMA did not include the identification and monitoring of a reference site. Therefore, we have chosen to define the success or failure of the LTBMP on CIDMMA using the same approach as Guilfoyle (2020b): we use long-term mean abundance and species richness trends of individual bird species and species groups using seasonal habitats. Successful creation of habitats for seasonal bird communities is assessed by determining whether abundance and richness trends show statistically significant increasing trends, or presence but stable trends (not significantly increasing or decreasing) during one or more seasons for species identified by the New England/Mid-Atlantic Region as regional species of concern. Success of the effort could be described as moderate or poor if regionally identified

priority species are either absent, present in low numbers, or show seasonally significant decreasing abundance or richness trends during the monitoring period.

## **1.4 Approach**

### **1.4.1 Procure and analyze population trends of the seasonal bird community**

The first step in developing this report involved procuring, compiling, and analyzing existing data on population trends of the bird community at the Craney Island over time. Data collected by USACE Norfolk District personnel, the CVWO Waterbirds Team, contractors, and Ruth A. Beck and others associated with the College of William and Mary, Williamsburg, VA, were procured with permission from the Cornell Lab of Ornithology, online database, eBird (<https://ebird.org/home>). We used the identical statistical approach to analyzing mean abundance and species richness trends as Guilfoyle et al. (2020b). Preliminary results are presented in this report, however, further consideration to data error and uncertainty will likely require additional future analyses. For example, the large differences in survey efforts in 2008 versus subsequent years, may require data from 2008 be dropped from any future analysis. For presenting preliminary results, data from 2008 are included, but trend results should be interpreted with caution. Future statistical analyses will also include an analysis into the effects of habitat availability on mean abundance and richness trends (Guilfoyle et al. 2020b). As presented by Guilfoyle et al. (2020b), we use a Bayesian community generalized linear mixed model approach with bird and species group abundance fitted into a negative binomial distribution, and species richness for each species group fitted into a Poisson distribution. We used the ‘jags UI’ 1.4.2 package (Kellner 2016) in the R 3.1.1 program (R Core Team 2014) to test for significant increasing or decreasing trends during the monitoring period. Significant seasonal population trends for abundance and species richness, positive, negative, or non-significant, were determined based on overlap of the logarithmic means and confidence intervals.

## **1.5 Scope**

This report targets USACE land managers or districts that manage or regulate coastal engineering projects or operations of various CDFs. However, the results of this effort may be of significant interest to all

biologists and land managers interested in habitat creation or restoration to benefit seasonal coastal waterbirds, shorebirds, waterfowl, and terrestrial landbirds. Results presented and methods described should be considered for monitoring on other USACE project lands or coastal engineering operations, and other state, federal, or private landowners with an interest and objectives to manage coastal lands for sustained seasonal use by populations of rare, sensitive, or regionally identified avian species of concern.

## 2 Study Area and Methods

### 2.1 Study area

The CIDMMA is located along the confluence of the James and Elizabeth Rivers, in the city of Portsmouth, just west of the city of Norfolk, Virginia (Figure 2-1). This CDF facility is a federally owned civil works project operated by the USACE, Norfolk District. This facility is the primary dredged material disposal facility in the Norfolk Harbor that receives dredged material during the dredging of navigation channels for the US Navy and the Virginia Port Authority. The facility is approximately 2,500 acres (1,011 ha) and extends from the mainland about 2 miles (3.2 km) north, 2 miles west, and 2 miles south. There are about 8 miles (12.9 km) of a primary containment dike serviced (additional containment dikes that separate the area into three subcontainment areas (Figure 2-1). The facility is surrounded by approximately 6 miles (9.7 km) of shoreline consisting of sand beaches, marshes, and hardened structures (Beck 2012).

Coastal habitats and vegetative communities on the Craney Island facility are likely very similar to conditions of coastal habitat measured in the city of Portsmouth (US Department of Commerce 1988). The most common type of habitat includes non-vegetated tidal wetlands that would include sand/mud mixed flats and sand or mud flats, and intertidal beach. Typical plants in these areas and emergent marsh areas would include saltmarsh chordgrass (*Spartina alterniflora*), saltbush (*Atriplex* spp.), saltmeadow hay (*Spartina patens*), marsh fleabane (*Pluchea odorata*), saltmarsh aster (*Symphytotrichum subulatum*), waterhemp (*Amaranthus tuberculatus*), seaside goldenrod (*Solidago sempervirens*), reedgrass (*Calamagrostis* spp.), black needlerush (*Juncus roemerianus*), saltmarsh bulrush (*Bolboschoenus robustus*), and other rushes (*Juncus* spp.). In addition to coastal wetland and beach areas, other areas have some scattered young forest and scrub areas. Typical plants in these areas include sycamore (*Platanus occidentalis*), red maple (*Acer rubrum*), sweetgum (*Liquidambar styraciflua*), ash species (*Fraxinus* spp.), honeysuckle (*Lonicera japonica*), trumpet creeper (*Campsis radicans*), poison sumac (*Toxicodendron vernix*), and bittersweet (*Celastrus scandens*). Other potential tree species include loblolly pine (*Pinus taeda*), oaks (*Quercus* spp.), American holly (*Ilex opaca*), wax myrtle (*Myrica cerifera*), and redbud (*Ceris canadensis*) (US Department of Commerce 1988). Cell hillside slopes vegetated by scattered to dense stands of Groundsel

(*Baccharis halimifolia*) and the roadside along the Portsmouth Landfill entrance has a large stand of black locust (*Robinia pseudoacacia*). Upper perimeter and cross-dikes roadsides are vegetated by white sweet clover (*Melilotus albus*) and Cocklebur (*Xanthium strumarium*). Substantial common reed (*Phragmites australis*) stands exist throughout the facility, and interior areas of the cells will support fall aster (*Symphotrichum* spp), Queen Anne's Lace (*Daucus carota*), and Dog-fennel (*Eupatorium capillifolium*) (Hamilton and Hall 2013).

Figure 2-1. Overview of the Craney Island Dredged Material Management Area, Portsmouth, Virginia.



## 2.2 Seasonal bird community monitoring

From 2008 to 2014, regularly conducted seasonal surveys generally occurred semimonthly on the CIDMMA, but on occasion, no surveys were conducted in some months, and occasionally, three or more surveys were conducted in a single month (Table 2-1). Since 2015, these surveys have occurred weekly. These surveys include a complete count of the area, which corresponds to an area search (Guilfoyle and Fischer 2007), where all birds detected by sight or sound were recorded and start and stop time data were recorded to determine the length of time for all surveys. Since surveys were intended to represent complete counts of all birds present, total time required to conduct the surveys varied depending on time of the year, current weather conditions, and number of birds present, among other factors. Surveys began in the early morning hours (generally before civil sunrise) and ended circa 1300-1400 hrs. Surveys were conducted during periods of relatively good weather; no surveys were conducted on days with heavy wind, rain, or other precipitation. All surveys were performed with a minimum of four experienced birders; usually, one would be responsible to record all observations and enter all data. All birders used high quality field binoculars and spotting scopes as necessary.

A total of 403 surveys were conducted during the monitoring period, with approximately 2,667.3 survey hours completed (Table 2-1). All birds detected by sight or sound within the CIDMMA boundaries were counted, whether they were flying or stationary. Survey boundaries were judged to be the boundary of the perimeter containment dikes on the east, north, and west side of the facility; south boundary was judged to be the southern perimeter access road.

We also identified priority species listed in ACJV (2008) to identify those birds detected on CIDMMA recognized as Highest, High, and Moderate Priority species for the New England/Mid-Atlantic Coast BCR.

**Table 2-1. Total number of survey days, and total survey hours for all birds detected on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008 to 2020.**

| Year  | Total number of days surveyed | Total hours of surveys |
|-------|-------------------------------|------------------------|
| 2008  | 5                             | 25.3                   |
| 2009  | 22                            | 104.6                  |
| 2010  | 21                            | 135.4                  |
| 2011  | 30                            | 187.5                  |
| 2012  | 22                            | 147.8                  |
| 2013  | 38                            | 248.6                  |
| 2014  | 35                            | 247.8                  |
| 2015  | 38                            | 259.8                  |
| 2016  | 43                            | 291.9                  |
| 2017  | 47                            | 330.1                  |
| 2018  | 44                            | 284.8                  |
| 2019  | 45                            | 310.4                  |
| 2020  | 13                            | 93.3                   |
| TOTAL | 403                           | 2,667.3                |

## **2.3 Initial mean abundance and species richness trends**

### **2.3.1 Mean abundance trends**

Surveys included in the analyses include 403 surveys (105 spring, 136 summer, 93 fall, and 69 winter). Seasons correspond to spring (March to May), summer (June - August), fall (September – November), and winter (December to February). These broad seasonal categories are intended to reflect specific seasonal changes in the seasonal life history phenology of these birds. The general seasonal pattern observed with most North American birds is that most species generally breed in the summer, migrate south during the fall (southward migration), and migrate north during the spring (northward migration) with several months generally spent in southern locations during the over-wintering season. This broad characterization can be highly variable within populations of many coastal waterbirds. Some may be migrating during February and March, breed in the spring, and begin their southward migration as early as July. Future efforts to analyze these data will attempt to correct the seasonal cut-off dates to reflect actual seasonal phenology of these birds more closely.

Also, in the current preliminary analyses, all birds are included, including birds identified only to species groups (e.g., Duck spp.) and identifications that may indicate a hybrid (e.g., Mallard X American Black Duck). Count data were only excluding for individual birds in which the identification could not be confirmed to species. Two exceptions include *Empidonax* spp. and *Calidris* spp., which represent a group of flycatchers and sandpipers, respectively, which can be difficult to identify in the field. In general, any species detected during fewer than 10% of surveys in any season will be eliminated in future analyses, as there simply are not enough data to estimate trends for those species. Therefore, the presented trend analyses only represent a preliminary analysis to provide broad-level trends on the CIDMMA, and results should be interpreted with caution.

The statistical trend analyses and associated equations are identical to the approach described in Guilfoyle et al. (2020b). For purposes of this report, and the preliminary nature of the trend analyses described, equations are not provided. For each species or species group in each season, a Bayesian community generalized linear mixed model was fixed with time treated as a fixed effect, and year as a random grouping variable. Bird abundance data were fitted to a negative binomial distribution. Community models estimate species-specific trends within a hierarchical framework where estimates are related through community-level hyperparameters (Kéry and Royle 2008; Zipkin et al. 2009). The full advantages of such an approach are detailed elsewhere (Kéry and Royle 2008; Royle and Dorazio 2008; Zipkin et al. 2009; Iknayan et al. 2014), but we chose this approach for two primary reasons. First, by borrowing information from more common species, the approach allows inclusion of rarer species in the analysis (Royle and Dorazio 2008; Zipkin et al. 2009; Iknayan et al. 2014). Second, we were interested in community hyperparameter estimates themselves to summarize overall group trends.

Trend significance was evaluated for individual species and species groups based on logarithmic values with 95% confidence intervals for each seasonal trend during the monitoring period. We determined significant trends based on whether the mean and confidence intervals overlapped with zero; those means and confidence intervals that are above zero indicate significantly increasing trends, while those means and confidence intervals below zero indicate significantly decreasing trends. Means and confidence intervals that overlap with zero indicate non-significant trends during the monitoring period (See Chapter 3 and Figures 3-10 – 3-27).

### 2.3.2 Mean species richness trends

The current analysis includes all 271 species recorded during the monitoring period that were then placed into one of 17 species groups (common names, scientific names, and 4-letter species codes for all birds detected in this effort are provided in Appendix A).

1. Blackbirds, Crows, and Jays
2. Dabbling Ducks
3. Diving and Sea Ducks
4. Flycatchers
5. Geese and Swans
6. Grassland Birds and Sparrows
7. Gulls, Terns, and Skimmers
8. Herons and Egrets
9. Other Nonpasserines
10. Other Passerines
11. Raptors and Vultures
12. Sandpipers and Plovers
13. Shorebirds
14. Swallows and Swifts
15. Thrushes
16. Warblers
17. Woodpeckers

For each species group, we fit a generalized linear mixed model where species richness values had a Poisson distribution, time was treated as a fixed effect, and year was a random grouping variable. The analysis was conducted in R 3.1.2 using the `glmer()` function in the `lme4` package (Bates et al. 2015). A summary of all species groups detected over the course of the 12-yr monitoring period (2008 to 2020) is provided, focusing only on mean species richness. For clarity on species richness, we generally only used data on detections identified to species. As for mean abundance models, exceptions will include the designation of birds in the genus, *Empidonax* and *Calidris*. As with abundance trends, significance of species trends was determined based on the overlap of the logarithmic means and confidence intervals as described above for mean abundance models (see Chapter 3 and Figure 3-28).

## 3 Results

### 3.1 Seasonal survey monitoring results

Over 1.5 million detections of 271 species were recorded during survey efforts on the CIDMMA over the 12-yr survey period (Appendix A). Eighty-four species are wintering species that breed farther north during the spring and summer; 33 species are present largely during the fall and spring migrations seasons; 27 species were Neotropical migrants that breed in the area, but undergo a long-distance migration to Central and South America during the winter months; 88 species are residents that occur year-round and some likely breed near or on site; and 50 are spring/summer breeding species that winter likely south of the study area. An additional 22 species are rare transients known to have irregular or only occasional occupancy in the area during the breeding, wintering, or migration seasons (Appendix A).

Numerous high priority species identified by the New England/Mid-Atlantic BCR 30 (Atlantic Coast Joint Venture 2008) were documented on the CIDMMA (Table 3-1). Of 134 ranked priority species in the region, the CIDMMA supported 102 of 134 (76%) recognized in the BCR, including 16 of 19 (84%) of Highest priority ranked species, 47 of 60 (78.3%) of High priority species, and 39 of 55 (71%) of Moderate priority species for BCR 30. Represented of these priority species found on the CIDMMA are shown in Figures 3-1 – 3-9.

Over the 2008 to 2020 survey period, the most commonly recorded species was the High Priority Species, Semipalmated Sandpiper (261,500; total cumulative annual detections), followed by Moderate Priority Species, Ruddy Duck (179, 692), unranked Northern Shoveler (145,476), *Calidris* ssp. Sandpipers (96,986), Double-crested Cormorant (76,016), Dunlin (68,934), Mallard (50,072), Red-winged Blackbird (47,168), Green-winged Teal (44,743), Lesser Yellowlegs (40, 910), European Starling (38,888), Ring-billed Gull (38,036), Herring Gull (28,221), Laughing Gull (25,636), Least Tern (25,071), Canvasback (24,945), and the American Avocet (24,568) (Appendix A).

The Semipalmated Sandpiper was the most common Neotropical Migrant or Migrant, followed by the Least Tern, Semipalmated Plover (14,560), Short-billed Dowitcher (12,337), Western Sandpiper (9,813), Chimney

Swift (9,558), Blue-winged Teal (3,058), and Caspian Tern (2,966) (Appendix A). The Mallard was the most common occurring Resident or Year-Round species, followed by the Red-winged Blackbird, European Starling, Herring Gull, and Killdeer (11,189) (Appendix A). The Ruddy Duck (179,682) was the most common winter species, followed by the Northern Shoveler (145,476), Dunlin (68,934), Green-winged Teal (44,743), Ring-billed Gull (38,036), Canvasback (24,945), Red-breasted Merganser (19,396), and the Gadwall (19,323) (Appendix A).

During the survey period, the bird community was most diverse during the winter and most abundant during the summer: 164 wintering species constituted 21.4% of total bird counts (total counts = 335,419 individuals), while 81 spring species constituted 26.6% of total birds (counts = 411,948 individuals); 138 summer species constituted 30.4% of total birds (counts = 470,586 individuals), and 118 fall species constituted 21.6% of all birds (counts = 331,524 individuals).

**Table 3-1. Total cumulative detections of the New England/Mid-Atlantic Coast Bird Conservation Region (BCR 30) Highest Priority, High Priority, and Moderate Priority Species of regional or local conservation concern (from Atlantic Coast Venture 2008) detected on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008 to 2020.**

| <b>Highest Priority Species</b>              |                              |                                    |                              |
|----------------------------------------------|------------------------------|------------------------------------|------------------------------|
| <b>Species</b>                               | <b>Cumulative Detections</b> | <b>Species</b>                     | <b>Cumulative Detections</b> |
| American Black Duck <sup>1,2,3</sup>         | 4,108                        | Prairie Warbler <sup>1</sup>       | 32                           |
| American Oystercatcher <sup>1</sup>          | 97                           | Red Knot <sup>3</sup>              | 74                           |
| American Woodcock <sup>1,2,3</sup>           | 4                            | Red-throat Loon <sup>2,3</sup>     | 1,992                        |
| (Atlantic) Brant <sup>2,3</sup>              | 13*                          | Ruddy Turnstone <sup>3</sup>       | 794                          |
| (Atlantic) Canada Goose <sup>2,3</sup>       | 11,235*                      | Saltmarsh Sparrow <sup>1,2,3</sup> | 3                            |
| Blue-winged Warbler <sup>1</sup>             | 1                            | Sanderling <sup>2,3</sup>          | 8,053                        |
| Gull-billed Tern <sup>1</sup>                | 772                          | Whimbrel <sup>3</sup>              | 36                           |
| Piping Plover <sup>1,2,e</sup>               | 18                           | Wood Thrush <sup>1</sup>           | 1                            |
| <b>High Priority Species</b>                 |                              |                                    |                              |
| American Golden Plover <sup>3</sup>          | 43                           | Least Tern <sup>1</sup>            | 25,071                       |
| Baltimore Oriole <sup>1</sup>                | 9                            | Long-tailed Duck <sup>2,3</sup>    | 4                            |
| Bay-breasted Warbler <sup>3</sup>            | 1                            | Mallard <sup>1,2,3</sup>           | 50,072                       |
| Black Scoter <sup>2,3</sup>                  | 82                           | Marbled Godwit <sup>3</sup>        | 37                           |
| Black-and-white Warbler <sup>1</sup>         | 30                           | Marsh Wren <sup>1</sup>            | 95                           |
| Black-bellied Plover <sup>2,3</sup>          | 2,333                        | Northern Bobwhite <sup>1,2</sup>   | 81                           |
| Brown Thrasher <sup>1</sup>                  | 942                          | Northern Flicker <sup>1,2,3</sup>  | 323                          |
| Buff-breasted Sandpiper <sup>3</sup>         | 20                           | Northern Gannet <sup>2,3</sup>     | 1,057                        |
| Bufflehead <sup>1,2,3</sup>                  | 8,266                        | Prothonotary Warbler <sup>1</sup>  | 1                            |
| (North Atlantic) Canada Goose <sup>2,3</sup> | 11,235*                      | Purple Sandpiper <sup>2,3</sup>    | 3                            |
| Canvasback <sup>2,3</sup>                    | 24,945                       | Scarlet Tanager <sup>1</sup>       | 4                            |

|                                          |        |                                           |         |
|------------------------------------------|--------|-------------------------------------------|---------|
| Chimney Swift <sup>1</sup>               | 9,558  | Semipalmated Sandpiper <sup>3</sup>       | 261,500 |
| Clapper Rail <sup>1</sup>                | 46     | Short-billed Dowitcher <sup>3</sup>       | 12,337  |
| Dunlin <sup>2,3</sup>                    | 68,934 | Solitary Sandpiper <sup>3</sup>           | 33      |
| Eastern Kingbird <sup>1</sup>            | 618    | Surf Scoter <sup>1,2,3</sup>              | 531     |
| Eastern Towhee <sup>1,2,3</sup>          | 947    | (Eastern) Tundra Swan <sup>2,3</sup>      | 885*    |
| Field Sparrow <sup>1,3</sup>             | 331    | White-rumped Sandpiper <sup>3</sup>       | 440     |
| Foster's Tern <sup>1,3</sup>             | 661    | White-winged Scoter <sup>2,3</sup>        | 6       |
| Glossy Ibis <sup>1</sup>                 | 145    | Willet <sup>1,2,3</sup>                   | 1,391   |
| Great-crested Flycatcher <sup>1</sup>    | 279    | Willow Flycatcher <sup>1</sup>            | 13      |
| Greater Scaup <sup>2,3</sup>             | 41     | Wilson's Phalarope <sup>3</sup>           | 441     |
| Greater Yellowlegs <sup>2,3</sup>        | 3,323  | Wilson's Plover <sup>1</sup>              | 1       |
| Horned Grebe <sup>2,3</sup>              | 9,246  | Worm-eating Warbler <sup>1</sup>          | 1       |
| Hudsonian Godwit <sup>3</sup>            | 9      | Yellow-throated Warbler <sup>1</sup>      | 3       |
| <b>Moderate Priority Species</b>         |        |                                           |         |
| American Avocet <sup>3</sup>             | 24,568 | Northern Pintail <sup>2,3</sup>           | 4,161   |
| American Bittern <sup>1,2,3</sup>        | 8      | Red-necked Phalarope <sup>3</sup>         | 760     |
| American Wigeon <sup>2,3</sup>           | 4,930  | Red Phalarope <sup>3</sup>                | 14      |
| Bald Eagle <sup>1,2,3</sup>              | 763    | Red-breasted Merganser <sup>2,3</sup>     | 19,396  |
| Black Skimmer <sup>1</sup>               | 569    | Red-headed Woodpecker <sup>1,2,3</sup>    | 5       |
| Black-crowned Night-Heron <sup>1,2</sup> | 252    | Royal Tern <sup>1</sup>                   | 18,186  |
| Brown-headed Nuthatch <sup>1,2</sup>     | 54     | Ruddy Duck <sup>2,3</sup>                 | 179,692 |
| Common Goldeneye <sup>1,2,3</sup>        | 18     | Seaside Sparrow <sup>1,2,3</sup>          | 39      |
| Common Tern <sup>1,3</sup>               | 379    | Sedge Wren <sup>1,2,3</sup>               | 1       |
| Gadwall <sup>1,2,3</sup>                 | 19,323 | Semipalmated Plover <sup>3</sup>          | 14,560  |
| Gray Catbird <sup>1</sup>                | 747    | Short-eared Owl <sup>2,3</sup>            | 5       |
| Green-winged Teal <sup>1,2,3</sup>       | 516    | Snowy Egret <sup>1,2</sup>                | 3,453   |
| Hooded Merganser <sup>1,2,3</sup>        | 44,743 | Sora <sup>1,3</sup>                       | 33      |
| Killdeer <sup>1,2,3</sup>                | 11,189 | Spotted Sandpiper <sup>1,3</sup>          | 2,267   |
| King Rail <sup>1,2</sup>                 | 1      | Tricolored Heron <sup>1</sup>             | 75      |
| Least Bittern <sup>1</sup>               | 29     | Upland Sandpiper <sup>1,3</sup>           | 6       |
| Least Sandpiper <sup>3</sup>             | 18,336 | Western Sandpiper <sup>3</sup>            | 9,813   |
| Lesser Yellowlegs <sup>2,3</sup>         | 40,910 | Wood Duck <sup>1,2,3</sup>                | 41      |
| Little Blue Heron <sup>1,2</sup>         | 10     | Yellow-crowned Night-Heron <sup>1,3</sup> | 107     |
| Nelson's Sparrow <sup>1,3</sup>          | 4      |                                           |         |

<sup>1</sup>Breeding population is designated or ranked for protection/conservation.

<sup>2</sup>Non-breeding or wintering population is ranked for protection/conservation.

<sup>3</sup>Migratory population is ranked for protection/conservation.

eSpecies is listed as federally endangered.

\*Species not identified to subspecies; therefore, presence and/or number of potential identified Priority Subspecies is unknown.

## 3.2 Examples of priority species

### 3.2.1 Highest priority species

Figure 3-1. The American Oystercatcher is a species documented on the Craney Island Dredged Material Management Area that has been ranked as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservations Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Rhododentries, Wikimedia Commons).



Figure 3-2. The Ruddy Turnstone is a wintering species documented on the Craney Island Dredged Material Management Area that has been ranked as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservations Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Kate Perez, Wikimedia Commons).



Figure 3-3. The Piping Plover is a federally listed endangered species that has been documented on the Craney Island Dredged Material Management Area and has been listed as a Highest Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008). (Photo Credit: Shutterstock.com, Wikimedia Commons).



### 3.2.2 High priority species

Figure 3-4. The Black-bellied Plover is a migratory species documented on the Craney Island Dredged Material Management Area that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture) (Photo Credit: Public Domain, Wikimedia Commons).



Figure 3-5. The Northern Bobwhite is a species documented on the Craney Island Dredged Material Management Area that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Uatrok77, Wikimedia Commons).



Figure 3-6. The Semipalmated Sandpiper is a migratory species documented on the Craney Island Dredged Material Management Areas that has been ranked as a High Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Venture 2008) (Photo Credit: USFWS Northeast Region, Wikimedia Commons).



### 3.2.3 Moderate priority species

Figure 3-7. The Royal Tern is a species documented on the Craney Island Dredged Material Management Areas that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Region Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: defaulder, Wikimedia Commons).



Figure 3-8. The Red-necked Phalarope is a migratory species documented on the Craney Island Dredged Material Management Areas that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: USFWS, Wikimedia Commons).



Figure 3-9. The Ruddy Duck is a wintering species documented on the Craney Island Dredged Material Management Area that has been ranked as a Moderate Priority Species in the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008) (Photo Credit: Public Domain, Wikimedia Commons).




### 3.3 Annual trends for species and species groups

Monitoring of avian populations is an essential tool in determining population trends and the overall health of specific bird populations. Currently, the data collected on the CIDMMA will need to be analyzed more thoroughly to identify specific statistical trends for any species' groups or individual species. It is beyond the scope of this report to present such an analysis; however, we provide a preliminary view of trends for individuals and species groups to provide some idea of the variation and trends among seasons present in the dataset. In future analyses, trends will be correlated with annual habitat availability, especially in the context of CIDMMA management due to application of the LTBMP and to determine if habitat quality and/or quantity provide any explanatory insights into why such trends may be occurring.

- Blackbirds/Crows/Jays:

This group was stable for all seasons during the monitoring period with no significant increase or decreases; most all individual species also showed no significant trends except the Red-winged Blackbird that had significant increases during the winter, spring, and fall (Figure 3-10).

Figure 3-10. Seasonal mean abundance trends for Blackbirds, Crows and Jays on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Dabbling Ducks:

This group was stable for all seasons during the monitoring period, with only the Gadwall showing significant increases in abundance trends during the winter and the Northern Shoveler showing significant increases during the summer (Figure 3-11).

- Diving and Sea Ducks:

This group was stable for all seasons during the monitoring period, with only the Red-breasted Merganser showing significant increases in abundance trends during the winter and spring (Figure 3-12).

Figure 3-11. Seasonal mean abundance trends for Dabbling Ducks on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.





Figure 3-12. Seasonal mean abundance trends for Diving and Sea Ducks on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Flycatchers:

No significant changes during the monitoring period for this group or any of the individual five species during any season (Figure 3-13).

- Geese and Swans:

No significant changes during the monitoring period for this group or any individual eight species during any season (Figure 3-14).

- Grassland birds and Sparrows:

No significant changes during the monitoring period for this group or 19 of 20 individual species during any season (Figure 3-15). The one exception includes the significant increase in abundance trends for the Savannah Sparrow during the fall (Figure 3-15).

Figure 3-13. Seasonal mean abundance trends for Flycatchers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.




Figure 3-14. Seasonal mean abundance trends for Geese and Swans on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.





Figure 3-15. Seasonal mean abundance trends for Grassland Birds and Sparrows on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Gulls/Terns/Skimbers:

No significant changes during the monitoring period for this group and 15 of 17 species; exceptions include a significant increase in abundance trends for the Ring-billed Gull during the winter and spring, and a significant decrease in trends for the Black Tern during the fall (Figure 3-16).

- Herons and Egrets:

No significant changes during the monitoring period for this group or for 10 of 11 individual species during any season; the one exception includes a significant increase for the Snowy Egret in the spring (Figure 3-17).

Figure 3-16. Seasonal mean abundance trends for Gulls, Terns and Skimmers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.





Figure 3-17. Seasonal mean abundance trends for Herons and Egrets on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Other Nonpasserines:

No significant changes during the monitoring period for this group or for 30 of 31 individual species during any season; the lone exception includes a significant decrease for the Northern Gannet in the winter (Figure 3-18).

- Other Passerines:

No significant changes during the monitoring period for this group or for 43 individual species; however, some of the 95% confidence intervals fall right on the 0-line (Figure 3-19).

- Pelicans and Cormorants:

No significant changes during the monitoring period for this group or for 3 of 4 individual species during any season; the Double-crested Cormorant is an exception showing significant increases in abundance trends during the winter and spring (Figure 3-20).

Figure 3-18. Seasonal mean abundance trends for Other Nonpasserines on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.




Figure 3-19. Seasonal mean abundance trends for Other Passerines on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.





Figure 3-20. Seasonal mean abundance trends for Pelicans and Cormorants on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Raptors and Vultures:

No significant changes during the monitoring period for this group or for 17 individual species during any season (Figure 3-21).

- Sandpipers and Plovers:

No significant changes during the monitoring period for this group or for 21 of 23 individual species during any season; exceptions include a significant decrease in abundance trends for *Calidris* ssp. during the winter, and a decrease for the Least Sandpiper during the fall (Figure 3-22).

Figure 3-21. Seasonal mean abundance trends for Raptors and Vultures on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.






Figure 3-22. Seasonal mean abundance trends for Sandpipers and Plovers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Shorebirds:

No significant changes during the monitoring period for this group or for 11 of 13 individual species during any season; the American Avocet is an exception showing weak (just barely above the 0-line) significant increases in abundance trends during all four seasons, and the Short-billed Dowitcher showing increases during the spring and summer (Figure 3-23). There has been a clear increase in the number of breeding pairs of Black-necked Stilt at CIDMMA. This facility is one of two known breeding sites for this species in Virginia.

Figure 3-23. Seasonal mean abundance trends for Shorebirds on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



- Swallows and Swifts:

No significant changes during the monitoring period for this group or for 7 of 9 individual species during any season; exceptions include a significant increase in abundance trends for Chimney Swift during the summer and fall, and a borderline decline for Cliff Swallow during the summer (Figure 3-24).

- Thrushes:

No significant changes during the monitoring period for this group or for six individual species during any season (Figure 3-25).

Figure 3-24. Seasonal mean abundance trends for Swallows and Swifts on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.





Figure 3-25. Seasonal mean abundance trends for Thrushes on the Craney Island Dredged Material Management Areas, Portsmouth, Virginia, 2008-2020.



- Warblers:

No significant changes during the monitoring period for this group or for 27 of 28 individual species during any season; the lone exception is a weak significant increase in abundance trends for the Pine Warbler during the spring (Figure 3-26).

- Woodpeckers:

No significant changes during the monitoring period for this group or for 7 individual species during any season (Figure 3-27).

Figure 3-26. Seasonal mean abundance trends for Warblers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.






Figure 3-27. Seasonal mean abundance trends for Woodpeckers on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



### 3.4 Species richness

During the monitoring period, most of the 17 species groups showed significant increases in mean species richness trends during all seasons, except the Flycatchers and Woodpeckers, which showed no significant changes during any season, and Shorebirds and Thrushes that showed significant decreases in mean species richness trends during the winter and fall, respectively (Figure 3-28).

Figure 3-28. Seasonal mean species richness for avian species groups on the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020.



## 4 Discussion

The CIDMMA supports a significant number and diversity of birds and bird species year-round. This report summarizes long-term monitoring of the CIDMMA for the approximately 12-yr period from 2008 to 2020. These data support prior research that shows habitats created from dredged material are often important for coastal bird species, and in some cases, such habitats may support a disproportionate proportion of a species' continental or regional population (Soots and Landin 1978, Guilfoyle et al. 2006, 2007). Confined Disposal Facilities in particular, have a long history of supporting large numbers of birds, even in largely urban and industrial ports and harbors (Guilfoyle et al. 2020a, b). In such cases, habitats created with dredged material may not just be beneficial but may be essential in the long-term persistence of a species. Currently, the USACE Norfolk District continues to perform dredged material deposition in the CIDMMA according to the LTBMP and personnel and contractors are continuing the regularly conducted seasonal monitoring efforts.

Results of this effort continue to show the importance of the CIDMMA for regional bird populations, including many species ranked at Highest Priority and High Priority by the New England/Mid-Atlantic Bird Conservation Region (Atlantic Coast Joint Venture 2008). During the monitoring period documented in this report, several species ranked as Highest Priority were observed on the CIDMMA in large numbers, including the American Black Duck, Red-throated Loon, and Sanderling. During the winter, the CIDMMA is also used in large numbers by the Ruddy Duck and the Northern Shoveler. The CIDMAA may occasionally be used by the endangered Piping Plover, including breeding individuals in 1989-1991 (Beck 2006). The CIDMMA also supports large numbers of High Priority species, including the Semipalmated Sandpiper, Mallard, and Dunlin. Large numbers of Moderate Priority Species have been detected on the CIDMMA, with counts of many species, including the American Avocet, Red-breasted Merganser, Royal Tern, Killdeer, Semipalmated Plover, Least Sandpiper, and Lesser Yellowlegs, numbering in the tens of thousands over the monitoring period. Not only are such areas important for many regional priority species, but by supporting moderately rare and/or sensitive species, areas such as the CIDMMA may act as critical links that serve to buffer populations from becoming critically rare in the future.

In addition, dredged material deposited on the CIDMMA creates sandy berms that constitute important breeding areas for several priority species listed by the Atlantic Coast Joint Venture (2008). Verified breeding efforts have been documented by species designated as Highest Priority species, including Piping Plover (last nested in 1997 [Beck 2006]), and the American Black Duck (presence of broods suggestive of breeding by this species onsite [Beck 2012]; High Priority Species, including the Least Tern and Mallard, and Moderate Priority Species, including the American Avocet (one nesting attempt in the 1970's), Bald Eagle and Killdeer. Non-ranked species, including the Black-necked Stilt and Osprey have also nested on or near the facility. Although Brown Pelicans have continued to increase in numbers at the facility, no nesting has been recorded to date.

Additional nesting by other species have been documented on other sites relatively close to the CIDMMA, which include Grandview Nature Preserve and the Hampton Road Bridge Tunnel South Island. Birds nesting at these sites include the following: Laughing Gull, Herring Gull, Great Black-backed Gull, Common Tern, Gull-billed Tern, Sandwich Tern, Royal Tern, Black Skimmer, and American Oystercatcher. These sites also manage and protect nesting birds by vegetation control, predator control, and placement of chick shelters to shield hatchlings from predators and mid-day heat. The proposed eastward expansion of the Craney Island facility provides an opportunity to build a permanent nesting area for birds outside the containment cells as proposed by Beck (2012). With proper management and monitoring, such a site would likely attract a similar suite of nesting birds to the Craney Island facility as observed at the Hampton Road Bridge Tunnel South Island. In addition, creation of areas with suitable access to shoreline and/or moist soil foraging site, or removal of riprap added to the Craney facility in 1997 (Beck 2006), may attract breeding Piping Plovers that have not nested on Craney Island since 1997.

Included in this report is a preliminary statistical seasonal mean abundance and species richness trend analysis for 271 species and 17 species groups. In general, these results reveal largely stable, unchanging mean abundance trends for most species groups and individual species. Some species show significant increases during some seasons, and very few reveal decreasing abundances in any season (see Figures 3-10 – 3-27). These results also reveal most species groups had significant increasing species richness for most seasons during the monitoring period (Figure 3-

28). These trend analyses, plus the documentation of site utilization by ranked priority species in the New England/Mid-Atlantic Bird Conservation Region on the Craney Island facility, show that the site is likely an important area that contributes to the local and regional stability of seasonal coastal bird populations. With the recent documentation of significant declines of North American birds (Rosenberg et al. 2019), including coastal shorebirds and sea birds, creation, management and monitoring of such sites becomes even more important for building long-term sustainable coastal bird populations. The USACE can play an important role contributing to supporting coastal bird populations through proactive management of CDFs (Calver et al. 2016; Guilfoyle et al. 2020a, b) and by employing best management practices during coastal engineering activities (Guilfoyle et al. 2019). Current management practices at the CIDMMA are consistent with EWN principles, providing benefits including enhancing habitat quality sufficient for multiple bird species and for recreational birding activities. Moreover, in these actions, the USACE can accrue benefits under the ESA Section 7(A)(1) conservation planning USFWS to lower costs of compliance, capitalize on collaborations and partnerships, and provide demonstrable benefits to listed species, regionally ranked priority species and other rare and sensitive species (Hartfield et al. 2017; Guilfoyle et al. 2022).

## References

- Atlantic Coast Joint Venture. 2008. *New England/Mid-Atlantic Coast bird conservation region (BCR 30) implementation plan*. Laurel, Maryland: US Fish and Wildlife Service. [https://acjv.org/BCR\\_30/BCR30\\_June\\_23\\_2008\\_final.pdf](https://acjv.org/BCR_30/BCR30_June_23_2008_final.pdf)
- Banks, C. J., T. J. Fredette, B. C. Suedel, and T. S. Bridges. 2013. *Implementing with nature within the Corps: A Workshop*. ERDC/TN DOER-R21. Vicksburg, MS: US Army Engineer Research and Development Center. <http://el.erdclib.usace.army.mil>.
- Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. "Fitting Linear Mixed-Effects Models Using lme4." *Journal of Statistical Software* 67:1-48.
- Beck, R. A. 2006. "Long-term bird use of the Craney Island Dredged Material Site." Pages 41-43, in Guilfoyle, M. P., R. A. Fischer, D. N. Pashley, and C. A. Lott, eds., *Summary of first regional workshop on dredging, beach nourishment, and birds on the South Atlantic Coast*. ERDC/TN-06-10. Vicksburg, MS.
- Beck, R. A. 2012. *Long-term Bird Management Plan for the Craney Island Dredged Material Management Area*. Prepared for the U.S. Army Corps of Engineers, Norfolk District. R.A. Beck Avian Research Consulting, Barhamsville, VA. 141 p.
- Bridges, T. S., E. M. Bourne, H. K. Kuzmitski, E. B. Moynihan, and B. C. Suedel. 2018. *Engineering with Nature: An Atlas*. Vicksburg, MS: U.S. Army Engineer Research and Development Center. <https://erdclib.usace.army.mil/xmlui/handle/11681/27929>.
- Brown, S., C. Hickey, B. Harrington, and R. Gill, eds. 2001. *The U.S. Shorebird Conservation Plan, 2<sup>nd</sup> ed.* Manomet, MA: Manomet Center for Conservation Sciences.
- Calver, J. S., M. P. Guilfoyle, R. A. Fischer, and E. L. Covington. 2016. *Birds of the Savannah Harbor Navigation Project, Dredged Material Disposal Areas, 1994-2012*. ERDC/EL TR-16-4. Vicksburg, MS: US Army Engineer Research and Development Center. [http://acwc.sdp.sirsi.net/client/en\\_US/search/asset/1048847](http://acwc.sdp.sirsi.net/client/en_US/search/asset/1048847)
- Carter, M. F., W. C. Hunter, D. N. Pashley, and K. V. Rosenberg. 2000. "Setting conservation priorities for landbirds in the United States: The Partners In Flight Approach." *Auk* 117:541-548.
- Environmental Protection Agency (EPA) and the U.S. Army Corps of Engineers (USACE). 2007. "Identify, planning, and financing beneficial use projects using dredged material: beneficial use planning manual." EPA842-B-07-001. Environmental Protection Agency and the U.S. Army Corps of Engineers, Washington, DC. 114 p. <https://nepis.epa.gov/Exe/ZyPDF.cgi/P100390N.PDF?Dockey=P100390N.PDF>
- Fischer, R. A., M. P. Guilfoyle, C. A. Lott, and D. N. Pashley. 2004. "Application of dredged material disposal for enhancement of bird habitat: Past and current efforts by the U.S. Army Corps of Engineers." Pages 95-103 in C. A. Fleming, ed., 3<sup>rd</sup> Missouri River and North American Piping Plover and Least Tern Symposium, Sioux City, NE.

- Fitzgerald, J. A., and D. N. Pashley. 2000. *Partner's In Flight Conservation Plan for the Ozark/Ouachitas (Physiographic Area 19)*. Washington, DC: American Bird Conservancy.
- Guilfoyle, M. P., R. A. Fischer, D. N. Pashley, and C. A. Lott (eds.). 2006. *Summary of first regional workshop on dredging, beach nourishment, and birds on the South Atlantic Coast*. ERDC/TN-06-10. Vicksburg, MS. <http://el.erdcd.usace.army.mil>.
- Guilfoyle, M. P., R. A. Fischer, D. N. Pashley, and C. A. Lott (eds.). 2007. *Summary of second regional workshop on dredging, beach nourishment, and birds on the North Atlantic Coast*. ERDC/TN-07-26. Vicksburg, MS. <http://el.erdcd.usace.army.mil>.
- Guilfoyle, M. P., and R. A. Fischer. 2007. *Implementing avian inventory and monitoring efforts on Corps of Engineers project lands*. EMRRP Technical Notes Collection (ERDC/TN EMRRP-SI-32). Vicksburg, MS: US Army Engineer Research and Development Center. <http://el.erdcd.usace.army.mil/emrrp/techtran.html>
- Guilfoyle, M. P., J. F. Jung, R. A. Fischer, and D. D. Dickerson. 2019. *Developing Best Management Practices for Coastal Engineering Projects That Benefit Atlantic Coast Shoreline-Dependent Species*. ERDC/TN EMRRP-SI-38. Vicksburg, MS: US Army Engineer Research and Development Center. <https://erdcd-library.erdcd.dren.mil/xmlui/handle/11681/33203>.
- Guilfoyle, M. P., J. S. Calver, M. E. Richards, and R. A. Fischer. 2020a. *Use of Engineering With Nature concepts on the Savannah Harbor Navigation Project, dredged material containment areas, Savannah, GA*. ERDC/TN EWN-20-1. Vicksburg, MS: US Army Engineer Research and Development Center. <https://erdcd-library.erdcd.dren.mil/xmlui/handle/11681/35353>
- Guilfoyle, M. P., J. S. Calver, S. S. Jackson, M. F. Richards, and R. A. Fischer. 2020b. *Benefits of Engineered Habitats to Seasonal Bird Communities on the Savannah Harbor Navigation Project, Dredged Material Containment Areas, 1994 – 2012*. DOER Technical Reports (DOER TR-20-2). Vicksburg, MS, US Army Engineer Research and Development Center. <https://erdcd-library.erdcd.dren.mil/xmlui/handle/11681/36039>
- Guilfoyle, M. P., P. Hartfield, R. A. Fischer, J. F. Jung, and K. A. Reine. *Implementing ESA Section 7(a)(1) Conservation Planning during USACE Coastal Engineering Projects*. EMRRP Technical Notes Collection (ERDC/TN EMRRP-SI-40). Vicksburg, MS: US Army Engineer Research and Development Center. <https://erdcd-library.erdcd.dren.mil/jspui/handle/11681/44845>.
- Hamilton, H., and G. Hall. 2013. "Wildlife Flowers and Grasses of Virginia's Coastal Plain. Virginia Native Plant Society, Botanical Research of Texas Press, Fort Worth, TX." *Botanical Miscellany*, No. 40. 288 pp.
- Hartfield, P., J. G. Brown, and R. A. Fischer. 2017. *The Role of Interagency Cooperation in the Conservation of Threatened and Endangered Species*. ERDC/TN DOER-40. Vicksburg, MS, US Army Engineer Research and Development Center. [https://doer.el.erdcd.dren.mil/notes\\_reports.html](https://doer.el.erdcd.dren.mil/notes_reports.html).

- Iknayan, K. J., M. W. Tingley, B. J. Furnas, and S. R. Beissinger. 2014. "Detecting Diversity: Emerging Methods to Estimate Species Diversity." *Trends in Ecology and Evolution* 29: 97–106.
- Kellner, K. 2016. "jagsUI: a wrapper around 'rjags' to streamline 'JAGS' analyses." *R package*. Version 1.4.2. <http://CRAN.R-project.org/package=jagsUI>.
- Kéry, M., and J. A. Royle. 2008. "Hierarchical Bayes Estimation of Species Richness and Occupancy in Spatially Replicated Surveys." *Journal of Applied Ecology* 45:589–598.
- Kuslan, J., M. J. Steinkamp, K. Parson, J. Capp, M. A. Cruz, M. Coulter, I. Davidson, L. Dickson, N. Edelson, R. Elliot, R. M. Erwin, S. Hatch, S. Kress, R. Milko, S. Miller, K. Mills, R. Paul, R. Phillips, J. E. Saliva, B. Sydeman, J. Trapp, J. Wheeler, and K. Wohl. 2002. *Waterbird Conservation for the Americas Conservation Plan, Version I*. Washington, DC: Waterbird Conservation for the Americas.
- North American Bird Conservation Initiative Committee. 2000. *North American Bird Conservation Initiative*. Arlington, VA: US Fish and Wildlife Service.
- Partners In Flight. 2014. Partners In Flight: Physiographic area plans. <http://www.partnersinflight.org/>. (3 March 2014).
- Pashley, D. N., C. J. Beardmore, J. A. Fitzgerald, R. P. Ford, W. C. Hunter, M. S. Morris, and K. V. Rosenberg. 2000. *Partners in Flight: Conservation of the Land Birds of the United States*. American Bird Conservancy, The Plans, VA.
- R Core Team. 2014. *R: a Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <http://www.R-project.org/>.
- Rosenberg, K. V., A. M. Dokter, P. J. Blancher, J. R. Sauer, A. C. Smith, P. A. Smith, J. C. Stanton, A. Panjabi, L. Helf, M. Parr, and P. P. Marra. 2019. "Decline of the North American Avifauna." *Science* 10.1126/science.aaw1313.
- Royle, J. A., and R. M. Dorazio. 2008. *Hierarchical Modeling and Inference in Ecology*. Boston, MA: Academic Press.
- Soots, R. F., Jr., and M. C. Landin. 1978, *Development and Management of Avian Habitat on Dredged Material Islands*. DMRP Technical Report DS-78-18. Vicksburg, MS: ERDC.
- USACE and USEPA (U.S. Environmental Protection Agency). 2003. *Great Lakes Confined Disposal Facilities*. Washington, DC: USACE and USEPA. [http://www.lrd.usace.army.mil/Portals/73/docs/Navigation/GL-CDF/GL\\_CDF.pdf](http://www.lrd.usace.army.mil/Portals/73/docs/Navigation/GL-CDF/GL_CDF.pdf).
- US Department of Commerce. 1988. *City of Portsmouth, Virginia: Coastal Zone Management Plan*. Final Report. Charleston, SC: National Oceanic and Atmospheric Administration, Coastal Services. <https://www.govinfo.gov/content/pkg/CZIC-ht-393-v8-p67-1988/html/CZIC-ht-393-v8-p67-1988.htm>

- US Fish and Wildlife Service. 1998. *North American Waterfowl Management Plan: Expanding the Vision, 1998 Update*. Arlington, VA: US Department of the Interior, Fish and Wildlife Service.
- Zipkin, E. F., A. DeWan, and J. A. Royle. 2009. "Impacts of Forest Fragmentation on Species Richness: A Hierarchical Approach to Community Modelling." *Journal of Applied Ecology* 46:815–822.

## Acronyms and Abbreviations

| Term   | Definition                                            |
|--------|-------------------------------------------------------|
| ACJV   | Atlantic Coast Joint Venture                          |
| BCR    | (Southeastern Coastal Plain) Bird Conservation Region |
| CDF    | Confined Disposal Facility                            |
| CIDMMA | Craney Island Dredged Material Management Area        |
| CVWO   | Coastal Virginia Wildlife Observatory                 |
| DMCA   | Dredged Material Containment Areas                    |
| EL     | Environmental Laboratory                              |
| ERDC   | Engineer Research and Development Center              |
| ESA    | U.S. Endangered Species Act                           |
| EST    | Eastern Standard Time                                 |
| EWN®   | Engineering With Nature                               |
| ICWW   | Intracoastal Waterway System                          |
| LTBMP  | Long-Term Bird Management Plan                        |
| NABCI  | North American Bird Conservation Initiative           |
| PIF    | Partners In Flight                                    |
| SHNP   | Savannah Harbor Navigation Project                    |
| USACE  | U.S. Army Corps of Engineers                          |
| USEPA  | U.S. Environmental Protection Agency                  |
| USFWS  | U.S. Fish and Wildlife Service                        |

## Unit Conversion Factors

| Multiply              | By             | To Obtain       |
|-----------------------|----------------|-----------------|
| acres                 | 4,046.873      | square meters   |
| acre-feet             | 1,233.5        | cubic meters    |
| angstroms             | 0.1            | nanometers      |
| atmosphere (standard) | 101.325        | kilopascals     |
| cubic feet            | 0.02831685     | cubic meters    |
| cubic inches          | 1.6387064 E-05 | cubic meters    |
| cubic yards           | 0.7645549      | cubic meters    |
| degrees Fahrenheit    | $(F-32)/1.8$   | degrees Celsius |
| feet                  | 0.3048         | meters          |
| hectares              | 1.0 E+04       | square meters   |
| inches                | 0.0254         | meters          |
| miles (U.S. statute)  | 1,609.347      | meters          |
| square feet           | 0.09290304     | square meters   |
| square inches         | 6.4516 E-04    | square meters   |
| square miles          | 2.589998 E+06  | square meters   |
| square yards          | 0.8361274      | square meters   |
| yards                 | 0.9144         | meters          |

---

## **Appendix A: Table of Counts of Birds Detected During Seasonal Surveys on the Craney Island Dredged Material Management Area, USACE Norfolk District, Portsmouth, Virginia, 2008-2020**

| Bird Species                     | Scientific Name                | Species Code | 2008 | 2009  | 2010  | 2011   | 2012  | 2013  | 2014  | 2015   | 2016   | 2017   | 2018   | 2019  | 2020  |
|----------------------------------|--------------------------------|--------------|------|-------|-------|--------|-------|-------|-------|--------|--------|--------|--------|-------|-------|
| Snow Goose <sup>W</sup>          | <i>Anser caerulescens</i>      | SNGO         | 0    | 0     | 0     | 0      | 0     | 0     | 5     | 0      | 0      | 1      | 0      | 0     | 0     |
| Brant <sup>W</sup>               | <i>Branta Bernicia</i>         | BRAN         | 0    | 0     | 0     | 1      | 0     | 0     | 0     | 0      | 5      | 3      | 0      | 4     | 0     |
| Canada Goose <sup>R</sup>        | <i>Branta canadensis</i>       | CAGO         | 0    | 231   | 1,756 | 417    | 203   | 213   | 1,442 | 827    | 1,770  | 982    | 1,434  | 1,431 | 521   |
| Tundra Swan <sup>W</sup>         | <i>Cygnus columbianus</i>      | TUSW         | 0    | 0     | 0     | 10     | 4     | 308   | 155   | 43     | 137    | 25     | 28     | 175   | 0     |
| Mute Swan <sup>T</sup>           | <i>Cygnus olor</i>             | MUSW         | 0    | 0     | 0     | 1      | 0     | 0     | 0     | 0      | 0      | 0      | 0      | 0     | 0     |
| Wood Duck <sup>YR</sup>          | <i>Aix sponsa</i>              | WODU         | 0    | 0     | 6     | 0      | 0     | 3     | 0     | 2      | 5      | 7      | 18     | 0     | 0     |
| Mallard <sup>R</sup>             | <i>Anas platyrhynchos</i>      | MALL         | 29   | 1,210 | 2,559 | 2,770  | 1,465 | 2,741 | 6,578 | 6,290  | 7,741  | 8,027  | 4,264  | 4,569 | 1,829 |
| Mottled Duck <sup>T</sup>        | <i>Anus fulvigula</i>          | MODU         | 0    | 0     | 0     | 0      | 0     | 0     | 0     | 0      | 0      | 0      | 2      | 0     | 0     |
| American Black Duck <sup>W</sup> | <i>Anas rubripes</i>           | AMBD         | 2    | 83    | 158   | 392    | 149   | 295   | 295   | 480    | 890    | 472    | 350    | 328   | 214   |
| Gadwall <sup>W</sup>             | <i>Mareca strepera</i>         | GADW         | 0    | 23    | 135   | 367    | 229   | 512   | 781   | 1,749  | 4,665  | 4,034  | 2,262  | 2,511 | 2,055 |
| Green-winged Teal <sup>W</sup>   | <i>Anas crecca</i>             | GWTE         | 0    | 522   | 955   | 1,138  | 729   | 544   | 2,699 | 3,095  | 13,271 | 10,234 | 3,306  | 5,544 | 2,706 |
| American Wigeon <sup>W</sup>     | <i>Mareca Americana</i>        | AMWI         | 0    | 82    | 231   | 311    | 242   | 473   | 392   | 784    | 723    | 682    | 463    | 270   | 278   |
| Eurasian Wigeon <sup>W</sup>     | <i>Anas penelope</i>           | EUWI         | 0    | 0     | 2     | 0      | 0     | 9     | 12    | 13     | 5      | 6      | 4      | 1     | 3     |
| Northern Pintail <sup>W</sup>    | <i>Anas acuta</i>              | NOPI         | 0    | 39    | 7     | 59     | 2,640 | 102   | 245   | 114    | 592    | 265    | 34     | 31    | 33    |
| Northern Shoveler <sup>W</sup>   | <i>Spatula clypeata</i>        | NOSH         | 42   | 3,260 | 1,397 | 10,802 | 9,693 | 9,218 | 8,392 | 12,668 | 21,983 | 34,944 | 20,962 | 8,067 | 4,048 |
| Blue-winged Teal <sup>M</sup>    | <i>Spatula discors</i>         | BWTE         | 1    | 718   | 40    | 267    | 463   | 252   | 79    | 168    | 560    | 290    | 147    | 38    | 36    |
| Canvasback <sup>W</sup>          | <i>Aythya valisineria</i>      | CANV         | 0    | 31    | 17    | 333    | 229   | 5,935 | 1,963 | 4,200  | 2,912  | 3,490  | 4,108  | 1,001 | 726   |
| Redhead <sup>W</sup>             | <i>Aythya americana</i>        | REDH         | 0    | 0     | 0     | 1      | 0     | 2     | 15    | 23     | 1      | 3      | 2      | 5     | 0     |
| Ring-necked Duck <sup>W</sup>    | <i>Aythya colaris</i>          | RNDU         | 0    | 0     | 18    | 12     | 6     | 83    | 21    | 36     | 114    | 45     | 33     | 113   | 2     |
| Greater Scaup <sup>W</sup>       | <i>Aythya marila</i>           | GRSC         | 0    | 0     | 0     | 0      | 0     | 6     | 1     | 0      | 0      | 1      | 5      | 26    | 2     |
| Lesser Scaup <sup>W</sup>        | <i>Aythya affinis</i>          | LESC         | 0    | 8     | 5     | 32     | 23    | 446   | 90    | 99     | 39     | 38     | 66     | 62    | 3     |
| Long-tailed Duck <sup>T</sup>    | <i>Clangula hyemalis</i>       | LTDU         | 0    | 0     | 0     | 0      | 0     | 0     | 0     | 0      | 0      | 0      | 0      | 1     | 0     |
| Surf Scoter <sup>W</sup>         | <i>Melanitta perspicillata</i> | SUSC         | 0    | 0     | 21    | 79     | 13    | 104   | 55    | 28     | 22     | 40     | 7      | 143   | 19    |
| Black Scoter <sup>W</sup>        | <i>Melanitta Americana</i>     | BLSC         | 0    | 0     | 0     | 1      | 2     | 11    | 8     | 4      | 1      | 1      | 47     | 5     | 2     |

|                                        |                              |      |    |     |       |       |       |        |        |        |        |        |        |        |       |
|----------------------------------------|------------------------------|------|----|-----|-------|-------|-------|--------|--------|--------|--------|--------|--------|--------|-------|
| White-winged Scoter <sup>W</sup>       | <i>Melanitta fusca</i>       | WWSC | 0  | 0   | 0     | 0     | 2     | 0      | 0      | 0      | 0      | 0      | 1      | 3      | 0     |
| Bufflehead <sup>W</sup>                | <i>Bucephala albeola</i>     | BUFF | 0  | 112 | 118   | 489   | 303   | 911    | 556    | 840    | 1,063  | 1,418  | 983    | 985    | 488   |
| Common Goldeneye <sup>W</sup>          | <i>Bucephala clangula</i>    | COGO | 0  | 0   | 0     | 0     | 0     | 2      | 0      | 0      | 3      | 2      | 3      | 8      | 0     |
| Hooded Merganser <sup>W</sup>          | <i>Lophodytes cucullatus</i> | HOME | 0  | 4   | 4     | 58    | 131   | 76     | 15     | 117    | 108    | 128    | 118    | 101    | 41    |
| Common Merganser <sup>W</sup>          | <i>Mergus merganser</i>      | COME | 0  | 0   | 0     | 0     | 0     | 0      | 0      | 5      | 0      | 3      | 0      | 4      | 0     |
| Red-breasted Merganser <sup>W</sup>    | <i>Mergus serrator</i>       | RBME | 0  | 125 | 28    | 169   | 638   | 403    | 396    | 851    | 1,936  | 5,334  | 877    | 4,750  | 3,889 |
| Ruddy Duck <sup>W</sup>                | <i>Oxyura jamaicensis</i>    | RUDU | 8  | 458 | 2,417 | 7,852 | 5,325 | 14,160 | 11,702 | 22,600 | 45,652 | 29,003 | 23,227 | 15,388 | 1,900 |
| Northern Bobwhite <sup>R</sup>         | <i>Colinus virginianus</i>   | NOBO | 0  | 1   | 3     | 13    | 7     | 7      | 7      | 3      | 8      | 2      | 15     | 14     | 1     |
| Wild Turkey <sup>YR</sup>              | <i>Meleagris gallopavo</i>   | WITU | 0  | 0   | 1     | 0     | 0     | 0      | 6      | 0      | 11     | 7      | 13     | 12     | 13    |
| Pied-billed Grebe <sup>R</sup>         | <i>Podilymbus podiceps</i>   | PBGR | 0  | 3   | 19    | 24    | 71    | 15     | 121    | 43     | 42     | 45     | 12     | 21     | 6     |
| Horned Grebe <sup>W</sup>              | <i>Podiceps auritus</i>      | HOGR | 0  | 97  | 300   | 185   | 269   | 560    | 356    | 603    | 1,005  | 2,093  | 1,659  | 1,321  | 798   |
| Eared Grebe <sup>T</sup>               | <i>Podiceps nigricollis</i>  | EAGR | 0  | 0   | 0     | 0     | 2     | 0      | 5      | 0      | 0      | 0      | 0      | 1      | 0     |
| Red-necked Grebe <sup>W</sup>          | <i>Podiceps grisegena</i>    | RNGR | 0  | 0   | 0     | 0     | 0     | 1      | 5      | 1      | 0      | 0      | 0      | 0      | 0     |
| Rock Pigeon <sup>YR</sup>              | <i>Columba livia</i>         | ROPI | 18 | 35  | 224   | 422   | 178   | 266    | 142    | 235    | 94     | 243    | 267    | 98     | 18    |
| Mourning Dove <sup>R</sup>             | <i>Zenaida macroura</i>      | MODO | 46 | 426 | 667   | 900   | 760   | 1,535  | 1,351  | 999    | 1,297  | 1,099  | 1,309  | 1,879  | 298   |
| Yellow-billed Cuckoo <sup>N</sup>      | <i>Coccyzus americanus</i>   | YBCU | 0  | 0   | 4     | 1     | 0     | 3      | 1      | 4      | 9      | 3      | 12     | 10     | 0     |
| Common Nighthawk <sup>N</sup>          | <i>Chordeiles minor</i>      | CONI | 8  | 5   | 6     | 4     | 12    | 60     | 12     | 7      | 33     | 65     | 25     | 20     | 3     |
| Chimney Swift <sup>M</sup>             | <i>Chaetura pelagica</i>     | CHSW | 5  | 92  | 264   | 227   | 111   | 214    | 953    | 733    | 1,162  | 2,063  | 1,149  | 2,547  | 38    |
| Ruby-throated Hummingbird <sup>S</sup> | <i>Archilochus colubris</i>  | RTHU | 0  | 1   | 12    | 7     | 5     | 17     | 20     | 16     | 0      | 26     | 51     | 42     | 11    |
| Sora <sup>SB</sup>                     | <i>Porzana Carolina</i>      | SORA | 0  | 0   | 0     | 3     | 4     | 1      | 11     | 2      | 4      | 5      | 0      | 1      | 5     |
| Clapper Rail <sup>YR</sup>             | <i>Rallus longirostris</i>   | CLRA | 0  | 0   | 0     | 1     | 1     | 1      | 0      | 0      | 8      | 19     | 12     | 4      | 0     |
| King Rail <sup>YR</sup>                | <i>Rallus elegans</i>        | KIRA | 0  | 0   | 0     | 0     | 0     | 0      | 1      | 0      | 0      | 0      | 0      | 0      | 0     |
| Virginia Rail <sup>YR</sup>            | <i>Rallus limicola</i>       | VIRA | 0  | 0   | 0     | 0     | 1     | 0      | 0      | 0      | 0      | 0      | 3      | 1      | 0     |
| Common Gallinule <sup>S</sup>          | <i>Gallinula galeata</i>     | COGA | 0  | 0   | 0     | 0     | 0     | 0      | 0      | 0      | 0      | 0      | 0      | 1      | 0     |

|                                      |                                |          |       |       |       |        |        |        |        |        |        |        |        |        |       |
|--------------------------------------|--------------------------------|----------|-------|-------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|
| American Coot <sup>R</sup>           | <i>Fulica Americana</i>        | AMCO     | 0     | 0     | 26    | 20     | 25     | 3      | 117    | 46     | 70     | 40     | 2      | 23     | 0     |
| Black-necked Stilt <sup>SB</sup>     | <i>Himantopus mexicanus</i>    | BNST     | 3     | 37    | 67    | 310    | 208    | 384    | 612    | 609    | 1,090  | 1,126  | 791    | 594    | 257   |
| American Avocet <sup>R</sup>         | <i>Recurvirostra americana</i> | AMAV     | 37    | 364   | 221   | 473    | 292    | 376    | 1,304  | 1,512  | 2,603  | 6,412  | 8,538  | 1,696  | 740   |
| American Oystercatcher <sup>YR</sup> | <i>Haematopus palliates</i>    | AMOY     | 0     | 11    | 4     | 9      | 1      | 10     | 9      | 16     | 7      | 4      | 7      | 1      | 18    |
| Black-bellied Plover <sup>W</sup>    | <i>Pluvialis squatarola</i>    | BBPL     | 27    | 116   | 60    | 100    | 74     | 212    | 116    | 746    | 253    | 331    | 179    | 106    | 13    |
| American Golden Plover <sup>M</sup>  | <i>Pluvialis dominica</i>      | AMGP     | 0     | 0     | 1     | 10     | 4      | 5      | 5      | 2      | 10     | 4      | 0      | 2      | 0     |
| Killdeer <sup>R</sup>                | <i>Charadrius vociferous</i>   | KILL     | 63    | 338   | 597   | 730    | 509    | 1,138  | 1,221  | 933    | 1,218  | 1,846  | 967    | 1,258  | 371   |
| Wilson's Plover <sup>T</sup>         | <i>Charadrius wilsonia</i>     | WIPL     | 0     | 0     | 1     | 0      | 0      | 0      | 0      | 0      | 0      | 0      | 0      | 0      | 0     |
| Semipalmated Plover <sup>M</sup>     | <i>Charadrius semipalmatus</i> | SEPL     | 27    | 597   | 767   | 415    | 374    | 850    | 1,775  | 1,725  | 2,337  | 2,750  | 1,142  | 1,421  | 380   |
| Piping Plover <sup>S</sup>           | <i>Charadrius melodus</i>      | PIPL     | 0     | 0     | 0     | 0      | 1      | 3      | 1      | 1      | 5      | 4      | 2      | 1      | 0     |
| Upland Sandpiper <sup>M</sup>        | <i>Bartramia longicauda</i>    | UPSA     | 0     | 1     | 1     | 0      | 2      | 0      | 1      | 0      | 0      | 0      | 1      | 0      | 0     |
| Whimbrel <sup>W</sup>                | <i>Numenius phaeopus</i>       | WHIM     | 3     | 0     | 0     | 0      | 0      | 8      | 1      | 6      | 1      | 3      | 4      | 4      | 1     |
| Hudsonian Godwit <sup>M</sup>        | <i>Limosa haemastica</i>       | HUGO     | 1     | 0     | 0     | 0      | 3      | 0      | 0      | 1      | 4      | 0      | 0      | 0      | 0     |
| Marbled Godwit <sup>W</sup>          | <i>Limosa fedoa</i>            | MAGO     | 2     | 1     | 0     | 7      | 0      | 7      | 9      | 3      | 6      | 1      | 0      | 1      | 0     |
| Ruddy Turnstone <sup>W</sup>         | <i>Arenaria interpres</i>      | RUTU     | 17    | 46    | 86    | 92     | 68     | 104    | 60     | 67     | 42     | 72     | 98     | 40     | 2     |
| Purple Sandpiper <sup>W</sup>        | <i>Calidris maritima</i>       | PUSA     | 0     | 0     | 0     | 8      | 0      | 0      | 0      | 0      | 0      | 0      | 2      | 1      | 0     |
| Calidris spp.                        | <i>Calidris spp.</i>           | CALIDRIS | 0     | 6,636 |       | 8,109  | 3,211  | 2,751  | 10,622 | 11,468 | 33,087 | 13,318 | 4,580  | 1,119  | 2,085 |
| Red Knot <sup>M</sup>                | <i>Calidris canutus</i>        | REKN     | 0     | 2     | 0     | 2      | 4      | 0      | 2      | 13     | 38     | 0      | 1      | 1      | 2     |
| Sanderling <sup>W</sup>              | <i>Calidris alba</i>           | SAND     | 89    | 462   | 311   | 848    | 246    | 819    | 412    | 923    | 1,138  | 899    | 795    | 691    | 425   |
| Semipalmated Sandpiper <sup>M</sup>  | <i>Calidris pusilla</i>        | SESA     | 3,135 | 3,391 | 7,843 | 21,572 | 30,015 | 31,634 | 35,371 | 19,992 | 13,447 | 27,468 | 38,671 | 23,869 | 5,092 |
| Western Sandpiper <sup>M</sup>       | <i>Calidris mauri</i>          | WESA     | 25    | 1,743 | 107   | 591    | 271    | 640    | 2,457  | 1,039  | 729    | 715    | 480    | 935    | 81    |
| Least Sandpiper <sup>W</sup>         | <i>Calidris minutilla</i>      | LESA     | 545   | 1,790 | 2,645 | 1,685  | 1,267  | 1,279  | 1,134  | 1,379  | 3,129  | 1,526  | 666    | 641    | 649   |

|                                      |                                     |      |     |       |       |       |       |       |       |       |       |        |        |       |        |
|--------------------------------------|-------------------------------------|------|-----|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|-------|--------|
| White-rumped Sandpiper <sup>M</sup>  | <i>Calidris fuscicollis</i>         | WRSA | 0   | 23    | 6     | 55    | 54    | 32    | 34    | 45    | 50    | 63     | 41     | 13    | 7      |
| Baird's Sandpiper <sup>T</sup>       | <i>Calidris bairdii</i>             | BASA | 0   | 0     | 0     | 0     | 0     | 2     | 0     | 3     | 2     | 1      | 0      | 0     | 2      |
| Dunlin <sup>W</sup>                  | <i>Calidris alpina</i>              | DUNL | 113 | 2,607 | 1,369 | 2,447 | 1,293 | 5,465 | 3,869 | 5,504 | 8,923 | 14,790 | 15,297 | 4,439 | 2,818  |
| Curlew Sandpiper <sup>T</sup>        | <i>Calidris ferruginea</i>          | CUSA | 0   | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 0      | 0      | 0     | 0      |
| Stilt Sandpiper <sup>M</sup>         | <i>Calidris himantopus</i>          | STSA | 12  | 260   | 60    | 99    | 127   | 97    | 452   | 541   | 150   | 390    | 585    | 123   | 109    |
| Pectoral Sandpiper <sup>M</sup>      | <i>Calidris melanotos</i>           | PESA | 1   | 84    | 21    | 28    | 27    | 28    | 20    | 70    | 117   | 68     | 22     | 19    | 69     |
| Buff-breasted Sandpiper <sup>T</sup> | <i>Calidris subruficollis</i>       | BBSA | 0   | 1     | 0     | 0     | 0     | 1     | 0     | 0     | 10    | 6      | 0      | 0     | 0      |
| Ruff <sup>T</sup>                    | <i>Calidris pugnax</i>              | RUFF | 0   | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 1      | 0      | 0     | 0      |
| Short-billed Dowitcher <sup>M</sup>  | <i>Limnodromus griseus</i>          | SBDO | 35  | 348   | 239   | 120   | 138   | 339   | 3,047 | 2,887 | 1,862 | 1,034  | 1,295  | 522   | 471    |
| Long-billed Dowitcher <sup>W</sup>   | <i>Limnodromus scolopaceus</i>      | LBDO | 0   | 0     | 1     | 3     | 0     | 0     | 0     | 42    | 14    | 5      | 6      | 6     | 0      |
| Wilson's Snipe <sup>W</sup>          | <i>Gallinago delicata</i>           | WISN | 0   | 0     | 2     | 0     | 1     | 13    | 4     | 4     | 14    | 17     | 7      | 7     | 1      |
| American Woodcock <sup>W</sup>       | <i>Scolopax minor</i>               | AMWO | 0   | 0     | 0     | 0     | 0     | 3     | 0     | 0     | 0     | 0      | 0      | 0     | 1      |
| Spotted Sandpiper <sup>SB</sup>      | <i>Actitis macularia</i>            | SPSA | 34  | 99    | 123   | 144   | 211   | 232   | 218   | 137   | 349   | 227    | 261    | 181   | 51     |
| Solitary Sandpiper <sup>M</sup>      | <i>Tringa solitaria</i>             | SOSA | 0   | 0     | 9     | 3     | 0     | 5     | 2     | 0     | 5     | 4      | 1      | 1     | 0      |
| Willet <sup>R</sup>                  | <i>Tringa semipalmata</i>           | WILL | 17  | 35    | 77    | 91    | 62    | 121   | 133   | 141   | 98    | 250    | 217    | 114   | 35     |
| Greater Yellowlegs <sup>W</sup>      | <i>Tringa melanoleuca</i>           | GRYE | 15  | 93    | 120   | 212   | 157   | 383   | 133   | 409   | 512   | 617    | 333    | 295   | 44     |
| Lesser Yellowlegs <sup>W</sup>       | <i>Tringa flavipes</i>              | LEYE | 483 | 2,200 | 2,350 | 2,183 | 1,566 | 1,973 | 4,838 | 3,149 | 5,301 | 7,402  | 5,371  | 3,076 | 1,018  |
| Wilson's Phalarope <sup>M</sup>      | <i>Phalaropus tricolor</i>          | WIPH | 0   | 19    | 22    | 24    | 34    | 36    | 37    | 67    | 55    | 91     | 22     | 21    | 10     |
| Red-necked Phalarope <sup>M</sup>    | <i>Phalaropus lobatus</i>           | RNPH | 13  | 30    | 15    | 51    | 13    | 16    | 43    | 34    | 376   | 143    | 14     | 0     | 12     |
| Red Phalarope <sup>W</sup>           | <i>Phalaropus fulicarius</i>        | REPH | 0   | 0     | 0     | 0     | 0     | 0     | 1     | 1     | 4     | 9      | 0      | 0     | 0      |
| Bonaparte's Gull <sup>W</sup>        | <i>Chroicocephalus Philadelphia</i> | BOGU | 0   | 0     | 0     | 4     | 35    | 5     | 30    | 3     | 12    | 12     | 11     | 38    | 6      |
| Laughing Gull <sup>R</sup>           | <i>Leucophaeus atricilla</i>        | LAGU | 388 | 2,273 | 1,399 | 5,059 | 870   | 1,272 | 1,281 | 5,178 | 2,042 | 3,807  | 1,108  | 586   | 373    |
| Ring-billed Gull <sup>W</sup>        | <i>Larus delawarensis</i>           | RBGU | 22  | 543   | 660   | 1,138 | 581   | 2,379 | 1,665 | 2,343 | 2,849 | 2,680  | 2,920  | 9,351 | 10,905 |
| Herring Gull <sup>YR</sup>           | <i>Larus argentatus</i>             | HEGU | 134 | 1,319 | 1,029 | 2,468 | 759   | 1,610 | 1,740 | 1,679 | 3,833 | 3,334  | 3,173  | 1,614 | 5,529  |

|                                       |                                  |      |     |       |       |       |       |       |       |       |        |        |       |       |       |
|---------------------------------------|----------------------------------|------|-----|-------|-------|-------|-------|-------|-------|-------|--------|--------|-------|-------|-------|
| Iceland Gull <sup>I</sup>             | <i>Larus glaucooides</i>         | ICGU | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 2      | 0      | 0     | 0     | 0     |
| Glaucous Gull <sup>W</sup>            | <i>Larus hyperboreus</i>         | GLGU | 0   | 0     | 0     | 8     | 0     | 0     | 0     | 0     | 0      | 0      | 0     | 0     | 0     |
| Lesser Black-backed Gull <sup>W</sup> | <i>Larus fuscus</i>              | LBBG | 2   | 49    | 4     | 56    | 24    | 32    | 55    | 35    | 56     | 38     | 45    | 39    | 86    |
| Great Black-backed Gull <sup>W</sup>  | <i>Larus marinus</i>             | GBBG | 60  | 1,098 | 670   | 631   | 305   | 632   | 699   | 474   | 559    | 685    | 529   | 424   | 254   |
| Least Tern <sup>N</sup>               | <i>Sternula antillarum</i>       | LETE | 309 | 1,877 | 2,213 | 4,143 | 1,772 | 2,874 | 1,978 | 2,071 | 1,511  | 2,545  | 1,484 | 1,827 | 467   |
| Black Tern <sup>M</sup>               | <i>Chidonias niger</i>           | BLTE | 1   | 227   | 1     | 638   | 17    | 49    | 49    | 44    | 40     | 102    | 4     | 7     | 0     |
| Common Tern <sup>M</sup>              | <i>Sterna hirundo</i>            | COTE | 26  | 11    | 9     | 70    | 124   | 7     | 14    | 22    | 81     | 9      | 3     | 2     | 1     |
| Forster's Tern <sup>W</sup>           | <i>Sterna forsteri</i>           | FOTE | 0   | 16    | 12    | 80    | 19    | 46    | 123   | 116   | 74     | 85     | 30    | 12    | 48    |
| Gull-billed Tern <sup>SB</sup>        | <i>Gelochelidon nilotica</i>     | GBTE | 18  | 24    | 52    | 73    | 61    | 58    | 14    | 35    | 72     | 156    | 101   | 88    | 20    |
| Sandwich Tern <sup>SB</sup>           | <i>Thalasseus sandvicensis</i>   | SATE | 0   | 0     | 2     | 4     | 0     | 2     | 33    | 8     | 2      | 0      | 0     | 2     | 1     |
| Royal Tern <sup>SB</sup>              | <i>Thalasseus maximus</i>        | ROYT | 159 | 1,170 | 714   | 358   | 426   | 2,142 | 2,144 | 1,145 | 2,502  | 1,665  | 1,632 | 3,537 | 592   |
| Caspian Tern <sup>M</sup>             | <i>Hydroprogne caspia</i>        | CATE | 54  | 500   | 184   | 226   | 204   | 494   | 145   | 249   | 190    | 100    | 310   | 150   | 160   |
| Black Skimmer <sup>SB</sup>           | <i>Rynchops niger</i>            | BLSK | 1   | 40    | 38    | 9     | 5     | 33    | 27    | 203   | 47     | 96     | 40    | 24    | 6     |
| Red-throated Loon <sup>W</sup>        | <i>Gavia Stellata</i>            | RTLO | 0   | 21    | 2     | 5     | 28    | 38    | 115   | 186   | 87     | 291    | 1,195 | 10    | 14    |
| Common Loon <sup>W</sup>              | <i>Gavia immer</i>               | COLO | 0   | 0     | 6     | 7     | 12    | 35    | 20    | 50    | 49     | 40     | 60    | 65    | 8     |
| Northern Gannet <sup>W</sup>          | <i>Morus bassanus</i>            | NOGA | 0   | 3     | 0     | 0     | 21    | 238   | 611   | 45    | 51     | 41     | 19    | 10    | 253   |
| Great Cormorant <sup>I</sup>          | <i>Phalacrocorax carbo</i>       | GRCO | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 2      | 0      | 0     | 0     | 0     |
| Double-crested Cormorant <sup>R</sup> | <i>Phalacrocorax auritus</i>     | DCCO | 45  | 607   | 1,037 | 1,706 | 434   | 2,085 | 5,447 | 3,781 | 13,804 | 26,085 | 5,469 | 7,018 | 8,498 |
| American White Pelican <sup>YR</sup>  | <i>Pelecanus erythrorhynchos</i> | AWPE | 0   | 14    | 0     | 0     | 0     | 0     | 0     | 0     | 6      | 0      | 0     | 0     | 0     |
| Brown Pelican <sup>R</sup>            | <i>Pelecanus occidentalis</i>    | BRPE | 32  | 1,416 | 138   | 606   | 269   | 526   | 475   | 914   | 1,013  | 1,567  | 1,016 | 825   | 192   |
| American Bittern <sup>W</sup>         | <i>Botaurus lentiginosus</i>     | AMBI | 0   | 0     | 0     | 0     | 0     | 0     | 3     | 2     | 2      | 1      | 0     | 0     | 0     |
| Least Bittern <sup>SB</sup>           | <i>Ixobrychus exilis</i>         | LEBI | 0   | 0     | 0     | 0     | 0     | 0     | 16    | 0     | 1      | 9      | 2     | 1     | 0     |

|                                          |                                 |      |    |     |     |     |     |     |     |     |     |     |     |     |    |
|------------------------------------------|---------------------------------|------|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|----|
| Great Blue Heron <sup>R</sup>            | <i>Ardea herodias</i>           | GBHE | 16 | 110 | 67  | 162 | 129 | 144 | 118 | 282 | 352 | 476 | 405 | 460 | 82 |
| Cattle Egret <sup>B</sup>                | <i>Bubulcus ibis</i>            | CAEG | 0  | 0   | 0   | 1   | 1   | 0   | 0   | 2   | 1   | 0   | 0   | 0   | 0  |
| Snowy Egret <sup>R</sup>                 | <i>Egretta thula</i>            | SNEG | 6  | 156 | 31  | 249 | 35  | 51  | 19  | 722 | 387 | 844 | 540 | 357 | 56 |
| Little Egret <sup>T</sup>                | <i>Egretta garzetta</i>         | LIEG | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0  |
| Great Egret <sup>R</sup>                 | <i>Ardea alba</i>               | GREG | 29 | 519 | 153 | 372 | 227 | 220 | 144 | 769 | 785 | 630 | 535 | 369 | 0  |
| Tricolored Heron <sup>YR</sup>           | <i>Egretta tricolor</i>         | TRHE | 0  | 0   | 0   | 7   | 0   | 1   | 2   | 33  | 5   | 6   | 14  | 6   | 0  |
| Little Blue Heron <sup>R</sup>           | <i>Egretta caerulea</i>         | LBHE | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 2   | 2   | 2   | 3   | 1   | 0  |
| Reddish Egret <sup>T</sup>               | <i>Egretta rufescens</i>        | REEG | 1  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0  |
| Green Heron <sup>SB</sup>                | <i>Butorides virescens</i>      | GRHE | 1  | 1   | 15  | 9   | 5   | 8   | 4   | 17  | 31  | 41  | 58  | 39  | 10 |
| Black-crowned Night-Heron <sup>R</sup>   | <i>Nycticorax nycticorax</i>    | BCNH | 0  | 0   | 0   | 0   | 3   | 2   | 1   | 34  | 36  | 79  | 58  | 34  | 5  |
| Yellow-crowned Night-Heron <sup>SB</sup> | <i>Nyctanassa violacea</i>      | YCNH | 5  | 5   | 0   | 7   | 2   | 3   | 6   | 25  | 8   | 13  | 23  | 10  | 0  |
| Glossy Ibis <sup>S</sup>                 | <i>Plegadis falcinellus</i>     | GLIB | 0  | 3   | 3   | 5   | 3   | 7   | 16  | 23  | 44  | 11  | 4   | 21  | 5  |
| White Ibis <sup>M</sup>                  | <i>Eudocimus albus</i>          | WHIB | 0  | 1   | 0   | 0   | 2   | 0   | 0   | 64  | 10  | 1   | 4   | 0   | 0  |
| Roseate Spoonbill <sup>T</sup>           | <i>Platalea ajaja</i>           | ROSP | 0  | 8   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 45  | 0   | 0  |
| Turkey Vulture <sup>YR</sup>             | <i>Cathartes aura</i>           | TUVU | 7  | 23  | 46  | 76  | 59  | 71  | 70  | 129 | 89  | 119 | 144 | 130 | 32 |
| Black Vulture <sup>YR</sup>              | <i>Coragyps atratus</i>         | BLVU | 0  | 0   | 0   | 3   | 0   | 2   | 4   | 5   | 3   | 12  | 0   | 2   | 2  |
| Osprey <sup>R</sup>                      | <i>Pandion haliaetus</i>        | OSPR | 45 | 124 | 195 | 151 | 121 | 206 | 170 | 245 | 310 | 334 | 356 | 333 | 43 |
| Mississippi Kite <sup>N</sup>            | <i>Ictinia mississippiensis</i> | MIKI | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 2   | 0   | 0   | 0  |
| Bald Eagle <sup>YR</sup>                 | <i>Haliaeetus leucocephalus</i> | BAEA | 0  | 5   | 13  | 24  | 22  | 56  | 55  | 56  | 76  | 75  | 100 | 215 | 56 |
| Northern Harrier <sup>W</sup>            | <i>Circus hudsonius</i>         | NOHA | 0  | 5   | 20  | 80  | 43  | 81  | 69  | 105 | 116 | 162 | 126 | 144 | 53 |
| Sharp-shinned Hawk <sup>W</sup>          | <i>Accipiter striatus</i>       | SSHA | 1  |     | 2   | 3   | 2   | 3   | 4   | 3   | 5   | 5   | 4   | 5   | 2  |
| Cooper's Hawk <sup>S</sup>               | <i>Accipiter cooperii</i>       | COHA | 2  | 3   | 7   | 12  | 6   | 30  | 19  | 24  | 32  | 30  | 20  | 31  | 2  |
| Red-shouldered Hawk <sup>YR</sup>        | <i>Buteo lineatus</i>           | RSHA | 0  | 1   | 1   | 1   | 0   | 1   | 4   | 1   | 11  | 6   | 6   | 4   | 1  |
| Red-tailed Hawk <sup>YR</sup>            | <i>Buteo jamaicensis</i>        | RTHA | 0  | 34  | 48  | 81  | 114 | 145 | 120 | 142 | 131 | 154 | 123 | 177 | 46 |

|                                       |                                   |           |   |    |    |    |    |    |    |    |    |    |    |     |    |
|---------------------------------------|-----------------------------------|-----------|---|----|----|----|----|----|----|----|----|----|----|-----|----|
| Short-eared Owl <sup>W</sup>          | <i>Asio flammeus</i>              | SEOW      | 0 | 0  | 0  | 1  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 1   | 0  |
| Great Horned Owl <sup>R</sup>         | <i>Bubo virginianus</i>           | GHOW      | 0 | 0  | 0  | 0  | 0  | 2  | 0  | 0  | 0  | 0  | 1  | 0   | 0  |
| Snowy Owl <sup>W</sup>                | <i>Bubo scandiacus</i>            | SNOW      | 0 | 0  | 0  | 0  | 0  | 3  | 3  | 0  | 0  | 0  | 0  | 0   | 0  |
| Eastern Screech-Owl <sup>YR</sup>     | <i>Megascops asio</i>             | EASO      | 0 | 0  | 0  | 0  | 0  | 3  | 0  | 0  | 0  | 0  | 0  | 0   | 1  |
| Belted Kingfisher <sup>YR</sup>       | <i>Megascops asio</i>             | BEKI      | 0 | 4  | 5  | 13 | 15 | 21 | 10 | 13 | 24 | 47 | 32 | 21  | 7  |
| Red-headed Woodpecker <sup>YR</sup>   | <i>Melanerpes erythrocephalus</i> | RHOW      | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 5   | 0  |
| Red-bellied Woodpecker <sup>R</sup>   | <i>Melanerpes carolinus</i>       | RBWO      | 0 | 4  | 18 | 28 | 30 | 49 | 61 | 64 | 78 | 74 | 81 | 135 | 17 |
| Yellow-bellied Sapsucker <sup>W</sup> | <i>Sphyrapicus varius</i>         | YBSA      | 0 | 0  | 1  | 0  | 0  | 1  | 0  | 4  | 1  | 2  | 4  | 7   | 0  |
| Hairy Woodpecker <sup>YR</sup>        | <i>Dryobates villosus</i>         | HAWO      | 0 | 0  | 0  | 0  | 1  | 1  | 0  | 0  | 2  | 0  | 12 | 6   | 2  |
| Downy Woodpecker <sup>YR</sup>        | <i>Dryobates pubescens</i>        | DOWO      | 0 | 6  | 1  | 10 | 14 | 39 | 42 | 66 | 54 | 52 | 76 | 124 | 28 |
| Northern Flicker <sup>YR</sup>        | <i>Colaptes auratus</i>           | NOFL      | 0 | 5  | 0  | 17 | 11 | 22 | 18 | 43 | 25 | 36 | 47 | 68  | 24 |
| Pileated Woodpecker <sup>YR</sup>     | <i>Dryocopus pileatus</i>         | PIWO      | 0 | 0  | 1  | 2  | 3  | 6  | 7  | 19 | 16 | 21 | 18 | 20  | 2  |
| American Kestrel <sup>WR</sup>        | <i>Falco sparverius</i>           | AMKE      | 0 | 12 | 7  | 30 | 11 | 35 | 31 | 27 | 21 | 27 | 29 | 44  | 13 |
| Merlin <sup>W</sup>                   | <i>Falco columbarius</i>          | MERL      | 0 | 1  | 0  | 1  | 0  | 6  | 2  | 2  | 2  | 4  | 7  | 4   | 0  |
| Peregrine Falcon <sup>W</sup>         | <i>Falco peregrinus</i>           | PEFA      | 1 | 11 | 3  | 12 | 12 | 17 | 21 | 26 | 29 | 30 | 22 | 14  | 2  |
| Olive-sided Flycatcher <sup>T</sup>   | <i>Contopus cooperi</i>           | OSFL      | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 1   | 0  |
| Eastern Wood-Pewee <sup>N</sup>       | <i>Contopus virens</i>            | EWPE      | 0 | 0  | 0  | 0  | 0  | 2  | 0  | 1  | 1  | 2  | 0  | 4   | 0  |
| Empidonax spp. <sup>N</sup>           | <i>Empidonax spp.</i>             | EMPIDONAX | 0 | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 1  | 0  | 0  | 1   | 0  |
| Willow Flycatcher <sup>N</sup>        | <i>Empidonax traillii</i>         | WIFL      | 3 | 0  | 0  | 0  | 0  | 0  | 11 | 0  | 0  | 1  | 0  | 1   | 0  |
| Eastern Phoebe <sup>W</sup>           | <i>Sayornis phoebe</i>            | EAPH      | 0 | 2  | 0  | 0  | 3  | 4  | 2  | 8  | 10 | 20 | 12 | 21  | 5  |
| Great Crested Flycatcher <sup>N</sup> | <i>Myiarchus crinitus</i>         | GCFL      | 5 | 1  | 10 | 9  | 11 | 16 | 31 | 27 | 35 | 33 | 43 | 51  | 12 |
| Eastern Kingbird <sup>N</sup>         | <i>Tyrannus tyrannus</i>          | EAKI      | 5 | 7  | 30 | 27 | 36 | 79 | 62 | 44 | 80 | 80 | 61 | 96  | 11 |

|                                            |                                   |      |    |     |       |       |     |       |     |       |     |       |     |     |     |
|--------------------------------------------|-----------------------------------|------|----|-----|-------|-------|-----|-------|-----|-------|-----|-------|-----|-----|-----|
| Scissor-tailed Flycatcher <sup>T</sup>     | <i>Tyrannus forficatus</i>        | STFL | 0  | 0   | 0     | 0     | 0   | 0     | 0   | 0     | 0   | 0     | 0   | 1   | 0   |
| White-eyed Vireo <sup>R</sup>              | <i>Vireo griseus</i>              | WEVI | 0  | 0   | 3     | 2     | 0   | 3     | 4   | 2     | 3   | 6     | 6   | 4   | 1   |
| Blue-headed Vireo <sup>W</sup>             | <i>Vireo solitaries</i>           | BHVI | 0  | 0   | 0     | 0     | 1   | 0     | 0   | 1     | 1   | 0     | 0   | 2   | 0   |
| Red-eyed Vireo <sup>N</sup>                | <i>Vireo olivaceus</i>            | REVI | 0  | 1   | 0     | 0     | 3   | 5     | 1   | 3     | 3   | 3     | 5   | 1   | 6   |
| Warbling Vireo <sup>N</sup>                | <i>Vireo gilvus</i>               | WAVI | 0  | 0   | 2     | 0     | 0   | 0     | 0   | 0     | 3   | 9     | 0   | 0   | 0   |
| Blue Jay <sup>YR</sup>                     | <i>Cyanocitta cristata</i>        | BLJA | 3  | 44  | 94    | 128   | 86  | 188   | 179 | 250   | 242 | 243   | 217 | 353 | 62  |
| Fish Crow <sup>R</sup>                     | <i>Corvus ossifragus</i>          | FICR | 5  | 6   | 50    | 42    | 36  | 323   | 70  | 651   | 198 | 172   | 277 | 133 | 28  |
| American Crow <sup>YR</sup>                | <i>Corvus brachyrhynchos</i>      | AMCR | 6  | 76  | 79    | 72    | 52  | 145   | 167 | 337   | 286 | 291   | 302 | 515 | 162 |
| Horned Lark <sup>YR</sup>                  | <i>Eremophila alpestris</i>       | HOLA | 1  | 11  | 32    | 119   | 51  | 102   | 27  | 68    | 165 | 88    | 33  | 47  | 6   |
| Purple Martin <sup>N</sup>                 | <i>Progne subis</i>               | PUMA | 15 | 82  | 132   | 159   | 95  | 122   | 146 | 94    | 182 | 227   | 276 | 167 | 45  |
| Tree Swallow <sup>W</sup>                  | <i>Agelaius bicolor</i>           | TRSW | 0  | 29  | 39    | 66    | 21  | 955   | 244 | 2,483 | 407 | 630   | 238 | 232 | 7   |
| Barn Swallow <sup>S</sup>                  | <i>Hirundo rustica</i>            | BARS | 89 | 565 | 1,965 | 1,088 | 345 | 1,052 | 971 | 698   | 834 | 1,491 | 609 | 480 | 129 |
| Northern Rough-winged Swallow <sup>N</sup> | <i>Stelgidopteryx serripennis</i> | NRWS | 2  | 3   | 7     | 21    | 2   | 33    | 19  | 35    | 33  | 51    | 68  | 49  | 1   |
| Bank Swallow <sup>S</sup>                  | <i>Riparia riparia</i>            | BANS | 0  | 8   | 51    | 27    | 11  | 11    | 10  | 10    | 28  | 45    | 2   | 2   | 0   |
| Cliff Swallow <sup>M</sup>                 | <i>Petrochelidon pyrrhonota</i>   | CLSW | 0  | 0   | 36    | 2     | 0   | 13    | 2   | 0     | 1   | 1     | 0   | 1   | 0   |
| Cave Swallow <sup>T</sup>                  | <i>Petrochelidon fulva</i>        | CASW | 0  | 0   | 11    | 0     | 0   | 0     | 0   | 24    | 0   | 1     | 0   | 0   | 0   |
| Tufted Titmouse <sup>YR</sup>              | <i>Baeolophus bicolor</i>         | TUTI | 0  | 6   | 10    | 18    | 17  | 36    | 37  | 40    | 72  | 127   | 138 | 147 | 35  |
| Carolina Chickadee <sup>R</sup>            | <i>Poecile carolinensis</i>       | CACH | 0  | 10  | 49    | 78    | 57  | 123   | 137 | 168   | 200 | 249   | 255 | 335 | 58  |
| Red-breasted Nuthatch <sup>W</sup>         | <i>Sitta canadensis</i>           | RBNU | 0  | 0   | 1     | 0     | 4   | 1     | 0   | 0     | 0   | 1     | 3   | 3   | 0   |
| White-breasted Nuthatch <sup>YR</sup>      | <i>Sitta carolinensis</i>         | WBNU | 0  | 0   | 1     | 0     | 4   | 0     | 1   | 0     | 0   | 1     | 0   | 0   | 0   |
| Brown-headed Nuthatch <sup>YR</sup>        | <i>Sitta pusilla</i>              | BHNU | 0  | 1   | 4     | 3     | 2   | 6     | 16  | 5     | 3   | 2     | 4   | 6   | 2   |
| Brown Creeper <sup>W</sup>                 | <i>Certhia Americana</i>          | BRCR | 0  | 0   | 0     | 0     | 0   | 0     | 0   | 0     | 0   | 2     | 5   | 5   | 0   |

|                                     |                                 |      |     |       |       |       |       |       |       |       |       |       |       |       |     |
|-------------------------------------|---------------------------------|------|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----|
| House Wren <sup>YR</sup>            | <i>Troglodytes aedon</i>        | HOWR | 2   | 10    | 20    | 6     | 11    | 26    | 30    | 27    | 38    | 26    | 33    | 69    | 8   |
| Winter Wren <sup>W</sup>            | <i>Troglodytes hiemalis</i>     | WIWR | 0   | 0     | 0     | 0     | 1     | 3     | 0     | 0     | 2     | 3     | 1     | 0     | 0   |
| Carolina Wren <sup>R</sup>          | <i>Thryothorus ludovicianus</i> | CARW | 3   | 27    | 78    | 90    | 107   | 218   | 195   | 242   | 356   | 470   | 401   | 561   | 148 |
| Marsh Wren <sup>R</sup>             | <i>Cistothorus palustris</i>    | MAWR | 0   | 2     | 0     | 2     | 4     | 4     | 12    | 18    | 15    | 17    | 7     | 11    | 3   |
| Sedge Wren <sup>W</sup>             | <i>Cistothorus platensis</i>    | SEWR | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 0   |
| Blue-gray Gnatcatcher <sup>SB</sup> | <i>Poliophtila caerulea</i>     | BGGN | 0   | 0     | 5     | 15    | 5     | 12    | 10    | 22    | 31    | 33    | 76    | 108   | 8   |
| Golden-crowned Kinglet <sup>W</sup> | <i>Regulus satrapa</i>          | GCKI | 0   | 0     | 0     | 0     | 1     | 6     | 3     | 5     | 0     | 12    | 7     | 24    | 9   |
| Ruby-crowned Kinglet <sup>W</sup>   | <i>Regulus calendula</i>        | RCKI | 0   |       | 3     | 7     | 2     | 11    | 8     | 23    | 19    | 78    | 35    | 66    | 29  |
| Eastern Bluebird <sup>YR</sup>      | <i>Sialia sialis</i>            | EABL | 0   | 2     | 29    | 48    | 41    | 60    | 101   | 103   | 67    | 162   | 165   | 306   | 67  |
| Gray-cheeked Thrush <sup>M</sup>    | <i>Catharus minimus</i>         |      | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0   |
| Veery <sup>M</sup>                  | <i>Catharus fuscescens</i>      | VEER | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 0   |
| Swainson's Thrush <sup>M</sup>      | <i>Catharus ustulatus</i>       | SWTH | 0   | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 1     | 1     | 0     | 0   |
| Hermit Thrush <sup>W</sup>          | <i>Catharus guttatus</i>        | HETH | 0   | 0     | 0     | 1     | 2     | 6     | 4     | 9     | 1     | 2     | 10    | 10    | 2   |
| Wood Thrush <sup>N</sup>            | <i>Hylocichla mustelina</i>     | WOTH | 0   | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 0     | 0     | 0     | 0     | 0   |
| American Robin <sup>YR</sup>        | <i>Turdus migratorius</i>       | AMRO | 16  | 62    | 187   | 375   | 160   | 369   | 541   | 775   | 670   | 219   | 235   | 538   | 67  |
| Gray Catbird <sup>R</sup>           | <i>Dumetella carolinensis</i>   | GRCA | 1   | 3     | 11    | 12    | 17    | 80    | 86    | 98    | 84    | 121   | 69    | 153   | 13  |
| Northern Mockingbird <sup>R</sup>   | <i>Mimus polyglottos</i>        | NOMO | 13  | 66    | 115   | 171   | 159   | 231   | 159   | 149   | 0     | 160   | 154   | 225   | 44  |
| Brown Thrasher <sup>R</sup>         | <i>Toxostoma rufum</i>          | BRTH | 2   | 3     | 44    | 66    | 22    | 69    | 78    | 131   | 109   | 123   | 101   | 167   | 27  |
| Sage Thrasher <sup>T</sup>          | <i>Oreoscoptes montanus</i>     | SATH | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 0     | 1     | 0   |
| European Starling <sup>YR</sup>     | <i>Sturnus vulgaris</i>         | EUST | 187 | 1,205 | 2,021 | 3,477 | 3,023 | 4,139 | 5,000 | 3,979 | 3,881 | 4,477 | 3,167 | 3,564 | 768 |
| Cedar Waxwing <sup>W</sup>          | <i>Bombycilla cedrorum</i>      | CEDW | 0   | 0     | 71    | 36    | 32    | 382   | 54    | 89    | 551   | 178   | 99    | 101   | 226 |
| House Sparrow <sup>YR</sup>         | <i>Passer domesticus</i>        | HOSP | 0   | 10    | 3     | 6     | 2     | 12    | 0     | 2     | 1     | 0     | 0     | 1     | 0   |
| American Pipit <sup>W</sup>         | <i>Anthus rubescens</i>         | AMPI | 0   | 0     | 0     | 2     | 0     | 0     | 1     | 1     | 0     | 3     | 0     | 0     | 0   |
| Purple Finch <sup>W</sup>           | <i>Haemorhous purpureus</i>     | PUFI | 0   | 0     | 0     | 0     | 0     | 0     | 0     | 1     | 0     | 0     | 0     | 0     | 0   |

|                                          |                                |      |    |    |     |     |     |     |     |     |     |     |     |     |     |
|------------------------------------------|--------------------------------|------|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| House Finch <sup>YR</sup>                | <i>Haemorhous mexicanus</i>    | HOFI | 9  | 25 | 79  | 171 | 104 | 168 | 144 | 101 | 164 | 134 | 278 | 283 | 52  |
| Pine Siskin <sup>W</sup>                 | <i>Spinus pinus</i>            | PISI | 0  | 0  | 2   | 0   | 0   | 0   | 0   | 1   | 22  | 0   | 0   | 0   | 0   |
| American Goldfinch <sup>W</sup>          | <i>Spinus tristis</i>          | AMGO | 14 | 37 | 172 | 129 | 115 | 108 | 165 | 228 | 247 | 259 | 427 | 468 | 44  |
| Snow Bunting <sup>W</sup>                | <i>Plectrophenas nivalis</i>   | SNBU | 0  | 0  | 19  | 62  | 32  | 0   | 4   | 5   | 13  | 96  | 2   | 0   | 18  |
| Ovenbird <sup>M</sup>                    | <i>Seiurus aurocapilla</i>     | OVEN | 0  | 0  | 1   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   |
| Worm-eating Warbler <sup>N</sup>         | <i>Helmitheros vermivorum</i>  | WEWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 0   |
| Northern Waterthrush <sup>M</sup>        | <i>Parkesia noveboracensis</i> | NOWA | 0  | 0  | 1   | 0   | 0   | 1   | 0   | 1   | 4   | 2   | 1   | 3   | 0   |
| Blue-winged Warbler <sup>M</sup>         | <i>Vermivora cyanoptera</i>    | BWWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   |
| Prothonotary Warbler <sup>N</sup>        | <i>Protonotaria citrea</i>     | PROW | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   |
| Black-and-white Warbler <sup>B</sup>     | <i>Mniotilta varia</i>         | BAWW | 11 | 0  | 0   | 0   | 1   | 0   | 2   | 5   | 4   | 7   | 2   | 6   | 3   |
| Tennessee Warbler <sup>M</sup>           | <i>Oreothlypis peregrine</i>   | TEWA | 0  | 0  | 0   | 0   | 0   | 1   | 0   | 0   | 0   | 0   | 0   | 1   | 0   |
| Orange-crowned Warbler <sup>W</sup>      | <i>Oreothlypis celata</i>      | OCWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 3   | 7   | 0   | 2   | 0   |
| Nashville Warbler <sup>M</sup>           | <i>Oreothlypis ruficapilla</i> | NAWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 0   |
| Common Yellowthroat <sup>N</sup>         | <i>Geothlypis trichas</i>      | COYT | 2  | 13 | 67  | 70  | 36  | 72  | 86  | 95  | 187 | 185 | 184 | 233 | 38  |
| Hooded Warbler <sup>N</sup>              | <i>Setophaga citrina</i>       | HOWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   |
| American Redstart <sup>N</sup>           | <i>Setophaga ruticilla</i>     | AMRE | 0  |    | 1   |     |     | 6   | 6   | 5   | 16  | 5   | 6   | 15  | 0   |
| Northern Parula <sup>YR</sup>            | <i>Setophaga Americana</i>     | NOPA | 0  | 0  | 1   | 0   | 2   | 1   | 0   | 5   | 5   | 15  | 8   | 25  | 0   |
| Yellow Warbler <sup>N</sup>              | <i>Setophaga petechial</i>     | YEWA | 0  | 0  | 3   | 0   | 0   | 14  | 2   | 8   | 27  | 9   | 4   | 9   | 0   |
| Chestnut-sided Warbler <sup>M</sup>      | <i>Setophaga pensylvanica</i>  | CSWA | 0  | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 0   |
| Magnolia Warbler <sup>M</sup>            | <i>Setophaga magnolia</i>      | MAWA | 0  | 0  | 0   | 0   | 0   | 1   | 1   | 0   | 0   | 1   | 1   | 7   | 1   |
| Black-throated Blue Warbler <sup>M</sup> | <i>Setophaga caeruleascens</i> | BTBW | 0  | 0  | 0   | 0   | 1   | 0   | 2   | 1   | 0   | 0   | 0   | 1   | 0   |
| Yellow-rumped Warbler <sup>W</sup>       | <i>Setophaga coronate</i>      | YRWA | 0  | 8  | 94  | 231 | 76  | 191 | 192 | 8   | 221 | 325 | 253 | 521 | 158 |

|                                           |                                  |      |    |     |     |     |     |     |     |     |     |     |     |     |     |
|-------------------------------------------|----------------------------------|------|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Black-throated Green Warbler <sup>M</sup> | <i>Setophaga virens</i>          | BTNW | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 0   | 0   | 0   |
| Bay-breasted Warbler <sup>M</sup>         | <i>Setophaga castanea</i>        | BBWA | 0  | 0   | 1   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   |
| Blackpoll Warbler <sup>M</sup>            | <i>Setophaga striata</i>         | BLPW | 0  | 0   | 0   | 1   | 0   | 2   | 0   | 0   | 9   | 7   | 0   | 5   | 3   |
| Pine Warbler <sup>YR</sup>                | <i>Setophaga pinus</i>           | PIWA | 1  | 0   | 10  | 5   | 5   | 16  | 30  | 25  | 70  | 56  | 88  | 140 | 28  |
| Yellow-throated Warbler <sup>N</sup>      | <i>Setophaga dominica</i>        | YTWA | 0  | 0   | 0   | 1   | 5   | 0   | 0   | 0   | 0   | 0   | 0   | 3   | 0   |
| Prairie Warbler <sup>N</sup>              | <i>Setophaga discolor</i>        | PRAW | 0  | 0   | 1   | 4   | 0   | 3   | 2   | 3   | 3   | 2   | 5   | 1   | 1   |
| Palm Warbler <sup>W</sup>                 | <i>Setophaga palmarum</i>        | PAWA | 0  |     | 1   | 25  | 16  | 1   | 8   | 10  | 41  | 60  | 22  | 125 | 31  |
| Wilson's Warbler <sup>T</sup>             | <i>Cardellina pusilla</i>        | WIWA | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   |
| Yellow-breasted Chat <sup>N</sup>         | <i>Icteria virens</i>            | YBCH | 0  | 1   | 2   | 5   | 2   | 8   | 14  | 8   | 11  | 11  | 12  | 18  | 1   |
| Eastern Towhee <sup>R</sup>               | <i>Pipilo erythrophthalmus</i>   | EATO | 0  | 0   | 19  | 43  | 42  | 78  | 107 | 130 | 123 | 125 | 120 | 131 | 29  |
| Field Sparrow <sup>W</sup>                | <i>Spizella pusilla</i>          | FISP | 0  | 3   | 10  | 7   | 1   | 4   | 39  | 41  | 52  | 41  | 49  | 77  | 7   |
| Chipping Sparrow <sup>S</sup>             | <i>Spizella passerine</i>        | CHSP | 0  | 2   | 3   | 10  | 2   | 6   | 15  | 11  | 6   | 6   | 5   | 21  | 4   |
| Clay-colored Sparrow <sup>T</sup>         | <i>Spizella pallida</i>          | CCSP | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 1   | 1   | 0   | 0   | 0   |
| Vesper Sparrow <sup>W</sup>               | <i>Poocetes gramineus</i>        | VESP | 0  | 0   | 0   | 0   | 1   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   |
| Lark Sparrow <sup>T</sup>                 | <i>Chondestes grammacus</i>      | LASP | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 1   | 0   | 0   |
| Savannah Sparrow <sup>W</sup>             | <i>Passerculus sandwichensis</i> | SAVS | 0  | 26  | 16  | 158 | 60  | 202 | 258 | 713 | 886 | 705 | 394 | 836 | 102 |
| Saltmarsh Sparrow <sup>W</sup>            | <i>Ammospiza caudacuta</i>       | SALS | 0  | 0   | 0   | 0   | 0   | 0   | 1   | 0   | 0   | 0   | 1   | 0   | 0   |
| Nelson's Sparrow <sup>W</sup>             | <i>Ammospiza nelsoni</i>         | NESP | 0  | 0   | 0   | 0   | 0   | 0   | 0   | 2   | 0   | 1   | 1   | 0   | 0   |
| Seaside Sparrow <sup>YR</sup>             | <i>Ammospiza maritima</i>        | SESP | 0  | 1   | 0   | 0   | 0   | 0   | 1   | 10  | 1   | 1   | 13  | 12  | 0   |
| Song Sparrow <sup>W</sup>                 | <i>Melospiza melodia</i>         | SOSP | 21 | 130 | 224 | 384 | 241 | 460 | 311 | 443 | 559 | 655 | 535 | 931 | 290 |
| Swamp Sparrow <sup>W</sup>                | <i>Melospiza georgiana</i>       | SWSP | 0  | 0   | 21  | 53  | 16  | 40  | 26  | 83  | 90  | 103 | 88  | 162 | 37  |
| Fox Sparrow <sup>W</sup>                  | <i>Passerella iliaca</i>         | FOSP | 0  | 0   | 0   | 2   | 1   | 1   | 3   | 17  | 0   | 5   | 3   | 9   | 3   |
| White-throated Sparrow <sup>W</sup>       | <i>Zonotrichia albicollis</i>    | WTSP | 0  | 0   | 23  | 75  | 25  | 103 | 45  | 219 | 213 | 188 | 191 | 449 | 100 |

|                                                                                                                                                                                                                                                                                                                                                                                              |                                      |      |      |       |       |       |       |        |        |        |        |        |        |        |       |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|------|------|-------|-------|-------|-------|--------|--------|--------|--------|--------|--------|--------|-------|
| White-crowned Sparrow <sup>W</sup>                                                                                                                                                                                                                                                                                                                                                           | <i>Zonotrichia leucophrys</i>        | WCSP | 0    | 1     | 0     | 0     | 25    | 3      | 4      | 1      | 4      | 4      | 1      | 23     | 22    |
| Dark-eyed Junco <sup>W</sup>                                                                                                                                                                                                                                                                                                                                                                 | <i>Junco hyemalis</i>                | DEJU | 0    | 0     | 6     | 13    | 12    | 6      | 1      | 8      | 4      | 14     | 4      | 4      | 0     |
| Summer Tanager <sup>N</sup>                                                                                                                                                                                                                                                                                                                                                                  | <i>Piranga rubra</i>                 | SUTA | 0    | 0     | 1     | 1     | 0     | 2      | 6      | 3      | 2      | 1      | 6      | 2      | 0     |
| Scarlet Tanager <sup>M</sup>                                                                                                                                                                                                                                                                                                                                                                 | <i>Piranga olivacea</i>              | SCTA | 0    | 0     | 0     | 0     | 0     | 1      | 1      | 0      | 2      | 0      | 0      | 0      | 0     |
| Northern Cardinal <sup>R</sup>                                                                                                                                                                                                                                                                                                                                                               | <i>Cardinalis cardinalis</i>         | NOCA | 9    | 34    | 102   | 154   | 114   | 204    | 183    | 266    | 375    | 338    | 445    | 564    | 110   |
| Dickcissel <sup>T</sup>                                                                                                                                                                                                                                                                                                                                                                      | <i>Spiz Americana</i>                | DICK | 0    | 0     | 0     | 0     | 0     | 2      | 1      | 0      | 0      | 3      | 0      | 0      | 0     |
| Rose-breasted Grosbeak <sup>M</sup>                                                                                                                                                                                                                                                                                                                                                          | <i>Pheucticus ludovicianus</i>       | RBGR | 0    | 0     | 0     | 0     | 0     | 0      | 0      | 0      | 1      | 0      | 0      | 0      | 0     |
| Blue Grosbeak <sup>N</sup>                                                                                                                                                                                                                                                                                                                                                                   | <i>Passerina caerulea</i>            | BLGR | 0    | 0     | 57    | 48    | 38    | 84     | 62     | 74     | 81     | 112    | 141    | 141    | 23    |
| Indigo Bunting <sup>N</sup>                                                                                                                                                                                                                                                                                                                                                                  | <i>Passerina cyanea</i>              | INBU | 0    | 3     | 28    | 25    | 19    | 55     | 72     | 38     | 75     | 79     | 95     | 75     | 6     |
| Eastern Meadowlark <sup>W</sup>                                                                                                                                                                                                                                                                                                                                                              | <i>Sturnella magna</i>               | EAME | 2    | 27    | 62    | 191   | 190   | 567    | 557    | 359    | 385    | 551    | 361    | 431    | 103   |
| Bobolink <sup>M</sup>                                                                                                                                                                                                                                                                                                                                                                        | <i>Dolichonyx oryzivorus</i>         | BOBO | 0    | 15    | 0     | 4     | 0     | 22     | 13     | 33     | 51     | 29     | 32     | 26     | 1     |
| Red-winged Blackbird <sup>R</sup>                                                                                                                                                                                                                                                                                                                                                            | <i>Agelaius phoeniceus</i>           | RWBL | 76   | 748   | 1,322 | 2,090 | 1,552 | 3,488  | 3,622  | 4,074  | 5,928  | 7,806  | 6,611  | 6,590  | 3,261 |
| Yellow-headed Blackbird <sup>T</sup>                                                                                                                                                                                                                                                                                                                                                         | <i>Xanthocephalus xanthocephalus</i> | YHBL | 0    | 2     | 1     | 0     | 0     | 0      | 0      | 0      | 0      | 0      | 1      | 0      | 0     |
| Common Grackle <sup>R</sup>                                                                                                                                                                                                                                                                                                                                                                  | <i>Quiscalus quiscula</i>            | COGR | 38   | 43    | 146   | 243   | 142   | 260    | 188    | 347    | 370    | 494    | 296    | 627    | 103   |
| Boat-tailed Grackle <sup>YR</sup>                                                                                                                                                                                                                                                                                                                                                            | <i>Quiscalus major</i>               | BTGR | 11   | 36    | 36    | 98    | 31    | 128    | 69     | 102    | 137    | 222    | 131    | 97     | 9     |
| Brown-headed Cowbird <sup>R</sup>                                                                                                                                                                                                                                                                                                                                                            | <i>Molothrus ater</i>                | BHCO | 1    | 470   | 656   | 157   | 47    | 255    | 72     | 111    | 158    | 60     | 668    | 263    | 28    |
| Orchard Oriole <sup>N</sup>                                                                                                                                                                                                                                                                                                                                                                  | <i>Icterus spurius</i>               | OROR | 2    | 5     | 35    | 27    | 27    | 56     | 42     | 19     | 34     | 54     | 18     | 26     | 2     |
| Baltimore Oriole <sup>YR</sup>                                                                                                                                                                                                                                                                                                                                                               | <i>Icterus galbula</i>               | BAOR | 0    | 0     | 0     | 0     | 0     | 0      | 0      | 3      | 3      | 1      | 0      | 2      | 0     |
| TOTAL                                                                                                                                                                                                                                                                                                                                                                                        |                                      |      | 6.8K | 45.1K | 46.1K | 94.5K | 78.6K | 120.5K | 140.3K | 151.5K | 236.5K | 259.1K | 193.4K | 145.3K | 73.1K |
| GRAND TOTAL = 1,590,674                                                                                                                                                                                                                                                                                                                                                                      |                                      |      |      |       |       |       |       |        |        |        |        |        |        |        |       |
| <sup>B</sup> = Breeds in the study area and is likely a short-distance migrant that winters elsewhere on the continent.<br><sup>N</sup> = Long-distance Neotropical migrant that breeds in the study area, but winters in Central/South America.<br><sup>M</sup> = Observed in the study area only during migration.<br><sup>R</sup> = Occurs year-round and likely nests in the study area. |                                      |      |      |       |       |       |       |        |        |        |        |        |        |        |       |

<sup>T</sup> = Transient with no record of regular occupancy during the breeding, wintering, or migration seasons.

<sup>W</sup> = Winters in the study area; likely a short-distance migrant that breeds north of the study area.

YR = Occurs year-round but no confirmed successful breeding in the study area.

WR = Winters in the study area and may breed occasionally.

RW = Rare, but has wintered in the study area.

S = Occurs only during the summer but no confirmed breeding on the study area.

RS = Rare, but has been detected during the summer on the study area.

SB = Occurs in the study area during the summer and breeds; remains in the vicinity but moves out of the study area during the winter.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              |                                                                         |                                                  |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------|------------------------------------------------------|----------------------------------------------|-------------------------------------------------------------------------|--------------------------------------------------|
| <b>1. REPORT DATE (DD-MM-YYYY)</b><br>September 2022                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                                        | <b>2. REPORT TYPE</b><br>Final                       |                                              | <b>3. DATES COVERED (From - To)</b>                                     |                                                  |
| <b>4. TITLE AND SUBTITLE</b><br><br>Birds of the Craney Island Dredged Material Management Area, Portsmouth, Virginia, 2008-2020                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |                                        |                                                      |                                              | <b>5a. CONTRACT NUMBER</b>                                              |                                                  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>5b. GRANT NUMBER</b>                                                 |                                                  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>5c. PROGRAM ELEMENT NUMBER</b>                                       |                                                  |
| <b>6. AUTHOR(S)</b><br><br>Michael P. Guilfoyle, Ruth A. Beck, Bill Williams, Shannon J. Reinheimer, Lyle D. Burgoon, Sam S. Jackson, Sherwin M. Beck, Burton C. Suedel, and Richard A. Fischer                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>5d. PROJECT NUMBER</b>                                               |                                                  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>5e. TASK NUMBER</b>                                                  |                                                  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>5f. WORK UNIT NUMBER</b>                                             |                                                  |
| <b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b><br><br>See reverse.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                        |                                                      |                                              | <b>8. PERFORMING ORGANIZATION REPORT NUMBER</b><br><br>ERDC/EL TR-22-15 |                                                  |
| <b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b><br><br>US Army Corps of Engineers<br>Washington, DC 20314-1000                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>                                 |                                                  |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                        |                                                      |                                              | <b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>                           |                                                  |
| <b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b><br>Approved for public release; distribution is unlimited.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                        |                                                      |                                              |                                                                         |                                                  |
| <b>13. SUPPLEMENTARY NOTES</b><br><br>Funding Account Code U4372823; AMSCO Code 086000                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                        |                                                      |                                              |                                                                         |                                                  |
| <b>14. ABSTRACT</b><br><br>This report presents the results of a long-term trend analyses of seasonal bird community data from a monitoring effort conducted on the Craney Island Dredged Material Management Area (CIDMMA) from 2008 to 2020, Portsmouth, VA. The USACE Richmond District collaborated with the College of William and Mary and the Coastal Virginia Wildlife Observatory, Waterbird Team, to conduct year-round semimonthly area counts of the CIDMMA to examine species presence and population changes overtime. This effort provides information on the importance of the area to numerous bird species and bird species' groups and provides an index to those species and group showing significant changes in populations during the monitoring period. We identified those species regionally identified as Highest, High, and Moderate Priority Species based on their status as rare, sensitive, or in need of conservation attention as identified by the Atlantic Coast Joint Venture (ACJV), Bird Conservation Region (BCR), New England/Mid-Atlantic Bird Conservation Area (BCR 30). Of 134 ranked priority species in the region, the CIDMMA supported 102 of 134 (76%) recognized in the BCR, including 16 of 19 (84%) of Highest priority ranked species, 47 of 60 (78.3%) of High priority species, and 39 of 55 (71%) of Moderate priority species for BCR 30. All bird count and species richness data collected were fitted to a negative binomial (mean abundance) or Poisson distribution (mean species richness) and A total of 271 species and over 1.5 million birds were detected during the monitoring period. Most all bird species and species groups showed stable or increasing trends during the monitoring period. These results indicate that the CIDMMA is an important site that supports numerous avian species of local and regional conservation concern throughout the year. |                                        |                                                      |                                              |                                                                         |                                                  |
| <b>15. SUBJECT TERMS</b><br>Craney Island (Va.) – Bird communities<br>Craney Island (Va.) – Bird populations                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                        | Environmental management<br>Environmental monitoring |                                              |                                                                         |                                                  |
| <b>16. SECURITY CLASSIFICATION OF:</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |                                        |                                                      | <b>17. LIMITATION OF ABSTRACT</b><br><br>SAR | <b>18. NUMBER OF PAGES</b><br><br>82                                    | <b>19a. NAME OF RESPONSIBLE PERSON</b>           |
| <b>a. REPORT</b><br><br>Unclassified                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <b>b. ABSTRACT</b><br><br>Unclassified | <b>c. THIS PAGE</b><br><br>Unclassified              |                                              |                                                                         | <b>19b. TELEPHONE NUMBER (include area code)</b> |

**7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)**

Environmental Laboratory  
US Army Engineer Research and Development Center  
3909 Halls Ferry Road  
Vicksburg, MS 39180-6199

Department of Biology  
College of William and Mary  
Williamsburg, Virginia 23185

Coastal Virginia Wildlife Observatory, Waterbird Team  
154 Lakewood Drive  
Williamsburg, Virginia 23185

US Army Corps of Engineers  
Norfolk District  
803 Front Street  
Norfolk, Virginia 23510

P.O. Box 270  
Barhamsville, Virginia 23011