

Distribution Statement

Distribution A: Public Release.

The views presented here are those of the author and are not to be construed as official or reflecting the views of the Uniformed Services University of the Health Sciences, the Department of Defense or the U.S. Government.



UNIFORMED SERVICES UNIVERSITY OF THE HEALTH SCIENCES

POSTGRADUATE DENTAL COLLEGE
ARMY POSTGRADUATE DENTAL SCHOOL
228 EAST HOSPITAL ROAD
FORT GORDON, GEORGIA 30905



THESIS APPROVAL PAGE FOR MASTER OF SCIENCE IN ORAL BIOLOGY

Title of Thesis: Influence of neodymium-doped yttrium aluminum garnet laser exposure time on cytokine secretion in lipopolysaccharide-challenged rat peripheral blood mononuclear cells

Name of Candidate: Sarah M. Vargas, DMD
Master of Science Degree
August 8, 2021

THESIS/MANUSCRIPT APPROVED:

DATE:

JOHNSON.THOMAS.
MICHAEL.1073956591

Digitally signed by
JOHNSON.THOMAS.MICHAEL.107
3956591
Date: 2021.08.08 09:44:26 -04'00'

20210808

Thomas M. Johnson
DEPARTMENT OF PERIODONTICS, ARMY POSTGRADUATE DENTAL SCHOOL
Committee Chairperson

LANCASTER.DOUGLAS.
S.DUNN.1120129313

Digitally signed by
LANCASTER.DOUGLAS.DUNN.11
20129313
Date: 2021.08.11 07:58:42 -04'00'

20210811

Douglas D. Lancaster
DEPARTMENT OF PERIODONTICS, ARMY POSTGRADUATE DENTAL SCHOOL
Committee Member

STANCOVEN.BRIAN.
WILLIAM.1239802733

Digitally signed by
STANCOVEN.BRIAN.WILLIAM.123
9802733
Date: 2021.08.12 07:11:05 -04'00'

20210812

Brian W. Stancoven
DEPARTMENT OF PERIODONTICS, ARMY POSTGRADUATE DENTAL SCHOOL
Committee Member

LINCICUM.ADAM.R
ANDAL.1139359891

Digitally signed by
LINCICUM.ADAM.RANDAL.113935
9891
Date: 2021.08.11 07:53:33 -04'00'

20210811

Adam R. Lincicum
DEPARTMENT OF PERIODONTICS, ARMY POSTGRADUATE DENTAL SCHOOL
Committee Member

Influence of neodymium-doped yttrium aluminum garnet laser exposure time on cytokine secretion in lipopolysaccharide-challenged rat peripheral blood mononuclear cells

by

Sarah M. Vargas, DMD

CPT, DC, USA

Thesis directed by:

Thomas M. Johnson, DMD, MS; COL, DC, USA

Professor, Department of Periodontics, Army Postgraduate Dental School

Thesis committee members:

Douglas D. Lancaster, DMD, MA; COL, DC, USA

Director, United States Army Advanced Education Program in Periodontics and Associate Professor, Department of Periodontics, Army Postgraduate Dental School

Brian W. Stancoven, DMD, MS; LTC(P), DC, USA

Assistant Professor, Department of Periodontics, Army Postgraduate Dental School

Adam R. Lincicum, DMD, MS; LTC, DC, USA

Assistant Professor, Department of Periodontics, Army Postgraduate Dental School

**Thesis submitted to the Faculty of the
Army Postgraduate Dental School
Postgraduate Dental College
Uniformed Services University of the Health Sciences
In partial fulfillment of the requirements for the degree of
Master of Science 2022**

ABSTRACT

Objective: The objective of this study was to assess the influence of neodymium-doped yttrium aluminum garnet (Nd:YAG) laser exposure time on proinflammatory cytokine/chemokine concentrations in lipopolysaccharide-stimulated rat peripheral blood mononuclear cells (PBMCs).

Materials and Methods: Rat PBMCs were cultured, then stimulated using various lipopolysaccharide (LPS) concentrations—0, 10, 100, or 1000 ng/ml. Cultures at each LPS stimulation level received Nd:YAG laser irradiation (3 or 5 W) over one of four exposure times—0 (control), 30, 45, or 60 s. Pulse duration and repetition rate remained constant at 100 μ s and 20 Hz, respectively. We completed magnetic microsphere immunoassays to compare concentrations of 27 inflammatory mediators in laser versus control groups.

Results: In control PBMC cultures, LPS stimulation increased secretion of cytokines and chemokines in a dose-dependent manner. In laser-irradiated cultures, we observed statistically significant reductions in concentrations of four proinflammatory cytokines/chemokines—tumor necrosis factor- α (TNF- α), macrophage inflammatory protein-1 α (MIP-1 α), macrophage inflammatory protein-2 (MIP-2), and interferon gamma-induced protein-10 (IP-10)—following laser exposures ranging from 30 to 60 s. The effect appeared more pronounced at longer exposure times (45 and 60 s). The lower power setting (3 W) resulted in greater variance in recorded cytokine levels.

Conclusions: Over a range of exposure times, Nd:YAG laser energy produced statistically significant reduction in secretion of 4 of 27 evaluated inflammatory mediators (15%), and the effect was more pronounced with longer exposures (45 and 60 s). Under the experimental conditions described, our findings suggest that Nd:YAG laser irradiation may affect secretion of cytokines and chemokines narrowly rather than broadly, producing statistically significant reductions in concentrations of a small subset of inflammatory mediators. Modifying irradiation parameters may alter the observed anti-inflammatory profile. Future studies should extend our experiments across a range of laser irradiation parameters and cell types.

KEYWORDS: Lasers; leukocytes, mononuclear; inflammation; cytokines; low-level light therapy; lipopolysaccharide

INTRODUCTION

The concept of utilizing light for medical purposes originated long ago. Ancient Egyptian, Greek, Roman, and Arab physicians utilized sunlight to treat a variety of human ailments, beginning as early as 1400 B.C.¹ However, modern phototherapy began to take form around the turn of the twentieth century. In 1903, Niels Ryberg Finsen received a Nobel Prize for his work demonstrating bactericidal and tissue-stimulating effects of concentrated light.² In 1960, Theodore Maiman introduced the first laser,³ and by the end of the same decade, Dr. Endre Mester demonstrated enhancement of a biologic process—hair growth—in mice irradiated with a ruby laser.⁴ Since then, researchers have associated laser irradiation with pain relief,⁵⁻¹¹ enhanced wound healing,^{5,13-17} bone¹⁸⁻²¹ and nerve regeneration,²²⁻²⁶ and reduced inflammation.^{5,27-38}

After Mester's early experiments, investigators quickly realized that progressively large laser exposures or "doses" did not always result in concomitant increases in the recorded biologic outcome—more was not necessarily better.⁶ In fact, laser irradiation may produce stimulatory or inhibitory responses, depending upon the applied laser parameters.² Of principal importance is the wavelength of the laser light, which determines the absorption profile within tissue.³⁹ Fluence (J/cm^2) and irradiance (W/cm^2) are also major determinants of the biologic response.⁴⁰ Other relevant and interrelated parameters include average power, peak power, pulse duration, pulse repetition rate, exposure time, spot size, target cell type, pulse energy, total energy applied, number of laser applications, and interval between laser applications.^{41,42}

Mester referred to the effect he observed as “laser biostimulation,”⁴ and low-level laser/light therapy (LLLT) is another commonly used term for the phenomenon.² In 2014, the North American Association for Photobiomodulation Therapy (NAALT) and the World Association for Laser Therapy (WALT) jointly affirmed “photobiomodulation” (PBM) as the most appropriate term. WALT defines PBM as induction of nonthermal photophysical and photochemical events in target cells or tissues, leading to physiological changes at various biologic scales.⁴³ Produced through nonthermal mechanisms, PBM responses are distinguishable from effects that result from simply heating tissue.² Typical fluence (energy density) values used in PBM range from 1 through 20 J/cm².⁴⁰ Above this threshold, thermal effects may predominate or partially account for the observed responses.

Modulation of the inflammatory response is one of the most consistent observations in PBM research, and this effect may account for many of the clinical benefits associated with laser therapy.²⁸ Various authors have reported reduction in proinflammatory cytokines such as interleukin (IL)-1 β and tumor necrosis factor (TNF)- α using infrared lasers.^{37,38,44} Infrared laser irradiation also appears to decrease cyclooxygenase (COX)-2 mRNA levels and prostaglandin synthesis in some cell types.³³ Given the number of laser parameters potentially influencing cell/tissue effects and the complexity of the immune response, additional research is necessary to clarify underlying PBM mechanisms and optimize therapeutic protocols.

OBJECTIVE

We aimed to evaluate the influence of a specific laser parameter (exposure time) on cytokine/chemokine secretion in stimulated peripheral blood mononuclear cells (PBMCs) irradiated with a neodymium-doped yttrium aluminum garnet (Nd:YAG) laser.

MATERIALS AND METHODS

Target Cells

Rat PBMCs (IQ Biosciences, Berkeley, California) were cultured and exposed to one of four lipopolysaccharide (LPS) concentrations—0, 10, 100, or 1000 ng/ml. We then transferred the cells to 96-well plates (2.0×10^4 cells per well).

Laser Irradiation Parameters

We used an Nd:YAG laser (1064 nm, Lightwalker AT, Fotona, Dallas, Texas) to irradiate cultures in laser treatment groups. A 300-micron optical fiber directed the laser beam perpendicularly to the plated cells from a distance of 1.8 mm, and a standardized support reliably reproduced the laser position. Cultures at each LPS stimulation level received Nd:YAG laser irradiation (3 or 5 W average power) at one of four exposure times—0 (control), 30, 45, or 60 s (Figure 1). Pulse duration and repetition rate remained constant at 100 μ s and 20 Hz, respectively. After irradiation, we incubated the cells for one hour at 37° C prior to analysis.

Evaluation of Cytokine/Chemokine Concentrations

We utilized a magnetic microsphere immunoassay (MAGPIX System, Luminex, Austin, Texas) to simultaneously analyze 27 cytokine/chemokine levels in each culture (Table 1). The immunoassay kit included color-coded magnetic microspheres tagged with fluorescent dyes covalently coupled with antibodies specific to cytokines of interest. After we subjected the microspheres to a culture, we introduced a reporter molecule labeled with a specific fluorescent dye. A light emitting diode (LED) excited the dyes, and a CCD camera detected the fluorescence from each microsphere. Median fluorescence intensities (MFIs) provided the basis for the sample analysis. The analysis software (MAGPIX System, Luminex) processed the images and determined cytokine/chemokine concentrations in picograms per milliliter (pg/ml) using standard curves.

Statistical Analysis

We performed each experiment (LPS concentration and laser treatment combination) in duplicate wells per condition and in two independent experiments. Data are presented as means $\pm$ SEM. Mean cytokine concentrations for treatment and control groups were compared using Student's t-tests. Differences were accepted as significant at the $p < 0.05$ level.

RESULTS

Four of 27 inflammatory mediators analyzed (15%) exhibited statistically significant reductions in concentration in the laser-irradiated compared with control cultures. As expected, we observed dose-dependent increases in cytokine/chemokine concentrations in the

control groups (Figure 2). We detected some statistically significant differences at each laser exposure (30, 45, and 60 s). However the anti-inflammatory effect was more pronounced at the higher exposure times (45 and 60 s) (Figure 3). Cytokine/chemokine concentrations recorded in cultures exposed to the lower power setting (3 W, 60 s exposure time) were not statistically different than corresponding concentrations in control cultures. Additionally, in the cultures receiving 3 W power exposure, cytokine/chemokine concentrations exhibited greater variation (standard deviation) in repeated experiments. Some cytokines exhibited a trend for increased concentrations following exposure to laser power of 3 W, although this trend did not reach statistical significance.

DISCUSSION

Our objective was to evaluate the effect of Nd:YAG laser exposure time on secretion of 27 inflammatory mediators in LPS-stimulated rat PBMCs. Although some statistically significant reductions in cytokine concentrations were noted at each evaluated exposure time, the longer exposures (45 s and 60 s) did produce more pronounced anti-inflammatory effects. Because general reduction in inflammation is a consistent finding in PBM studies,²⁸ an expectation of extensive laser influence on a large proportion of the evaluated proteins may have been reasonable. On the contrary, we observed statistically significant reductions in only four proinflammatory cytokines/chemokines. It is possible that the power levels we applied exceeded ideal for producing anti-inflammatory effects. Notably, our experiments were incapable of establishing optimal parameters to minimize proinflammatory cytokine levels. Since numerous PBM outcomes of interest exhibit biphasic responses,⁴⁰ it is possible that a

different fluence value (J/cm^2) may have affected the evaluated inflammatory mediators more broadly and to a greater degree.

Most of our experiments utilized an average laser power of 5 W. We selected this power level based on the findings of a previous study in our lab. Utilizing an experimental design similar to that of the present study, Bunting and colleagues assessed cytokine concentrations after irradiating cultures for 30 s with one of four power values—0 (control), 5, 7.5, or 10 W.⁴⁵ Bunting found in cultures receiving 7.5- and 10-W laser exposures, cytokine concentrations were reduced below detection limits.⁴⁵ The maximum exposure time used in the present investigation (60 s) was twice the exposure time Bunting applied.⁴⁵ Consistent with Bunting's findings, we noted a less pronounced anti-inflammatory effect, as well as higher standard deviation, in the group receiving the lower power value (3 W). Additionally, our observation of a trend for increased concentrations of some cytokines in the 3 W group may suggest a biphasic response related to laser power. Our findings, in the context of Bunting's work, indicate that higher power (7.5 to 10 W) over a shorter exposure period (30 s) or moderate power (5 W) over a longer period (45 to 60 s) may produce superior suppression of some proinflammatory cytokines.

CONCLUSIONS

Under the experimental conditions described, our findings suggest that Nd:YAG laser irradiation may affect secretion of cytokines and chemokines narrowly rather than broadly, producing statistically significant reductions in concentrations of a small subset of the evaluated inflammatory mediators. A more pronounced anti-inflammatory effect occurred with longer

laser exposures (45 and 60 s). Modifying irradiation parameters may alter the observed anti-inflammatory profile. Future studies should extend our experiments across a range of laser irradiation parameters and cell types.

REFERENCES

1. Gusain P, Paliwal R, Joga R, Gupta N, Singh V. Ancient light therapies: a boon to medical science. *Sci Cult* 2016;82:231-236.
2. Anders JJ, Lanzafame RJ, Arany PR. Low-level light/laser therapy versus photobiomodulation therapy. *Photomed Laser Surg* 2015;33:183-184.
3. Townes CH. The first laser. In: Garwin L, Lincoln T, eds. *A Century of Nature: Twenty-One Discoveries that Changed Science and the World*. London: Nature; 2010:107.
5. Mester E, Szende B, Gartner P. The effect of laser beams on the growth of hair in mice. *Radiobiol Radiother* 1968;9:621–626.
5. Carroll JD, Milward MR, Cooper PR, Hadis M, Palin WM. Developments in low level light therapy (LLLT) for dentistry. *Dent Mater* 2014;30:465-475.
6. Enwemeka CS. Laser biostimulation of healing wounds: specific effects and mechanisms of action. *J Orthop Sports Phys Ther* 1988;9:333-338.
7. Suter VG, Sjölund S, Bornstein MM. Effect of laser on pain relief and wound healing of recurrent aphthous stomatitis: a systematic review. *Lasers Med Sci* 2017;32:953-963.
8. Qamruddin I, Alam MK, Mahroof V, Fida M, Khamis MF, Husein A. Effects of low-level laser irradiation on the rate of orthodontic tooth movement and associated pain with self-ligating brackets. *Am J Orthod Dentofacial Orthop* 2017;152:622-630.
9. Jang H, Lee H. Meta-analysis of pain relief effects by laser irradiation on joint areas. *Photomed Laser Surg* 2012;30:405-417.

10. Chow RT, Johnson MI, Lopes-Martins RA, Bjordal JM. Efficacy of low-level laser therapy in the management of neck pain: a systematic review and meta-analysis of randomised placebo or active-treatment controlled trials. *Lancet* 2009;374:1897–908.
11. Bjordal JM, Lopes-Martins RA, Joensen J, et al. A systematic review with procedural assessments and meta-analysis of low level laser therapy in lateral elbow tendinopathy (tennis elbow). *BMC Musculoskelet Disord* 2008;9:75. doi:10.1186/1471-2474-9-75.
12. Mester E, Nagylucskay S, Doklen A, Tisza S. Laser stimulation of wound healing. *Acta Chir Acad Sci Hung* 1976;17:49–55.
13. Suzuki R, Takakuda K. Wound healing efficacy of a 660-nm diode laser in a rat incisional wound model. *Lasers Med Sci* 2016;31:1683-1689.
14. Solmaz H, Ulgen Y, Gulsoy M. Photobiomodulation of wound healing via visible and infrared laser irradiation. *Lasers Med Sci* 2017;32:903-910.
15. Yu W, Naim JO, Lanza fame RJ. Effects of photostimulation on wound healing in diabetic mice. *Lasers Surg Med* 1997;20:56-63.
16. Bryant GL, Davidson JM, Ossoff RH, Garrett CG, Reinisch L. Histologic study of oral mucosa wound healing: a comparison of a 6.0-to 6.8- μ m pulsed laser and a carbon dioxide laser. *Laryngoscope* 1998;108:13-17.
17. Fujimoto K, Kiyosaki T, Mitsui N, et al. Low-intensity laser irradiation stimulates mineralization via increased BMPs in MC3T3-E1 cells. *Lasers Surg Med* 2010;42:519-526.
18. Ueda Y, Shimizu N. Effects of pulse frequency of low-level laser therapy (LLLT) on bone nodule formation in rat calvarial cells. *J Clin Laser Med Surg* 2003;21:271–277.

19. Karoussis IK, Kyriakidou K, Psarros C, Lang NP, Vrotsos IA. Nd:YAG laser radiation (1.064 nm) accelerates differentiation of osteoblasts to osteocytes on smooth and rough titanium surfaces in vitro. *Clin Oral Implants Res* 2017;28:785-790.
20. Arisu HD, Türköz E, Bala O. Effects of Nd:YAG laser irradiation on osteoblast cell cultures. *Lasers Med Sci* 2006;21:175-180.
21. Aleksic V, Aoki A, Iwasaki K, Takasaki AA, Wang CY, Abiko Y, Ishikawa I, Izumi Y. Low-level Er: YAG laser irradiation enhances osteoblast proliferation through activation of MAPK/ERK. *Lasers Med Sci* 2010;25:559-569.
22. Rochkind S. Stimulation effect of laser energy on the regeneration of traumatically injured peripheral nerves. *Morphogen Regen* 1978;83:25-27.
23. Gigo-Benato D, Geuna S, de Castro Rodrigues A, et al. Low-power laser photobiomodulation enhances nerve repair after end-to-side neurorrhaphy: a double-blind randomized study in the rat median nerve model. *Lasers Med Sci* 2004;19:57-65.
24. Matsushita HI, Kakami KA, Ito A, et al. Effect on the action potential of the low power Nd: YAG laser as irradiated directly to the nerve. *Aichi Gakuin Dent Sci* 1989;2:19-28.
25. Orchardson R, Peacock JM, John Whitters C. Effect of pulsed Nd:YAG laser radiation on action potential conduction in isolated mammalian spinal nerves. *Lasers Surg Med* 1997;21:142-148.
26. Midamba ED, Haanaes HR. Low reactive-level 830 nm GaAlAs diode laser therapy (LLLT) successfully accelerates regeneration of peripheral nerves in human. *Laser Ther* 1993;5:125-129.

27. Giannelli M, Bani D, Tani A, et al. In vitro evaluation of the effects of low-intensity Nd:YAG laser irradiation on the inflammatory reaction elicited by bacterial lipopolysaccharide adherent to titanium dental implants. *J Periodontol* 2009;80:977-984.
28. Hamblin MR. Mechanisms and applications of the anti-inflammatory effects of photobiomodulation. *AIMS Biophys* 2017;4:337-361.
29. Yamaura M, Yao M, Yaroslavsky I, et al. Low level light effects on inflammatory cytokine production by rheumatoid arthritis synoviocytes. *Lasers Surg Med* 2009;41:282–290.
30. Hwang MH, Shin JH, Kim KS, et al. Low level light therapy modulates inflammatory mediators secreted by human annulus fibrosus cells during intervertebral disc degeneration in vitro. *Photochem Photobiol* 2015; 91:403–410.
31. Lim W, Choi H, Kim J, et al. Anti-inflammatory effect of 635 nm irradiations on in vitro direct/indirect irradiation model. *J Oral Pathol Med* 2015;44:94–102.
32. Choi H, Lim W, Kim I, et al. Inflammatory cytokines are suppressed by light-emitting diode irradiation of *P. gingivalis* LPS-treated human gingival fibroblasts: inflammatory cytokine changes by LED irradiation. *Lasers Med Sci* 2012; 27:459–467.
33. Sakurai Y, Yamaguchi M, Abiko Y. Inhibitory effect of low-level laser irradiation on LPS-stimulated prostaglandin E2 production and cyclooxygenase-2 in human gingival fibroblasts. *Eur J Oral Sci* 2000;108:29–34.
34. Nomura K, Yamaguchi M, Abiko Y. Inhibition of interleukin-1 β production and gene expression in human gingival fibroblasts by low-energy laser irradiation. *Lasers Med Sci* 2001;16:218–223.

35. Funk JO, Kruse A, Kirchner H. Cytokine production after helium neon laser irradiation in cultures of human peripheral blood mononuclear cells. *J Photochem Photobiol* 1992. 16:347-355.
36. Pourzarandian A, Watanabe H, Ruwanpura SMPM, Aoki A, Noguchi K, Ishikawa I. Er:YAG laser irradiation increases prostaglandin E2 production via the induction of cyclooxygenase-2 mRNA in human gingival fibroblasts. *J Periodontal Res* 2005;40:182-186.
37. Eltas A, Orbak R. Effect of 1,064-nm Nd:YAG laser therapy on GCF IL-1beta and MMP-8 levels in patients with chronic periodontitis. *Lasers Med Sci* 2012; 27: 543–550.
38. Gomez C, Dominguez A, Garcia-Kass AI, Garcia-Nunez JA. Adjunctive Nd:YAG laser application in chronic periodontitis: clinical, immunological, and microbiological aspects. *Lasers Med Sci* 2011; 26: 453–463.
39. Coluzzi DJ, Convissar RA, Roshkind DM. Laser Fundamentals. In: Convissar RA, ed. *Principles and Practice of Laser Dentistry, 2nd ed.* Amsterdam: Elsevier; 2015:12-26.
40. Huang YY, Chen ACH, Carroll JD, Hamblin MR. Biphasic dose response in low level light therapy. *Dose Response* 2009;7:358-383.
41. Choi AY, Reddy CM, McGary RT, Hill RB, Swenson DT, Seibel P, Hoag JM, Berridge JP, Johnson TM. Adjunctive Nd: YAG laser irradiation for ridge preservation and immediate implant procedures: a consecutive case series. *Clin Adv Periodontics* 2019;9:125-134.
42. Johnson TM, Jusino MA. Management of an immediate implant horizontal defect using freeze-dried bone allograft and a neodymium: yttrium-aluminum-garnet laser. *Clin Adv Periodontics* 2017;7:175-181.

43. Robijns J, Lodewijckx J, Mebis J. Photobiomodulation therapy for acute radiodermatitis. *Curr Opin Oncol* 2019;31:291-298.
44. Aoki A, Mizutani K, Schwarz F, et al. Periodontal and peri-implant wound healing following laser therapy. *Periodontol 2000* 2015;68:217-269.
45. Bunting M. Influence of neodymium-doped yttrium aluminum garnet laser irradiation on pro-inflammatory cytokine concentrations in lipopolysaccharide-challenged peripheral blood mononuclear cells: an in vitro study. [Thesis]. Fort Gordon, GA: Postgraduate Dental College, Uniformed Services University of the Health Sciences; 2021.

FIGURES




Figure 1. Experimental design. In all groups, we held pulse duration and repetition rate constant at 100 s and 20 Hz, respectively. We completed each experiment (LPS concentration and laser exposure combination) in duplicate.







Figure 2. Control cultures for inflammatory mediators exhibiting statistically significant concentration reductions. In control cultures, cytokine/chemokine and lipopolysaccharide (LPS) concentrations correlated appropriately.




Figure 3. Normalized mean concentrations of cytokines/chemokines exhibiting statistically significant differences compared with controls.

TABLES

Table 1. Inflammatory mediators evaluated by magnetic microsphere immunoassay

Tumor necrosis factor- α (TNF- α)
Macrophage inflammatory protein-1 α (MIP-1 α)
Macrophage inflammatory protein-2 (MIP-2)
Interferon gamma-induced protein 10 (IP-10)
Granulocyte colony-stimulating factor (G-CSF)
Eotaxin
Granulocyte-macrophage colony-stimulating factor (GM-CSF)
Interleukin-1 α (IL-1 α)
Leptin
Interleukin-4 (IL-4)
Interleukin-1 β (IL1- β)
Interleukin-2 (IL-2)
Interleukin-6 (IL-6)
Epidermal growth factor (EGF)
Interleukin-13 (IL-13)
Interleukin-10 (IL-10)
Interleukin-12p70 (IL-12p70)
Interferon (IFN)
Interleukin-5 (IL-5)
Interleukin-17 α (IL-17 α)
Interleukin-18 (IL-18)
Monocyte chemoattractant protein-1 (MCP-1)
Growth-regulated oncogene/keratinocyte chemoattractant (GRO/KC)
Vascular epithelial growth factor (VEGF)
Fractalkine
Liposaccharide-Induced CXC chemokine (LIX)