

**Naval Information
Warfare Center**



PACIFIC

TECHNICAL REPORT 3346
MAY 2024

Summary Report on the 2023 Collaborative Beaked Whale Cruise off Baja California, Mexico

E. Elizabeth Henderson
NIWC Pacific

Gustavo Cárdenas Hinojosa
Rodrigo Huerta Patiño
Jasivi Arcos Díaz
CONANP

Sergio Martínez Aguilar
UABCS

Robert Pitman
Oregon State University

Jay Barlow
Unaffiliated

DISTRIBUTION STATEMENT A: Approved for public release.
Distribution is unlimited.

Naval Information Warfare Center (NIWC) Pacific
San Diego, CA 92152-5001

This page is intentionally blank.

TECHNICAL REPORT 3346
MAY 2024

Summary Report on the 2023 Collaborative Beaked Whale Cruise off Baja California, Mexico

E. Elizabeth Henderson
NIWC Pacific

Gustavo Cárdenas Hinojosa
Rodrigo Huerta Patiño
Jasivi Arcos Díaz
CONANP

Sergio Martínez Aguilar
UABCS

Robert Pitman
Oregon State University

Jay Barlow
Unaffiliated

DISTRIBUTION STATEMENT A: Approved for public release.
Distribution is unlimited.

Administrative Notes:

This report was approved through the Release of Scientific and Technical Information (RSTI) process in September 2023 and formally published in the Defense Technical Information Center (DTIC) in May 2024.



NIWC Pacific
San Diego, CA 92152-5001

NIWC Pacific
San Diego, California 92152-5001

P.M. McKenna, CAPT, USN
Commanding Officer

M. J. McMillan
Executive Director

ADMINISTRATIVE INFORMATION

The work described in this report was performed by the Environmental Readiness Branch (56720) that performed the work of the ISR Division, Naval Information Warfare Center (NIWC) Pacific, San Diego, CA. Funding was provided by Naval Sea Systems Command (NAVSEA).

Released by
Mark Xitco, Division Head
Reconnaissance and Interdiction

Under authority of
Nicole Stone, Department Head
ISR Department

ACKNOWLEDGMENTS

We would like to acknowledge the crew of the Azteca for their hard work during the field effort, and Shannon Rankin, Annette Henry, and Brittany Hancock-Hanser at Southwest Fisheries Science Center (SWFSC) and John Atkins at Ocean Instruments, Ltd for the use of their equipment, and Armando Jaramillo and Diana López for their assistance tracking the Drifting Acoustic Spar Buoy Recorder (DASBR).

This is a work of the United States Government and therefore is not copyrighted. This work may be copied and disseminated without restriction.

The citation of trade names and names of manufacturers is not to be construed as official government endorsement or approval of commercial products or services referenced in this report.

Editor: MRM

EXECUTIVE SUMMARY

In September 2023, a third multi-week expedition was conducted off the coast of Baja California, Mexico, to relocate the unidentified species of beaked whales encountered in 2020 as well as look for the beaked whale (BW) 43. Daily, non-systematic searches were carried out close to shore off Bahia San Quintín. Unfortunately, weather conditions became too poor to observe and to survey further offshore of Baja. Only two beaked whale sightings occurred; one of possible Cuvier's beaked whales (*Ziphius cavirostris*) and one of Baird's beaked whales (*Berardius bairdii*). Both species were also detected acoustically, in addition to one detection of BW43 echolocation pulses. In addition to the beaked whales, many other species were also encountered, including killer whales (*Orcinus orca*), short-beaked common dolphins (*Delphinus delphis*), Pacific white-sided dolphins (*Lagenorhynchus obliquidens*), bottlenose dolphins (*Tursiops truncatus*), blue whales (*Balaenoptera musculus*), Bryde's whales (*Balaenoptera edeni*), humpback whales (*Megaptera novaeangliae*), and gray whales (*Eschrichtius robustus*).

This page is intentionally blank.

ACRONYMS

BW	beaked whale
DASBR	Drifting acoustic spar buoy recorder
GPS	Global Positioning System
US Navy	United States Navy
SWFSC	Southwest Fisheries Science Center

This page is intentionally blank.

CONTENTS

EXECUTIVE SUMMARY	V
ACRONYMS.....	VII
1. INTRODUCTION.....	1
2. METHODS	3
3. RESULTS	5
3.1 SIGHTING DATA.....	6
3.2 ACOUSTIC DATA.....	8
4. DISCUSSION.....	11

FIGURES

Figure 1. <i>Azteca</i> vessel track for September 6-16 2023.	5
Figure 2: Distribution of the sightings observed during September 2023.	7
Figure 3. DASBR tracks (blue), beaked whale sightings (red), and acoustic detections from Cuvier’s beaked whale (yellow), Baird’s beaked whale (magenta), BW43 (green), and <i>Kogia</i> (cyan) in September 2023.	10

TABLES

Table 1. Sighting information for all cetacean encounters. Beaked whale sightings are in bold.....	6
Table 2. Drifting acoustic recorder deployment and recovery times (PDT) and durations.....	8
Table 3. Acoustic detection times and locations for beaked whales and <i>Kogia</i>	9

This page is intentionally blank.

1. INTRODUCTION

In 2018, a series of acoustic recordings were made using Drifting Acoustic Spar Buoy Recorders (DASBRs) along the Baja California, Mexico, peninsula during the Southwest Fisheries Science Center (SWFSC) California Current Ecosystem Survey (Simonis, et al. 2020). Several detections of the beaked whale (BW) echolocation pulse called BW43 were made. It is unknown what species makes this pulse, but it's hypothesized to be Perrin's beaked whale (*Mesoplodon perrini*; Baumann-Pickering, et al. 2013), which has never been identified at sea and has only been described from a handful of strandings in southern and central California (Dalebout, et al. 2002). These findings led to an expedition in 2020 to acoustically and visually link and identify the BW43 species (Barlow, et al. 2021; Henderson et al. 2021). However, the expedition found a potentially new species of beaked whale that produces a similar but unique echolocation pulse, termed the BWB pulse (Barlow, et al. 2021). Another expedition was conducted in 2021 in the same offshore area, as well as closer to shore near Bahia San Quintín where deep basins with complex bathymetry are found relatively close to shore (Henderson, et al. 2022).

During this expedition, multiple sightings were made of Cuvier's beaked whale (*Ziphius cavirostris*) and at least one *Mesoplodon* that could not be identified to species; acoustic detections were also made of both Cuvier's beaked whale and BW43 echolocation pulses. This effort was still unable to link a *Mesoplodon* species to the BW43 or BWB pulses. Here, we report on a third expedition to the same region off Baja California, Mexico, in September 2023, during which we attempted to locate both *Mesoplodon* species and get photographs, acoustic detections, and DNA samples to confirm the species' identifications. This work supports the U.S. Navy as this region is part of the greater Navy Southern California Range Complex. For the U.S. Navy to conduct accurate impact assessments as part of their Environmental Impact Statement analyses, it is critical to know what species are present in the area, their population sizes and distributions.

This page is intentionally blank.

2. METHODS

A collaborative expedition was planned for September 6-20, 2023, on board the charter vessel the *Azteca*, based out of the Ensenada Harbor. The vessel surveyed in deep waters towards the nearshore region west of San Quintín. The focus of the effort remained there, as no effort could be conducted further offshore due to poor weather conditions. DASBRs were deployed repeatedly throughout the area in order to acoustically survey the region. When weather conditions became too poor for visual observations, the vessel worked closer to shore and anchored off Bahía San Quintín. Five DASBRs were available for deployment with GPS trackers and radio frequency location tags that could be deployed for periods of several days, each with a single-channel SoundTrap recorder (made by Ocean Instruments <http://www.oceaninstruments.co.nz/>) set to record at 576 kHz, continuously.

Seven visual observers conducted continuous visual observations from sunrise (at approximately 06:30 PDT) to sunset (at approximately 19:00 PDT). Observers rotated through four positions monitoring 360 degrees around the ship in half-hour shifts at each location, with the other observers on standby. The front position was equipped with a pair of mounted 25 x 150 mm bigeye binoculars that were used to scan the water in calm conditions and for species identification of distant sightings. When an individual or group of marine mammals was observed, the start time, start latitude and longitude, species, an estimate of group size (including a minimum, maximum, and best size estimate), group behavior, and any other behavioral observations were recorded on a tablet computer using a custom-made Access database. Photographs were taken of all individuals when possible.

For all species other than beaked whales, once the species and group size had been confirmed and photographed, the sighting was terminated with a final time and position update. For beaked whale sightings, a DASBR would be deployed after the first or second sighting. Due to the long intervals between surfacings for beaked whales (5 to 30 minutes), the vessel would remain in the area of the initial sighting of beaked whales for at least 30 minutes, with all observers helping to visually search the area in order to locate the whales upon their next surfacing. A beaked whale sighting was terminated if a group was not resighted after at least 60 minutes of monitoring. A 7 m panga was available to approach beaked whale groups to obtain biopsies and additional photographs when conditions allowed.

This page is intentionally blank.

3. RESULTS

Due to poor weather conditions for much of the planned period, a shortened expedition took place from September 6-16, 2023. The *Azteca* initially transited directly to the area off Bahía San Quintín, where deep basins and complex bathymetry occur only 22 km from shore (Figure 1). However, due to poor weather conditions that inhibited beaked whale visual sightings, the *Azteca* spent four days surveying this area before seeking shelter in Bahía San Quintín for a day (September 10, 2023). During these first four days of effort, two groups of beaked whales (Baird's beaked whales and likely Cuvier's beaked whales) were observed, but the vessel was never able to approach closely for photographs or biopsies. Both species were also detected acoustically on the DASBRs. In addition, a group of killer whales (*Orcinus orca*) was encountered, which was an unusual species for this area.

While in Bahía San Quintín, three gray whales (*Eschrichtius robustus*) were observed in the shallow waters of the bay (Figure 2). Photographs confirmed that at least one of the whales had been previously observed in the bay (Cárdenas Hinojosa, unpublished data). The next two days were spent back on the water searching the deep basins off Bahía San Quintín and deploying DASBRs in the area. The vessel then returned to the bay for another day (September 13, 2023) due to bad weather and refueling. Two more days of surveys in the nearshore deep waters were conducted, with no additional beaked whale sightings, although one acoustic detection of the BW43 echolocation pulses was obtained. Ultimately, the weather prohibited additional survey effort in the area and the expedition was terminated. The full survey effort, including DASBR tracks, beaked whale sightings, and acoustic detections, is depicted in Figure 3.




Figure 1. *Azteca* vessel track for September 6-16 2023.

3.1 SIGHTING DATA

There was a total of 42 cetacean sightings during the expedition (Table 1); only two of these were beaked whales (Figures 2 and 3). One of the beaked whale sightings was likely Cuvier’s beaked whales (UnIDBW). Although, no photographs were obtained, so species could not be confirmed. However, the other was a group of Baird’s beaked whales (Bba). This group was observed for two hours and four surfacings, with shallow dive periods of 15 to 25 minutes between sightings. However, the panga was unable to reach the group to obtain photographs usable for individual identification before they initiated their deep foraging dive. Other species observed included killer whales (Oo), common dolphins (*Delphinus delphis*, Dd or Dsp), Pacific white-sided dolphins (*Lagenorhynchus obliquidens*, Lo), bottlenose dolphins (*Tursiops truncatus*, Tt), blue whales (*Balaenoptera musculus*, Bm), Bryde’s whales (*Balaenoptera edeni*, Be), gray whales (*Eschrichtius robustus*, Er), humpback whales (*Megaptera novaeangliae*, Mn), and several unidentified large whales (UnMyst, UnLgCet) and unidentified dolphins (UnDelph) (Table 1). Most of the sightings were encountered along the continental slope and in deep waters ranging from 1,000 to 2,000 m (Figure 2).

Table 1. Sighting information for all cetacean encounters. Beaked whale sightings are in bold.

Date	Start Time	End Time	Start Dec Lat	Start Dec Long	Species	Est Size Best
06-Sep-23	15:43:22	15:53:21	30.5804	-116.722	Dd	250
07-Sep-23	6:58:54	6:59:53	30.2951	-115.98	UnDelph	30
07-Sep-23	7:56:56	10:05:35	30.1993	-116.075	Oo	5
07-Sep-23	8:41:36		30.2313	-116.092	Dd	550
07-Sep-23	11:00:37	11:23:14	30.1555	-116.137	Be	1
07-Sep-23	11:43:49	11:50:46	30.1502	-116.128	Be	1
08-Sep-23	7:09:45		30.2442	-116.086	Dd	30
08-Sep-23	7:27:30		30.221	-116.116	Be	1
08-Sep-23	9:12:10	9:15:06	30.0514	-116.077	Dd	100
08-Sep-23	9:53:22	9:54:41	29.977	-116.036	Lo	20
08-Sep-23	11:41:52	11:43:05	29.8693	-116.084	Dd	15
08-Sep-23	12:38:51		29.926	-116.176	UnDelph	
08-Sep-23	13:12:23	14:25:52	29.9553	-116.237	UnIDBW	1
08-Sep-23	14:10:43		29.9359	-116.246	Be	1
08-Sep-23	15:08:20	16:51:59	29.9965	-116.215	Bba	10
08-Sep-23	17:17:48	17:18:32	30.0834	-116.21	UnLgCet	1
08-Sep-23	17:37:48	17:58:26	30.1029	-116.192	Bm	1
08-Sep-23	18:01:01		30.1165	-116.172	Dd	100
08-Sep-23	18:04:09		30.122	-116.167	UnMyst	1
08-Sep-23	18:05:39		30.1246	-116.165	Be	1
08-Sep-23	18:12:41	18:13:01	30.137	-116.154	UnMyst	1
08-Sep-23	18:13:17	18:15:43	30.1381	-116.153	Be	1
09-Sep-23	6:47:29		30.2588	-116.076	Lo	18
09-Sep-23	7:40:44	7:41:09	30.1803	-116.172	UnMyst	1
09-Sep-23	13:51:13	13:54:45	30.3876	-116.324	Dd	20
09-Sep-23	15:45:23	15:47:15	30.3981	-116.305	Dd	30

Table 1. Sighting information for all cetacean encounters. Beaked whale sightings are in bold. (Continued)

Date	Start Time	End Time	Start Dec Lat	Start Dec Long	Species	Est Size Best
09-Sep-23	15:54:03	15:54:57	30.3797	-116.31	Dsp	50
09-Sep-23	16:37:39	16:39:15	30.2908	-116.319	Dsp	30
11-Sep-23	7:13:06	7:18:03	30.3672	-116.133	UnDelph	100
11-Sep-23	13:43:44	13:45:41	30.0222	-116.065	Bm	2
12-Sep-23	7:04:46	7:05:01	30.3745	-116.228	UnDelph	8
12-Sep-23	14:26:13	14:26:29	30.0597	-116.189	Be	1
13-Sep-23	9:12:01	9:53:27	30.3828	-115.988	Er	1
14-Sep-23	7:28:55	7:31:36	30.4074	-115.98	Tt	20
14-Sep-23	7:46:09	8:00:03	30.3847	-115.99	Er	2
14-Sep-23	10:24:47	10:25:16	30.1193	-116.175	UnDelph	5
15-Sep-23	7:17:53	7:18:35	30.2421	-116.098	Dd	50
16-Sep-23	7:20:41	7:28:26	30.9398	-116.498	Lo	20
16-Sep-23	7:38:12		30.9392	-116.54	Dd	100
16-Sep-23	7:48:47	7:54:15	30.9461	-116.556	Be	1
16-Sep-23	12:45:24	12:50:04	31.4626	-116.828	Dd	30
16-Sep-23	16:43:53	16:43:59	31.8431	-116.625	Mn	1




Figure 2: Distribution of the sightings observed during September 2023.

3.2 ACOUSTIC DATA

Drifting acoustic recorders were deployed eight times, for durations of 40 mins – 3.3 days (Table 2). Beaked whales were acoustically detected on five of the eight drifts (Figure 3). Cuvier’s beaked whale echolocation pulses were detected 15 times, Baird’s beaked whales were detected twice, and BW43 pulses were detected once; there was also one detection of *Kogia* sp. (Table 3). No detections occurred in conjunction with a beaked whale sighting, so no sightings or detections could be directly linked as being the same animal(s). Many detections of dolphins occurred; however, their times and locations were not noted. An abundance of dolphin echolocation clicks, primarily at night, impeded our ability to detect beaked whale echolocation pulses at those times. Some beaked whale acoustic detections may have been missed for that reason.

Table 2. Drifting acoustic recorder deployment and recovery times (PDT) and durations.

DASBR Drift #	DASBR #	Deployment Time	Recovery Time	Recording Duration (days)
1	1	9/6/23 18:48	9/7/23 14:05	0.8
2	1	9/8/23 08:40	9/9/23 11:19	1.1
3	4	9/8/23 13:45	9/8/23 14:25	0.04
4	4	9/8/23 16:51	9/9/23 10:05	0.7
5	1	9/11/23 09:07	9/12/23 09:12	1.0
6	3	9/11/23 09:41	9/14/23 16:30	3.3
7	2	9/11/23 12:10	9/14/23 14:08	3.1
8	1	9/14/23 10:35	9/15/23 09:59	1.0

Table 3. Acoustic detection times and locations for beaked whales and *Kogia*.

Drift	Det ID	Species	Start Time (PST)	End Time (PST)	Lat	Long	# clicks
1	1	ZC	9/7/2023 4:57	9/7/2023 4:58	30.30948	-116.397	3
1	2	ZC	9/7/2023 5:40	9/7/2023 5:43	30.30141	-116.401	10
1	3	ZC	9/7/2023 8:09	9/7/2023 8:14	30.28865	-116.396	14
2	1	ZC	9/8/2023 13:50	9/8/2023 14:03	30.09893	-116.172	22
2	2	BB	9/8/2023 17:09	9/8/2023 18:07	30.10348	-116.2	231
5	1	ZC	9/11/2023 11:12	9/11/2023 11:33	30.36733	-116.365	18
5	2	ZC	9/11/2023 13:57	9/11/2023 13:57	30.36831	-116.353	11
6	4	BB	9/12/2023 20:30	9/12/2023 20:40	30.10032	-116.404	173
6	6	ZC	9/13/2023 20:34	9/13/2023 20:56	29.7468	-116.44	43
6	7	BW43	9/14/2023 13:03	9/14/2023 13:19	29.57528	-116.356	11
6	8	<i>Kogia</i>	9/14/2023 5:41	9/14/2023 5:42	29.6256	-116.442	450
7	1	ZC	9/11/2023 15:29	9/11/2023 15:49	30.13164	-116.176	62
7	2	ZC	9/11/2023 22:34	9/11/2023 22:50	30.1429	-116.209	18
7	3	ZC	9/12/2023 1:29	9/12/2023 1:38	30.14138	-116.225	11
7	4	ZC	9/12/2023 11:00	9/12/2023 11:04	30.18391	-116.281	6
7	5	ZC	9/12/2023 13:32	9/12/2023 13:50	30.19074	-116.284	9
7	6	ZC	9/12/2023 18:54	9/12/2023 18:59	30.1577	-116.3	23
7	7	ZC	9/13/2023 19:09	9/13/2023 19:10	29.97599	-116.371	3
7	8	ZC	9/14/2023 2:58	9/14/2023 3:19	29.89012	-116.421	19




Figure 3. DASBR tracks (blue), beaked whale sightings (red), and acoustic detections from Cuvier's beaked whale (yellow), Baird's beaked whale (magenta), BW43 (green), and Kogia (cyan) in September 2023.

4. DISCUSSION

This expedition was the third in a series designed to locate and identify the beaked whale species that produce BW43 and BWB echolocation pulses. The region off Baja California, Mexico, was chosen based on the high number of acoustic detections made by SWFSC in 2018 (Simonis, et al. 2020). These beaked whale surveys have all been conducted in the fall as that was historically the period of best weather for the area. Unfortunately, in all three years the weather has proven to be a challenge, as the excellent conditions necessary for visually locating beaked whales have not been found. However, the three years of surveys have started to provide some information on these and other cetacean species in the area.

First, the deep waters just offshore of Bahía San Quintín do appear to be an important habitat for Cuvier's beaked whales. There were at least four confirmed sightings and 20 acoustic detections that have occurred in that area over three years, with two additional sightings off Ensenada. For all beaked whale sightings, there has been a sighting per day of effort rate of 0.32 over 38 days of effort in four years, with a far higher acoustic detection rate of 1.32 detections per day of effort. Furthermore, satellite-tagged Cuvier's beaked whales have been tracked through that nearshore, deep-water habitat as they traveled south from the Channel Islands (e.g., Schorr, et al. 2020). The bathymetry in that area is complex, with a number of seamounts and ridges that are known to provide good foraging habitat for beaked whales. There were also sightings of a few unidentified *Mesoplodon* species in the same area in 2021, including one acoustic detection of BW43 echolocation pulses (Henderson et al. 2022). However, there may be some niche partitioning occurring between different species of beaked whales as more acoustic detections of BW43 and BWB echolocation pulses have occurred further offshore along the ridge to the west of San Quintín Basin (Barlow, et al. 2021; Henderson, et al. 2021, 2022). It may be that the *Mesoplodon* species in this region prefer deeper waters or a steeper slope than is found on the eastern side of San Quintín Basin.

Second, no beaked whales of any species were seen or acoustically detected during a short, three-day survey effort in June 2022 (Cárdenas Hinojosa and Henderson, unpublished data). While that survey was brief, this may begin to provide some insight about beaked whales seasonal presence in this area. Future work at other times of the year will help to determine any seasonality that may exist in beaked whale distributions in this region. During an ongoing effort using a panga to survey for beaked whales at San Quintín and Ensenada, two Cuvier's beaked whales encounters occurred in winter (December 2022 and February 2023) offshore of Ensenada, and another occurred in the summer (July 2023) offshore of San Quintín. Furthermore, a group of Baird's beaked whales were also sighted in the same area in July 2023 (Cárdenas Hinojosa, unpublished data).

Third, a number of Bryde's whales have been observed in this region over the last three years of surveys. In addition to the nine sightings on the longer surveys, during the ongoing effort with panga, 16 sightings of this species were recorded in San Quintín Basin, mainly in July and August 2023 (Cárdenas Hinojosa, unpublished data). While Bryde's whales are widely distributed globally in tropical and sub-tropical waters, they are one of the less well-known rorquals, and are often found far offshore where visual surveys are rare (e.g., Helble, et al. 2016). The proximity of this basin to shore and the possible frequent occurrence of this species may provide an opportunity to study these animals more closely.

Fourth, in addition to the lesser-known Bryde's whales, blue whales and humpback whales have also been sighted repeatedly in the waters west of northern Baja California, Mexico, during this effort. Six blue whales were observed on the longer surveys, while 12 sightings occurred during the panga effort. Similarly, 19 sightings of humpback whales occurred during the longer surveys in the fall, while two sightings were made via panga in May and July, likely of members of the Mexican population. During the sighting in May the humpback whales were observed to be feeding. Sightings and photographs in this region further understanding of seasonal occurrence and migration behavior and help provide insight into connectivity between populations found in southern California and Mexican waters (Jorge Urbán and Anelio Aguayo 1987; Mate, et al. 1999; Calambokidis, et al. 2000, 2009).

Finally, at least three gray whales were observed in the shallow waters of Bahia San Quintín. One of these whales had also been observed in May, July, and August 2023 in the bay (Cárdenas Hinojosa, unpublished data), and at least one gray whale was observed in the bay in November 2021 (Henderson, et al. 2022). Eastern North Pacific gray whales migrate seasonally along the west coast of North America, foraging from the Pacific Northwest to the Bering Sea in the summer and breeding in the warm lagoon waters of Mexico in the winter (Calambokidis, et al. 2002; Rojas-Bracho, et al. 2003). The gray whales in Bahia San Quintín may be taking advantage of the safety of the shallow waters to rest along their migration and are probably foraging opportunistically because the body conditions of the whales were not emaciated. Alternatively, the bay may be an over-summering habitat for some whales. More photo-ID work throughout the year will provide more information on when and how gray whales are using the bay.

The DASBRs with vertical arrays of two hydrophones (used on our previous surveys) were not available on loan for this survey. Although our effort this year showed that DASBRs with single-channel Soundtrap recorders can be effective for detecting beaked whales and were faster to deploy and retrieve, these were less effective at detecting beaked whales when many dolphin echolocation clicks were present. Previously, by having two hydrophones at 100-150 m depth and by estimating bearing angles to sound sources, beaked whale echolocation pulses were more conspicuous because they were all from below the hydrophones' depths whereas the vast majority of dolphin clicks were coming from above. If possible, vertical arrays with two hydrophones should be included in future acoustic surveys of beaked whales. In the future, improved software may allow better detection of beaked whales in the presence of dolphins and could be used to reanalyze our recordings to detect more beaked whales.

While the waters off northern Baja California, Mexico appear to be a region utilized by a remarkable diversity of beaked whales (including *Ziphius*, *Berardius* and potentially five species of *Mesoplodon* [*M. densirostris*, *M. peruvianus*, *M. ginkgodens*, *M. perrini*, and *M. hotaula*]), the weather conditions (e.g., Beaufort sea state at 3 or higher, swell height 1-2 m) in the area have made visual surveys difficult. Additional future efforts may need to be conducted from a larger vessel that can allow for around-the-clock acoustic survey with a towed array, even in rough waters, to identify where best to search for BW43 and BWB beaked whales when weather conditions are amenable.

REFERENCES

- Barlow, J., Cárdenas-Hinojosa, G., Henderson, E.E., Breese, D., López-Arzate, D., Hidalgo Pla, E., and B.L. Taylor. 2021. Unique morphological and acoustic characteristics of beaked whales (*Mesoplodon* sp.) off the west coast of Baja California, Mexico. *Marine Mammal Science*.
- Baumann-Pickering, S., McDonald, M.A., Simonis, A.E., Solsona Berga, A., Merkens, K.P., Oleson, E.M., Roch, M.A., Wiggins, S.M., Rankin, S., Yack, T.M. and Hildebrand, J.A., 2013. Species-specific beaked whale echolocation signals. *The Journal of the Acoustical Society of America*, 134(3), pp.2293-2301.
- Calambokidis, J., Steiger, G.H., Rasmussen, K., Urbán, J., Balcomb, K.C., Salinas, M., Jacobsen, J.K., Baker, C.S., Herman, L.M., Cerchio, S. and Darling, J.D., 2000. Migratory destinations of humpback whales that feed off California, Oregon and Washington. *Marine Ecology Progress Series*, 192, pp.295-304.
- Calambokidis, J., Darling, J.D., Deecke, V., Gearin, P., Goshō, M., Megill, W., Tombach, C.T., Goley, D., Toropova, C. and Gisborne, B., 2002. Abundance, range and movements of a feeding aggregation of gray whales (*Eschrichtius robustus*) from California to southeastern Alaska in 1998. *J. Cetacean Res. Manage.*, 4(3), pp.267-276.
- Calambokidis, J., Barlow, J., Ford, J.K., Chandler, T.E. and Douglas, A.B., 2009. Insights into the population structure of blue whales in the Eastern North Pacific from recent sightings and photographic identification. *Marine Mammal Science*, 25(4), pp.816-832.
- Dalebout, M.L., Mead, J.G., Baker, C.S., Baker, A.N. and van Helden, A.L., 2002. A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine mammal science*, 18(3), pp.577-608.
- Helble, T.A., Henderson, E.E., Ierley, G.R. and Martin, S.W., 2016. Swim track kinematics and calling behavior attributed to Bryde's whales on the Navy's Pacific Missile Range Facility. *The Journal of the Acoustical Society of America*, 140(6), pp.4170-4177.
- Henderson, E.E., Barlow, J., Cardenas-Hinososa, G., Lopez-Arzate, D.C., Breese, D. and Hidalgo Pla, E., 2021. FY20 Summary Report on the Collaborative Beaked Whale Cruise Off Baja California, Mexico. Naval Information Warfare Center Pacific Technical Report 3249.
- Henderson, E.E., Cardenas-Hinososa, Huerta Patino, R., D.C., Breese, D. Pusser, T., Pitman, R. 2022, and Barlow, J. Summary Report on the Second Collaborative Beaked Whale Cruise Off Baja California, Mexico. Naval Information Warfare Center Pacific Technical Report 3271.
- Jorge Urbán, R. and Anelio Aguayo, L., 1987. Spatial and seasonal distribution of the humpback whale, *Megaptera novaeangliae*, in the Mexican Pacific. *Marine Mammal Science*, 3(4), pp.333-344.
- Mate, B.R., Lagerquist, B.A. and Calambokidis, J., 1999. Movements of North Pacific blue whales during the feeding season off southern California and their southern fall migration 1. *Marine Mammal Science*, 15(4), pp.1246-1257.
- Rojas-Bracho, L., Perez-Cortes, H., Gomez-Gallardo, A., Swartz, S.L., Ludwig, S. and Brownell Jr, R.L., 2003. A review of gray whales (*Eschrichtius robustus*) on their wintering grounds in Mexican waters. *J. Cetacean Res. Manage.*, 5(3), pp.281-295.

Schorr, G.S., Rone, B.K., Falcone, E.A., Keene, E.L., Sweeney, D.A. 2019. Cuvier's beaked whale and fin whale survey at the Southern California Offshore Anti-Submarine Warfare Range (SOAR). Final Report to the US Navy Pacific Fleet Integrated Comprehensive Monitoring Program, Award No. N66604-18-P-2187 and Interim Report to the Cooperative Agreement Studies Unit, Award No. N62473-19-2-0025. 54 p.

Simonis, A.E., Trickey, J.S., Barlow, J., Rankin, S., Urbán, J., Rojas-Bracho, L. and Moore, J.E., 2020. Passive Acoustic Survey of Deep-Diving Odontocetes in the California Current Ecosystem 2018. SWFSC Report.

INITIAL DISTRIBUTION

84310	Technical Library/Archives	(1)
56720	E. Henderson	(1)

Defense Technical Information Center		
Fort Belvoir, VA 22060-6218		(1)

Naval Sea Systems Command		(1)
---------------------------	--	-----

This page is intentionally blank.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-01-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to Department of Defense, Washington Headquarters Services Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) May 2024		2. REPORT TYPE Final		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Summary Report on the Second Collaborative Beaked Whale Cruise off Baja California, Mexico				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
6. AUTHORS E. Elizabeth Henderson NIWC Pacific				Gustavo Cardenas-Hinojosa Rodrigo Huerta Patino Sergio Martínez Aguilar Jasivi Arcos Díaz CONANP	
				Jay Barlow Unaffiliated	
				Robert Pitman Oregon State University	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NIWC Pacific 53560 Hull Street San Diego, CA 92152-5001				8. PERFORMING ORGANIZATION REPORT NUMBER TR-3346	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Sea Systems Command, Headquarters - SEA 04RE WNY Bldg. 197, Room 1E-2094 Phone: 202-781-3735				10. SPONSOR/MONITOR'S ACRONYM(S) NAVSEA	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT DISTRIBUTION STATEMENT A: Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT In September 2023, a third multi-week expedition was conducted off the coast of Baja California, Mexico to relocate the unknown species of beaked whales encountered in 2020 as well as look for the BW43 beaked whale. Daily non-systematic searches were carried out close to shore off Bahia San Quintín. Only two beaked whale sightings occurred, one of possible Cuvier's beaked whales (<i>Ziphius cavirostris</i>) and one of Baird's beaked whales (<i>Berardius bairdi</i>). Both species were also detected acoustically, in addition to one detection of BW43 echolocation pulses. In addition to the beaked whales, many other species were also encountered, including short-beaked common dolphins (<i>Delphinus delphis</i>), Pacific white-sided dolphins (<i>Lagenorhynchus obliquidens</i>), bottlenose dolphins (<i>Tursiops truncatus</i>), blue whales (<i>Balaenoptera musculus</i>), Bryde's whales (<i>Balaenoptera edeni</i>), and gray whales (<i>Eschrichtius robustus</i>).					
15. SUBJECT TERMS Beaked whales, BW43, BWB, echolocation pulses					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			E. Elizabeth Henderson
U	U	U	SAR	30	19b. TELEPHONE NUMBER (Include area code) (619) 553-5067

This page is intentionally blank.

This page is intentionally blank.

DISTRIBUTION STATEMENT A: Approved for public release. Distribution is unlimited.

Naval Information Warfare Center Pacific (NIWC) Pacific
San Diego, CA 92152-5001