

AD/A-004 668

THE HIGH PRESSURE NERVOUS SYNDROME
DURING HUMAN DEEP SATURATION AND
EXCURSION DIVING

A. J. Bachrach, et al

Naval Medical Research Institute
Bethesda, Maryland

1973

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE

**Best
Available
Copy**

AD 10-004668

NAVAL MEDICAL RESEARCH INSTITUTE
BETHESDA, MARYLAND 20014

UNCLASSIFIED

THE HIGH PRESSURE NERVOUS SYNDROME DURING HUMAN DEEP
SATURATION AND EXCURSION DIVING

MEDICAL RESEARCH PROGRESS REPORT

A. J. BACHRACH and P.B. BENNETT

1973

8

19

M4306.03-2112 Report No. 4

THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE:
ITS DISTRIBUTION IS UNLIMITED

Reprinted from
Forvarsmedicin 9 (3), 1973

BUREAU OF MEDICINE AND SURGERY
(NAVY)
WASHINGTON, D.C. 20372

The High Pressure Nervous Syndrome (HPNS) is a condition found in deep diving in excess of 400 feet (13 ATA) whilst breathing oxygen-helium. It is characterized by tremors, an increase in theta activity (4-7 c/sec) in the electroencephalogram (EEG) accompanied by a depression of faster activities and if sufficiently severe, lapses of consciousness. In animals,

convulsions occur, but to date these have not been seen in man. The HPNS is a complex phenomenon comprised of effects, some of which are a function of the hydrostatic pressure. Others seem due to a complex combination of the two. This paper will describe briefly some of the relevant factors responsible for the HPNS in saturation and excursion diving.

Reproduced from
best available copy.



Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
US Department of Commerce
Springfield, VA. 22151

DD FORM 1473

574 0102-014-700

UNCLASSIFIED

Security Classification

AD A004668

The High Pressure Nervous Syndrome during Human Deep Saturation and Excursion Diving

A. J. RACFRACH & P. B. BENNETT

The High Pressure Nervous Syndrome (HPNS) is a condition found in deep diving in excess of 400 feet (13 ATA) whilst breathing oxygen-helium. It is characterized by tremors, an increase in theta activity (4-7 c/sec) in the electroencephalogram (EEG) accompanied by a depression of faster activities and if sufficiently severe, lapses of consciousness (1, 4, 5, 11). In animals, convulsions occur, but to date these have not been seen in man (7).

The HPNS is a complex phenomenon comprised of effects, some of which correlate with the rate of compression and others which are a function of the hydrostatic pressure. Others seem due to a complex combination of the two. This paper will describe briefly some of the relevant factors responsible for the HPNS in saturation and excursion diving.

Saturation dives

Tremors

In 1960, tremor of the hands, arms and torso were reported (2) in subjects compressed at 40-50 fms to 600 and 800 ft and quantified by an Ball bearing test which requires subjects to pick up ball bearings with tweezers and place them in a hole of the same diameter. Rabinowitz, Gumpel, Smith and Schreiner (12) in a study of 1961, also in 1965, observed one or two subjects had slight tremor of both hands and arms, and the other, tremor of the hands. Rabinowitz (12, 19) used a "tremograph" to measure tremor, tremors with a frequency of 5-8 c/sec, according to the upper extremities, were observed in the lower jaw. Will power was used to suppress the tremors to some extent and they appeared more commonly in dry land than dives, more than in the wet, when the rise in temperature was accompanied by the rise in temperature.

finger. This has permitted also a study of the frequency of the tremor. A useful classification of tremor is that of Brumlik and Yap (8) into Rest Tremor, Postural Tremor and Intentional Tremor which includes normal and abnormal expressions of these.

All normal tremors, on frequency analysis, show a large frequency component between 8-12 c/sec whereas Parkinsons and cerebellar disease has a rest peak frequency of 3-8 c/sec. A postural tremor of 8-12 c/sec, as found with the HPNS, is found also in alcoholism and thyrotoxicosis and during the shivering of cold.

Whilst tremor may not be very incapacitating, it is an important early sign of the HPNS and may be the first warning that the rate of compression for the depth desired is too fast, before other more serious HPNS changes are seen, such as in the electroencephalogram (EEG). To prevent tremors, the deeper the depth the slower must be compression.

Electroencephalogram

Measurement of the EEG with out on-line frequency analysis may show little change on visual appraisal unless the HPNS is severe. Graphic representation of on-line EEG frequency analysis, as in the 1500 ft. British experiment (4) especially at depths of 600 ft and greater, with the eyes open, show, however, there is a compression induced rise in theta activity (4-6 c/sec) which once elicited continues for 6 hours, regardless of the fact that compression has ceased, and then falls to more normal values over a further 12 hours. In addition, there may well be a reduction of the remaining EEG activity.

This depression of the electrical activity of the brain is seen also in man, compressed at 1500 ft, during the first 12 hours of saturation, and in the first 12 hours of saturation.



Performance

Performance decrement tests are not particularly sensitive in identifying the presence of HPNS except perhaps for the ball bearing test. Intellectual tests usually are unaffected. However, when the HPNS is more severe, as during compressions of 40-90 ft/min to 600 and 800 ft, then mental impairment does occur accompanied by dizziness, nausea and sometimes vomiting (2).

The performance decrement normally is the result of the hand tremors and so the reverse of inert gas narcosis with psychomotor tasks most affected and intellectual performance less or not at all (5).

Using tremor, EEG and performance as indicators of the HPNS, the rate of compression in deep saturation oxygen-helium dives has been reduced significantly to ameliorate the signs and symptoms of HPNS found with early 100 ft/min rates (2, 3), to such values as 16.7 ft/min with stages to 1500 ft (4, 5), to the technique of a slower and slower compression speed also with stages to permit adaptation. In the French Sagittaire II experiment to 1640 ft the compression decreased to a final rate of 14 ft/hour. Thus compression profiles to great depths now look like those of decompression.

Excursions

Now it has been observed that decompression from excursion dives permits quite large excursions from a saturation depth without the need for stops (9) and in the same way excursion dives may provide one way to dive deep with fast compressions but without undue HPNS. The technique involves slow compression to a saturation depth, as described previously, followed by a fast compression some hours later to the work depth some 150 ft, or so deeper at rates which would not be possible from the surface. With such techniques it may be possible to avoid the phenomenon of the "excursion" which has been reported after excursions from 1000 ft to 1400 ft. Whether or not HPNS would result with standard rates of compression of 50 ft/min, however, has yet to be ascertained.

Bennett and Towse (6) studied rates of 100

ft/min with 20/80 oxygen/helium on EEG, tremor and performance at depths to 300 ft and noted no HPNS. However, other experiments with compression rates of a similar magnitude to 300 ft, 400 ft and 500 ft with 10/90 oxygen-helium did show some decrement in psychomotor tests as evidenced by the occurrence of tremors (3). Other studies indicated that 50 ft/min to 600 ft breathing 5/95 oxygen-helium will evoke HPNS (1) as indeed will the slower rate of 16.7 ft/min as shown in the first stage of compression to 1500 ft and also in the subsequent compressions after 24-hour stages from 600 ft to 1000 ft, 1000 ft to 1300 ft and 1300 ft to 1500 ft (4, 5) and in direct compression to 1000 ft (10). Indeed even rates as slow as 3.5 ft/min to 800 ft as in the experiment at New London (14) elicited the classic EEG changes of HPNS at 400 ft and deeper.

During the latter study, excursions were made from 800 ft to 1112 ft and 800 ft to 1550 ft at 27 and 28 ft/min respectively. These resulted in weakness and slight tremors but no serious increase in the EEG changes already elicited. A slower rate of 17 ft/min a day later did not produce tremor. Nor did the very similar rate of 16.7 ft/min used by Bühlman et al. (10) in excursions from 1000 ft to 1150 ft. However, the results of the RNPL 1500 ft dive suggest that in some individuals, at least, a rate of 16.7 ft/min from 1000 ft to only 1100 ft and 1300 ft to 1400 ft will cause EEG changes and tremor (4, 5).

Deep Work 1000

In January 1973, at Duke University Hyperbaric Facility during a coordinated experiment by many groups including Harbor Branch, Oceanengineering Inc. and University of Florida, known as Deep Work 1000, six men were exposed to 870 ft breathing 0.15 ATA O₂ and the remainder helium.

Decompression to 870 ft was very slow, being 1 ft/min to 450 ft with stops of 1 hour 15 minutes at 450 ft, 350 ft, and 420 ft. The night was spent at 450 ft and compression started again next day at 10 ft/min to 600 ft where a six hour 15 minute stop was made prior to compression to 870 ft at 2 ft/min.

PERCENTAGE CHANGES IN THE EEG
SUBJECT G. B. EYES OPEN

PERCENTAGE EEG CHANGE




Fig. 2. Decreases in EEG activity on compression to 870 ft and 1000 ft.

ally in the 60-80% range, but often nearer 100% in both subjects. Subject G. B. (Fig. 2) showed only increases from baseline during the 10' interval, while subject G. L. did not. Subject D. showed slight increases in alpha (6-13 cps) and delta (2-4 cps) frequencies in isolated instances in the eyes closed con-

dition at 870 ft; no such increases were ever seen in subject G. B.'s profiles. Delta frequencies in general appeared to be more resistant than the other bands to the depressive effects at depths below 100 ft.

Sleep EEG measurements were made in one subject at 870 ft and 76 ft. These indicated a

Table 1. Performance Efficiency During Deep Work 1000 After Excursion to 1000 ft at 50 ft/min

Test		Pre-dive Test-3	870 ft (Pre-excursion)	Arrival at 1000 ft 50 ft/min	1 3/4 hrs at 1000 ft 50 ft/min
Ball	Mean	15.67	17.68	16.33	18.00
Bearing	S.D.	±3.79	±1.53	±3.51	±3.61
Peg	Mean	28.67	33.67	30.00	32.67
Board	S.D.	±3.26	±7.23	±4.36	±6.11
Visual	Mean	47.00	48.00	49.3	40.33
Analogy	S.D.	±6.25	±6.56	±5.69	±2.08
Arith.	Mean	11.00	8.67	8.00	10.67
Correct	S.D.	±5.57	±8.96	±8.71	±8.08

decrease in Stage 3 from 5.6% of total at 76 ft. to 2.4% at 870 ft and an abolition of Stage 4 at 870 ft compared with 16.3% at 76 ft. Total sleep time and REM sleep were equal over the two nights. An increase of 8% in Stage 2 at 870 ft means that this stage was increasing at the expense of the deeper sleep stages.

Performance — The ball bearing test indicated a slight but not significant fall of efficiency only during each of the initial tests on arrival at depth, regardless of the rate of compression. None of the other tests, including the underwater tape assembly and hand tool task indicated a performance decrement (Table 1).

The time estimation task has been shown effective in the analysis of hyperbaric behavior (10, 13).

At times during the 870 ft saturation, marked disruption of the timing behavior was observed in all subjects. When this occurred the 10-20 sec time frame was consistently overestimated with estimation errors increasing by as much as 10-12 sec. Some of the results were very good and accurate but in other cases poor timing accuracy was observed. The accuracy of the 50 ft per min rate of compression.

Previous research (15) at 300 ft or less has shown that shifts in such timing behavior are to a certain extent dependent on the gas mixture breathed. Compressed air causes an underestimate of time, neon an overestimate and no change with helium. The overestimate in the present study is interesting since it is in a direction opposite to inert gas narcosis. More research is needed to confirm these findings.

Overall these results permit the conclusion that a standard rate of compression of 60 ft/min from a saturation depth of 870 ft to 1000 ft is unlikely to precipitate incapacitating HPNS. Nevertheless extrapolation of this data is not justified for it would seem that the deeper the saturation depth, the slower must be the rate of compression.

It is pertinent that in the 1500 ft RNPL dive, confusion and a sense of impending unconsciousness were experienced by Bevan, at 1535 ft during recompression from 1170 ft in an attempt to relieve vestibular decompression sickness in the other subject (4,5). Although he had spent three days deeper than 1000 ft, he was unable to tolerate, even with a very slow rate of compression which with stages totalled 8 1/2 hours, an excursion of 365 ft from 1170 ft.

The data to date, however, is too small with too few subjects to permit firm statements and it is evident that much more research is required in this critical area, with careful measurement of the PIINS to identify the operationally optimum rates of compression both for deep saturation and excursion diving.

A detailed report of HPNS studies during Deep Work 1000 will appear elsewhere by Bennett, P. B., Bachrach, A.J., Findli, A., Wilcox, R. and Walsch, J.M.

This work was partially supported by BUMED Grant No. N00014-67-A-0251-0022. + N4370, 73, 118.

References

1. BACHRACH, A.J. & P.B. BENNETT. Tremor in diving. *Aerospace Med.* 44:613-23, 1973.
2. BENNETT, P.B. *Behavioral Aspects of Deep Diving*. Report No. 1, 1973. (Available from the Naval Submarine Medical Research Committee, Underwater Physiology Sub-Committee Report No. 251, 1964).

3. BENNETT, P.B. & A.N. DOSSSETT. The effects of oxygen-helium at great depths. Medical Research Council, R.N. Personnel Research Committee. Underwater Physiology Sub-Committee Report No. 960 1967.
4. BENNETT, P.B. & E.J. TOWSE. The High Pressure Nervous Syndrome during a simulated oxygen-helium dive to 1500 ft. EEG, Clin. Neurophysiol. 31:583-93, 1971a.
5. BENNETT, P.B. & E.J. TOWSE. Performance efficiency of men breathing oxygen-helium at depths between 100 ft. and 1500 ft. Aerospace Med. 42:1147-56, 1971b.
6. BENNETT, P.B. & E.J. TOWSE. Electroencephalogram, tremors and mental performance during exposure to air or oxygen-helium at 100 ft, 200 ft, and 500 ft. Report 3/72. R.N. Physiological Laboratory, Alverstoke, England.
7. BRAUER, R.W., R.O. WAY, M.R. JORDAN & D.E. PARRISH. Experimental studies on the High Pressure Nervous Syndrome in various mammalian species. In: Underwater Physiology - Proc. Fourth Symposium on Underwater Physiology, edited by C.J. Lambertsen. New York and London: Academic Press, 1971, 471-500.
8. BRUMLIK, J. & C.B. YAP. Normal Tremor: A Comparative Study. Springfield, Illinois: Charles C. Thomas, 1970.
9. BORNEMANN, R.C. Helium-oxygen saturation excursion diving for the U.S. Navy. In: Underwater Physiology - Proc. Fourth Symposium on Underwater Physiology, edited by C.J. Lambertsen. New York and London: Academic Press, 1971, 529-36.
10. BÜHLMANN, A.A., H. MATTHIAS, G. OVERATH, P.B. BENNETT, D.H. ELLIOTT & S.P. GRAY. Saturation exposures of 31 ATA in an oxygen-helium atmosphere with excursions to 36 ATA. Aerospace Med. 41:391-402, 1970.
11. FRUCTUS, X.R., R.W. BRAUER & R. NAQUET. Physiological effects observed in the course of simulated deep chamber dives to a maximum of 36.5 atm. in helium-oxygen atm. In: Underwater Physiology - Proc. Fourth Symposium on Underwater Physiology, edited by C.J. Lambertsen. New York and London: Academic Press, 1971, 545-50.
12. HAMILTON, R.W., J.B. MACINNIS, A.D. NOBLE & H.R. SCHREINER. Saturation Diving at 650 ft. Tech. Memorandum. BI 411. Ocean Systems, Inc. New York, 1966.
13. KINSEY, J.S., G.L. BEFAY & S.M. LURIA. A study of the effects of nitrogen narcosis during saturation diving. In: Underwater Physiology, 705. U.S. Navy Submarine Medical Research Center, London, 1972.
14. PROCTOR, E.D., G.L. BEFAY & M. LURIA. SCHALLER & H. van den ENDE. Electroencephalographic changes during saturation excursion dives to a simulated maximum depth of 1600 ft. Aerospace Med. 43:567-77, 1972.
15. THOMAS, J.B., J.M. WALSH & A.J. BACHRACH. D.P. THOMAS. Different behavioral effects of nitrogen, helium and neon at increased pressures. (Abstract) In: Proc. Fifth Symposium on Underwater Physiology, Treport, British Bahamas, 1972, p. 97.
16. WALSH, J.M. & A.J. BACHRACH. Tiring behavior in the assessment of adaptation to nitrogen narcosis. U.S. Naval Medical Research Institute Report No. 2, Project M4 396, 93-2049D. Bethesda, Maryland, 1971.
17. WALSH, J.M. & A.J. BACHRACH. Behavioral adaptation to hyperbaric gas narcosis. J. Comp. Physiol. Psych. In press.
18. ZALTSMAN, G.L. Physiological Principles of a Sojourn of a Human in Conditions of Raised Pressure of the Gaseous Medium. Leningrad. (English translation.) Foreign Technology Division, Wright-Patterson Air Force Base, Ohio. AD 655, 360, 1967.
19. ZALTSMAN, G.L. (Ed.) Hyperbaric Epilepsy and Narcosis. 1-265. Sechenov Institute of Evolutionary Physiology and Biochemistry USSR Academy of Sciences Leningrad.

Dr. A.J. Bachrach
Behavioral Sciences Department
U.S. Naval Medical Research Institute
Bethesda, Maryland 20014
U.S.A.

Dr. P.B. Bennett
Department of Anesthesiology and
Biomedical Engineering
Duke University Medical Center
Durham, North Carolina 27710
U.S.A.