



ADA 028376

⑨ Progress Report, No. 2, 15 Jun - 14 Jul 76, ⑪ 29 Jul 1976

⑫ 10 p.

June 15 to July 14, 1976

⑥ DESENSITIZATION OF EXPLOSIVE MATERIALS.

⑩ By: Howard M. Peters and John M. Guimont

Prepared for:

Scientific Officer, Director Vehicle Technology Program  
Assistant Chief for Technology  
Office of Naval Research  
800 North Quincy Street  
Arlington, Virginia 22217

Attention: LT Richard S. Miller, Code 211

⑮ Contract Number N00014-76-C-0810  
⑬ (SRI-Project) FYU-5374

Approved:

M. E. Hill  
M. E. Hill, Director  
Chemistry Laboratory

DDC  
RECEIVED  
AUG 17 1976  
A

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

1473  
332 500  
AB

## INTRODUCTION AND SUMMARY

The objective of our work is to determine the generality of an observation that substitution of fluorine for hydrogen in an explosive material leads to desensitization. Substitution of the  $-\text{CH}_2-$  group in FEFO,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{O}]_2\text{CH}_2$ , with a  $-\text{CF}_2-$  group to give DFF,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{O}]_2\text{CF}_2$ , reduces the sensitivity to impact and low velocity detonation (LVD). In addition to having reduced sensitivity, DFF has higher density and lower melting point than FEFO.<sup>1</sup> To generalize our hypothesis that the substitution of fluorine for hydrogen will desensitize an explosive, first we are preparing formals analogous to FEFO and DFF and determining their sensitivity characteristics to be followed by fluorine analogs of explosive nitrate esters, heterocyclics and other classes.

In this report period,  $[\text{C}(\text{NO}_2)_3\text{CH}_2\text{O}]_2\text{CF}_2$ , compound 8,<sup>†</sup> was prepared for comparison with  $[\text{C}(\text{NO}_2)_3\text{CH}_2\text{O}]_2\text{CH}_2$ , compound 7. Also, the synthesis of HADDF,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{OCH}_2\text{OCH}_2]_2\text{C}(\text{NO}_2)_2$ , compound 13, was begun for comparison of the chemical and physical properties with ADDF,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{OCF}_2\text{OCH}_2]_2\text{C}(\text{NO}_2)_2$ , compound 14.

<sup>1</sup>J. M. Guimont and H. M. Peters, "Desensitization of Explosive Materials", Monthly Report No. 1, July 2, 1976.

<sup>†</sup>Numbers refer to compounds listed in Table 1.

|                                 |   |
|---------------------------------|---|
| COPIES FROM                     |   |
| ATIS                            | White Section <input checked="" type="checkbox"/> |
| DDC                             | Buff Section <input type="checkbox"/>             |
| UNANNOUNCED                     | <input type="checkbox"/>                          |
| JUSTIFICATION                   |   |
| <i>Not in file</i>              |   |
| BY                              |   |
| DISTRIBUTION/AVAILABILITY CODES |   |
| Dist.                           | AVAIL. and/or SPECIAL                             |
| A                               |   |

Table 1

## LINEAR FORMALS AND DIFLUOROFORMALS

| (RO) <sub>2</sub> CH <sub>2</sub> (Formals) |     | (RO) <sub>2</sub> CF <sub>2</sub> (Fluoroformals)   |              |   |
|---|-----|---|--------------|---|
| Compound                                    | No. | Structure   | Compound No. | Structure   |
| FEFO  | 1   | [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CH <sub>2</sub> *  | DFD          | [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CF <sub>2</sub> *  |
| DNPF  | 3   | [H <sub>3</sub> CC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CH <sub>2</sub> *                                     | NPFF         | [H <sub>3</sub> CC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CF <sub>2</sub> *                                     |
| TDPF  | 5   | [CF <sub>3</sub> OCH <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CH <sub>2</sub> *                    | OTT          | [CF <sub>3</sub> OCH <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CF <sub>2</sub> *                    |
| TEFO  | 7   | [C(NO <sub>2</sub> ) <sub>3</sub> CH <sub>2</sub> O] <sub>2</sub> CH <sub>2</sub> *   | †            | [C(NO <sub>2</sub> ) <sub>3</sub> CH <sub>2</sub> O] <sub>2</sub> CF <sub>2</sub> *   |
| †   | 9   | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> C(NO <sub>2</sub> )F <sub>2</sub> *                 | MFF          | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> C(NO <sub>2</sub> )F <sub>2</sub> *                 |
| †   | 11  | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> CF <sub>3</sub> *                                   | TMFF         | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> CF <sub>3</sub> *                                   |
| †   | 13  | [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> ] <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> *  | ADDF         | [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> OCH <sub>2</sub> ] <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> *  |
| †   | 15  | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> F* | HTD          | FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> OCH <sub>2</sub> CF <sub>2</sub> OCH <sub>2</sub> C(NO <sub>2</sub> ) <sub>2</sub> F* |


\* Site of change.

† Not previously reported.

## DISCUSSION OF RESULTS

### Comparison of $[\text{C}(\text{NO}_2)_3\text{CH}_2\text{O}]\text{CF}_2$ , compound 8, with $[\text{C}(\text{NO}_2)_3\text{CH}_2\text{O}]\text{CH}_2$ , compound 7 (TEFO)

We prepared bis(trinitroethyl) difluoroformal, compound 8, by the route shown in Eqs. 1 and 2.



The carbonate, 8b, prepared during the previous report period, was fluorinated in a high pressure reactor using sulfur tetrafluoride with anhydrous hydrofluoric acid as catalyst and solvent. The results of the five trials are shown in Table 2. Based on our previous experience with sulfur tetrafluoride, we expected the reaction temperature to be the most critical parameter. The results of trials 2, 4, and 5 showed that reaction temperature is important but that other parameters may be equally so. Since the conditions of trial 5 produced about 75% conversion to the desired product, 8, we plan to force the reaction to completion by repeating trial 5 at 105°C on a larger scale using a longer reaction time.

A complete physical property determination of compound 8 is in progress.

### Comparison of ADDF with HADDF

We also began the synthesis of the hydrocarbon analog (HADDF, compound 13) of ADDF for comparison of sensitivity properties. In one approach, fluorodinitroethylchloromethyl ether was treated with 2,2-dinitropropane-1,3-diol in the presence of stannic chloride as catalyst, Eq. 3.




Table 2




| Trial | Carbonate<br>(g/mmol) | HF<br>mmol | SF <sub>4</sub><br>mmol | Temp.<br>°C | Time<br>hrs | Product<br>(g) | Remarks #                                   |
|-------|-----------------------|------------|-------------------------|-------------|-------------|----------------|---|
| 1     | 10/48                 | 100        | 157                     | 90          | 20          | 9.5            | Starting material recovered                 |
| 2     | 10/48                 | 1100       | 102                     | 90          | 20          | 9.2            | Starting material recovered                 |
| 3     | 10/48                 | 1300       | 74                      | 120         | 20          | 9.2            | Starting material plus small amount product |
| 4     | 10/48                 | 1300       | 102                     | 120         | 120         | 0.8            | Primarily decomposition                     |
| 5     | 10/48                 | 1100       | 194                     | 105         | 72          | 9.4            | 75% Product, 25% starting material          |

\*Product identification is based on ir and nmr analyses.

The mixture of reaction products is currently undergoing separation and identification.<sup>2</sup>

Similarly in a second approach, 1,3-bis(chloromethyl)-2,2-dinitropropyl ether was treated with fluorodinitroethanol Eq. 4.



Again, a mixture of products was obtained. Although bis(fluorodinitroethyl) formal,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{O}]_2\text{CH}_2$ , and 5,5-dinitro-1,3-dioxane,  $(\text{NO}_2)_2\text{C}(\text{CH}_2\text{OCH}_2)_2$ , have been identified as reaction products of both attempted reactions, the presence of Compound 13 has not yet been confirmed. The product analyses of the two reactions, Eq. 3 and 4, have not been completed and are continuing because we believe that some of compound 13 was formed and can be isolated. Other workers have obtained mixtures from chloromethyl ether reactions with similar electronegatively substituted alcohols.<sup>3</sup>

---

<sup>2</sup>G. W. Lawrence, L. E. Kayser, and H. G. Adolph, "ALWT - High Energy Plasticizer and Binder Synthesis", Final Report, February 15, 1976.

<sup>3</sup>H. G. Adolph and M. J. Kamlet, J. Org. Chem., 34, 45 (1969).

## FUTURE WORK

During the next report period we plan to characterize and prepare in quantity  $[\text{C}(\text{NO}_2)_3\text{CH}_2\text{O}]_2\text{CF}_2$ , compound 8, and begin testing its sensitivity for comparison with compound 7 (impact, HVD, LVD).

We also plan to complete the characterization of the products of the chloromethyl ether reactions, Eq. 3 and 4, and to continue our investigation of the synthesis of compound 13,  $[\text{FC}(\text{NO}_2)_2\text{CH}_2\text{OCH}_2\text{OCH}_2]_2\text{C}(\text{NO}_2)_2$ .

DISTRIBUTION LIST

| Address   | DODAAD<br>Code | No. Copies |
|---|----------------|------------|
| Office of Naval Research<br>800 North Quincy Street<br>Arlington, Virginia 22217                                  |                |            |
| Scientific Officer  | N09014         | 1          |
| Administrative Contracting Officer  | S0507A         | 1          |
| Director, Naval Research Laboratory<br>Attn: Code 2627<br>Washington, D.C. 20375                                  | N00173         | 6          |
| Office of Naval Research<br>Department of the Navy<br>Arlington, Virginia 22217<br><br>Attn: Code 1G21P           | N00014         | 6          |
| Defense Documentation Center<br>Bldg. 5, Cameron Station<br>Alexandria, Virginia 22314                            | S47031         | 12         |
| Office of Naval Research<br>Room 447<br>760 Market Street<br>San Francisco, Ca. 94102<br><br>Attn: Phillip Miller | N62887         | 1          |

| REPORT DOCUMENTATION PAGE   |                       | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM   |                       |
|---|-----------------------|---|-----------------------|
| 1. REPORT NUMBER<br>SRI-5374 - #2   | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER   |                       |
| 4. TITLE (and Subtitle)<br>Desensitization of Explosive Materials   |                       | 5. TYPE OF REPORT & PERIOD COVERED<br>Technical Monthly<br>June 15 to July 14, 1976 |                       |
| 7. AUTHOR(s)<br>J. M. Guimont and H. M. Peters  |                       | 6. PERFORMING ORG. REPORT NUMBER<br>N/A   |                       |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Stanford Research Institute<br>333 Ravenswood Avenue<br>Menlo Park, CA 94025   |                       | 8. CONTRACT OR GRANT NUMBER(s)<br>Contract No. N00014-76-C-0810                     |                       |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Attn Code 211<br>Office of Naval Research<br>800 Quincy Street<br>Arlington, VA 22217  |                       | 10. PROGRAM ELEMENT, PROJECT, TASK<br>AREA & WORK UNIT NUMBERS<br>N/A               |                       |
| 14. MONITORING AGENCY NAME & ADDRESS (if diff. from Controlling Office)<br>N/A  |                       | 12. REPORT DATE<br>July 29, 1976  | 13. NO. OF PAGES<br>7 |
|   |                       | 15. SECURITY CLASS. (of this report)<br>Unclassified                                |                       |
| 16. DISTRIBUTION STATEMENT (of this report)<br>Attached   |                       | 15a. DECLASSIFICATION/DOWNGRADING<br>SCHEDULE<br>N/A                                |                       |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)<br>N/A   |                       |   |                       |
| 18. SUPPLEMENTARY NOTES<br>N/A  |                       |   |                       |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br>Chemical Desensitization<br>Fluorodinitroethyl formal (FEFO)<br>Fluorodinitroethyl difluoroformal (DFF)<br>Trinitroethyl formal (TEFO)  |                       |   |                       |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br>The objective of our work is to determine the generality of an observation that substitution of fluorine for hydrogen in an explosive material leads to desensitization. Substitution of the -CH <sub>2</sub> - group in FEFO, [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CH <sub>2</sub> , with a -CF <sub>2</sub> - group to give DFF, [FC(NO <sub>2</sub> ) <sub>2</sub> CH <sub>2</sub> O] <sub>2</sub> CF <sub>2</sub> , reduces the sensitivity to impact and low velocity detonation (LVD). In addition to having reduced sensitivity, DFF has higher density and lower melting point than FEFO. To |                       |   |                       |

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19. KEY WORDS (Continued)

Trinitroethyl difluoroformal (TEDFO)  
Replacement C-H by C-F  
Reduced low velocity detonation (LVD)  
High Velocity Detonation (HVD)  
Impact Sensitivity  
ADDF versus -CH<sub>2</sub>- ADDF

20 ABSTRACT (Continued)

generalize <sup>the</sup> ~~our~~ hypothesis that the substitution of fluorine for hydrogen will desensitize an explosive, first ~~we are preparing~~ formals analogous to FEFO and DFF\* and determining their sensitivity characteristics. ~~to be followed by~~ fluorine analogs of explosive nitrate esters, heterocyclics and other classes. ↑

\* are being prepared

xx This research will continue on