

AD-A031 221

NAVAL RESEARCH LAB WASHINGTON D C
ELECTRON IMPACT RATE COEFFICIENTS FOR THE LOW LYING METASTABLE --ETC(U)
SEP 76 A W ALI


F/G 4/1

UNCLASSIFIED

NRL-MR-3371

NL

1 of 1
ADA031221



END

DATE
FILMED

11 - 76

AD A031221

(12) AC

NRL Memorandum Report 3371

Electron Impact Rate Coefficients for the Low Lying Metastable States of O, O⁺, N and N⁺

A. W. ALI

Plasma Physics Division

September 1976

DDC
OCT 26 1976
C

This work was supported by the Defense Nuclear Agency under Subtask S99QAXHD010,
work unit 87, work unit title, Reaction Rate Studies of Disturbed E and F Region.



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 3371	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ELECTRON IMPACT RATE COEFFICIENTS FOR THE LOW LYING METASTABLE STATES OF O, O ⁺ , N AND N ⁺		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem, 3
7. AUTHOR(s) A. W. Ali	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D.C. 20375		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Defense Nuclear Agency Washington, D.C. 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem H02-27D Subtask S99QAXHD010
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1976
		13. NUMBER OF PAGES 19
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. (16) NRL-H02-27D, DNA-NWED-QAXH		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) (17) DDIP		
18. SUPPLEMENTARY NOTES This work was supported by the Defense Nuclear Agency under Subtask S99QAXHD010, work unit 87, work unit title, Reaction Rate Studies of Disturbed E and F Region.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Disturbed atmosphere Rate coefficients Low lying metastable states O, O ⁺ , N, N ⁺		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A set of electron impact excitation rate coefficients for the low lying metastable states of O, N, O ⁺ , and N ⁺ are presented. These rates are obtained using recent and more accurate cross sections. These species play an important role in the emission, deionization processes and communication problems of a disturbed atmosphere. N(+)		

251 950
hpg

CONTENTS

INTRODUCTION 1

NITROGEN ATOM LOW LYING METASTABLE STATES 2

OXYGEN ATOM LOW LYING METASTABLE STATES 2

THE COLLISION STRENGTHS FOR THE LOW LYING
METASTABLE STATES OF N^+ AND O^+ 3

REFERENCES 5

ACQUISITION NO.	INDEXED	<input checked="" type="checkbox"/>
FILE	FILED	<input type="checkbox"/>
DATE	DATE	<input type="checkbox"/>
BY	BY	
DATE	DATE	
BY _____/ARL/CHY/STP		
DATE	DATE	
A		

Electron Impact Rate Coefficients for the
Low Lying Metastable States of O, O⁺, N and N⁺

INTRODUCTION

The atmosphere, from the sea level to the F-region of the ionosphere can be ionized by many external forces. These disturbing forces include, the lightning discharges, the passage of charged particle beams, the passage of laser beams, the solar flares, the cosmic rays and the atmospheric nuclear bursts. The ionization of the atmosphere under these external forces results in the creation of a large number of excited atomic and molecular species and their corresponding ions which also are in excited states. In addition, the free electrons ejected under the ionization force possess enough kinetic energy to alter the distribution of these species.

Among the atomic and atomic ions species of the disturbed atmosphere, the following excited metastable states play an important role in the chemistry of the atmosphere. These are: O(¹D), O(¹S), O⁺(²D), O⁺(²P), N(²D), N(²P), N⁺(¹D) and N⁺(¹S). These metastable states lie close, within few eV, to the ground state of their respective species. Their densities are controlled to a large degree by electron excitation and de-excitation processes in addition to recombination, charge exchange, and ion-molecule rearrangements (for the ionic species only).

In order to calculate the deionization process of such a disturbed atmosphere, its emission, the conductivity of a disturbed channel and other parameters, one must know the electron temperature and a wide range of inelastic collision cross sections. The collision cross sections of electrons with the species mentioned above, and the relevant rate coefficients, are essential for calculations of relevant parameters.

The rate coefficients and the collision strengths for the inelastic processes of the free electrons with these species have been reported.^{1,2}

Note: Manuscript submitted September 3, 1976.

However, due to the availability of better cross sections we present a new set of rate coefficients for excitations among these low lying metastable states.

NITROGEN ATOM LOW LYING METASTABLE STATES

The low lying metastable states of nitrogen are $N(^2D)$ and $N(^2P)$. The rate coefficients for the electron impact excitations of $N(^4S) - N(^2D)$, $N(^4S) - N(^2P)$ and $N(^2D) - N(^2P)$, reported earlier¹ were based on cross sections calculated by Henry et al.³ These calculations, however, neglected the polarization of the target atom during the collision, incomplete allowance for short range correlations and omissions of higher lying configurations. A recent electron-nitrogen atom scattering calculations by Berrington, et al⁴ includes all these effects. We have elected these cross sections to obtain the relevant rates. Figures 1-3 show these cross sections and are compared with those of Henry et al.³ and Ormonde, et al.⁵

We have utilized these current cross sections⁴ to obtain, in the usual manner,⁶ the relevant electron impact excitation rate coefficients. These rates are given numerically in Table 1 and are shown graphically in Fig. 4. Comparison of these rates with the rates reported previously¹ show that the current rates are significantly smaller for $N(^4S) - N(^2D)$ at low T_e . This is due obviously to the difference in the cross sections (see Fig. 1). It should be also stated that, the resonance structure in the cross section $N(^2D) - N(^2P)$ (see Fig. 3), was ignored in obtaining the current relevant corresponding rate.

OXYGEN ATOM LOW LYING METASTABLE STATES

The low lying metastable states of oxygen are $O(^1D)$ and $O(^1S)$. The rate coefficients for the electron impact excitations of $O(^3P) - O(^1D)$, $O(^3P) - O(^1S)$ and $O(^1D) - O(^1S)$ reported earlier¹ were based on the cross sections calculated by Henry, et al.³ However, the current relevant rate coefficients reported in this section are based on the recent, more accurate, cross sections calculated by Thomas and Nisbet⁷. These cross sections are shown in Fig. 5 along with Henry et al.³ and Vo Ky Lan et al.⁸ It is obvious from this figure that there is very little change

in the cross section for $O(^3P) - O(^1S)$, therefore we shall retain the rate calculated earlier.¹ However, the changes are obvious near threshold and above for $O(^3P) - O(^1D)$, and above threshold for $O(^1D) - O(^1S)$. The new rate coefficients for the low lying oxygen metastable states are given in Table 2 and are shown in Fig. 6.

THE COLLISION STRENGTHS FOR THE LOW LYING METASTABLE STATES OF N^+ AND O^+

The metastable, low lying, excited states of N^+ are $N^+(^1D)$ and $N^+(^1S)$. Those for O^+ are $O^+(^2D)$ and $O^+(^2P)$. The collision strength for most of these states in O^+ and N^+ have been calculated by Henry, et al.³ and more recently for O^+ by Czyzak, et al.⁹ For oxygen ion, the results of Refs. (3) and (9) are in close agreement (within 10%). As for N^+ , Henry et al.³ results are in good agreement with those calculated by Saraph, et al.¹⁰ Therefore, for calculational purposes one may select the collision strengths, for O^+ and N^+ , as given by Henry et al.³ These values are given in Table 3.

In electron ion collision, the cross section is finite at threshold and so is the collision strength. This latter varies slowly as a function of incident electron energy over ranges of interest. Therefore, one can utilize these collision strengths to obtain the relevant electron impact excitation or de-excitation rate coefficients.

The de-excitation rate coefficient from level j to i ($j > i$) is¹¹

$$Y_{ji} = \frac{8.63 \times 10^{-6} \nu(i, j)}{g_j \sqrt{T}} \quad (1)$$

with g_j being the statistical weight of level j and

$$\nu(j, i) = \int_0^{\infty} \Omega(j, i) \text{Exp} \left(\frac{E}{kT} \right) d \left(\frac{E}{kT} \right) \quad (2)$$

In Eq. (2) E is the electron energy, T is the electron temperature in $^{\circ}K$ and $\Omega(j, i)$ is the collision strength. Equation (2) represents the

averaging of the cross section and the electron velocity over a Maxwellian velocity distribution. When the collision strength is constant i.e. it does not depend on E then Eq. (1) is used to obtain the de-excitation rate coefficient. The excitation rate coefficient can be obtained via the detailed balancing.

Using the collision strengths given in Table 3 and Eq. (1), the corresponding de-excitation rate coefficients are given in Table 4.

REFERENCES

1. A. W. Ali and A. D. Anderson, "Low-Energy Electron Impact Rate Coefficients for some Atmospheric Species," NRL Report 7432 (1972).
2. A. W. Ali, "The Physics and the Chemistry of Two NRL Codes for the Disturbed E and F Regions," NRL Report 7578 (1973).
3. R. J. W. Henry, P. G. Burke and A. L. Sinfailam, Phys. Rev. 178, 218 (1969).
4. K. A. Berrington, P. G. Burke and W. D. Robb, J. Phys. B: Atom. Mol. Phys. 8, 2500 (1975).
5. S. Ormonde, K. Smith, B. W. Torres, and A. R. Davies, Phys. Rev. A8, 262 (1973).
6. A. W. Ali and A. D. Anderson, NRL Report 7282 (1971).
7. L. D. Thomas and R. K. Nisbet, Phys. Rev. A11, 170 (1975).
8. Vo Ky Lan, N. Feautrier, M. Le Dourneuf, and Van Regemorter, J. Phys. B5, 1506 (1972).
9. S. J. Czyzak, T. K. Krueger, P. de A. P. Martins, H. E. Saraph and M. J. Seaton, Mon. Not. R. Astron. Soc. 148, 361 (1970).
10. H. E. Saraph, M. J. Seaton, and J. Shemming, Proc. Roy. Soc. (London) 89, 27 (1966).
11. M. J. Seaton, in "Advances in Atomic and Molecular Physics," Bates and Estermann eds. Vol. 4, Academic Press, New York (1968).

TABLE 1

Electron Impact Excitation Rate Coefficients for the Low
Lying States of Nitrogen Atom

T_e (eV)	$^4S - ^2D$	$^4S - ^2P$	$^2D - ^2P$
0.1	8.0 (- 20)	2.6 (- 25)	2.10 (- 14)(*)
0.2	1.54 (- 14)	2.03 (- 17)	1.16 (- 11)
0.3	1.25 (- 12)	1.11 (- 14)	9.78 (- 11)
0.5	4.2 (- 11)	1.47 (- 12)	6.0 (- 10)
0.7	1.98 (- 10)	1.63 (- 11)	1.16 (- 9)
1.0	6.38 (- 10)	8.91 (-11)	2.05 (- 9)
1.2	1.08 (- 9)	1.73 (- 10)	2.56 (- 9)
1.5	1.52 (- 9)	3.38 (- 10)	3.24 (- 9)
2.0	2.27 (- 9)	6.54 (- 10)	4.1 (- 9)
3.0	3.31 (- 9)	1.23 (- 9)	5.18 (- 9)
5.0	4.50 (- 9)	1.99 (- 9)	6.2 (- 9)
7.0	5.26 (- 9)	2.36 (- 9)	6.5 (- 9)
10.0	5.85 (- 9)	2.4 (- 9)	6.5 (- 9)
15.0	5.97 (- 9)	2.49 (- 9)	6.1 (- 9)
20.0	4.96 (- 9)	2.28 (- 9)	5.47 (- 9)

(*) Numbers in parenthesis indicate the power of ten by which the entries are multiplied.

TABLE 2

Electron Impact Excitation Rate Coefficients for
the Low Lying States of Oxygen Atom

T_e (eV)	$^3P - ^1D$	$^3P - ^1S$	$^1D - ^1S$
0.1	1.92 (- 18)	1.78 (- 28)	2.25 (- 19)(*)
0.2	5.28 (- 14)	1.96 (- 19)	1.76 (- 14)
0.3	1.76 (- 12)	2.10 (- 16)	8.07 (- 13)
0.5	3.28 (- 11)	6.04 (- 14)	1.52 (- 11)
0.7	1.21 (- 10)	7.25 (- 13)	5.4 (- 11)
1.0	3.43 (- 10)	4.93 (- 12)	1.38 (- 10)
1.2	5.20 (- 10)	1.06 (- 11)	1.97 (- 10)
1.5	7.94 (- 10)	2.32 (- 11)	2.76 (- 10)
2.0	1.21 (- 9)	5.15 (- 11)	3.98 (- 10)
3.0	1.84 (- 9)	1.16 (- 10)	5.30 (- 10)
5.0	2.52 (- 9)	2.21 (- 10)	7.16 (- 10)
7.0	2.73 (- 9)	2.8 (- 10)	7.76 (- 10)
10.0	2.80 (- 9)	3.3 (- 10)	7.5 (- 10)
15.0	2.58 (- 9)	3.8 (- 10)	6.9 (- 10)
20.0	2.23 (- 9)	3.7 (- 10)	6.4 (- 10)

(*) Numbers in parenthesis indicate the power of ten by which the entries are multiplied.

TABLE 3

Collision Strengths for the Electron Impact Excitations
of the Low Lying Metastable States of O^+

<u>Transition</u>	<u>Collision Strength</u>
$O^+(^4S) - O^+(^2D)$	1.57
$O^+(^4S) - O^+(^2P)$	0.475
$O^+(^2D) - O^+(^2P)$	1.77
$N^+(^3P) - N^+(^1D)$	2.98
$N^+(^3P) - N^+(^1S)$	0.395
$N^+(^1D) - N^+(^1S)$	0.41

TABLE 4

De-excitation Rate Coefficients for the Low Lying
Metastable States of O^+ and N^+ . T_e is in units of eV.

<u>Transition</u>	<u>g_j</u>	<u>De-excitation Rate Coefficient</u>
$^2D - ^4S$	10	$\frac{1.26 \times 10^{-8}}{\sqrt{T_e}}$
$^2P - ^4S$	6	$\frac{6.33 \times 10^{-9}}{\sqrt{T_e}}$
$^2P - ^2D$	6	$\frac{2.36 \times 10^{-8}}{\sqrt{T_e}}$
$^1D - ^3P$	5	$\frac{4.77 \times 10^{-8}}{\sqrt{T_e}}$
$^1S - ^3P$	1	$\frac{3.16 \times 10^{-8}}{\sqrt{T_e}}$
$^1S - ^1D$	1	$\frac{3.28 \times 10^{-8}}{\sqrt{T_e}}$




Fig. 1 - Electron impact excitation cross section for $N(4S) - N(2D)$. Solid curve Ref. (4), dashed curve Ref. (3), and circles Ref. (5).




Fig. 2 - Electron impact excitation cross section for $N(4S) - N(2)$. Solid curve Ref. (4), dashed curve, Ref. (3), and circles Ref. (5).




Fig. 3 - Electron impact excitation cross section for $N(2D) - N(2)$.
 Solid curve Ref. (4), dashed curve Ref. (3) and circles Ref. (5).




Fig. 4 — Excitation rate coefficients for the low lying metastable states of nitrogen atom as a function of the electron temperature




Fig. 5 — Electron impact excitation cross sections for low lying metastable states of oxygen. Solid curve Ref. (7), dashed curve Ref. (8), and circles Ref. (3).




Fig. 6 — Electron impact excitation rate coefficient for $O(^3P) - O(^1D)$, $O(^3P) - O(^1S)$ and $O(^1D) - O(^1S)$ as a function of the electron temperature

Distribution List

Director
Defense Nuclear Agency
Washington, D. C. 20305
Attn: RAAE, Dr. Harold C. Fitz
RAAE, Dr. Charles A. Blank
RAAE, Mr. Paul B. Fleming
STTL, Technical Library
STSI, (Archives)

Defense Documentation Center
Attn: IC
Cameron Station
Alexandria, VA 22314

Director
U.S. Army Ballistic Research Laboratories
Aberdeen Proving Ground, MD 21005
Attn: AMXBR-CA Franklin Niles

Commander
Harry Diamond Laboratories
2800 Powder Mill Road
Adelphi, MD 20783
Attn: AMXDU-NP

Commander
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, MD 20910
Attn: Code 121, Navy NUC Programs Office

Institute for Defense Analysis
400 Army-Navy Drive
Arlington, VA. 22202
Attn: Dr. E. Bauer
Dr. H. Wolfhard

AF Geophysical Laboratory, AFSC
L. G. Hanscom Field
Bedford, MA 01730
Attn: Dr. John S. Garing
Dr. J. Ulwick
Dr. K. S. W. Champion
Dr. Alva T. Stair
Dr. R. E. Huffman

R&D Associates
Attn: Dr. F. Gilmore
P.O. Box 3580
Santa Monica, CA 90403

Science Applications, Inc.
Attn: Dr. D. Hamlin
P.O. Box 2451
LaJolla, CA 92037

AF Weapons Laboratory, AFSC
Kirtland AFB, NM 87117
Attn: DYT, Maj. Don Mitchell
DYT, Lt. Davis Goetz
SUL, Technical Library

Photometrics, Inc.
Attn: Dr. Irving Kofsky
442 Marrett Road
Lexington, MA 02173

Director
Defense Advanced Research Projects
Agency
Architect Building
1400 Wilson Blvd.
Arlington, VA 22209
Attn: LTCOL W. Whitaker
STO, Capt. J. Justice
Maj. G. Canavan

Director
Naval Research Laboratory
Attn: Dr. T. Coffey
Mr. J. D. Brown
Dr. S. Ossakow
Dr. A. W. Ali (20 copies)
Technical Library, Code 2627

General Electric Company
Tempo-Center for Advanced Studies
816 State Street (P.O. Drawer QQ)
Santa Barbara, CA 93102
Attn: Warren S. Knapp
Art Feryok

General Research Corporation
Attn: Technical Info. (Dr. John Ise)
P.O. Box 3587
Santa Barbara, CA 93105

Mission Research Corporation
735 State Street
Santa Barbara, CA 93101
Attn: Dr. D. Archer
Dr. M. Scheibe