

AD-A032 543

GEORGETOWN UNIV WASHINGTON D C DEPT OF CHEMISTRY
ORGANIC DERIVATIVES OF HETEROPOLY MOLYBDATES.(U)


F/G 7/3

UNCLASSIFIED

NOV 76 M T POPE, C O QUICKSALL, W KWAK
TR-4

N00014-75-C-0945
NL

| OF |
AD
A032543



END
DATE
FILMED
1-77

AD A032543

Office of Naval Research

Contract No. N00014-75A-C-0945

Task No. NR 053-576

Technical Report No.4

Organic Derivatives of Heteropoly Molybdates

by


M.T. Pope*, C.O. Quicksall, W. Kwak, L.M. Rajdović,
J.K. Stalick, K.M. Barkigia and T.F. Scully
Department of Chemistry, Georgetown University,
Washington, D.C.

Prepared for Publication

in

Proceedings of the Second International Conference
on the Chemistry and Uses of Molybdenum

See 1473



Reproduction in whole or in part is permitted for any purpose
of the United States Government.

Approved for public release; distribution unlimited.

401215

OB

M.T. Pope*, C.O. Quicksall, W. Kwak, L.M. Rajković,
J.K. Stalick, K.M. Barkigia and T.F. Scully

Department of Chemistry, Georgetown University,
Washington, D.C. 20057 USA

A survey is given of recent work that characterizes several examples of heteropoly molybdates with covalently attached organic groups. The structures of complexes with the following stoichiometries, $(\text{RP})_2\text{Mo}_3\text{O}_{11}^{3-}$, $(\text{RAS})_2\text{Mo}_3\text{O}_{11}^{3-}$, $(\text{RAS})_4\text{Mo}_{12}\text{O}_{46}^{8-}$, and $\text{R}_2\text{AsMo}_3\text{O}_{11}\text{H}^{2-}$ further illustrate the prevalence of cis MoO_6 groups in polymolybdates, and they also appear to be governed by the dimensions of the $\text{RP}(\text{As})\text{O}_3^{2-}$ and R_2AsO_2^- heterogroups. Novel features of the new complexes include optical activity based on the oxometallate structure of $(\text{RP})_2\text{Mo}_3\text{O}_{11}^{3-}$, neutral zwitterionic species, and a localized non-acidic proton in $\text{R}_2\text{AsMo}_3\text{O}_{11}\text{H}^{2-}$.

Heteropoly molybdate(VI) and tungstate(VI) anions are typically 'ionic' oxometallate clusters of high symmetry, large size, and high ionic weight¹. They are usually the anions of strong acids, they form salts that are soluble in a variety of aqueous and nonaqueous solvents, and they may

also undergo a series of reversible electron transfer reactions to yield mixed valence (V, VI) "heteropoly blues". It has been customary to represent the structures of heteropoly anions as assemblages of metal-oxygen polyhedra which share corners, edges and occasionally faces^{1,2}, and to invoke electrostatic arguments ("Pauling's Rules") to rationalize bond-lengths and angles. However, in spite of the almost identical ionic radii of Mo^{6+} and W^{6+} , there are very few instances of isostructural heteropoly molybdates and tungstates. In general, molybdates tend to adopt structures in which each octahedrally coordinated molybdenum has two cis terminal oxo-oxygens (a notable exception is the Keggin structure for $\alpha\text{-PMo}_3\text{O}_{11}$) whereas tungstates seem to favor structures in which each metal atom has a single terminal oxygen atom. Only anions with the latter structural feature undergo reduction to give heteropoly blues³.

We have recently started to explore the possibility of synthesizing heteropoly complexes that would contain, in addition to the ionic oxometallate cluster, one or more covalently-attached organic groups. Such complexes would represent an entirely new class of compounds that would be expected to display novel combinations of properties, and would lead to applications in areas as diverse as organic synthesis, polymer chemistry, X-ray and electron microscope

ACCESSION for

NTS White Section ☒

DOC Draft Section ☐

UNCLASSIFIED ☐

JUSTIFICATION

BY

DISTRIBUTION / AVAILABLE TO

DATE

APPROVED

1

labelling, and catalysis.

There are several possible routes to complexes of this type, and we summarize here our recent work characterizing new heteropoly molybdates in which the organic groups are directly bound to the heteroatoms. Four classes of such complexes have so far been identified.

1. Pentamolydbisorganophosphonates, $(\text{RP})_2\text{Mo}_5\text{O}_{21}^{1-}$, and Hexamolydbisorganocarsonates, $(\text{RAS})_2\text{Mo}_5\text{O}_{21}^{2-}$

The 2:5 phosphonate⁴ and 2:6 arsonate⁵ complexes are rapidly formed when molybdate and the appropriate organic

anion, RPO_3^{2-} or RASO_3^{2-} , are mixed at pH 4-5. Both types of complexes are characterized by an absorption band at

ca 250 nm, by slight downfield nmr shifts of the R-group protons compared with the uncomplexed acids, and by IR

spectra. Salts of the phosphonate (2:5) complexes with

$\text{R} = \text{H}, \text{CH}_3, \text{C}_2\text{H}_5, \text{C}_6\text{H}_5, \text{C}_2\text{H}_4\text{NH}_2^+$ and $\text{p-CH}_2\text{C}_6\text{H}_4\text{NH}_2^+$ have been

isolated, and the crystal structures of $(\text{NH}_4)_2[(\text{CH}_3\text{P})_2\text{Mo}_5\text{O}_{21}] \cdot$

$5\text{H}_2\text{O}$ (I) and $\text{Na}[(\text{CH}_3)_2[(\text{H}_3\text{NC}_2\text{H}_4\text{P})_2\text{Mo}_5\text{O}_{21}]] \cdot 5\text{H}_2\text{O}$ (II) were

determined⁶. The anions in both structures contain a ring

of five edge- and corner-shared MoO_4 octahedra capped

above and below by the RPO_3 tetrahedra as shown in Figure 1.

The same structural arrangement has been found in the

molybdophosphate anions $\text{H}_n\text{P}_2\text{Mo}_5\text{O}_{21}^{(4-n)-}$ which may be viewed

as inorganic analogues ($\text{R} = \text{O}^{2-}, \text{OH}^-$) of the complexes

described here.⁷

The structure illustrated in Figure 1 is chiral. Compound I crystallizes in space group $P2_1$, and the polarity of the crystal employed in the structure analysis was determined⁶. Although racemization of the optical isomers is expected to be very rapid in hydrolytic solvents, it has been possible to precipitate a diastereomeric brucine salt of $(\text{C}_6\text{H}_5\text{P})_2\text{Mo}_5\text{O}_{21}^{1-}$ which is soluble and stable in acetonitrile. The solution in acetonitrile shows a negative Cotton effect for the 250-nm chromophore of the heteropoly anion as well as the expected optical rotatory dispersion of the brucine present⁸.

The arsonate (2:6) complexes, $\text{R} = \text{CH}_3, \text{C}_6\text{H}_5$, and $\text{pC}_6\text{H}_4\text{NH}_2$, have the structure illustrated in Figure 2. The ring of six as opposed to five MoO_4 octahedra is almost certainly a consequence of the larger size of arsenic over phosphorus. Recently, Pettersson⁹ has proposed an analogous structure for the anion $\text{As}_2\text{Mo}_5\text{O}_{21}^{6-}$ ($= (\text{OAs})_2\text{Mo}_5\text{O}_{21}^{6-}$). Several possibilities for derivatization of the 2:5 and 2:6 complexes may be envisaged, although the heteropoly structures are stable only at pH 2.5-5. Examples of novel zwitterionic species with the 2:5 structure are $[(\text{H}_3\text{NC}_2\text{H}_4\text{P})_2\text{Mo}_5\text{O}_{21}]^-$ (compound II) and the neutral complex $\{[(\text{en})_2\text{Co}(\text{H}_2\text{O})\text{OP})_2\text{Mo}_5\text{O}_{21}]\}$ prepared from $\text{cis}[\text{Co}(\text{en})_2(\text{H}_2\text{O})(\text{HPO}_4)]^+$.

2. Dodecamolybdotetrakisorganoarsonates, $(RAs)_4Mo_{12}O_{46}^{2-}$

In acidic solutions (pH ca 0.5) the 2:6 arsonate complexes are converted to 4:12 species $(RAs)_4Mo_{12}O_{46}^{2-}$. The 4:12 anions with $R = OH^-$ were first prepared by Debray in 1874¹¹, and shown to be tetrameric by Contant in 1973¹². The crystal structure of the ammonium salt, $(NH_4)_4[(HOAs)_4Mo_{12}O_{46}] \cdot 4H_2O$ has recently been reported¹³ and the anion structure is shown in Figure 3. The anion has tetrahedral symmetry and may be viewed as an "inside out" Keggin structure with bridging $HOAsO_3^{2-}$ groups. The central cavity, which appears to be empty, has an estimated radius of 1.43 Å. We have isolated organic derivatives with the 4:12 stoichiometry and $R = CH_3, C_2H_5OH, C_6H_5$, and $pC_6H_4NH_2^+$. Although a crystal structure determination of one of these derivatives is not yet complete, IR and UV spectra of the new complexes are very similar to those of the Debray salt.

3. Tetramolybdodiodiorganoarsinates, $R_2AsMo_4O_{11}H^{2-}$

The 2:5, 2:6, and 4:12 complexes incorporate tetrahedral heteroatoms that have one of the four possible bridging oxygens replaced by an organic group. Organoarsinates, $R_2AsO_3^-$, have two such binding positions blocked. Complexes of molybdate(VI) with $R_2AsO_3^-$ anions are formed at pH 4-5 and have the stoichiometry $R_2AsMo_4O_{11}H^{2-}$. The guanidinium

salt of the dimethyl derivative appears to be the same as a compound first prepared by Rosenheim and Bilecki¹⁴ and formulated by them as $(CN_3H_6)_2H[As(OH)_2(CH_3)_2(Mo_2O_7)_2]$. The crystal structure of $(CN_3H_6)_2[(CH_3)_2AsMo_4O_{11}H] \cdot H_2O$ shows the anion as in Figures 4 & 5¹⁵. This is only the second example (after dodecamolybdocerate¹⁶, $CeMo_{12}O_{48}^{2-}$) of a polyanion structure that involves face-sharing of MO_6 octahedra. As might be expected the molybdenum atoms are displaced from the centres of their respective octahedra towards the unshared edges: metal-metal separations are 3.36 and 3.16 Å through shared edges and faces respectively. The compact structure of the anion is dictated by the steric requirements of the $(CH_3)_2AsO_3^-$ group which suffers very little distortion from 'tetrahedral' symmetry (OAsO angle 106.6°).

The complexes are stable at pH 2-6 in aqueous solution and, with appropriate counterions, in organic solvents. All of the salts prepared contain two cations per anion and must be formulated with an extra proton. Direct evidence for the proton is provided by IR and NMR spectra of anhydrous salts such as $(Bu_4N)_2[(CH_3)_2AsMo_4O_{11}H]$. The IR spectra show a narrow line at 3615 cm^{-1} and the NMR spectra (in methylene chloride) a singlet with the appropriate intensity at ca 2.2 ppm. The proton is not titratable but

undergoes rapid exchange in protic solvents. It is almost certainly located on the unique oxygen that bridges the four molybdenum atoms (see Figure 5) since this is likely to be the most basic. Furthermore the x-ray data for the guanidinium salt show a tightly bound water of hydration 2.8 Å from this particular oxygen.

The structures of the new heteropoly complexes further exemplify the prevalent structural feature for polymolybdates (VI), namely each molybdenum atom with two cis unshared oxo-oxygens³. Consequently, none of these complexes undergoes reversible one or two-electron reductions to heteropoly blue species, but all are irreversibly reduced to molybdenum(V) at the dropping mercury and wax-impregnated graphite electrodes. The structures are 'rational' in the sense that the dimensions of the heterogroups are virtually unchanged upon complexation. The nonbonded O.....O distances in PO_4^{3-} (2.53 Å) and AsO_4^{3-} (2.86 Å) match very closely the appropriate distances in the four heteropoly structures. A dialkylphosphinate, R_2PO_2^- , analogue of the 1:4 complex (Figure 4), could not be synthesized, presumably because the short OPO bridge would further compress the already closely packed structure. However, it may be noted that the dimensions of SO_3^{2-} and SeO_3^{2-} (O.....O = 2.41 and 2.61 Å respectively¹⁷) are appropriate for the 2:5

structure and both $\text{S}_2\text{Mo}_2\text{O}_{11}^{4-}$ and $\text{Se}_2\text{Mo}_2\text{O}_{11}^{4-}$ complexes are well characterized.¹⁸


Acknowledgment. We thank the Office of Naval Research for support of this research through contracts N00014-69-A-0220-0009 and N00014-75-C-0945.



References

1. T.J.R. Weakley, Structure and Bonding, 1974, 18, 131
2. H.T. Evans, Jr., Perspectives in Structural Chem., 1971, 4, 1
3. M.T. Pope, Inorg.Chem., 1972, 11, 1973
4. W. Kwak, M.T. Pope and T.F. Scully, J.Amer.Chem.Soc., 1975, 97, 5735
5. W. Kwak, L.M. Rajković, M.T. Pope, J.K. Stalick and C.O. Quicksall, submitted for publication in Inorg.Chem.
6. J.K. Stalick and C.O. Quicksall, Inorg.Chem., in the press
7. R. Strandberg, Acta Chem.Scand., A, 1973, 27, 1004; B. Hedman, ibid., 1973, 27, 3335; J. Fischer, L. Ricard, and P. Toledano, J.Chem.Soc., Dalton Trans., 1974, 941
8. W. Kwak and M.T. Pope, Unpublished data
9. L. Pettersson, Acta Chem.Scand., A, 1975, 29, 677
10. W. Kwak and M.T. Pope, Inorg.Chem., in the press
11. H. Debray, Compt.rend., 1874, 78, 1411
12. R. Contant, Bull.Soc.Chim.France, 1973, 3277
13. T. Nishikawa and Y. Sasaki, Chemistry Letters, 1975, 1185
14. A. Rosenheim and R. Bilecki, Chem.Ber., 1913, 46, 539
15. K. Barkigia, L.M. Rajković, M.T. Pope and C.O. Quicksall, J.Amer.Chem.Soc., 1975, 97, 4146
16. D.D. Dexter and J.V. Silverton, J.Amer.Chem.Soc., 1968, 90, 3589
17. A.F. Wells, 'Structural Inorganic Chemistry', Clarendon Press, Oxford, 4th edition, p.585
18. A. Rosenheim, Z.anorg.Chem., 1894, 7, 181; R. Lautie, Bull.Soc.Chim.France, 1935, [5], 1, 105; A. Rosenheim and L. Krause, Z.anorg.Chem., 1921, 118, 184; K.Y. Matsumoto, M. Kato and Y. Sasaki, Bull.Chem.Soc.Japan, 1976, 49, 106

Figure Captions

- Figure 1. The structure of $(\text{CH}_3\text{P})_2\text{Mo}_5\text{O}_{21}^{4-}$ (Stalick and Quicksall) showing arrangement of edge- and corner-shared MoO_6 octahedra. The circles represent the phosphorus atoms and one of the methyl groups.
- Figure 2. The structure of $(\text{CH}_3\text{As})_2\text{Mo}_6\text{O}_{24}^{4-}$ (Kwak *et al.*) The circles represent the arsenic atoms and the methyl groups.
- Figure 3. The structure of $(\text{HOAs})_4\text{Mo}_{12}\text{O}_{46}^{4-}$ (Nishikawa and Sasaki). The free vertices of the AsO_4 tetrahedra (shaded) are presumed to be protonated.
- Figure 4. The structure of $(\text{CH}_3)_2\text{AsMo}_4\text{O}_{15}\text{H}^{2-}$ (Barkigia *et al.*), showing edge- and face-shared MoO_6 octahedra and $(\text{CH}_3)_2\text{As}$ group (circles).
- Figure 5. Another view of the structure of $(\text{CH}_3)_2\text{AsMo}_4\text{O}_{15}\text{H}^{2-}$ showing the probable location of the proton.





DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Department of Chemistry Georgetown University Washington, D.C. 20057		2a. REPORT SECURITY CLASSIFICATION unclassified	
		2b. GROUP	
3. REPORT TITLE Organic Derivatives of Heteropoly Molybdates.			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical rept.			
5. AUTHOR(S) (First name, middle initial, last name) Michael T./Pope, Carl O./Quicksall, Wonsuk/Kwak, L.M./Rajdović, Judith K./Stalick, K.M. Baskic-a and T.F. Scully			
6. REPORT DATE November 1976 11 Nov 76		7a. TOTAL NO. OF PAGES 16	7b. NO. OF REFS 18
8a. CONTRACT OR GRANT NO. 15 N00014-759-C-0945		9a. ORIGINATOR'S REPORT NUMBER(S) Technical Report No. 4 14 TR-4	
b. PROJECT NO.		9b. OTHER REPORT NUMBER(S) (Any other numbers that may be assigned this report) 12 10 p.	
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Chemistry Branch Office of Naval Research Arlington, Va. 22217	

13. ABSTRACT

A survey is given of recent work that characterizes several examples of heteropoly molybdates with covalently-attached organic groups. The structures of complexes with the following stoichiometries, $(RP)_2Mo_5O_{21}^{4-}$, $(RAS)_2Mo_6O_{24}^{4-}$, $(RAS)_4Mo_{12}O_{46}^{4-}$, and $R_2AsMo_4O_{15}H^{2-}$ further illustrate the prevalence of cis MoO_6 groups in polymolybdates, and the also appear to be governed by the dimensions of the $RP(As)O_3^{2-}$ and R_2AsO_2 heterogroups. Nine features of the new complexes include optical activity based on the oxometallate structure of $(RP)_2Mo_5O_{21}^{4-}$, neutral zwitterionic species, and a localized non-acidic proton in $R_2AsMo_4O_{15}H^{2-}$.

14.

KEY WORDS

LINK A

LINK B

LINK C

ROLE

WT

ROLE

WT

ROLE

WT

Heteropoly complexes

Organic derivatives

Molybdophosphonates

Molybdoarsonates

Survey