

AD-A032 681

BROWN UNIV PROVIDENCE R I DIV OF ENGINEERING
A NOTE ON THE ALGEBRAIC MATRIX RICCATI EQUATION. (U)
1976 A E PEARSON, W H KWON

F/G 12/1

UNCLASSIFIED

AFOSR-TR-76-1154

AF-AFOSR-2793-75

NL

[OF]
AD
A032681



END

DATE
FILMED
1-77

AD A032681

AFOSR - TR - 76 - 1154

A Note on the Algebraic Matrix Riccati Equation*

2

by

W. H. Kwon and A. E. Pearson

Division of Engineering
and
Lefschetz Center for Dynamical Systems
Brown University
Providence, Rhode Island 02912

Abstract

The results in [1,2] for the matrix Lyapunov equation are extended to the case of an algebraic matrix Riccati equation. Some errors in [1,2] are pointed out by a counter example. The estimations obtained in this note are shown to be exact for certain cases. Similar results are possible for the discrete algebraic matrix Riccati equation.

See Form 1473

Approved for public release;
distribution unlimited.

DDC
RECEIVED
DEC 3 1976
A.

* This research was supported in part by the Air Force Office of Scientific Research under Grant AFOSR-75-2793E.

COPY AVAILABLE TO DDC DOES NOT
PERMIT FULLY LEGIBLE PRODUCTION

I. Introduction

Lower bounds of the solution K to the matrix Lyapunov equation

$$A'K + KA + Q = 0 \quad (1)$$

have been obtained in [1,2] in terms of $\{A,Q\}$. Equation (1) is related to the linear constant homogeneous system

$$\dot{x}(t) = Ax(t), \quad x(t_0) = x_0. \quad (2)$$

The well known result is that the system (2) is asymptotically stable if and only if for each positive definite matrix Q there exists a positive definite solution K to (1). We consider in this note the algebraic matrix Riccati equation

$$A'K + KA - KBB'K + Q = 0 \quad (3)$$

which is related to the linear constant dynamical system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad x(t_0) = x_0. \quad (4)$$

The corresponding result is that the system (4) is stabilizable if and only if for each positive definite matrix Q there exists a positive definite solution K to (3). We will obtain lower bounds of the solution to (3) in this note. Results for (3) should coincide with those of (1) when $B = 0$. Bounds for the finite time solution of the matrix differential Riccati equation are given in [4,5] for continuous systems, and in [6,7] for discrete systems, in terms of the controllability and observability matrices. These are compared with the results in this note for some special cases. Applications of bounds for (1) are listed in [1,2] and similar applications for bounds of (3) are possible.

II. Main Results

In the following, $|x|$ denotes the Euclidean vector norm and $\|C\|$ the matrix norm induced by the Euclidean norm, i.e.,

$$\|C\| = \sup_{|x|=1} |Cx| = \max_i \lambda_i^{1/2}(C'C) \quad (5)$$

where $\lambda_i(M)$ denotes an eigenvalue of a matrix M . For a nonnegative definite matrix S it holds that $\|S\| = \sup_{|x|=1} x'Sx = \lambda_{\max}(S)$ and $\inf_{|x|=1} x'Sx = \lambda_{\min}(S)$.

Theorem 1. Assume that Q is a symmetric positive definite matrix. The positive definite solution matrix K to the algebraic matrix equation (3) has the following bounds:

$$K \geq \frac{\lambda_{\min}(Q)}{\|A\| + \{\|A\|^2 + \|BB'\| \lambda_{\min}(Q)\}^{1/2}} I \quad (6)$$

$$\|K\| \geq \frac{\lambda_{\max}(Q)}{\|A\| + \{\|A\|^2 + \|BB'\| \lambda_{\max}(Q)\}^{1/2}} \cdot \quad (7)$$

Proof: Pre- and post-multiplying (3) by x' and x respectively yields

$$x'KBB'Kx - 2x'A'Kx - x'Qx = 0,$$

which can be expressed as

$$x'KBB'Kx + 2|x'A'Kx| - x'Qx \geq 0. \quad (8)$$

From the Schwarz Inequality we have

$$|x'A'Kx| \leq |Ax| |Kx| \quad (9)$$

$$x'KBB'Kx \leq |BB'Kx| |Kx| \leq \|BB'\| |Kx|^2. \quad (10)$$

Combining (8), (9), and (10) yields

$$\|BB'\| \cdot |Kx|^2 + 2|Ax| \cdot |Kx| - x'Qx \geq 0. \quad (11)$$

Since $|Kx|$ is nonnegative, it should satisfy

$$|Kx| \geq \frac{1}{\|BB'\|} \{-|Ax| + \{|Ax|^2 + \|BB'\| x'Qx\}^{1/2}\} = \frac{x'Qx}{|Ax| + \{|Ax|^2 + \|BB'\| x'Qx\}^{1/2}} \cdot \quad (12)$$

The last equality in (12) follows from the following identity

$$\{-a + \sqrt{a^2 + bc}\} \{a + \sqrt{a^2 + bc}\} = bc.$$

Dividing both sides of (12) by $|x|$ yields

$$\frac{|Kx|}{|x|} \geq \frac{\frac{x'Qx}{|x|^2}}{\frac{|Ax|}{|x|} + \left\{ \frac{|Ax|^2}{|x|^2} + \|BB'\| \frac{x'Qx}{|x|^2} \right\}^{1/2}} \geq \frac{\frac{x'Qx}{|x|^2}}{\|A\| + \left\{ \|A\|^2 + \|BB'\| \frac{x'Qx}{|x|^2} \right\}^{1/2}}. \quad (13)$$

For $a > 0$ and $b > 0$, it can be shown that the function

$$f(x) = \frac{x}{a + \sqrt{a^2 + bx}} \quad (14)$$

is monotonically increasing with respect to x for $x \geq 0$. Thus we have

$$K \geq \lambda_{\min}(K)I = \inf_x \frac{|Kx|}{|x|} I \geq \frac{\lambda_{\min}(Q)}{\|A\| + \{ \|A\|^2 + \|BB'\| \lambda_{\min}(Q) \}^{1/2}} I \quad (15)$$

and

$$\|K\| = \lambda_{\max}(K) \geq \frac{\lambda_{\max}(Q)}{\|A\| + \{ \|A\|^2 + \|BB'\| \lambda_{\max}(Q) \}^{1/2}}. \quad (16)$$

This completes the proof.

By taking $B = 0$, the results in (6) and (7) coincide with those of [1,2]. The results in (6) and (7) are sharp in the sense that its estimations are exact in some cases as shown in the following example. It is also demonstrated in the example that the estimations in Theorem 1 are better than those of [4,8] for some special cases of $Q > 0$.

Example: (a) Let $A = -I$, $B = I$, $Q = 3I$. Then $\|A\| = \|BB'\| = 1$. From (6) and (7), $K \geq I$ and $\|K\| \geq 1$. The exact solution to (3) is $K = I$. This shows that the estimations in (6) and (7) are exact in this case. In [4] it is suggested that $K \geq \left(\frac{2}{3(1-e^{-2T})} + \frac{e^{2T}-1}{2} \right)^{-1}$ for any $T > 0$, which implies $K \geq \frac{3}{4}(\sqrt{3}-1)I$. Also in [8] it is suggested that $K \geq \frac{3}{4}I$.

(b) Let $A = 0$, $B = bI$, and $Q = q^2I$. From (6) and (7) $K \geq \frac{q}{b}I$ and $\|K\| \geq \frac{q}{b}$. The exact solution to (3) is $K = \frac{q}{b}I$. Thus the estimations of (6) and (7) in this case are exact. It is suggested in [4] that $K \geq \left(-\frac{1}{2} + b^2T \right)^{-1}I$ for

any $T \geq 0$, which implies $K \geq \frac{q}{2b} I$.

By a slight modification the results in Theorem 1 can be applied to the discrete algebraic matrix Riccati equation

$$K = \Phi'K\Phi - \Phi'KB(I + B'KB)^{-1}B'K\Phi + Q, \quad (17)$$

which can be transformed to

$$K\Phi^{-1}BB'K + K\Phi^{-1} - (Q\Phi^{-1}BB' + \Phi')K - Q\Phi^{-1} = 0. \quad (18)$$

The corresponding result can be given as follows.

Theorem 2. Assume that Φ is nonsingular, Q nonnegative definite, and $(Q\Phi^{-1} + \Phi'^{-1}Q)$ nonsingular. The positive definite solution matrix K to the discrete algebraic matrix Riccati equation (17) has the following bounds:

$$K \geq 2 \frac{\gamma_{\min}(Q\Phi^{-1})}{\|G\| + \{\|G\|^2 + 4\|BB'\Phi'^{-1}\| \gamma_{\min}(Q\Phi^{-1})\}^{1/2}} I \quad (19)$$

$$\|K\| \leq 2 \frac{\gamma_{\max}(Q\Phi^{-1})}{\|G\| + \{\|G\|^2 + 4\|BB'\Phi'^{-1}\| \gamma_{\max}(Q\Phi^{-1})\}^{1/2}} \quad (20)$$

where

$$G = \Phi^{-1} - \Phi - BB'\Phi'^{-1}Q \quad (21)$$

$$\gamma_{\min}(C) = \frac{1}{2} \min_i |\lambda_i(C + C')| \quad (22)$$

$$\gamma_{\max}(C) = \frac{1}{2} \max_i |\lambda_i(C + C')| \quad (23)$$

The results in Theorem 2 can be obtained from the fact that $\inf_{|x|=1} |x'Cx| = \gamma_{\min}(C)$ and $\sup_{|x|=1} |x'Cx| = \gamma_{\max}(C)$. By taking $B = 0$ the results in Theorem 2 apply to the discrete Lyapunov matrix equation, $\Phi'K\Phi - K = -Q$. It is also true that the estimations in (19) and (20) are exact for some special cases and thus better than those estimations in [7] in certain cases.

III. Correction to [2,1]

In this section we use the same notations as [2,1]. Since the equation (17)

in [2],

$$\|(A')^{-1}PA^{-1}\|_2 = \frac{\beta_n}{\sigma_1}, \quad (24)$$

is not generally true, Theorem 2 in [2] which claims

$$\alpha_n \geq \frac{\beta_n}{2\sigma_1^{1/2}} \quad (25)$$

is invalid.

Counter example: Let $P = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ and $A = \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$. Then $A'^{-1}PA^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$,

$\|A'^{-1}PA^{-1}\|_2 = 1$, $\beta_n = 2$, and $\sigma_1 = 1$. Thus the inequality (24) does not hold. The solution matrix Q of the Lyapunov equation is $Q = \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$. Thus $\alpha_n = \frac{1}{2}$ which shows that the relation (25) is invalid.

The relation (25) also appeared in [1] and should be changed to $\alpha_n \geq \frac{\beta_n}{2\sigma_n^{1/2}}$, which can be obtained from (7) by taking $B = 0$.

ACCESSION OF	
RTIS	With Section <input checked="" type="checkbox"/>
DOC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. END. OF SPECIAL
A	

References

- [1] E. Y. Shapiro, "On the Lyapunov matrix equation," IEEE Trans. Automat. Contr., vol. AC-19, pp. 594-596, October 1974.
- [2] J. J. Montemayer and B. F. Womack, "Comments on 'On the Lyapunov matrix equation'", IEEE Trans. Automat. Contr., vol. AC-20, pp. 814-815, December 1975.
- [3] R. S. Bucy, "Global Theory of the Riccati equation," J. of Computer and System Science: 1, pp. 349-361, 1967.
- [4] R. S. Bucy, "The Riccati equation and its bounds," J. of Computer and System Sciences; 6, pp. 343-353, 1972.
- [5] R. Canales, "A lower bound on the performance of optimal regulators," IEEE Trans. Automat. Contr., vol. AC-15, pp. 409-415, 1970.
- [6] J. J. Deyst, Jr., and C. F. Price, "Conditions for asymptotic stability of the discrete minimum-variance linear estimator," IEEE Trans. Automat. Contr., vol. AC-13, pp. 702-705, 1968.
- [7] J. J. Deyst, Jr., "Corrections for the above paper," IEEE Trans. Automat. Contr., vol. AC-18, 1973.
- [8] R. W. Brockett, *Finite Dimensional Linear Systems*, John Wiley & Sons, Inc., New York, 1970.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

19 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER	
18 AFOSR - TR - 76 - 1154			
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED	
6 A NOTE ON THE ALGEBRAIC MATRIX RICCATI EQUATION.		7 Interim rept.	
7. AUTHOR(s)		6. PERFORMING ORG. REPORT NUMBER	
10 W. H. /Kwon A. E. /Pearson		8. CONTRACT OR GRANT NUMBER(s)	
		15 VAF - AFOSR W-2793-75	
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Brown University Division of Engineering Providence, Rhode Island 02912		61102F 9769-01 17 p1	
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE	
Air Force Office of Scientific Research (NM) Bolling AFB, DC 20332		11 1976	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES	
12 8 p.		6	
		15. SECURITY CLASS. (of this report)	
		UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)			
Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)			
The results for the matrix Lyapunov equation are extended to the case of an algebraic matrix Riccati equation. Some errors are pointed out by a counter example. The estimations obtained in this note are shown to be exact for certain cases. Similar results are possible for the discrete algebraic matrix Riccati equation.			