

AD-A032 690

NORTH CAROLINA UNIV AT CHAPEL HILL DEPT OF STATISTICS F/G 12/1  
ON THE EXPONENTIAL BOUNDEDNESS OF STOPPING TIMES OF INVARIANT S--ETC(U)  
JUL 76 H ROOTZEN, G SIMONS N00014-75-C-0P09  
MIMEO SER-1076 NL

UNCLASSIFIED

1 of 1  
ADA032690



END  
DATE  
FILMED  
2 - 77

AD A 0 32690

(12) jr

On the Exponential Boundedness of Stopping Times of Invariant SPRT's

by

Holger Rootzén<sup>1</sup>

*Department of Mathematical Statistics  
University of Lund*

*and  
Department of Statistics  
University of North Carolina at Chapel Hill*

and

Gordon Sirons<sup>2</sup>

*Department of Statistics  
University of North Carolina at Chapel Hill*

Institute of Statistics Mimeo Series #1076

July, 1976

DDC  
NOV 30 1976  
RECEIVED

**COPY AVAILABLE TO DDC DOES NOT  
PERMIT FULLY LEGIBLE PRODUCTION**

- <sup>1</sup> Research sponsored in part by the Office of Naval Research under Contract N00014-75-C-0309.
- <sup>2</sup> Research supported by the National Science Foundation under Contract MPS75-07556.

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) 6 On the Exponential Boundedness of Stopping Times of Invariant SPRT's.		5. TYPE OF REPORT & PERIOD COVERED 9 TECHNICAL rept.
7. AUTHOR(s) 10 Holger/Rootzen & Gordon/Simons		6. PERFORMING ORG. REPORT NUMBER 14 Mimeo Series - 1076
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of North Carolina Chapel Hill, North Carolina 27514		8. CONTRACT OR GRANT NUMBER(s) 15 N00014-75-C-0809
11. CONTROLLING OFFICE NAME AND ADDRESS Statistics & Probability Program Office of Naval Research, Attn: Dr. McDonald Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12/10 p.
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 11 July 1976
		13. NUMBER OF PAGES 7
		18. SECURITY CLASS. (of this report) UNCLASSIFIED
		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for Public Release: Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Invariance, sequential probability ratio test, exponential boundedness, likelihood ratio, stopping time		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  It is shown, under conditions which include invariant sequential probability ratio tests, that the stopping time is always exponentially bounded when the null or alternative hypothesis holds, except in a trivial instance.		

mt

On the Exponential Boundedness of Stopping Times of Invariant SPRT's

Short Title: Exponential Boundedness of Stopping Times

by

Holger Rootzén<sup>1</sup>      University of Lund and University of North Carolina  
Gordon Simons<sup>2</sup>      University of North Carolina


Abstract: It is shown, under conditions which include invariant sequential probability ratio tests, that the stopping time is always exponentially bounded when the null or alternative hypothesis holds, except in a trivial instance.

AMS Classification: Primary 62L10  
Secondary 60G17, 60G40

Key Words and Phrases: Invariance, sequential probability ratio test, exponential boundedness, likelihood ratio, stopping time

<sup>1</sup> Research sponsored in part by the Office of Naval Research under Contract N00014-75-C-0809.

<sup>2</sup> Research supported by the National Science Foundation under Contract MPS75-07556.



This paper is primarily concerned with showing that, for invariant sequential probability ratio tests, the stopping time is exponentially bounded under the null and alternative hypotheses, with the exception of trivial situations. This work is the by-product of a largely unsuccessful attempt to obtain general results under non-model distributions. Since we view the lack of these general results as a major deficiency in the theory of sequential analysis, we shall suggest what we believe are reasonable conjectures and shall discuss some of the past literature on the subject.

Let  $(\Omega, \mathcal{B})$  be a measurable space and  $\{\mathcal{B}_n, n \geq 1\}$  be a nondecreasing sequence of sub- $\sigma$ -fields of  $\mathcal{B}$ . A stopping time  $N$  adapted to  $\{\mathcal{B}_n, n \geq 1\}$ <sup>3</sup> is said to be *exponentially bounded* for a family of probability measures  $\mathcal{R}$  on  $(\Omega, \mathcal{B})$  if for each probability measure  $R \in \mathcal{R}$  there exist constants  $c > 0$  and  $\rho < 1$  such that  $R(N > n) < c\rho^n$ ,  $n = 1, 2, \dots$ . Such a condition implies that  $N$  is finite a.s. ( $R$ ) and that  $N$  has a moment generating function in some neighborhood of the origin for each  $R \in \mathcal{R}$ .

Let  $P$  and  $Q$  be elements in  $\mathcal{R}$  and let  $L_n$  denote the  $P - Q$  likelihood ratio for  $\mathcal{B}_n$ .<sup>4</sup> Sequential probability ratio tests are defined in terms of a stopping variable  $N$  of the general form

<sup>3</sup> Here,  $N$  is said to be adapted to  $\{\mathcal{B}_n, n \geq 1\}$  if  $[N = 1, 2, \dots \text{ or } \infty] = \Omega$  and  $[N = n] \in \mathcal{B}_n$  for  $n = 1, 2, \dots$ .

<sup>4</sup> For a precise definition of this concept see Eisenberg, Ghosh and Simons (1976). Essentially,  $L_n$  is the Radon-Nikodym derivative  $dQ/dP$  relative to  $\mathcal{B}_n$ .

- (1)  $N =$  the first  $n \geq 1$  such that  $L_n \notin (A, B)$   
 $= \infty$  if no such  $n$  exists,

where  $0 < A < B < \infty$ . At this level of generality, it is easy to construct examples for which  $N$  is exponentially bounded for  $\{P, Q\}$  and others for which it is not. All that is known, in general, is derived from the following set of inequalities, described by Eisenberg, Ghosh and Simons (1976):

$$AP(N > n) \leq Q(N > n) \leq BP(N > n), \quad n = 1, 2, \dots$$

It follows immediately that

- (a)  $P(N < \infty) = 1$  iff  $Q(N < \infty) = 1$ ,  
 (b) for each  $r > 0$ ,  $\int N^r dP < \infty$  iff  $\int N^r dQ < \infty$ , and  
 (c)  $N$  is exponentially bounded for  $\{P\}$  iff it is exponentially bounded for  $\{Q\}$ .

Now suppose  $X_1, X_2, \dots$  is a sequence of random elements on  $(\Omega, \mathcal{B})$  (to the statistician, "potential data") and  $\mathcal{B}_n = \sigma(X_1, \dots, X_n)$  (the  $\sigma$ -field generated by  $X_1, \dots, X_n$ ). If  $X_1, X_2, \dots$  is an i.i.d. sequence under  $P$  and  $Q$ , then  $\{\log L_n, n \geq 1\}$  is a random walk under both  $P$  and  $Q$ . It follows that  $N$  is exponentially bounded for  $\{P, Q\}$  unless  $L_1 = 1$  a.s.  $(P, Q)$ . This is a well-known result due to Stein (1946). We shall now describe a similar result for invariant sequential probability ratio tests.

Consider the following mathematical structure:

- (i)  $\mathcal{P}$  and  $\mathcal{Q}$  are two disjoint families of probability measures on  $(\Omega, \mathcal{B})$ .
- (ii)  $X_1, X_2, \dots$  is an i.i.d. sequence of random elements for each probability measure  $R \in \mathcal{P} \cup \mathcal{Q}$ .
- (iii) For each  $n \geq 1$ ,  $\mathcal{B}_n = \sigma(T_n)$  where  $T_n$  is a statistic which is a symmetric function of  $X_1, \dots, X_n$ .
- (iv) For each  $n \geq 1$ ,  $T_k$  is  $\mathcal{B}_n$ -measurable for  $k = 1, 2, \dots, n$ .  
(I.e.,  $\mathcal{B}_1 \subset \mathcal{B}_2 \subset \dots$ .)
- (v) For each  $n \geq 1$ , the probability space  $(\Omega, \mathcal{B}_n, R)$  is the same for every  $R \in \mathcal{P}$ , and is the same for every  $R \in \mathcal{Q}$ .

This is the structure that one encounters when invariant sequential probability ratio tests are being considered. If one wishes to test the (composite) hypothesis that the true probability measure  $R$  belongs to  $\mathcal{P}$  against the (composite) alternative that it belongs to  $\mathcal{Q}$ , the following sequential probability ratio test suggests itself: Let  $P \in \mathcal{P}$  and  $Q \in \mathcal{Q}$  be chosen in an arbitrary manner, and let  $L_n$  be the  $P - Q$  likelihood ratio for  $\mathcal{B}_n$  ( $n \geq 1$ ). Because of property (v), each possible pair  $(P, Q)$  leads to the same sequence of likelihood ratios  $L_1, L_2, \dots$ . Let  $N$  be the stopping variable described in (1) and choose  $P$  or  $Q$  according as  $N < \infty$  and  $L_N \leq A$  or  $N < \infty$  and  $L_N \geq B$ , respectively. (If  $N = \infty$ , no decision is made.) We have the following theorem:

THEOREM *Either  $L_n = 1$  a.s. ( $\mathcal{P} \cup \mathcal{Q}$ ) for  $n \geq 1$ , or  $N$  is exponentially bounded for  $\mathcal{P} \cup \mathcal{Q}$ .*

PROOF In view of (c) and (v),  $N$  is exponentially bounded for  $P \cup Q$  iff it is exponentially bounded for  $\{P\}$ . Likewise,  $L_n = 1$  a.s. ( $P \cup Q$ ) for  $n \geq 1$  iff  $P(L_n = 1) = 1$  for  $n \geq 1$ . Suppose the latter is false, i.e., that for some  $k$ ,  $P(L_k = 1) < 1$ . For simplicity, we shall assume that  $k$  can be taken to be unity. If a larger value of  $k$  is required, the proof we shall give that  $N$  is exponentially bounded for  $\{P\}$  can be easily modified. Express  $T_1$  as  $t(X_1)$  and  $L_1$  as  $\ell(X_1)$  and observe that  $EL_1 \leq 1$  and (since  $P(L_1=1) < 1$ ) that  $E \log L_1 < 0$ . Set  $B'_n = \sigma(t(X_1), \dots, t(X_n))$  and observe that  $L'_n = \prod_{i=1}^n \ell(X_i)$  is the  $P - Q$  likelihood ratio for  $B'_n$  in view of condition (ii) ( $n \geq 1$ ). Moreover, conditions (iii) and (iv) imply that  $B'_n \subset B_n$  and, hence,  $E^{B'_n} L'_n \leq L'_n$ <sup>5</sup> for  $n \geq 1$ . Thus, for  $\rho < 1$ , to be chosen later,

$$P^{B'_n}(L'_n > A, L'_n \leq \rho^n) \leq A^{-1} I_{[L'_n \leq \rho^n]} E^{B'_n} L'_n \leq A^{-1} \rho^n$$

and, hence,

$$\begin{aligned} P(N > n) &\leq P(L'_n > A) = E P^{B'_n}(L'_n > A) \\ &\leq A^{-1} \rho^n + P(L'_n > \rho^n). \end{aligned}$$

Thus it suffices to show, for properly chosen  $\rho < 1$  and  $\rho' < 1$ , that

$$P(L'_n > \rho^n) \leq (\rho')^n.$$

Since  $EL_1 \leq 1$ ,  $E(L_1)^t \leq 1$  for  $0 < t \leq 1$ . In fact, if  $\rho$  is chosen so that  $E \log L_1 < \log \rho < 0$ , then there exists a small positive  $t_0$  such that

<sup>5</sup> Equality holds if  $P$  and  $Q$  are equivalent probability measures on  $B_n$ .

$$\rho' \equiv E e^{t_0(\log L_1 - \log \rho)} < 1.$$

Then

$$\begin{aligned} P(L'_n > \rho^n) &= P\left(\prod_{i=1}^n \ell(X_i) > \rho^n\right) \\ &\leq \rho^{-nt_0} E\left[\prod_{i=1}^n \ell(X_i)\right]^{t_0} \\ &= \rho^{-nt_0} E^n(L_1)^{t_0} = (\rho')^n. \quad \square \end{aligned}$$

We should point out that it is always possible that  $L_n = 1$  a.s. ( $P \cup Q$ ) for  $n \geq 1$ . (This is the case, for instance, if  $T_n \equiv 1$  for  $n \geq 1$ .) However, this possibility does not detract from the strength of the theorem. For if  $L_n = 1$  a.s. ( $P \cup Q$ ) for  $n \geq 1$ , then, for each  $n \geq 1$ , the probability spaces  $(\Omega, \mathcal{B}_n, R)$ ,  $R \in P \cup Q$ , are all the same and there is no information available for distinguishing between the two families  $P$  and  $Q$ . In practice, one would never consider proposing such a test.

A close examination of our proof reveals that conditions (ii) and (iii) are stronger than necessary. We shall not try to state the weakest possible conditions, but simply make the observation that the argument works because the  $\sigma$ -fields  $\mathcal{B}_n$ ,  $n \geq 1$ , grow at a sufficiently fast rate under conditions (ii) and (iii). Note also that the proof shows exponential boundedness under  $P$  for the "single boundary" stopping time

$$\begin{aligned} N^* &= \text{the first } n \geq 1 \text{ such that } L_n \geq A, \\ &= \infty \text{ if no such } n \text{ exists.} \end{aligned}$$

It is reasonable to ask whether our theorem has an extension to probability measure  $R \notin P \cup Q$  under which  $X_1, X_2, \dots$  is an i.i.d.

sequence. It is possible to extend it to other probability measures within a monotone likelihood ratio class (cf., A. Ifram (1965)) but this amounts to a very restrictive improvement. R. Wijsman (1970) has found an interesting example for which  $R(N<\infty) = 1$  but  $N$  is not exponentially bounded under  $R$ . This would seem to put to rest any hope of finding a general extension of our theorem. However, it must be remembered that a  $P - Q$  likelihood ratio is uniquely defined only up to a  $P$  and  $Q$  equivalence, and, in Wijsman's example, the probability measure  $R$  is *orthogonal* to the probability measures in  $P \cup Q$  on the  $\sigma$ -fields  $\mathcal{B}_n$ . Indeed there exist other versions of the likelihood ratios  $L_n$ , appropriate to his example (albeit less natural from a topological viewpoint), for which  $N$  is exponentially bounded under  $R$ . Recently, Wijsman (1976a) has been working with some examples for which  $R$  is dominated (on each  $\mathcal{B}_n$ ) by the probability measures in  $P \cup Q$ . (In such a case, the likelihood ratios  $L_n$  are uniquely specified up to an  $R$ -equivalence.) For these, again,  $R(N<\infty) = 1$  but  $N$  is not exponentially bounded under  $R$ . We believe it is significant that the probability measures  $R$ , in his examples, are *not* equivalent to the probability measures in  $P \cup Q$  (on each  $\mathcal{B}_n$ ). For one thing, there exist examples, within the context of conditions (i) - (v), for which  $R$  is dominated by but is not equivalent to the probability measures in  $P \cup Q$ , with the property that  $R(N=\infty) = 1$ , and yet the condition " $R(L_n=1) = 1$  for  $n \geq 1$ " does not hold. However, on the basis of our studies, we are prepared to conjecture that  $R(N<\infty) = 1$ , whenever  $R$  and the probability measures in  $P \cup Q$  are equivalent on  $\sigma(\mathcal{X}_1)$  and the probability spaces  $(\Omega, \mathcal{B}_n, P)$  and  $(\Omega, \mathcal{B}_n, Q)$  are distinct

for some  $n \geq 1$ ,  $P \in \mathcal{P}$ ,  $Q \in \mathcal{Q}$ . Indeed we suspect that these conditions guarantee that  $N$  is exponentially bounded under  $R$ . Here, we are assuming the structure imposed by conditions (i) - (v).

We do not want to leave the reader with the impression that there are no results of a general nature concerning  $R \downarrow P \cup Q$ . R. Berk (1970) has obtained results for what he calls *parametric* sequential probability ratio tests and R. Wijsman (1976b) has recently obtained a theorem concerned with exponential boundedness. Their results exploit the fact that, in many useful examples of invariant sequential probability ratio tests, the log-likelihood ratios  $\{\log L_n, n \geq 1\}$  become a random walk asymptotically (under  $R$ ) as  $n \rightarrow \infty$ . This is not always the case. (There are practical examples to the contrary.) Therefore, it appears that some substantially new approaches are called for.

#### REFERENCES

- [1] Berk, Robert (1970). Stopping times of SPRTs based on exchangeable models. *Ann. Math. Statist.* 41, 979-990.
- [2] Eisenberg, Ghosh and Simons (1976). Properties of generalized sequential probability ratio tests. *Ann. Statist.* 4, 237-251.
- [3] Ifram, Adnan (1965). On the asymptotic behavior of densities with applications to sequential analysis. *Ann. Math. Statist.* 36, 615-637.
- [4] Stein, Charles (1946). A note on cumulative sums. *Ann. Math. Statist.* 17, 498-499.
- [5] Wijsman, Robert (1972). Examples of exponentially bounded stopping time of invariant sequential probability ratio tests when the model may be false. *Proc. Sixth Berkeley Symp. Math. Statist. Prob.* 1, 109-128.
- [6] \_\_\_\_\_ (1976a). Obstructive distributions in a sequential rank-order test based on Lehmann alternatives. Unpublished manuscript.
- [7] \_\_\_\_\_ (1976b). A general theorem with applications on exponentially bounded stopping time, without moment conditions. Unpublished manuscript.