

AD-A033 057

SYDNEY UNIV (AUSTRALIA) DEPT OF AERONAUTICAL ENGINEERING F/G 20/4
NUMERICAL STUDIES IN GAS DYNAMICS.(U)
OCT 76 G A BIRD

UNCLASSIFIED

AFOSR-TR-76-1178


AF-AFOSR-2336-72
NL

| OF |
AD
A033057
M
M
M



END

DATE
FILMED
1 - 77



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS - 1963 - A

AFOSR - TR - 76 - 1178

J

12

FINAL SCIENTIFIC REPORT

RESEARCH GRANT AFOSR-72-2336

NUMERICAL STUDIES IN GAS DYNAMICS

ADA 033057

Submitted to the

Air Force Office of Scientific Research

by the

Department of Aeronautical Engineering

University of Sydney

Sydney, N.S.W. Australia.

14th October, 1976.

DDC
FORM 12
DEC 7 1976
REGISTRATION
C

Principal Investigator: G.A. Bird,
Professor of Aeronautical Engineering.

COPY AVAILABLE TO DDC DOES NOT
PERMIT FULLY LEGIBLE PRODUCTION

This document summarizes work supported by Grant AFOSR-72-2336 over the complete grant period April 15, 1972 to October 14, 1976 and describes the work carried out during the period from April 15, 1974.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

Handwritten initials or mark in a circle.

RESEARCH REPORT

7200330700

APPROVED FOR RELEASE
DATE 10-1-80
BY [illegible]

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is approved for public release IAW AFR 190-12 (7b). Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

DATA MONITORING
[illegible text]

1. REVIEW OVER COMPLETE GRANT PERIOD 1972-1976

1.1 Major Developments

Most of the research has been associated with the further development and application of the direct simulation Monte Carlo method for the modeling of gas flows at the molecular level. A monograph¹ which includes a full description of the method, together with a number of new applications, has been written and published during this period. Acknowledgment was made generally of the support provided by the AFOSR during the whole period over which the method has been developed and specifically to the current grant with respect to some new applications that appear in the book.

A significant development has been the demonstration that the computations are best carried out on a dedicated minicomputer rather than on a large machine in a computing center. The probability of this was proposed for preliminary investigation in 1973 and this led to a proposal to the Australian Research Grants Committee for a PDP 11/40 system. This submission was successful and the computer was installed early in 1975. Since this time, it has run twenty four hours a day and seven days a week without an operator and the improvement in cost-effectiveness has been found to be near two orders of magnitude. The reduction in computing costs has meant that small disturbance flows, which had previously been thought to be inaccessible to the method, can now be studied. The first demonstration of this capability was for the classical surface slip or Kramers problem². The flow profile was studied at Mach numbers as low as 0.05, with the computer runs being sufficiently long to reduce the statistical scatter to one part in a thousand.

An additional consequence of the practicability of very long runs is that molecular simulation can now be considered as an alternative to finite difference methods for the computation of some continuum flows. This point was first demonstrated³ by lengthy runs on a large computer and a more recent study⁴ has further demonstrated the advantages of the dedicated minicomputer. The advantages of the molecular simulation over the finite difference continuum representation relate to the absence of stability problems and the ease of incorporation of complex flow boundaries. The molecular approach appears to be most advantageous for problems involving complex shock wave interactions, but having inviscid boundaries. On the other hand, the method has been used⁵ to study a viscous problem at sufficiently low Knudsen number that the conclusions should be valid in continuum flow.

The most recent extension of the method has been to reacting gas flows. Particular attention has been paid to the dissociation-recombination reaction in a diatomic gas. The model permits the analytical determination of the equilibrium state¹ and it has now been applied to the reaction zone behind a strong shock wave. Specific cases have been computed

2. REPORT ON PERIOD 1975-1976.

The final extension of the Grant was from April 15, 1975 to April 14, 1976 but, due to a period of absence of the Principal Investigator, this was extended to October 14, 1976. Four topics were included in the proposal and the work done on each of these is described below.

2.1 Collision Rates and Collisional Energy Distribution within Shock Waves.

This work led to publication 7 in the list of Section 1.2. The direct simulation Monte Carlo method was used to determine the collision rate and the distribution of relative translational energy between colliding molecules within strong shock waves. The results were compared with the corresponding theoretical quantities in an equilibrium gas. The objective was to determine whether the region of marked translational nonequilibrium within the wave needs to be taken into account in the interpretation of reaction rate experiments in shock tubes. It was found that a correction should be applied to measured relaxation distances behind shock waves. The magnitude of this correction is of the order of ten times the translational shock thickness.

2.2 Calculations with Quantum Mechanics Cross-sections.

Dr. A.U. Chatwani of the Max-Planck Institut für Strömungsforschung spent six weeks in the Aeronautical Engineering Department and incorporated quantum theory collision cross-sections into the existing simulation program for a steady spherical expansion. This program had been used to study the onset of translational non-equilibrium, and this generally occurs at very low temperatures where quantum effects are significant. Dr. Chatwani continued the work after his return to Germany and carried out a thorough study of terminal Mach numbers, both for real gases and the idealised models. This work was presented at the Tenth International Symposium on Rarefied Gas Dynamics in a paper entitled "Monte Carlo Simulation of Nozzle Beam Expansions" and is to be published in the proceedings.

2.3 The Onset of Mach Reflection

The direct simulation Monte Carlo method was used to study the transition from regular to Mach reflection of oblique two-dimensional steady shock waves. Particular attention was paid to the region of Mach number and deflection angle in which both types of reflection are theoretically possible. The results indicated that regular reflection can occur in the region where both types of reflection are possible. The study confirmed that the molecular approach can provide a practical alternative to finite difference and finite element methods for the computation of some continuum flows, especially when a dedicated mini-computer is used instead of a large machine in a general computing center. A paper was presented at the AIAA 9th Fluid and Plasma Dynamics Conference.

2.4 The Dissociation-Recombination Reaction in a Shock Tube

The direct simulation Monte Carlo method has been extended to cover molecular dissociation and recombination reactions. The model also includes internal degrees of freedom with separate relaxation times for rotation and vibration. The model is based on a classical collision theory analysis that provides an expression for the equilibrium state and allows the appropriate simulation parameters to be inferred from thermodynamic data for real gases. The model was first tested by the computation of the constant volume relaxation of a homogeneous gas sample. A finite difference procedure was then used to extend this program to cover the dissociation relaxation zone behind strong normal shock waves. A number of nitrogen cases were computed and the results were in agreement with existing experimental data. A paper has been submitted to the Journal of Computational Physics.

In addition to the work on the above topics, the work on the incompressible Kramers problem (reference 9 in the list of Section 1.2) was carried out during this period.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 18 AFOSR - TR - 76 - 1178	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NUMERICAL STUDIES IN GAS DYNAMICS	5. TYPE OF REPORT & PERIOD COVERED FINAL scientific rept 15 Apr 72 - 14 October 76	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) G A BIRD	8. CONTRACT OR GRANT NUMBER(s) 35 VAF - AFOSR - 2336 - 72	
9. PERFORMING ORGANIZATION NAME AND ADDRESS UNIVERSITY OF SYDNEY DEPARTMENT OF AERONAUTICAL ENGINEERING N S W 2006 AUSTRALIA	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 681307 9783/106 61102F	
11. CONTROLLING OFFICE NAME AND ADDRESS AIR FORCE OFFICE OF SCIENTIFIC RESEARCH/NA BLDG 410 BOLLING AIR FORCE BASE, D C 20332	12. REPORT DATE July 1974	13. NUMBER OF PAGES 5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) RAREFIED GAS FLOWS FORCES ON SMALL PARTICLES MONT CARLO REACTING GAS FLOWS SIMULATION NUMERICAL METHODS SHOCK WAVES		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A survey is made of the overall development of the direct simulation Monte Carlo method, over the period 1972 to 1976. This is a method for the computation of gas flows at the molecular level. The major advances have related to the extension of the method to cover some continuum flows, small disturbance flows and reacting gas flows. All these require very extensive computation and dedicated mini-computers have been shown to be one or two orders of magnitude more cost-effective than large general purpose machines in conventional computer centers.		