

BRL R 1952

ADA 035070

BRL

12⁵

REPORT NO. 1952

CONVEX INTERPOLATION OF CONVEX DATA

Walter O. Egerland

January 1977

Approved for public release; distribution unlimited.

COPY AVAILABLE TO DDC DOES NOT
PERMIT FULLY LEGIBLE PRODUCTION

DDC
REFINED
FEB 1 1977
D

USA BALLISTIC RESEARCH LABORATORIES
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating
or sponsoring activity is prohibited.

Additional copies of this report may be obtained
from the National Technical Information Service,
U.S. Department of Commerce, Springfield, Virginia
22151.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER BRL 12-10p. 1952	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CONVEX INTERPOLATION OF CONVEX DATA	5. TYPE OF REPORT & PERIOD COVERED BRL Report	
7. AUTHOR(s) Walter O. Egerland	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory Aberdeen Proving Ground, MD 21005	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Materiel Development & Readiness Command 5001 Eisenhower Avenue Alexandria, VA 22333	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE JANUARY 1977	
	13. NUMBER OF PAGES 11 (12 10p.)	
	15. SECURITY CLASS. (of this Report) UNCLASSIFIED	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Convex Interpolation Nonlinear Systems of Equations Antitone Operator Brouwer's Fixed Point Theorem		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (ams) This report contains the mathematical basis of an interpolation technique that constructs a smooth convex interpolant, called an H-Spline, for convex data on the real line. It is shown that an H-Spline always exists and is unique.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

1

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

050 750

TABLE OF CONTENTS

	Page
I. INTRODUCTION	5
II. DEFINITIONS.	5
III. EXISTENCE AND UNIQUENESS PROOF	6
REFERENCES	10
DISTRIBUTION LIST.	11

ACCESSION BY	
DTIC	White Section ☒
DTIC	Black Section ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

DDC
RECEIVED
 FEB 1 1977
 D

I. INTRODUCTION

In various interpolation problems of experimental and pointwise computer-constructed data the convexity and smoothness (continuous differentiability to the second order) of the interpolant is either implied by the nature of the underlying process or expressly desired. It is well known that classical (Lagrange, Hermite, etc.) and ordinary spline interpolation procedures do not in general produce convex interpolants for convex data and more often than not introduce spurious oscillations between data points. We introduce here a technique for interpolation and display of convex data without such drawbacks. Its basis is provided by an explicit solution of the convex two-point Hermite interpolation problem by a special rational function. To meet the smoothness requirement, a piecewise rational function, called an H-Spline, is constructed. The feasibility of the construction is equivalent to the existence and uniqueness of a solution of a nonlinear system of equations which may be brought into the fixed point form $x = Fx$. Here x is an n -dimensional vector whose dimension depends on the number of data points and whose components represent linear functions of the slopes to be assigned at the internal data points. Section II contains the basic definitions and preparatory remarks. A proof for the existence and uniqueness of an H-Spline for any finite number of data points, based on Brouwer's Fixed Point Theorem and the antitonicity of the operator F , is given in Section III. The general mathematical background for our considerations is covered in^{1,2,3}. A companion report⁴ describes in detail the actual construction of H-Splines on a computer.

II. DEFINITIONS

We begin with the following definitions:

Definition. A table $T: (a_i, y_i), i = 0, 1, \dots, n+1$, such that $a = a_0 < a_1 < \dots < a_n < a_{n+1} = b$ and $s_0 < s_1 < \dots < s_{n-1} < s_n$, where $s_i = (y_{i+1} - y_i)/(a_{i+1} - a_i), i = 0, 1, \dots, n$, is called a convex table.

Definition. Let R_2^1 be the class of rational functions of order 2 with at most one finite pole. Given a strictly monotone increasing sequence of real numbers $a_0, a_1, \dots, a_{n+1}$, an H-Spline with joints $a_i, i = 0, 1, \dots, n+1$, is a function $H(x)$ defined for $a = a_0 < x < a_{n+1} = b$ that satisfies the following three conditions:

- ¹ Collatz, L., Functional Analysis and Numerical Mathematics, (pp. 350-361), Academic Press, New York and London, 1966.
- ² Ortega, J.M. and Rheinboldt, W.C., Iterative Solution of Nonlinear Equations in Several Variables, (pp. 432-446), Academic Press, 1970.
- ³ Rall, L. B., Nonlinear Functional Analysis and Applications, (pp. 16-18), Academic Press, 1971.
- ⁴ Egerland, W. O. and Wisniewski, H. L., "Convex Interpolation with H-Splines," to appear.

a. In each interval (a_i, a_{i+1}) for $i = 0, 1, \dots, n$, $H(x)$ is given by some $h_i \in R_2^1$.

b. $H(x) \in C^2[a, b]$, i.e., $H(x)$ is twice continuously differentiable on $[a, b]$.

c. $H''(x) > 0$, $a < x < b$, i.e., $H(x)$ is strictly convex on $[a, b]$.

Since $h_i(x) = (A_{0i} + A_{1i}x + A_{2i}x^2)/(B_{0i} + B_{1i}x)$, $a_i < x < a_{i+1}$, $A_{2i} \neq 0$, $H(x)$, as a plane curve, is represented by an arc of an hyperbola (or a parabola in case $B_{1i} = 0$) between adjacent joints. Hence the name H-Spline was suggested.

Definition. Let ℓ and r be positive numbers. A function $f(x)$ solves the "convex two-point Hermite interpolation problem" on the interval $[a, b]$ if (1) $f(a) = 0$, $f(b) = 0$, (2) $f'(a) = -\ell$, $f'(b) = r$, and (3) $f''(x) > 0$, $a < x < b$.

According to this definition, the function

$$B(x) = B(x; a, b, \ell, r) = \frac{\ell r (x-a)(x-b)}{\ell(x-a) + r(b-x)}$$

solves the convex two-point Hermite interpolation problem for arbitrary a, b , $a < b$, and positive ℓ and r . $B(x)$ is the simplest H-Spline. We note that

$$B''(x) = \frac{2\ell^2 r^2 (b-a)^2}{[\ell(x-a) + r(b-x)]^3} \quad (1)$$

III. EXISTENCE AND UNIQUENESS PROOF

With the definitions given in Section II, the following theorem holds:

Theorem. Given a convex table $T: (a_i, y_i)$, $i = 0, 1, \dots, n+1$, and endconditions y'_0, y'_{n+1} , $y'_0 < s_0$, $s_n < y'_{n+1}$, there exists a unique H-Spline such that $H(a_i) = y_i$, $i = 0, 1, \dots, n+1$, $H'(a_0) = y'_0$, and $H'(a_{n+1}) = y'_{n+1}$.

Proof. Let y'_i , $i = 1, 2, \dots, n$, be a sequence of slopes such that $y'_0 < s_0 < y'_1 < s_1 < \dots < s_{n-1} < y'_n < s_n < y'_{n+1}$ and consider the function

$$\tilde{H}(x) = L_i(x) + B_i(x), \quad a_i \leq x \leq a_{i+1}, \quad i = 0, 1, \dots, n, \quad (2)$$

where $L_i(x) = y_i + s_i(x - a_i)$ and $B_i(x) = B(x; a_i, a_{i+1}, s_i - y'_i, y'_{i+1} - s_i)$. $\tilde{H}(x)$ is strictly convex on $[a, b]$, $\tilde{H}(a_i) = y_i$, $\tilde{H}'(a_i) = y'_i$, and $\tilde{H}(x) \in C^1[a, b]$. $\tilde{H}(x)$ is an H-Spline if and only if the continuity requirements

$$B''_{i-1}(a_i) = B''_i(a_i) > 0, \quad i = 1, 2, \dots, n, \quad (3)$$

are satisfied. Using (1) and setting

$$y'_i = x_i s_{i-1} + (1 - x_i) s_i, \quad i = 1, 2, \dots, n,$$

$$K_1 = \left(\frac{a_1 - a_0}{a_2 - a_1} \right)^{1/2} \left(\frac{s_0 - y'_0}{s_2 - s_1} \right)^{1/2}$$

$$K_i = \left(\frac{a_i - a_{i-1}}{a_{i+1} - a_i} \right)^{1/2} \left(\frac{s_{i-1} - s_{i-2}}{s_{i+1} - s_i} \right)^{1/2}, \quad i = 2, \dots, n-1,$$

$$K_n = \left(\frac{a_n - a_{n-1}}{a_{n+1} - a_n} \right)^{1/2} \left(\frac{s_{n-1} - s_{n-2}}{y'_{n+1} - s_n} \right)^{1/2},$$

(3) is equivalent to the existence of a solution of the system

$$x_1 = \frac{(1 - x_2)^{1/2}}{(1 - x_2)^{1/2} + K_1} = f_1(x_2)$$

$$x_i = \frac{(1 - x_{i+1})^{1/2}}{(1 - x_{i+1})^{1/2} + K_i x_{i-1}^{1/2}} = f_i(x_{i-1}, x_{i+1}), \quad i = 2, \dots, n-1$$

$$x_n = \frac{1}{1 + K_n x_{n-1}^{1/2}} = f_n(x_{n-1}) \quad (4)$$

in the open cube $I_n: 0 < x_i < 1, i = 1, 2, \dots, n$. This, in turn, is equivalent to the existence of a fixed point in I_n of the mapping $F: I \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ where for $x^T = (x_1, x_2, \dots, x_n)$ Fx is represented by the column vector

$$Fx = \begin{pmatrix} f_1(x) \\ \vdots \\ f_i(x) \\ \vdots \\ f_n(x) \end{pmatrix}.$$

The mapping F has the following properties:

- (P₁) F is continuous on I_n .
- (P₂) $FI_n \subset I_n$.
- (P₃) F is antitone on I_n .
- (P₄) If $v, w \in I_n$ and $v \leq w$, then $v = Fw$ and $w = Fv$ imply $v = w$.

(P₁) and (P₂) are obvious, and (P₃) follows from the mean value theorem and $\partial f_i(x)/\partial x_j \leq 0$, $i, j = 1, \dots, n$, $x \in I_n$. To prove (P₄), we observe that the constants K_i , $i = 1, \dots, n$, can be expressed by the components of v and w as follows:

$$K_1 = \frac{1 - v_1}{v_1} (1 - w_2)^{1/2} = \frac{1 - w_1}{w_1} (1 - v_2)^{1/2}$$

$$K_i = \frac{1 - v_i}{v_i} \left(\frac{1 - w_{i+1}}{w_{i-1}} \right)^{1/2} = \frac{1 - w_i}{w_i} \left(\frac{1 - v_{i+1}}{v_{i-1}} \right)^{1/2}, \quad (6)$$

$$i = 2, \dots, n-1,$$

$$K_n = \frac{1 - v_n}{v_n} w_{n-1}^{-1/2} = \frac{1 - w_n}{w_n} v_{n-1}^{-1/2}$$

The equality of the product of the left sides with that of the right sides in (6) yields, after cancellation,

$$\left(\frac{1 - v_1}{1 - w_1} \right)^2 \prod_{i=2}^n \frac{1 - v_i}{1 - w_i} = \left(\frac{v_n}{w_n} \right)^2 \prod_{i=1}^{n-1} \frac{v_i}{w_i}, \quad (7)$$

a contradiction unless $v_i = w_i$, $i = 1, \dots, n$, i.e., $v = w$.

We show next that F has exactly one fixed point in I_n . First, let $\delta_0 > 0$ be such that $\delta_0 < \delta_1 = \min (m^2(1+m)^{-2}; (1+M)^{-2}; 1/2)$, where $m = \min (K_1, \dots, K_n)$ and $M = \max (K_1, \dots, K_n)$. If $x \in I_n(\delta_0)$: $\delta_0 < x_i < 1 - \delta_0$, $i = 1, \dots, n$, then it is easy to verify that the components of F satisfy the inequalities

$$\delta_0 < f_i(x) < 1 - \delta_0, \quad i = 1, \dots, n. \quad (8)$$

$I_n(\delta_0)$ is compact and convex, F is continuous on $I_n(\delta_0)$, and, by (8), $F I_n(\delta_0) \subset I_n(\delta_0)$. Therefore, we may apply Brouwer's fixed point theorem to conclude that F has at least one fixed point x^* in $I_n(\delta_0)$. Furthermore, by the antitonicity of F , the iterations

$$\begin{aligned} v^{k+1} &= F w^k \\ w^{k+1} &= F v^k \end{aligned} \quad k = 0, 1, \dots \quad (9)$$

with initial points $(v^0)^T = (\delta_0, \dots, \delta_0)$, $(w^0)^T = (1 - \delta_0, \dots, 1 - \delta_0)$ define iterates such that $([1], [3], [4])$

$$v^0 < v^1 < \dots < v^{k+1} < w^{k+1} < \dots < w^1 < w^0. \quad (10)$$

The limits $\lim_{k \rightarrow \infty} v^{k+1} = v$ and $\lim_{k \rightarrow \infty} w^{k+1} = w$ exist, $v \leq w$, x^* is contained in the order interval $\langle v, w \rangle$, and, by the continuity of F , we have $v = F w$ and $w = F v$ from (9). Hence, in view of (5) - (P_4) , $v = w = x^*$ is the only fixed point of F in $I_n(\delta_0)$. Since the argument can be repeated with an arbitrary positive $\delta < \delta_0$ and $I_n(\delta_0) \subset I_n(\delta)$, it follows that F has no other fixed point in I_n besides x^* . The unique H-Spline correspondence to $(x^*)^T = (x_1^*, x_2^*, \dots, x_n^*)$ is given by $H(x) = \tilde{H}(x)$ in (2) with $y_i^1 = x_i^* s_{i-1} + (1 - x_i^*) s_i$, $i = 1, \dots, n$. This completes the proof of the theorem.

NOTE:

An outline for the proof of the existence and uniqueness of an H-Spline for a given convex table was presented at the Twentieth Conference of Army Mathematicians at the U.S. Army Natick Laboratories, Natick, Massachusetts, May 1974.

REFERENCES

1. Collatz, L., Functional Analysis and Numerical Mathematics, (pp. 350-361), Academic Press, New York and London, 1966.
2. Ortega, J. M. and Rheinboldt, W. C., Iterative Solution of Nonlinear Equations in Several Variables, (pp. 432-446), Academic Press, 1970.
3. Rall, L. B., Nonlinear Functional Analysis and Applications, (pp. 16-18), Academic Press, 1971.
4. Egerland, W. O. and Wisniewski, H. L., "Convex Interpolation with H-Splines," to appear.

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: DDC-TCA Cameron Station Alexandria, VA 22314	2	Commander US Army Mobility Equipment Research & Development Command ATTN: Tech Docu Cen, Bldg. 315 DRSME-RZT Fort Belvoir, VA 22060
1	Commander US Army Materiel Development and Readiness Command ATTN: DRCDMA-ST 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander US Army Armament Command Rock Island, IL 61202
1	Commander US Army Aviation Systems Command ATTN: DRSAB-E 12th and Spruce Streets St. Louis, MO 63166	1	Commander US Army Frankford Arsenal ATTN: SARFA-FCV, Stan Goodman Philadelphia, PA 19137
1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	Commander US Army Harry Diamond Labs ATTN: DRXDO-TI 2800 Powder Mill Road Adelphi, MD 20783
1	Commander US Army Electronics Command ATTN: DRSEL-RD Fort Monmouth, NJ 07703	1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SA White Sands Missile Range NM 88002
1	Commander US Army Missile Command ATTN: DRSMI-R Redstone Arsenal, AL 35809	1	Director The Johns Hopkins University Applied Physics Lab Johns Hopkins Road Laurel, MD 20810
1	Commander US Army Tank Automotive Development Command ATTN: DRDTA-RWL Warren, MI 48090	1	University of Wisconsin Math Research Center Madison, WI 53706
			<u>Aberdeen Proving Ground</u> Marine Corps Ln Ofc Dir, USAMSAA