

AD-A035 264

OKLAHOMA UNIV HEALTH SCIENCES CENTER OKLAHOMA CITY
LEUKOCYTE RESPONSE AND HYPOGLYCEMIA: CO-DETERMINANTS IN THE ALT--ETC(U)
DEC 76 G L WHITE, L T ARCHER, B K BELLER

F/G 6/5

N00014-76-C-0229

UNCLASSIFIED

TR-114

NL

1 OF 1
AD-A
035 264



END
DATE
FILMED
3-11-77
NTIS

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

AD-A035 264

LEUKOCYTE RESPONSE AND HYPOGLYCEMIA
CO-DETERMINANTS IN THE ALTERED SURVIVAL
TO SUPERLETHAL ENDOTOXIC SHOCK

OKLAHOMA UNIVERSITY HEALTH SCIENCES
CENTER, OKLAHOMA CITY

13 DECEMBER 1976

ADA 035264

OFFICE OF NAVAL RESEARCH

Contract N00014-76-C-0229

Project NR 105-516

TECHNICAL REPORT NO. 114

LEUKOCYTE RESPONSE AND HYPOGLYCEMIA:
CO-DETERMINANTS IN THE ALTERED SURVIVAL TO SUPERLETHAL ENDOTOXIC SHOCK

Gary L. White, Linda T. Archer, Beverly K. Beller,
Donald D. Holmes, and Lerner B. Hinshaw

Prepared for Publication

in

Circulatory Shock

University of Oklahoma Health Sciences Center
Departments of Pathology and Physiology & Biophysics
Oklahoma City, Oklahoma

13 December 1976

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Distribution of this report is unlimited

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161



OFFICE OF NAVAL RESEARCH

Contract N00014-76-C-0229

Project NR 105-516

TECHNICAL REPORT NO. 114

LEUKOCYTE RESPONSE AND HYPOGLYCEMIA:
CO-DETERMINANTS IN THE ALTERED SURVIVAL TO SUPERLETHAL ENDOTOXIC SHOCK

Gary L. White, Linda T. Archer, Beverly K. Beller,
Donald D. Holmes, and Lerner B. Hinshaw

Prepared for Publication

in

Circulatory Shock

University of Oklahoma Health Sciences Center
Departments of Pathology and Physiology & Biophysics
Oklahoma City, Oklahoma

13 December 1976

Reproduction in whole or in part is permitted for
any purpose of the United States Government

Distribution of this report is unlimited

io

ADVISORY	
NTS	With Section <input checked="" type="checkbox"/>
DSG	Diff Section <input type="checkbox"/>
MANUSCRIPT	<input type="checkbox"/>
IDENTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY GROUP	
Dist.	AVAIL. INFO. SPEC.
A	

ABSTRACT

A prime goal of this laboratory has been to develop animal shock models more closely approximating septic shock in man. Anesthesia in dogs has been found to depress leukocyte concentration, body temperature, and central nervous system activities. This laboratory has documented a progressively developing hypoglycemia associated with systemic hypotension, hepatosplanchnic pathology and death in endotoxin-shocked dogs. Recent data documented accelerated uptake of glucose in blood following endotoxin, with certain components of the buffy coat responsible for the increased uptake. The present study explored the use of the awake dog as an improved shock model and assayed a possible protective role of leukocytes against the lethal effects of endotoxin. Experiments were conducted on unanesthetized healthy dogs with initial WBC counts between 7,000 and 20,000/mm³. Dogs were divided into paired groups: saline controls (Group I) and endotoxin experimentals (Group II). Group II animals (N=5) were injected intravenously with sublethal doses of E. coli endotoxin on Days 1 and 2, LD₁₀₀ on Day 3, and 2 X LD₁₀₀ on Day 4. The control group (N=5) received equal volumes of saline on Days 1, 2 and 3, but on Day 4 received an identical superlethal dose of endotoxin (2 X LD₁₀₀). Data document that the awake dog becomes febrile and exhibits initial leukopenia with subsequent marked leukocytosis in response to endotoxin. Lethal hypoglycemia is not seen in animals demonstrating leukocytosis on the day of superlethal endotoxin challenge, while animals with normal leukocyte counts die with low glucose concentrations (mean, 40 mg%). Results suggest that leukocytosis and sustained gluconeogenic function are important co-determinants of survivability to endotoxin shock.

KEY WORDS: endotoxin, leukocyte, febrile response, awake dog, glucose concentration, leukocytosis, phagocytosis, gluconeogenesis, shock model

The development of animal models for the study of endotoxin shock more closely approximating the state of septic shock in man has long been the goal of clinically oriented research laboratories. A persistent problem has been the adverse effects of surgical levels of anesthesia which question the validity of the majority of animal models for comparative clinical application. The use of anesthesia in dogs has been found to depress the white blood cell concentration (1), body temperature (2), and central nervous system activities (3).

Progressively developing hypoglycemia associated with systemic hypotension, hepatosplanchnic pathology and death has been observed in the dog subjected to endotoxin shock (4) and the cause has been proposed to be a depressant effect of endotoxin on liver function, specifically gluconeogenesis (5-10) combined with increased glucose need (11,12). Recent studies have demonstrated accelerated uptake of glucose in the blood following endotoxin, with certain components of the buffy coat responsible for the increased uptake (11,13). Endotoxin is efficiently phagocytized by polymorphonuclear leukocytes (14,15), and an associated increase in glucose utilization by neutrophils has been reported to occur (16,17).

The purpose of the present study was to explore the use of the awake dog as an improved shock model and to assay a possible protective role of leukocytes against the lethal effects of endotoxin. The unanesthetized dog model was utilized since it was considered to more closely parallel the pathophysiological alterations observed in human septic shock. Results reveal that the awake dog becomes febrile and exhibits initial leukopenia with subsequent leukocytosis in response to sublethal, lethal and superlethal doses of endotoxin. Lethal hypoglycemia is not seen in animals demonstrating leukocytosis at time of endotoxin challenge. Data are consistent with the view that leukocytosis and sustained liver gluconeogenic function are important co-determinants of

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) UNIVERSITY OF OKLAHOMA HEALTH SCIENCES CENTER OKLAHOMA CITY, OKLAHOMA		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP UNCLASSIFIED	
3. REPORT TITLE LEUKOCYTE RESPONSE AND HYPOGLYCEMIA: CO-DETERMINANTS IN THE ALTERED SURVIVAL TO SUPERLETHAL ENDOTOXIC SHOCK			
4. DESCRIPTIVE NOTES (Type of report and, inclusive dates) Technical Report			
5. AUTHOR(S) (First name, middle initial, last name) Gary L. White, Linda T. Archer, Beverly K. Beller, Donald D. Holmes, and Lerner B. Hinshaw			
6. REPORT DATE 13 December 1976		7a. TOTAL NO. OF PAGES 26	7b. NO. OF REFS 41
8a. CONTRACT OR GRANT NO. N00014-76-C-0229		9a. ORIGINATOR'S REPORT NUMBER(S) 114	
b. PROJECT NO. NR 105-516			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT Distribution of this report is unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Office of Naval Research	
13. ABSTRACT A prime goal of this laboratory has been to develop animal shock models more closely approximating septic shock in man. Anesthesia in dogs has been found to depress leukocyte concentration, body temperature, and central nervous system activities. This laboratory has documented a progressively developing hypoglycemia associated with systemic hypotension, hepatosplanchnic pathology and death in endotoxin-shocked dogs. Recent data documented accelerated uptake of glucose in blood following endotoxin, with certain components of the buffy coat responsible for the increased uptake. The present study explored the use of the awake dog as an improved shock model and assayed a possible protective role of leukocytes against the lethal effects of endotoxin. Experiments were conducted on unanesthetized healthy dogs with initial WBC counts between 7,000 and 20,000/mm ³ . Dogs were divided into paired groups: saline controls (Group I) and endotoxin experimentals (Group II). Group II animals (N=5) were injected intravenously with sublethal doses of <i>E. coli</i> endotoxin on Days 1 and 2, LD ₁₀₀ on Day 3, and 2XLD ₁₀₀ on Day 4. The control group (N=5) received equal volumes of saline on Days 1, 2 and 3, but on Day 4, received an identical superlethal dose of endotoxin (2XLD ₁₀₀). Data document that the awake dog becomes febrile and exhibits initial leukopenia with subsequent marked leukocytosis in response to endotoxin. Lethal hypoglycemia is not seen in animals demonstrating leukocytosis on the day of superlethal endotoxin challenge, while animals with normal leukocyte counts die with low glucose concentrations (mean, 40 mg%). Results suggest that leukocytosis and sustained gluconeogenic function are important co-determinants of survivability to endotoxin shock.			

survivability to endotoxin. The unanesthetized animal administered endotoxin was found to be a useful model for septic shock in man.

METHODS

Experiments were designed to follow the changes of the peripheral white blood cell (WBC) counts, rectal temperatures and blood glucose values during sublethal, lethal, and superlethal doses of E. coli endotoxin (Difco, Detroit) without the influence of anesthetics. Ten awake mongrel adult dogs of random sex, selected for freedom of clinical signs of disease, were used in the present study. Each dog was screened for microfilaria of heartworms and eliminated if positive and was treated for intestinal parasites. Animals were allowed a stabilization period of 3 to 6 weeks prior to use in the study. Only dogs with initial WBC counts between 7,000 and 20,000/mm³ and hematocrits exceeding 37% were utilized.

Dogs were divided into paired saline control and endotoxin experimental groups (Groups I and II, respectively). The endotoxin group received sublethal doses of endotoxin of 0.003 mg/kg body weight on days 1 and 2 (i.e., 1/1,000 LD₁₀₀), 3 mg/kg on day 3 (i.e., LD₁₀₀) and a challenge dose of endotoxin of 6 mg/kg on day 4 (i.e., 2 X LD₁₀₀). The control group received equal volumes of saline on days 1, 2 and 3, but on day 4 received a 6 mg/kg challenge dose of endotoxin (i.e., 2 X LD₁₀₀). The LD₁₀₀ of E. coli endotoxin (3 mg/kg) had been previously established in a group of approximately 25 dogs. White blood cell counts, differential WBC, rectal temperatures and blood glucoses were measured initially (i.e., before endotoxin or saline injection) and then each 60 to 90 minutes for 8 1/2 hours during each of the 4 days of observation. Animals surviving injection of endotoxin were sacrificed following a 30-day observation period.

The WBC counts were measured with an automatic particle counter (Coulter Z_P; Hialeah, Florida) and the differential WBC by microscopic

examination of blood smears stained with Wrights stain (100 cells counted). Blood glucose concentrations were determined using a Beckman glucose analyzer (Beckman Instruments; Fullerton, Calif.) with an accuracy of ± 3 mg%, and rectal temperatures were obtained using a Tele-Thermometer probe (Yellow Springs Instruments; Yellow Springs, Ohio). The blood samples were obtained by venipunctures of either the cephalic or saphenous veins, then placed in vacutainers containing ethylenediamine-tetraacetic acid (EDTA; Becton, Dickinson, and Co.), and immediately placed on ice. The injection of saline or endotoxin was by the intravenous route utilizing either the cephalic or saphenous vein. The results were analyzed using the t test for paired or unpaired data.

RESULTS

Leukocyte, temperature and blood glucose alterations for days 1, 2 and 3 for both endotoxin experimental and saline control groups

Results from Figure 1 reveal significant increases ($p < 0.05$) in rectal temperature at all time intervals between 60 and 510 minutes post-endotoxin injection for days 1 and 3 and from 60 through 240 minutes for day 2 when contrasted with the control group receiving saline instead of endotoxin. The initial mean WBC counts for the endotoxin group and saline control group were $12,840/\text{mm}^3$ and $12,540/\text{mm}^3$, respectively. Figure 2 demonstrates significant elevations of WBC count after endotoxin administration on days 1, 2 and 3 at 420 and 510 minutes ($p < 0.05$) when compared with the saline control group.

Alterations in blood glucose concentration are arrayed in Figure 3. There were no significant changes in the blood glucose on days 1 and 2, but

on day 3 lower blood glucose values were observed at 240 through 420 minutes ($p < 0.05$).

Leukocyte, body temperature and blood glucose responses to superlethal challenge of endotoxin

Figure 4 presents a comparison of changes in WBC count, rectal temperature and blood glucose that occurred on day 4, when both the experimental endotoxin and the saline control groups were administered 6 mg/kg of endotoxin ($2 \times LD_{100}$). The endotoxin experimental group had an initial WBC count of $39,000/\text{mm}^3$ at zero time on day 4, which is significantly elevated ($p < 0.001$) when compared with its simultaneous time for the saline control group. Initial blood glucose values for the endotoxin group were not significantly different from either zero time on day 1 for the same group or zero time on day 4 for the saline group. WBC count was lowest ($p < 0.01$) for both groups at approximately $5,000 \text{ WBC}/\text{mm}^3$ at 60 minutes post-endotoxin. From 60 through 510 minutes post-endotoxin, the WBC count progressively rose to near control values for the endotoxin group and to above control values for the saline group. On day 4 after injection of endotoxin, a significant difference in WBC count between the two groups occurred at 120 minutes ($p < 0.05$). Significant febrile responses are seen with both groups from 60 through 180 minutes when compared with zero time ($p < 0.05$). Comparing the two groups on day 4, a markedly elevated ($p < 0.005$) rectal temperature was observed at 330 minutes in the saline group. The mean glucose concentration significantly fell ($p < 0.01$) from 95 mg% at zero time on day 4 to 67 mg% at 120 minutes for the endotoxin experimental group. At 60 minutes after endotoxin injection, for the saline control group mean glucose concentration rose from 91 mg% to 108 mg%, and then progressively declined below the zero time control values until death. When comparing the two groups on day 4, the 60 and 120 minute measurements show that mean blood glucose for the experi-

mental group was lower ($p < 0.02$) than the saline control group. Blood glucose concentration in the endotoxin experimental group steadily rose from 120 to 330 minutes, where it stabilized near control values.

Changes in differential leukocyte counts in endotoxin experimental and saline control groups

The responses of the total WBC, lymphocytes and mature and immature neutrophils are shown in Tables 1 and 2. The progressive leukocytosis ($p < 0.005$) seen in the endotoxin group (Group II) on days 2 through 4 can be attributed to the increased numbers of mature neutrophils ($p < 0.05$). In these animals, absolute leukocyte and lymphocyte counts remained constant days 1 through 4, while mature neutrophils increased ($p < 0.05$) on day 2 and immature neutrophils decreased ($p < 0.025$) on days 2 and 3. On day 4 in Groups I and II, the marked leukopenia ($p < 0.02$) observed at 60 minutes post-endotoxin appeared to be a function of the depressed ($p < 0.02$) mature neutrophil counts while depression of the immature neutrophils also occurred in the saline control group.

Mortality rates following superlethal challenge of *E. coli* endotoxin

Both groups received a 6 mg/kg ($2 \times LD_{100}$) *E. coli* endotoxin after zero time values on day 4. In the endotoxin experimental group (Group II), 100% of the animals ($N=5$) survived the $2 \times LD_{100}$ endotoxin challenge (30-day survival). In the saline control group all dogs ($N=5$) died within an average of 6 hours when challenged with $2 \times LD_{100}$ endotoxin (see Table 3).

DISCUSSION

The unanesthetized animal model

The purpose of the present study was to explore the use of the unanesthetized dog as a model for septic shock and to determine the leukocyte and febrile response and changes in blood glucose concentrations. Results from the present experiments reveal a rapid leukopenia followed by a leukocytosis

in response to sublethal and lethal, as well as superlethal injections, of endotoxin. Results showed that animals receiving daily sublethal injections of endotoxin survive a superlethal challenge of endotoxin and are permanent survivors. Animals not receiving sublethal injections are very susceptible to the adverse effects of endotoxin, including the full pathophysiological impact of shock, and die in an average time of 6 hours. The unanesthetized animal model described in this study appears to be well suited for application and comparison of findings with septic shock in man.

Response of the white blood cell to endotoxin

Mulholland and Cluff reported that endotoxin causes granulocytes to adhere to the capillary endothelial cells and then later leave the circulation and move into the tissue (18). These findings corroborate the data in the present study and would explain the early leukopenia seen post-endotoxin. Leukocytosis associated with endotoxin is reported to occur via entry of new leukocytes from the bone marrow into the circulation (19). The circulating neutrophils have been demonstrated to be the key cellular components in the clearance of bacterial organisms from the blood of dogs (20). Blood from endotoxin-treated rabbits has been shown to possess a significant phagocytic capacity (18) and leukocytes were found to detoxify endotoxin in vitro in rabbit serum (21). Glucose has been demonstrated to be the substrate essential for phagocytosis by polymorphonuclear leukocytes (22).

Animals in the experimental group exhibited a marked leukocytosis ($39,000/\text{mm}^3$) after endotoxin by day 4, and the increased white cell count can be mainly accounted for by increased numbers of neutrophils. In the present study all of the animals in the endotoxin-pretreated group survived the massive endotoxin challenge while every animal in the control group succumbed to the identical challenge dose. Since Balis et al. (14) and Cline et al. (15) have documented that neutrophils phagocytize endotoxin, the increased numbers of neutrophils

seen in the present study may have effected an efficient and rapid clearance of endotoxin leading to the 100% survivability results. Efficient removal of endotoxin was further suggested in the present experimental group since the animals exhibited minimal clinical signs of endotoxin administration and were eating and drinking normally within 9 hours post-endotoxin.

Blood glucose changes after endotoxin: Role of hypoglycemia

Recent studies in dogs administered endotoxin have documented the development of hypoglycemia concomitant with the systemic hypotension, hepatosplanchnic pathology and death (4). Hypoglycemia has been attributed to depressed glucose production due to defective hepatic function (5-10), while accelerated uptake of glucose by the blood has recently been reported (13). Early hypoglycemia seen in the endotoxin-pretreated animals of the present study in response to the $2 \times LD_{100}$ of endotoxin may suggest an early increased glucose uptake in response to leukocyte phagocytosis.

Hypoglycemia has been reported to play a significant role in the pathophysiology of endotoxin shock (8,11-13). Recent studies from this laboratory have documented a positive correlation between levels of blood glucose and survival in endotoxin shock (12,13). Further, the early hyperglycemia seen in various shock states has been attributed to sympathetic-stimulated glycogenolysis as well as alpha-adrenergic suppression of beta cell insulin release in the pancreas (8,23). Clearly, the "unprotected" saline group responded with a significant hyperglycemia during the first hour after superlethal endotoxin challenge, suggesting a greater stress to this group. At 5 1/2 hours post-endotoxin, the significantly decreased blood glucose level of the saline animals suggests an impairment of liver gluconeogenesis capacity, as has been reported by Groves *et al.* and Filkins *et al.* in endotoxin shock (7,8).

The recovery from hypoglycemia observed in the endotoxin experimental group indicates that the ability of the liver to perform gluconeogenesis in response

to superlethal endotoxin challenge has been protected. Normoglycemia was observed in the "protected" animals in contrast to hypoglycemia in the saline control dogs at 5 1/2 hours post-endotoxin. The lethal hypoglycemia previously observed in the laboratory in canine endotoxin shock was progressive but animals reached lethal levels of 12 mg% at 5 hours post-endotoxin. The sharp contrast observed in blood glucose concentration between the two groups and the 100% survival versus 100% mortality seen in the two groups support a liver-sparing role of the circulating neutrophil.

Temperature response to endotoxin

Data from the present experiments in unanesthetized dogs show significant increases in rectal temperature with a mean range from +1.4 to 3.2°C above control values. The rise in temperature in the endotoxin "protected" group corresponds with the time when blood glucose concentrations in these animals were decreasing, suggesting that fever may add an additional drain to the available glucose. Studies have shown that the neutrophil can release a pyrogen which will produce a febrile response in rabbits when administered endotoxin (24). Other reports found evidence that endotoxin produces fever by direct action upon the brain (25). It is known that anesthetized dogs in endotoxin shock become progressively hypothermic; while major trauma patients as well as septic animals and man spike fevers in the disease course (26,27). The awake dog thus appears to be a more valid model for application to the clinical septic state, since marked febrile responses were seen following both sub- and superlethal injections of E. coli endotoxin. Furthermore, leukopenia, followed by leukocytosis concomitant with the febrile responses, appears to more closely approximate changes seen in the septic patient.

Role of the RES and white blood cell in endotoxin shock

(a) Tolerance. Review of the literature describes the elicitation of a tolerant state following sublethal injections of endotoxins (28-30). Greisman

and Hornick have suggested possible mechanisms by which man develops tolerance to bacterial endotoxins and have separated the tolerant condition into two phases (30). Early tolerance appears to be mediated by a non-antibody mechanism entailing a transiently occurring refractory state of mainly the hepatic macrophages (Kupffer cells), while late tolerance appears to be mediated by anti-endotoxin antibodies directed against both O and common core antigens. At present there is no clear consensus of opinion on the role of the reticulo-endothelial system (RES) and the peripheral cellular components on increased tolerance and resistance to endotoxin. Beeson (31) found that an increased resistance to endotoxin occurs because of increased functional capacity of the RES to remove bacterial toxins unassociated with antibody formation. Tolerance to endotoxin involves acceleration of the degrading and detoxifying systems which exist in circulating blood as well as other factors (28,30,32,33) and it is not transferred passively with serum from resistant animals (32). It was not the purpose of this laboratory to re-evaluate the mechanisms involved in tolerance but rather to try to determine the importance of leukocytes in the total host-defense mechanism. Leukocytes have been reported to exert a significant role in phagocytosis, although serum and opsonins may also perform key roles (21,34). It is known that active phagocytosis requires increased substrate in the form of glucose (16,17), but it may be the most effective defense against the hepatic dysfunction observed in shock. A study utilizing radioactively-labelled endotoxin traced its passage from the plasma to the buffy coat and finally to the liver during the early clearance period associated with leukopenia, fever and diarrhea (35). If phagocytosis in the buffy coat were sufficiently effective, it seems that the liver could be spared the added stress of endotoxin detoxification (6) and the possible depression of gluconeogenic function from the action of endotoxin (7-9). Fukuda and Akiyama (36) propose that

it is not an increased RES activity but a greater stability of carbohydrate metabolism that provides the tolerance to endotoxin. Again in the present study, the return of blood glucose concentration to normal in the "protected" group reveals a possible association of leukocytosis with preserved liver gluconeogenic function, thus melding increased RES (circulating and sessile) function and normal carbohydrate metabolism.

(b) Therapy. This study suggested the protective influences of leukocytosis stimulated by sublethal injections of endotoxin. Since survivability seemed to be associated with leukocytosis, there are possible therapeutic implications regarding the role of the white blood cell in endotoxin shock. Recent reports have described beneficial effects of transfused white blood cells as a treatment for septicemia in neutropenic patients (37) and dogs (38,39). Granulocyte transfusions markedly reduced the number of circulating Pseudomonas aeruginosa bacteria in dogs, as contrasted to non-transfused controls (39). Leukopenic dogs in E. coli septicemia when given leukocyte transfusion had prolonged periods of sterile blood cultures and extended survival times compared with leukopenic septic controls (38). Graw et al. (37) reported a significant increase in survival rate in patients with gram-negative sepsis when transfused with granulocytes compared with septic patients. Administration of hypertonic glucose has resulted in an increased clearance of E. coli from the blood of dogs (40), suggesting that supplying substantial metabolic support for accelerated leukocyte activity may enhance survival in septic shock. By using the awake dog as an improved shock model and by studying the febrile and leukocyte response to varying doses of endotoxin, we hoped to observe a combination of clinical signs that would be a clue as to the appropriate time for initiating treatment via activation of natural host-defense mechanisms or supplying the host's first natural line of defense (41) against infection-- leukocytes via transfusion.

REFERENCES

1. Usenik, E. A., and Cronkite, E. P. 1965. Effect of barbituate anesthetic on leukocyte in normal and splenectomized dogs. *Anesth. Analg.* 44:167.
2. Lumb, W. V., and Jones, E. W. 1973. *Veterinary anesthesia*, p. 602. Lea and Febiger, Philadelphia.
3. Sharpless, S. K. 1975. Hypnotics and sedatives. I. The barbituates. Page 106 in L. S. Goodman and A. Gilman, eds. *The pharmacological basis of therapeutics*. McMillan Publishing Co., Inc., New York.
4. Hinshaw, L. B., Peyton, M. D., Archer, L. T., Black, M. R., Coalson, J. J., and Greenfield, L. J. 1974. Prevention of death in endotoxin shock by glucose administration. *Surg. Gynec. Obstet.* 139:851.
5. Brunson, J. G., Schulz, D. M., Angevine, D. M., Imrie, S. T., and Douglas, B. H. 1966. Evaluation of therapeutic agents. Morphologic changes and survival data in dogs subjected to endotoxin during the shock tour. *J. Okla. State Med. Assoc.* 59:479.
6. DiLuzio, N. R., Olcay, I., Holper, K., Drapanas, T., and Trejo, R. 1975. The relationship of reticuloendothelial dysfunction and endotoxemia to survival after hepatic ischemia injury in the baboon. *Circ. Shock* 2:77.
7. Filkins, J. P., and Cornell, R. P. 1974. Depression of hepatic gluconeogenesis and the hypoglycemia of endotoxin shock. *Am. J. Physiol.* 227:778.
8. Groves, A. C., Woolf, L. I., O'Regan, P. J., Beach, C., Hasinoff, C., and Sutherland, W. H. 1974. Impaired gluconeogenesis in dogs with E. coli bacteremia. *Surgery* 76:533.

9. Harken, A. H., Lillo, R. S., and Hufnagel, H. V. 1975. Direct influence of endotoxin on cellular respiration. *Surg. Gynec. Obstet.* 140:858.
10. McFadzean, A. J. S., and Yeung, R. T. T. 1965. Hypoglycaemia in suppurative pancholangiitis due to Clonorchis sinensis. *Trans. Royal Soc. Trop. Med. Hygiene* 59:179.
11. Archer, L. T., Benjamin, B., Lane, M. M., and Hinshaw, L. B. 1976. Renal gluconeogenesis and increased glucose utilization in shock. *Am. J. Physiol.* 231:872.
12. Peyton, M. D., Hinshaw, L. B., Greenfield, L. J., and Elkins, R. C. 1975. Hypoglycemic effects of endotoxin in eviscerated dogs. *Surg. Gynec. Obstet.* 141:727.
13. Hinshaw, L. B., Beller, B. K., Archer, L. T., and Benjamin, B. 1976. Hypoglycemic response of blood to live Escherichia coli organisms and endotoxin. *J. Surg. Res.* 21:141.
14. Balis, J. U., Rapport, E. S., Gerber, L., and Buddingh, F. 1974. Continuous endotoxemia in rhesus monkeys as a clinically relevant model for shock lung. *Am. J. Path.* 74:90a.
15. Cline, M. J., Melmon, K. L., Davis, W. C., and Williams, H. E. 1968. Mechanism of endotoxin interaction with human leucocytes. *Brit. J. Haematol.* 15:539.
16. Sbarra, A. J., and Karnovsky, M. L. 1959. The biochemical basis of phagocytosis. I. Metabolic changes during ingestion of particles by polymorphonuclear leukocytes. *J. Biol. Chem.* 234:1355.
17. Stjernholm, R. L., and Manak, R. C. 1970. Carbohydrate metabolism in leukocytes. XIV. Regulation of pentose cycle activity and glycogen metabolism during phagocytosis. *J. Reticuloendo. Soc.* 8:550.

18. Mulholland, J. H., and Cluff, L. E. 1964. The effect of endotoxin upon susceptibility to infection: The role of the granulocyte. Page 211 in M. Landy and W. Braun, eds. Bacterial endotoxins, proceedings of a symposium. Quinn & Boden Co., Inc., Rahway, New Jersey.
19. Herion, J. C., Walker, R. I., Herring, W. B., and Palmer, J. G. 1965. Effects of endotoxin and nitrogen mustard on leukocyte kinetics. Blood 25:522.
20. Postel, J., Schloerb, P. R., and Furtado, D. 1975. Pathophysiologic alterations during bacterial infusions for the study of bacterial shock. Surg. Gynec. Obstet. 141:683.
21. Rutenburg, S. H., Schweinburg, F. B., and Fine, J. 1960. In vitro detoxification of bacterial endotoxin by macrophages. J. Exptl. Med. 112:801.
22. Cohn, Z. A., and Morse, S. I. 1960. Functional and metabolic properties of polymorphonuclear leucocytes. II. The influence of a lipopolysaccharide endotoxin. J. Exptl. Med. 111:689.
23. Cryer, P. E., Coran, A. G., Sode, J., Herman, C. M., and Horwitz, D. L. 1972. Lethal Escherichia coli septicemia in the baboon: Alpha-adrenergic inhibition of insulin secretion and its relationship to the duration of survival. J. Lab. Clin. Med. 79:622.
24. Gillman, S. M., Bornstein, D. L., and Wood, W. B., Jr. 1961. Studies on the pathogenesis of fever. J. Exptl. Med. 114:729.
25. Freedman, H. H. 1960. Further studies on passive transfer of tolerance to pyrogenicity of bacterial endotoxin. The febrile and leukopenic responses. J. Exptl. Med. 112:619.

26. Roe, C. F., and Kinney, J. M. 1965. The caloric equivalent of fever. II. Influence of major trauma. *Ann. Surg.* 161:140.
27. Clowes, G. H. A., Jr., O'Donnell, T. F., Jr., and Ryan, N. T. 1975. Abnormalities of energy metabolism in sepsis and endotoxemia. Page 248 in B. Urbaschek, R. Urbaschek, and E. Neter, eds. *Gram-negative bacterial infections*. Springer-Verlag, New York/Wien.
28. Chedid, L., Parant, M., Boyer, F., and Skarnes, R. C. 1964. Non-specific host responses in tolerance to the lethal effect of endotoxins. Page 500 in M. Landy and W. Braun, eds. *Bacterial endotoxins, proceedings of a symposium*. Quinn & Boden Co., Inc., Rahway, New Jersey.
29. Greisman, S. E., and Hornick, R. B. 1976. Endotoxin tolerance. Page 43 in R. F. Beers, Jr. and E. G. Bassett, eds. *The role of immunological factors in infectious, allergic, and autoimmune processes*. Raven Press, New York.
30. Petersdorf, R. G., and Shulman, J. A. 1964. The role of tolerance in the action of bacterial endotoxins. Page 482 in M. Landy and W. Braun, eds. *Bacterial endotoxins, proceedings of a symposium*. Quinn & Boden Co., Inc., Rahway, New Jersey.
31. Beeson, P. B. 1947. Tolerance to bacterial pyrogens. II. Role of the reticuloendothelial system. *J. Exptl. Med.* 86:39.
32. Cluff, L. E. 1953. Studies of the effect of bacterial endotoxins on rabbit leukocytes. II. Development of acquired resistance. *J. Exptl. Med.* 98:349.
33. Herring, W. B., Herion, J. C., Walker, R. I., and Palmer, J. G. 1963. Distribution and clearance of circulating endotoxin. *J. Clin. Invest.* 42:79.

34. Weinstein, R. J., and Young, L. S. 1976. Neutrophil function in gram-negative rod bacteremia. *J. Clin. Invest.* 58:190.
35. Braude, A. I., Carey, F. J., and Zalesky, M. 1955. Studies with radioactive endotoxin. II. Correlation of physiologic effects with distribution of radioactivity in rabbits injected with lethal doses of *E. coli* endotoxin labelled with radioactive sodium chromate. *J. Clin. Invest.* 34:858.
36. Fukuda, T., and Akiyama, S. 1963. Mechanisms of endotoxin tolerance in relation to carbohydrate metabolism. *Japan. J. Physiol.* 13:486.
37. Graw, R. G., Jr., Herzig, G., Perry, S., and Henderson, E. S. 1972. Normal granulocyte transfusion therapy. Treatment of septicemia due to gram-negative bacteria. *New Engl. J. Med.* 287:367.
38. Epstein, R. B., Clift, R. A., and Thomas, E. D. 1969. The effect of leukocyte transfusions on experimental bacteremia in the dog. *Blood* 34:782.
39. Epstein, R. B., Waxman, F. J., Bennett, B. T., and Andersen, B. R. 1974. *Pseudomonas* septicemia in neutropenic dogs. I. Treatment with granulocyte transfusions. *Transfusion* 14:51.
40. Heggers, J. P., Robson, M. C., Jennings, P. B., and Fariss, B. L. 1976. Effects of glucose therapy on experimental *Escherichia coli* septicemia. *J. Surg. Res.* 20:33.
41. Stossel, T. P. 1972. Phagocytosis: The department of defense. *New Engl. J. Med.* 286:776.

Table 1. Effects of *E. coli* endotoxin on leukocyte, mature neutrophil, lymphocyte and immature neutrophil counts^a

	Control ^b								
	Day 1	Day 2	Day 3	Day 4	+1	+2	+4	+5-6	+7
WBC									
Mean	12540	12840	13620	13540	4900	7140	19660	53200	8500
±SE	2499	2151	1882	1630	1441	2142	6888	26179	--
N	5	5	5	5	5	5	5	3	1
paired "t" ^c		NS	NS	NS	<0.01	<0.005	NS	NS	NS
Mature Neutrophils									
Mean	6784	7846	7152	7557	2143	3870	11039	23143	396
±SE	1783	1546	493	1317	636	1607	4221	9720	--
N	5	5	5	5	5	5	5	3	1
paired "t" ^c		<0.05	NS	NS	<0.02	<0.05	NS	NS	NS
Lymphocytes									
Mean	2878	2279	4190	2995	2314	2280	5400	12135	2635
±SE	571	324	697	543	828	471	1762	8597	--
N	5	5	5	5	5	5	5	3	1
paired "t" ^c		NS	NS	NS	NS	NS	NS	NS	NS

(Table 1, continued)

Immature Neutrophils

Mean	1113	790	542	985	233	269	2115	6833	340
±SE	202	177	175	120	44	211	1450	3237	--
N	5	5	5	5	5	5	5	3	1
paired "t"	<0.025	<0.025	<0.025	NS	<0.005	<0.01	NS	NS	NS

^a Saline pretreated control group (Group I). On Days 1, 2 and 3 saline was administered intravenously in volumes equal to the endotoxin given to experimental animals. On Day 4, 2 X LD₁₀₀ (6 mg/kg) E. coli endotoxin was injected. All values represent mean±SE.

^b Initial measurements in each group on four consecutive days. Day 4 mean±SE serves as the control measurement for the time designations of hours.

^c p values represent paired comparisons from Day 1 control values through the time designations of days. Through the time designations of hours, p values are paired from Day 4 control value.

Table 2. Effects of *E. coli* endotoxin on leukocyte, mature neutrophil, lymphocyte and immature neutrophil counts^a

	Control ^b						Hrs			
	Day 1	Day 2	Day 3	Day 4	+1	+2	+4	+5-6	+8-9	
WBC										
Mean	12840	27680	26720	39000	5120	14440	25280	30900	33880	
±SE	908	2978	3211	2735	985	1710	5107	5609	4268	
N	5	5	5	5	5	5	5	5	5	
paired "t" ^{nc}	<0.01	<0.01	<0.02	<0.001	<0.001	<0.005	<0.05	NS	NS	
unpaired "t" nd	NS	<0.005	<0.01	<0.001	NS	<0.05	NS	NS	NS	
Mature Neutrophils										
Mean	6801	20138	17215	26790	2360	6881	10704	12973	17219	
±SE	1320	2196	4256	2223	724	1647	1573	4183	1305	
N	5	5	5	5	5	5	5	5	5	
paired "t" ^c	<0.005	<0.005	<0.05	<0.001	<0.001	<0.005	<0.01	NS	<0.02	
unpaired "t" ^c	NS	<0.005	<0.05	<0.001	NS	NS	NS	NS	NS	
Lymphocytes										
Mean	3661	3649	4628	3086	1650	2912	6964	5606	7221	
±SE	451	959	1290	1223	366	980	4155	3721	4566	
N	5	5	5	5	5	5	5	5	5	

(Table 2 continued)

	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
paired "t"											
unpaired "t"	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Immature Neutrophils											
Mean	888	272	3016	7939	528	3653	6154	11650	9724		
±SE	348	1481	995	4635	146	563	2044	3816	4406		
N	5	5	5	5	5	5	5	5	5	5	5
paired "t"	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
unpaired "t"	NS	NS	<0.05	NS	NS	<0.001	NS	NS	NS	NS	NS

^aEndotoxin pretreated experimental group (Group II); *E. coli* endotoxin administered intravenously Days 1 and 2, 1/1,000 LD₁₀₀ (0.003 mg/kg); Day 3, LD₁₀₀ (3 mg/kg); and Day 4, 2 X LD₁₀₀ (6 mg/kg). All values represent mean±SE.

^bInitial measurements in each group on four consecutive days. Day 4 mean±SE serves as the control measurement for the time designations of hours.

^cp values represent paired comparisons from Day 1 control values through time designations of days. Through the time designations of hours, p values are paired from Day 4 control value.

^dp values represent an unpaired comparison from saline pretreated control group.

Table 3. Lethality response to superlethal doses of E. coli endotoxin^a

	<u>Injected</u>	<u>Day 4</u>	
		<u>Died</u>	<u>Lived</u>
Saline pretreated controls (Group I) ^b	N=5	5 within +7 hours	0
Endotoxin pretreated experimentals (Group II) ^c	N=5	0	5 for 30 days

^a2 X LD₁₀₀ endotoxin (6 mg/kg) injected intravenously on Day 4.

^bOn Days 1, 2 and 3 saline was administered intravenously in volumes equal to the endotoxin given to experimental animals.

^cE. coli endotoxin administered intravenously; Days 1 and 2, 1/1,000 LD₁₀₀ (0.003 mg/kg) and Day 3, LD₁₀₀ (3 mg/kg).

FIGURE LEGENDS

- Figure 1. Effect of intravenous administration of E. coli endotoxin on rectal temperature (T_r - °C) in dogs. Results (mean±SE) from 10 dogs are depicted in two groups: saline controls (n=5) and endotoxin experimentals (n=5). Experimental animals received sublethal doses of E. coli endotoxin; 1/1,000 LD₁₀₀ on Days 1 and 2 and LD₁₀₀ on Day 3. The control group received equal volumes of saline on Days 1, 2 and 3. P values represent an unpaired comparison between control and experimental groups.
- Figure 2. Effect of intravenous administration of E. coli endotoxin on white blood cell concentration (mm^3) in dogs. Results (mean±SE) from 10 dogs are depicted in two groups: saline controls (n=5) and endotoxin experimentals (n=5). (See Figure 1 for details of experiments.)
- Figure 3. Effect of intravenous administration of E. coli endotoxin on blood glucose concentration (mg%) in dogs. Results (mean±SE) from 10 dogs are depicted in two groups: saline controls (n=5) and endotoxin experimentals (n=5). (See Figure 1 for details of experiments.)
- Figure 4. Effect of intravenous pretreatment with E. coli endotoxin on rectal temperature (°C), white blood cell concentration, and blood glucose concentration following 2 X LD₁₀₀ E. coli endotoxin (6 mg/kg). Results (mean±SE) from 10 dogs are depicted; 5 from experimental group pretreated with sublethal and lethal doses of endotoxin ** and 5 control animals receiving saline injections in place of endotoxin *. Both groups received 2 X LD₁₀₀ endotoxin on day of challenge (Day 4). P values represent an unpaired comparison between control and experimental group.




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5