

AD-A041 035

BALLISTIC RESEARCH LABS ABERDEEN PROVING GROUND MD
CONFIDENCE BOUNDS FOR THE GENERAL LINEAR MODEL. (U)
MAY 77 M S TAYLOR, J R MOORE

F/G 12/1

UNCLASSIFIED

BRL-1986

NL


| OFF |

AD
A041035



END

DATE
FILMED
7-77



RLR 1986

BRL

12

AD

AD A 041 035

REPORT NO. 1986

CONFIDENCE BOUNDS FOR THE GENERAL
LINEAR MODEL

Malcolm S. Taylor
J. Richard Moore

May 1977

DDC
JUN 29 1977
C

Approved for public release; distribution unlimited.

AU NO.
DDC FILE COPY.

USA ARMAMENT RESEARCH AND DEVELOPMENT COMMAND
USA BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating
or sponsoring activity is prohibited.

Additional copies of this report may be obtained
from the National Technical Information Service,
U.S. Department of Commerce, Springfield, Virginia
22151.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|---|---|
| 1. REPORT NUMBER BRL REPORT NO. 1986 | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) CONFIDENCE BOUNDS FOR THE GENERAL LINEAR MODEL | 5. TYPE OF REPORT & PERIOD COVERED FINAL / repl. | |
| 7. AUTHOR(s) Malcolm S. Taylor J. Richard Moore | 6. PERFORMING ORG. REPORT NUMBER | |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory Aberdeen Proving Ground, MD 21005 | 8. CONTRACT OR GRANT NUMBER(s) | |
| 11. CONTROLLING OFFICE NAME AND ADDRESS US Army Materiel Development and Readiness Command 5001 Eisenhower Avenue Alexandria, Virginia 22333 | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1T161102AH43 | |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) BRL-1986 | 12. REPORT DATE MAY 1977 | |
| | 13. NUMBER OF PAGES 17 | |
| | 15. SECURITY CLASS. (of this report) UNCLASSIFIED | |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) general linear model confidence bounds regression data analysis X beta | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper, for the general linear model $Y = XB + e$, we consider the construction of confidence bounds about the entire regression line. To accomplish this we exploit a powerful theorem of Scheffé. A procedure often encountered is one in which a set of confidence intervals about $E(y x)$ or prediction intervals for future observations are determined and then the end points are connected in such a fashion as to describe an envelope. The belief is that what has been accomplished is precisely what Scheffé's theorem allows one to do. → next page | | |

DDC
RECEIVED
JUN 02 1977
C

$E(y, x)$ 050450

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

cont

→ In addition, we present some extensions concerning confidence bounds about combinations of regression lines and suggest a useful application of these results. Specifically, we propose to use the confidence bounds about the difference of regression lines to make a quantitative assessment of when and where independent sets of data characterizing the same phenomena are in agreement or disagreement.

2

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE OF CONTENTS

| | Page |
|---|------|
| 1. INTRODUCTION | 5 |
| 2. CONFIDENCE REGION FOR A POLYNOMIAL | 7 |
| 3. THE TWO SAMPLE CASE | 9 |
| 4. THE k-SAMPLE CASE | 12 |
| REFERENCES | 16 |
| DISTRIBUTION LIST | 18 |

ACCESSION for

NTIS White Section

DBS Buff Section

UNANNOUNCED

JUSTIFICATION

DISTRIBUTION/AVAILABILITY CODES

AVAIL. AND/OR SPECIAL

A

1. INTRODUCTION

It is appropriate at the onset that we devote a few paragraphs to the introduction of the general linear hypothesis model of full rank. We want to consider uncorrelated observations y_1, y_2, \dots, y_n that satisfy the relation

$$y_i = \sum_{j=1}^p x_{ij} \beta_j + e_i, \quad i = 1, 2, \dots, n \quad (1.1)$$

and are linear in the unknown parameters $\beta_1, \beta_2, \dots, \beta_p$ with known coefficients x_{ij} and random term e_i satisfying

$$E(y_i) = \sum_{j=1}^p x_{ij} \beta_j,$$

and

$$\text{Var}(y_i) = \sigma^2.$$

In other words, the random term e_i is a random variable with expected value $E(e_i)$ equal to zero and unknown variance $\text{Var}(e_i)$ equal to σ^2 .

The problem, in its most general sense, involves determining point and interval estimates of several quantities of interest of the model and the testing of various statistical hypotheses.

For compactness of notation and ease of manipulation let

$$Y = \begin{pmatrix} y_1 \\ y_2 \\ \cdot \\ \cdot \\ y_n \end{pmatrix}, \quad X = \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1p} \\ x_{21} & x_{22} & \cdots & x_{2p} \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ x_{n1} & x_{n2} & \cdots & x_{np} \end{pmatrix}, \quad \beta = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \cdot \\ \cdot \\ \beta_p \end{pmatrix}, \quad e = \begin{pmatrix} e_1 \\ e_2 \\ \cdot \\ \cdot \\ e_n \end{pmatrix};$$

then we can write the system of relations (1.1) as

$$Y = X\beta + e$$

and proceed to define the general linear hypothesis model of full rank as follows:

Definition 1.1. The model $Y = X\beta + e$ where Y is a random observed vector, e is a random vector, X is an $n \times p$ matrix of known fixed quantities, and β is a vector of unknown parameters is called the general linear hypothesis model of full rank, provided the rank of X is equal to p where $p \leq n$.

In the present inquiry we restrict our consideration to the normal theory case, which means the random vector e , already satisfying $E(e) = 0$ and $\text{cov}(e) = \sigma_i^2$, will, in addition, be assumed to be normally distributed.

The problem most frequently addressed is that of estimating the unknown parameters β_j on the basis of the observations y_i . These estimates of β_j , denoted by $\hat{\beta}_j$, are functions of y_i ; and, as such, are themselves random variables about which confidence intervals can be constructed. These ideas are fully developed in a number of text-books.^{1,2} A point not so widely expounded is that the usual frequency interpretation of a confidence interval based on a single sample y_1, y_2, \dots, y_n holds only for a single coefficient β_j ; if the same data are used to determine confidence intervals for both β_i and β_j , $i \neq j$, the probability is not $1-\alpha$ that the confidence intervals thus constructed will simultaneously contain β_i and β_j . The complexity is advanced by the fact that the interval estimates are not independent; so, in general, only a single confidence statement can be made from a single set of observations.

It is not our intent here to address this problem directly; such an inquiry falls into the general area of simultaneous confidence intervals. It is our intent, however, to consider a ramification of this problem: namely, the construction of a confidence envelope about the entire regression line. We will, in addition, provide some results concerning confidence envelopes about combinations of regression lines and implications of their use.

Toward this end consider the following definition due to Bose:³

-
- 1 Graybill, F. A., An Introduction to Linear Statistical Models, Volume I, McGraw-Hill Book Company, Inc., New York, 1961.
 - 2 Rao, C. R., Linear Statistical Inference and Its Applications, John Wiley & Sons, Inc., New York, 1965.
 - 3 Bose, R. C., "The Fundamental Theorem of Linear Estimation", Proceedings of the 31st Indian Science Congress, 1944, pp. 2-3.

Definition 1.2. A parametric function ψ is called an estimable function if it has an unbiased linear estimate, i.e., if there exists an n -vector a of constant coefficients such that $E(a'y) = \psi$.

If L is a p -dimensional space of estimable functions with basis $\{\psi_1, \psi_2, \dots, \psi_p\}$ and $\hat{\psi}$ is the least squares estimate of $\psi \in L$, then we have the following theorem due to Scheffé⁴.

Theorem 1.1. Under the general linear hypothesis model (normal case) the probability is $1 - \alpha$ that simultaneously for all $\psi \in L$

$$\hat{\psi} - S\hat{\sigma}_\psi \leq \psi \leq \hat{\psi} + S\hat{\sigma}_\psi$$

where the constant $S = \{pF_\alpha(p, n-r)\}^{1/2}$ and rank $X = r$.

The implications of this theorem are far reaching; and in this article we will exploit a single facet, albeit an important and useful one. To facilitate this we need to be aware of the fact that since least squares estimates $\hat{\beta}$ are BLUE, the elements of the vector β of the general linear model of full rank form a basis of a space L of estimable functions which includes polynomials as a special case.

2. CONFIDENCE REGION FOR A POLYNOMIAL

To determine a confidence region for a polynomial with observational equations

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \dots + \beta_{p-1} x_i^{p-1} + e_i, \quad i = 1, 2, \dots, n$$

in the model $Y = X\beta + e$, the $n \times p$ matrix $X = (x_{ij})$ of known constant coefficients takes the form

4 Scheffé, H., The Analysis of Variance, John Wiley & Sons, Inc., New York, 1959.

$$X = \begin{pmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{p-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{p-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & x_n & x_n^2 & \dots & x_n^{p-1} \end{pmatrix} .$$

The least squares estimate of β is given by $\hat{\beta} = (X'X)^{-1}X'Y$. If we choose $\psi_i = \beta_i$, $i = 0, 1, \dots, p-1$, then $\{\psi_i\} = \{\beta_i\}$ is a set of linearly independent estimable functions which forms a basis for the space L . For any value x_0 denote $X'_0 = (1, x_0, \dots, x_0^{p-1})$. Clearly, $E(y_0) = X'_0\beta \in L$ and has least squares estimate

$$X'_0\hat{\beta} = X'_0(X'X)^{-1}X'Y = \sum_{i=1}^n a_i y_i$$

where the coefficients a_i are the elements of the $1 \times n$ vector $X'_0(X'X)^{-1}X'$. Thus

$$\begin{aligned} \sum_{i=1}^n a_i^2 &= X'_0(X'X)^{-1}X'[X'_0(X'X)^{-1}X']' \\ &= X'_0(X'X)^{-1}X'X(X'X)^{-1}X_0 \\ &= X'_0(X'X)^{-1}X_0 \end{aligned}$$

so that $\sigma_{\psi}^2 = \sigma^2 X'_0(X'X)^{-1}X_0$ with unbiased estimate $s^2 X'_0(X'X)^{-1}X_0$.

From Scheffé's theorem we can assert with probability $1-\alpha$ that simultaneously for all $\psi \in L$ and, in particular, $X'_0\beta \in L$

$$X'_0\hat{\beta} - S\hat{\sigma}_{\psi} \leq X'_0\beta \leq X'_0\hat{\beta} + S\hat{\sigma}_{\psi}$$

where $S = [pF_{\alpha}(p, n-p)]^{1/2}$.

As an illustration, suppose the paired data (1.20, 0.34), (1.37, 0.94), (1.38, 0.99), (1.65, 1.58), (1.71, 2.08), (1.82, 2.25) are characterized by the quadratic $y = -0.31x^2 + 3.97x - 3.95$ over the interval of interest, $1 \leq x \leq 2$. The 95% confidence region for the entire true line is given by

$$X'_0 \hat{\beta} - (5.28) \hat{\sigma}_\psi \leq X'_0 \beta \leq X'_0 \hat{\beta} + (5.28) \hat{\sigma}_\psi$$

as shown in Figure 1.

Grubbs⁵ showed that for the case $y = \beta_0 + \beta_1 x$ the confidence bounds resulting from Scheffé's theorem are

$$\hat{\beta}_0 + \hat{\beta}_1 x \pm [2F_\alpha(2, n-2)]^{1/2} S \left(\frac{1}{n} + \frac{n(x-\bar{x})^2}{A_{xx}} \right)^{1/2} \quad (2.1)$$

where $S = \left(\frac{1}{n-2} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \right)^{1/2}$ and $A_{xx} = n \sum x_i^2 - [\sum x_i]^2$.

Note that the value x appearing in (2.1) is not limited to an x_i which appears in the observations (x_i, y_i) , $i = 1, 2, \dots, n$.

3. THE TWO-SAMPLE CASE

Suppose two independent sets of data have given rise to two characterizations of the same phenomenon so that we are now confronted with what is, in essence, two models:

$$Y_1 = X_1 \beta_1 + e_1, \text{ an } n_1 \times p_1 \text{ problem,}$$

and

$$Y_2 = X_2 \beta_2 + e_2, \text{ an } n_2 \times p_2 \text{ problem.}$$

We can still represent this situation as $Y = X\beta + e$ where now

⁵ Grubbs, F. E., *Linear Statistical Regression and Functional Relations*, BRL Report No. 1842, November 1975. (AD #A018651)




FIGURE 1

95 % CONFIDENCE LIMITS FOR A LINEAR FUNCTION

$$Y = \begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix}, \quad X = \begin{pmatrix} X_1 & | & 0 \\ \hline 0 & | & X_2 \end{pmatrix}, \quad \beta = \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}, \quad e = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}.$$

The least squares estimate is given, as before, by $\hat{\beta} = (X'X)^{-1}X'Y$.

Consider now the difference of two polynomials $y_1^* - y_2^* \in L$ with LS estimate $X_1^{*'}\hat{\beta}_1 - X_2^{*'}\hat{\beta}_2$ where $X_i^{*'} = (1, x_i, \dots, x_i^{p_i-1})$. Rewriting,

$$\begin{aligned} X_1^{*'}\hat{\beta}_1 - X_2^{*'}\hat{\beta}_2 &= \begin{pmatrix} X_1^{*'} & | & -X_2^{*'} \end{pmatrix} \begin{pmatrix} \hat{\beta}_1 \\ \hat{\beta}_2 \end{pmatrix} = \begin{pmatrix} X_1^{*'} & | & -X_2^{*'} \end{pmatrix} (X'X)^{-1}X'Y \\ &= \sum_{i=1}^{n_1+n_2} a_i y_i \end{aligned}$$

where the coefficients a_i are the elements of the $1 \times (n_1+n_2)$ vector $X^{*'}(X'X)^{-1}X'$. Thus,

$$\sum a_i^2 = X^{*'}(X'X)^{-1}X'[X^{*'}(X'X)^{-1}X']' = X^{*'}(X'X)^{-1}X'X(X'X)^{-1}X^* = X^{*'}(X'X)^{-1}X^*.$$

$$\text{Now } X^{*'} = \begin{pmatrix} X_1^{*'} & | & -X_2^{*'} \end{pmatrix} \text{ and } (X'X)^{-1} = \begin{pmatrix} (X_1'X_1)^{-1} & | & 0 \\ \hline 0 & | & (X_2'X_2)^{-1} \end{pmatrix}$$

$$\text{so } X^{*'}(X'X)^{-1}X^* = X_1^{*'}(X_1'X_1)^{-1}X_1^* + X_2^{*'}(X_2'X_2)^{-1}X_2^*.$$

As before, $\text{Var}(\hat{\psi})$ has the unbiased estimate $\hat{\sigma}_{\hat{\psi}}^2 = s^2 X^{*'}(X'X)^{-1}X^*$,

where s^2 is now the pooled estimate of the variance, and with probability $1 - \alpha$ simultaneously for all $\psi \in L$

$$X_1^* \hat{\beta}_1 - X_2^* \hat{\beta}_2 - S \hat{\sigma}_\psi \leq y_1^* - y_2^* \leq X_1^* \hat{\beta}_1 - X_2^* \hat{\beta}_2 + S \hat{\sigma}_\psi$$

$$\text{with } S = \left[(p_1 + p_2) F_\alpha [p_1 + p_2, n_1 + n_2 - (p_1 + p_2)] \right]^{1/2}.$$

To illustrate one of the most useful potential applications of this result consider the situation where we are presented with two sets of data, $\{(x_{11}, y_{11}), (x_{12}, y_{12}), \dots, (x_{1n}, y_{1n})\}$, $\{(x_{21}, y_{21}), (x_{22}, y_{22}), \dots, (x_{2m}, y_{2m})\}$, collected from the same process; and we want to say something about the similarity or dissimilarity of the two descriptions. Suppose each set is fitted with a quadratic; and we construct the confidence bound about the difference $y_1^* - y_2^*$, as shown in Figure 2. Over the region $(1.65 \leq x \leq 2.05)$, where the confidence bounds cover the line $y = 0$, we will say the two descriptions are consistent, although the associated probability level cannot be attached without qualification and interpretation.

The extension of Grubbs result (2.1) to this case is direct; the bounds take the form

$$X_1^* \hat{\beta}_1 - X_2^* \hat{\beta}_2 \pm \left[(p_1 + p_2) F_\alpha (p_1 + p_2, n_1 + n_2 - p_1 - p_2) \right]^{1/2} \cdot S \cdot \left[\frac{1}{n_1} + \frac{1}{n_2} + \frac{n_1(X^* - \bar{X}_1)}{A_{xx}^1} + \frac{n_2(X^* - \bar{X}_2)}{A_{xx}^2} \right]^{1/2}$$

where S^2 is the pooled estimate of variance and A_{xx}^i is computed from the i -th data set.

4. THE k-SAMPLE CASE

The straightforward generalization to k sets of data proceeds as follows:

$$\begin{pmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_k \end{pmatrix} = \begin{pmatrix} X_1 & 0 & \dots & 0 \\ 0 & X_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & X_k \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_k \end{pmatrix} + \begin{pmatrix} e_1 \\ e_2 \\ \vdots \\ e_k \end{pmatrix}$$




FIGURE 2

95 % CONFIDENCE LIMITS FOR THE DIFFERENCE BETWEEN TWO QUADRATIC FUNCTIONS

and

$$\hat{\beta} = \begin{pmatrix} \hat{\beta}_1 \\ \hline \hat{\beta}_2 \\ \hline \cdot \\ \hline \hat{\beta}_k \end{pmatrix} = \begin{pmatrix} (X_1'X_1)^{-1}X_1'Y_1 \\ \hline (X_2'X_2)^{-1}X_2'Y_2 \\ \hline \cdot \\ \hline (X_k'X_k)^{-1}X_k'Y_k \end{pmatrix}$$

Now for $X^* = \left(X_1^* \mid \dots \mid X_k^* \right)$ where $X_i^* = \left(1, x, \dots, x^{p_i-1} \right)$

we can write $\hat{y}_i^* = X_i^{*'} \hat{\beta}_i = X_i^{*'} (X_i'X_i)^{-1} X_i'Y_i$, $i = 1, 2, \dots, k$.

Suppose we now consider the estimate of $\sum c_i y_i^*$:

$$\sum c_i \hat{y}_i^* = \sum c_i X_i^{*'} \hat{\beta}_i = \sum c_i X_i^{*'} (X_i'X_i)^{-1} X_i'Y_i. \text{ Rewriting,}$$

$$\sum c_i X_i^{*'} \hat{\beta}_i = \left(c_1 X_1^{*'} \mid \dots \mid c_k X_k^{*'} \right) (X'X)^{-1} X'Y = \sum_{i=1}^{n_1 + \dots + n_k} a_i y_i$$

where the coefficients a_i are the elements of the $1 \times \sum n_i$ vector $CX^* (X'X)^{-1} X'$.

Thus,

$$\begin{aligned} \sum a_i^2 &= CX^* (X'X)^{-1} X' [CX^* (X'X)^{-1} X']' \\ &= CX^* (X'X)^{-1} X' X (X'X)^{-1} CX^* \\ &= CX^* (X'X)^{-1} CX^*. \end{aligned}$$

Now

$$CX^*{}' = \left(c_1 X_1^*{}' \mid \dots \mid c_k X_k^*{}' \right) \text{ and } (X'X)^{-1} = \begin{pmatrix} (X_1'X_1)^{-1} & & \\ & \ddots & \\ & & (X_k'X_k)^{-1} \end{pmatrix}$$

so $CX^*{}'(X'X)^{-1}CX^* = \sum_{i=1}^k c_i^2 X_i^*{}'(X_i'X_i)^{-1}X_i^*$. The confidence region now assumes

the form

$$\sum c_i X_i^*{}' \hat{\beta}_i - S \left(sCX^*{}'(X'X)^{-1}CX^* \right)^{1/2} \leq \sum c_i y_i^* \leq \sum c_i X_i^*{}' \hat{\beta}_i + S \left(sCX^*{}'(X'X)^{-1}CX^* \right)^{1/2};$$

$$\text{with } S = \left(\sum p_i \cdot F_{\alpha}(\sum p_i, \sum n_i - \sum p_i) \right)^{1/2}.$$

For the linear case we obtain

$$\begin{aligned} CX^*{}'(X'X)^{-1}CX^* &= \sum_{i=1}^k c_i^2 X_i^*{}'(X_i'X_i)^{-1}X_i^* \\ &= \sum_{i=1}^k c_i^2 \left(\frac{1}{n_i} + n_i (x_{i0} - \bar{x}_i)^2 \right), \end{aligned}$$

and the two sample case (Section 3) is obtained by setting $c_1 = 1$ and $c_2 = -1$.

REFERENCES

1. Graybill, F. A., An Introduction to Linear Statistical Models, Volume I, McGraw-Hill Book Company, Inc., New York, 1961.
2. Rao, C. R., Linear Statistical Inference and Its Applications, John Wiley & Sons, Inc., New York, 1965.
3. Bose, R. C., "The Fundamental Theorem of Linear Estimation", Proceedings of the 31st Indian Science Congress, 1944, pp. 2-3.
4. Scheffé, H., The Analysis of Variance, John Wiley & Sons, Inc., New York, 1959.
5. Grubbs, F. E., "Linear Statistical Regression and Functional Relations", BRL Report No. 1842, November 1975. (AD #A018651)

DISTRIBUTION LIST

| <u>No. of Copies</u> | <u>Organization</u> | <u>No. of Copies</u> | <u>Organization</u> |
|--------------------------|---|--------------------------|--|
| 12 | Commander Defense Documentation Center ATTN: DDC-TCA Cameron Station Alexandria, VA 22314 | 2 | Commander US Army Mobility Equipment Research & Development Command ATTN: Tech Docu Cen, Bldg. 315 DRSME-RZT Fort Belvoir, VA 22060 |
| 1 | Commander US Army Materiel Development and Readiness Command ATTN: DRCDMA-ST 5001 Eisenhower Avenue Alexandria, VA 22333 | 1 | Commander US Army Armament Research and Development Command ATTN: DRDAR-TD Dover, NJ 07801 |
| 1 | Commander US Army Aviation Systems Command ATTN: DRSV-E 12th and Spruce Streets St. Louis, MO 63166 | 1 | Commander US Army Armament Materiel and Readiness Command Rock Island, IL 61202 |
| 1 | Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035 | 1 | Commander US Army Harry Diamond Labs ATTN: DRXDO-TI 2800 Powder Mill Road Adelphi, MD 20783 |
| 1 | Commander US Army Electronics Command ATTN: DRSEL-RD Fort Monmouth, NJ 07703 | 1 | Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SA White Sands Missile Range NM 88002 |
| 1 | Commander US Army Missile Research and Development Command ATTN: DRDMI-R Redstone Arsenal, AL 35809 | | <u>Aberdeen Proving Ground</u> Marine Corps Ln Ofc Dir, USAMSAA |
| 1 | Commander US Army Tank Automotive Development Command ATTN: DRDTA-RWL Warren, MI 48090 | | |