

AD A 041149

NPS55-77-6

NAVAL POSTGRADUATE SCHOOL
Monterey, California



MEANS AND VARIANCES OF STOCHASTIC VECTOR PRODUCTS
WITH APPLICATIONS TO RANDOM LINEAR MODELS

by

Gerald G. Brown


and

Herbert C. Rutemiller

February 1977

Approved for public release; distribution unlimited

AD NO. _____
DDC FILE COPY




NAVAL POSTGRADUATE SCHOOL
Monterey, California

Rear Admiral Isham Linder
Superintendent

Jack R. Borsting
Provost


Reproduction of all or part of this report is authorized.

This report was prepared by:




Gerald G. Brown
Associate Professor
Departments of Computer Science
and Operations Research

Reviewed by:



Michael G. Sovereign, Chairman
Department of Operations Research

Released by:



Robert R. Fossum
Dean of Research

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-77-6	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) MEANS AND VARIANCES OF STOCHASTIC VECTOR PRODUCTS WITH APPLICATIONS TO RANDOM LINEAR MODELS.		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Gerald G. Brown and Herbert C. Rutemiller		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12/237
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		12. REPORT DATE February 1977
		13. NUMBER OF PAGES 18
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Random Linear Models, Stochastic Programming, Chance Constrained Linear Programming, Tchebycheff Inequalities, Joint Tchebycheff Bounds, Dependent Stochastic Vector Products, Moments of Dependent Stochastic Vector Products		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Many mathematical models in operations research require computation of products of vectors whose elements are random variables. Unfortunately, analytic results for functions of interest are only obtained through highly restrictive, often unrealistic, choices of prior densities for the vectors' elements. Often, an investigation is performed by discretizing the random variables at point-quantile levels, or by outright simulation. This paper addresses the problem of characterizing the inner product of two stochastic vectors with		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

arbitrary multivariate densities. Expressions for means and variances of vector products are obtained, and used to make Tchebycheff-type probability statements. Included are applications to stochastic programming models.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

D D C
RECEIVED
JUL 8 1977
RECEIVED
D

MEANS AND VARIANCES OF STOCHASTIC VECTOR PRODUCTS
WITH APPLICATIONS TO RANDOM LINEAR MODELS

ACCESSION FOR	
NTIS	Write Section ☒
DDC	Off Section ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION AVAILABILITY CODES	
DISC. AVAIL. CODES	

A

by

Gerald G. Brown
Naval Postgraduate School
Monterey, California

and

Herbert C. Rutemiller
California State University
Fullerton, California

Many mathematical models in operations research require computation of products of vectors whose elements are random variables. Unfortunately, analytic results for functions of interest are only obtained through highly restrictive, often unrealistic, choices of prior densities for the vectors' elements. Often, an investigation is performed by discretizing the random variables at point-quantile levels, or by outright simulation. This paper addresses the problem of characterizing the inner product of two stochastic vectors with arbitrary multivariate densities. Expressions for means and variances of vector products are obtained, and used to make Tchebycheff-type probability statements which can accommodate and exploit stochastic dependence. Included are applications to stochastic programming models.

1. INTRODUCTION

Mathematical models in operations research often require the calculation of quantities which are sums of products of random variables. (For example, see the realistic view of stochastic linear programming taken by Tintner [15]). Sums of products of random variables are the generic structural elements fundamental to linear models exhibiting complex randomness.

In the interest of tractable mathematics, it is commonly assumed either that the random variables may be treated as constants, or replaced by their expectations. Even where these simplifying assumptions are not made, severe parametric assumptions, e.g., normality, pairwise independence, etc., are usually required to achieve analytic probability statements concerning the function of interest. Other approaches use weighted sums of uniform, exponential or other variates in approximation schemes, or use characteristic functions of component densities to carry out the indicated transformation and attempt inversion of the result. Finally, Monte Carlo solutions to this problem may be achieved, but are of limited value due to the parametric specificity and high cost of results. For excellent examples of these methods see Jagannathan [7]; King, Sampson, and Simms [8]; Miller and Wagner [9]; Searle [12]; and Sengupta [13, 14].

This paper addresses the problem of determining the mean and variance of sums of products of random variables in the most general linear case, i.e., arbitrary correlated multivariate densities for all components in the model. The results obtained are useful for making non-parametric Tchebycheff-type probability statements. Particular examples are given which use these probability statements as chance constraints in stochastic linear programming.

Simplifications of this model for moments of weighted sums of random variables have been presented by several authors. For instance, the highly successful heuristic given by Graves and Whinston [5] for the quadratic assignment problem is based on such a derivation.

2. NOTATION AND DEVELOPMENT

Let b and c be two random n -element column vectors. We denote the expectation of these vectors as

$$E \begin{pmatrix} b \\ c \end{pmatrix} = \mu = \begin{pmatrix} \mu_b \\ \mu_c \end{pmatrix} .$$

The symbols $\tilde{b}$ and $\tilde{c}$ will indicate the deviation of each random vector from its expectation. The variance-covariance matrix will be signified by

$$E \begin{pmatrix} \tilde{b} \\ \tilde{c} \end{pmatrix} \begin{pmatrix} \tilde{b} \\ \tilde{c} \end{pmatrix}' = \Sigma = \begin{pmatrix} \Sigma_{bb} & \Sigma_{bc} \\ \Sigma_{cb} & \Sigma_{cc} \end{pmatrix} ,$$

while the symbol Σ^* will signify this matrix with b and c interchanged.

Let q be a random scalar variable, defined as the inner product of these random vectors:

$$q = b'c. \quad (1)$$

Expressions for the mean, $E(q)$, and variance, $V(q)$, in terms of the moments of b and c are derived in the appendix. They are:

$$E(q) = \mu_b' \mu_c + \text{Tr } \Sigma_{bc}, \quad (2)$$

where $\text{Tr } \Sigma_{bc}$ indicates the trace, or sum of the diagonal elements of Σ_{bc} ;

$$\begin{aligned} V(q) = & \mu' \Sigma \mu + 2\mu_b' E(\tilde{c}\tilde{c}'\tilde{b}) + 2\mu_c' E(\tilde{b}\tilde{b}'\tilde{c}) \\ & + E(\tilde{b}'\tilde{c}\tilde{c}'\tilde{b}) - (\text{Tr } \Sigma_{bc})^2. \end{aligned} \quad (3)$$

Equations (2) and (3) represent the most general outcome, where any element of b may be correlated with any other element of b or c . Two special cases occur frequently. First, suppose that the elements of b are correlated with one another, the elements of c are correlated with one another, but that b and c are stochastically independent. Then

$$\Sigma = \begin{pmatrix} \Sigma_{bb} & 0 \\ 0 & \Sigma_{cc} \end{pmatrix},$$

and

$$E(q) = \mu_b' \mu_c, \quad (4)$$

$$V(q) = \mu_b' \Sigma_{cc} \mu_b + \mu_c' \Sigma_{bb} \mu_c + \text{Tr}(\Sigma_{bb}\Sigma_{cc}). \quad (5)$$

Secondly, we note that, if all elements of both matrices are stochastically independent, Σ is a diagonal matrix, and computation of (5) is greatly simplified.

An assumption frequently made is that elements of vectors b and c have a multivariate normal distribution. If we adopt this parametric condition, then, even in the general case where b and c are correlated, (3) is greatly simplified. For multivariate normal distributions, $E(\tilde{b}\tilde{b}'\tilde{c}) = E(\tilde{c}\tilde{c}'\tilde{b}) = 0$, and all higher moments are expressible in terms of the first two moments [1, p. 38]. Equation (3) becomes

$$V(q) = \mu'\Sigma\mu + \text{Tr}(\Sigma_{bb}\Sigma_{cc}) + \text{Tr}(\Sigma_{bc}^2) . \quad (6)$$

3. AN ESTIMATION EXAMPLE

A job shop is bidding on the manufacture of units which will consume two raw materials and skilled labor. Let b be the vector of material and labor requirements per unit of product, and c the vector of respective costs per unit of b . Then the total manufacturing cost per unit, q , may be computed with equation (1). Due to unknown factors in the manufacturing process (e.g., proportion of scrap units produced) the elements of b are random variables. Further, although the skilled labor cost is fixed by contract, other elements of c are random variables dependent upon commodity prices and other factors.

Historical cost accounting data are available for similar products in the form of job orders and associated material requisitions and labor charges, and are used as prior

information to determine the means and variance-covariance matrices of b and c as follows:

$$\mu_b = \begin{pmatrix} 4.2 \text{ units of I} \\ 6.8 \text{ units of II} \\ 4.8 \text{ hours of labor} \end{pmatrix}$$

$$\mu_c = \begin{pmatrix} \$2.15 \text{ per unit of I} \\ \$1.50 \text{ per unit of II} \\ \$6.20 \text{ per hour} \end{pmatrix}$$

$$\Sigma_{bb} = \begin{pmatrix} 0.7 & 0.8 & 0.2 \\ 0.8 & 1.2 & 0.3 \\ 0.2 & 0.3 & 0.5 \end{pmatrix}$$

$$\Sigma_{cc} = \begin{pmatrix} 0.6 & 0.1 & 0.0 \\ 0.1 & 0.2 & 0.0 \\ 0.0 & 0.0 & 0.0 \end{pmatrix}$$

$$\Sigma_{bc} = 0$$

Prices of the two raw materials are positively correlated, as are the amounts consumed and the labor expended. In this example, there is no correlation between commodity prices and quantities consumed during manufacture, so (4) and (5) are the appropriate equations.

$$\begin{aligned}
 E(q) &= \mu'_b \mu_c \\
 &= \$48.99
 \end{aligned}$$

$$\begin{aligned}
 V(q) &= \mu'_c \Sigma_{bb} \mu_c + \mu'_b \Sigma_{cc} \mu_b + \text{Tr}(\Sigma_{bb} \Sigma_{cc}) \\
 &= 41.23 + 25.54 + 0.82 \\
 &= 67.59
 \end{aligned}$$

Hence, the expected cost for one completed unit is \$48.99 with a standard deviation of \$8.22.

The (unknown) probability distribution of q depends upon the mathematical form of the multivariate density functions of b and c , and the transformation, equation (1). However, the moments calculated above are exact, and we may use them to make non-parametric statements about the distribution of q . Two illustrations follow.

Tchebycheff's Inequality

Regardless of the underlying distributions, at least $1 - 1/k^2$ of the observations of q will be within k standard deviations of $E(q)$; e.g., there is a 75 percent probability that the price will lie in the interval \$32.55-65.43.

Camp-Meidell Inequality

Some mild assumptions about the distribution of q lead to much tighter bounds [2]. If we assume that the distribution is unimodal, with the mode occurring at $E(q)$, then at least $1 - 1/2.25k^2$ of the observations will lie within k standard

deviations of the mean; e.g., there is an 89 percent probability that the price will lie in the interval \$32.55-65.43.

For more general treatments of Tchebycheff-type bounds, see [4] and [9].

4. LINEAR PROGRAMMING APPLICATIONS

The chance constrained programming model of Charnes and Cooper [3] presents interesting applications of these techniques. A deterministic linear program may be stated

$$\begin{aligned} \text{Max} \quad & c'x \\ \text{s.t.} \quad & Ax \leq b \end{aligned} \quad (7)$$

with A an $m \times n$ constant matrix of technological coefficients, c a vector of constant prices and b a vector of constant resource levels. If we change the sense of b to that of a vector of resource levels realized from a multivariate density function, $f(b)$, then a reformulation of (7) yields

$$\begin{aligned} \text{Max} \quad & c'x \\ \text{s.t.} \quad & \Pr[Ax \leq b] \geq \alpha \end{aligned} \quad (8)$$

where α is a set of probabilities of individually satisfying each of the m constraints. This problem may be solved by using the density function asserted for b to find a set of bounds, $\beta(\alpha, f(b))$, such that

$$\Pr[b \geq \beta] \geq \alpha \quad ,$$

and, if $f(b)$ is free of terms involving x , reformulating (8) to the "deterministic equivalent" form

$$\begin{aligned} \text{Max} \quad & c'x \\ \text{s.t.} \quad & Ax \leq \beta. \end{aligned} \quad (9)$$

Alternatively, let $\dot{\alpha}$ be the scalar joint probability that solutions of (9) will ultimately prove simultaneously feasible for all constraints. If the variates b are independent,

$$\dot{\alpha} = \prod_{j=1}^m \alpha_j, \text{ but if the terms in } b \text{ are correlated,}$$

extraction of $\dot{\alpha}$ from $f(b)$ may, or may not, be possible or practical depending on the mathematical form of $f(b)$, the dimensionality of b , and so forth.

If $f(b)$ is unknown or mathematically uncooperative we may use the moments of the vector b to specify β nonparametrically with Tchebycheff's inequalities. For instance

$$\Pr[b_j < \mu_j - k_j \sqrt{\sigma_{jj}} = \beta_j] = 1 - \alpha_j \leq \frac{1}{k_j^2}, \quad j = 1, 2, \dots, m$$

Also, a frequent and reasonable simplification is

$$k_1 = \dots = k_m = k.$$

Further, we can specify a bound for $\dot{\alpha}$ by using a multivariate Tchebycheff inequality.

$$\Pr[b_j < \mu_j - k_j \sqrt{\sigma_{jj}} = \beta_j; j=1, \dots, m] = 1 - \dot{\alpha} \leq \sum_{j=1}^m \frac{1}{k_j^2}; \quad (10)$$

or, for $k_1 = \dots = k_m = k$

$$\Pr[b < \mu - k((\text{diag } \Sigma))^{1/2} = \beta] = 1 - \dot{\alpha} \leq \frac{m}{k^2} . \quad (11)$$

Here, $(\text{diag } \Sigma)^{1/2}$ denotes a column vector with components $\sqrt{\sigma_{jj}}$.

Another Tchebycheff inequality presented by Olkin and Pratt [11] can be used to specify a bound for $\dot{\alpha}$. If we use Π to represent the correlation matrix of b so that

$$\pi_{jk} = \frac{\sigma_{jk}}{\sqrt{\sigma_{jj}\sigma_{kk}}} ,$$

and simplify the presentation by choosing the constants

$k_1 = \dots = k_m = k$, then

$$\begin{aligned} \Pr[b < \mu - k((\text{diag } \Sigma))^{1/2} = \beta] &= 1 - \dot{\alpha} \\ &\leq \frac{1}{k^2 m^2} \left[\sqrt{\rho} + \sqrt{(m^2 - \rho)(m - 1)} \right]^2 , \end{aligned} \quad (12)$$

with

$$\rho = e' \Pi e, \text{ where } e \text{ is a vector of 1's.}$$

5. A LINEAR PROGRAMMING EXAMPLE

To illustrate, using b from the example in section 3,

$$\rho = (1 \quad 1 \quad 1) \begin{pmatrix} 1.0 & 0.87 & 0.34 \\ 0.87 & 1.0 & 0.39 \\ 0.34 & 0.39 & 1.0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 6.20 ;$$

$$1 - \dot{\alpha} \leq \frac{2.62}{k^2} \text{ by use of (12) .}$$

We note that this is a better bound than the $3/k^2$ produced by (11).

For $k = 3$ we find

$$\hat{\alpha} \geq 0.71 ;$$

$$\beta = \mu - k(\text{diag } \Sigma)^{1/2} = \begin{pmatrix} 4.2 - 3(.84) \\ 6.8 - 3(1.10) \\ 4.8 - 3(.71) \end{pmatrix} = \begin{pmatrix} 1.68 \\ 3.50 \\ 2.67 \end{pmatrix} .$$

If (7) is further generalized to allow both b and c to be random vectors, possibly correlated, we can easily incorporate the expectation of the objection function in a deterministic equivalent.

Alternately, following the development of [3]

$$x = Db ,$$

with D an $n \times m$ matrix, we reformulate (7) to

$$\begin{aligned} \text{Max} \quad & E[c'x] \\ \text{s.t.} \quad & \Pr[Ax \leq b] \geq \alpha \\ & x = Db , \end{aligned} \tag{13}$$

which by substitution becomes

$$\begin{aligned} \text{Max} \quad & E[c'Db] \\ \text{s.t.} \quad & \Pr[ADb \leq b] \geq \alpha . \end{aligned}$$

Using the notation and results of section 2, we derive for the objective function:

$$E[c'Db] = \mu_c' D\mu_b + \text{Tr}\{D\Sigma_{bc}\} .$$

The constraint set may be restated

$$\begin{aligned} \Pr[ADb \leq b] &= \Pr[ADb - b \leq 0] \\ &= \Pr[(AD - I)b \leq 0] \geq \alpha . \end{aligned}$$

Thus, using the moments of b it follows [1, p. 24] that the vectors of expectations and variances are

$$\begin{aligned} E[(AD - I)b] &= (AD - I)\mu_b = \delta(D) ; \\ V[(AD - I)b] &= \text{diag} \{ (AD - I)\Sigma_{bb} (AD - I)' \} \\ &= \gamma(D) . \end{aligned}$$

The procedure, then, is as follows:

- (1) Specify an acceptable α_i , the probability of feasibility for each constraint.
 - (2) Use (10), (11), or (12) to generate the set of k 's and $\dot{\alpha}$.
- A set of slack variables, v , is introduced. Adopting the symbol '*' to represent congruent multiplication of two commensurate matrices [6, p. 549 ff], the deterministic equivalent programming model for (13) becomes

$$\begin{aligned} \text{Max} \quad & \mu_c' D \mu_b + \text{Tr}\{D \Sigma_{bc}\} \\ \text{s.t.} \quad & v - \delta(D) \leq 0 \\ & v * v + k * k * \{ \delta(D) * \delta(D) - \gamma(D) * \gamma(D) \} \leq 0 \\ & v \geq 0. \end{aligned} \tag{14}$$

Finally, a solution, D^* , to this problem can be used to make nonparametric interval estimates for the value of the optimal objective function using equation (3) or simplifications (5) or (6), much as in the numerical example in the preceding section.

This variance can be used as the objective function in another model proposed by [3]

$$\begin{aligned} \text{Min} \quad & V[c'x] \\ \text{s.t.} \quad & \Pr[Ax \leq b] \geq \alpha \\ & x = Db \end{aligned}$$

and its deterministic equivalent which follows the development of (14).

A "partitioned" stochastic programming model considers the problem

$$\begin{aligned} \text{Max} \quad & c'x + d'y \\ \text{s.t.} \quad & Ax + By \leq b; \end{aligned}$$

where y is a set of external, uncontrollable prior random variables, d a vector of random prices, and B a random technological coefficient matrix. By using the moments of y , d , B , and b , the problem may be restated

$$\begin{aligned} \text{Max} \quad & c'x + c_0 \\ \text{s.t.} \quad & Ax \leq b_0, \end{aligned}$$

with the moments of b_0 and c_0 specified exactly, and solved as previously shown.

6. CONCLUSION

Estimation of stochastic vector, or matrix, products may be virtually impossible by use of multivariate density functions and statistical transformations without severe and unrealistic simplifying assumptions about the random variates involved. Simulation can lead to approximations for a given problem, but at high computation cost and specificity for results.

However, an approach using multivariate moments, which are available from either empirical data or analytic densities, leads to exact statements about the mean and variance of the vector products and useful Tchebycheff-type probability estimates. These estimates are very helpful in generalizing the mathematical models of operations research.

- [1] Anderson, T. An Introduction to Multivariate Statistical Analysis. New York: John Wiley and Sons, 1958.
- [2] Camp, B. H. "A New Generalization of Tchebycheff's Statistical Inequality." Bulletin of the American Mathematical Society. Vol. 28 (1922), p. 427.
- [3] Charnes, A. and Cooper, W. "Deterministic Equivalents for Optimizing and Satisficing under Chance Constraints." Operations Research, Vol. 11, No. 1, pp. 18-39 (1963).
- [4] Godwin, H. J. "On Generalization of Tchebycheff's Inequality." Journal of the American Statistical Society, Vol. 50 (1955), p. 923.
- [5] Graves, G. and Whinston, A. "An Algorithm for the Quadratic Assignment Problem." Management Science, Vol. 17, No. 7, pp. 453-471 (1970).
- [6] Howard, R. Dynamic Probabilistic Systems, Vol. 1. New York: John Wiley and Sons, 1971.
- [7] Jagannathan, R. "Chance Constrained Programming with Joint Constraints." Operations Research, Vol. 22, No. 2, pp. 358-372 (1974).
- [8] King, E., Sampson, C., and Simms, L. "An Alternative to Monte Carlo Sampling in Statistical Models." Management Science, Vol. 21, No. 6 (February, 1975), p. 649.
- [9] Mallows, C. C. "Generalization's of Tchebycheff's Inequalities." Journal of the Royal Statistical Society, Series B, Vol. XVIII, No. 2, 1956, p. 139.

- [10] Miller, B. and Wagner, H. "Chance-Constrained Programming with Joint Constraints." Operations Research, Vol. 3, pp. 930-945.
- [11] Olkin, I. and Pratt, J. "A Multivariate Tchebycheff Inequality." Annals of Mathematical Statistics, Vol. 29, pp. 226-234 (1958).
- [12] Searle, S. R. Linear Models. New York: John Wiley and Sons, 1971.
- [13] Sengupta, J. Stochastic Programming Methods and Applications. New York: North-Holland Publishing Company, 1973.
- [14] Sengupta, J. "Chance-Constrained Linear Programming with Chi-Square Type Deviates." Management Science, Vol. 19, No. 3, pp. 337-349 (1972).
- [15] Tintner, G. "A Note on Stochastic Linear Programming." Econometrica, Vol. 38, No. 2, pp. 490-495 (1960).

APPENDIX

Derivation of Expressions for the Mean and Variance
of the Inner Product of Two Vectors

$$q = b'c ;$$

$$q = (\tilde{b} + \mu_b)'(\tilde{c} + \mu_c) ;$$

$$q = \tilde{b}'\tilde{c} + 2\tilde{b}'\mu_c + 2\mu_b'\tilde{c} + \mu_b'\mu_c .$$

Noting that $E(\tilde{b}) = E(\tilde{c}) = 0$;

$$E(q) = E(\tilde{b}'\tilde{c}) + \mu_b'\mu_c ;$$

$$E(q) = \mu_b'\mu_c + \text{Tr } \Sigma_{bc} ;$$

$$V(q) = E(q^2) - [E(q)]^2 ;$$

$$V(q) = E(b'c)^2 - (\mu_b'\mu_c)^2 - 2\mu_b'\mu_c \text{Tr } \Sigma_{bc} - (\text{Tr } \Sigma_{bc})^2 .$$

In terms of moments about the mean,

$$E(b'c)^2 = E[(\tilde{b} + \mu_b)'(\tilde{c} + \mu_c)]^2 ;$$

$$\begin{aligned} E(b'c)^2 = E[& \tilde{b}'\tilde{c}\tilde{c}'\tilde{b} + \mu_b'\tilde{c}\tilde{c}'\mu_b + \mu_c'\tilde{b}\tilde{b}'\mu_c + (\mu_b'\mu_c)^2 + 2\mu_b'\tilde{c}\tilde{c}'\tilde{b} \\ & + 2\mu_c'\tilde{b}\tilde{b}'\tilde{c} + 2\mu_b'\mu_c\tilde{b}'\tilde{c} + 2\mu_b'\tilde{c}\tilde{b}'\mu_c + 2\mu_c'\tilde{b}\mu_b'\mu_c \\ & + 2\mu_b'\tilde{c}\mu_b'\mu_c] . \end{aligned}$$

We note that

$$E(\tilde{c}\tilde{c}') = \Sigma_{cc} ;$$

$$E(\tilde{b}\tilde{b}') = \Sigma_{bb} ;$$

$$E(\tilde{b}'\tilde{c}) = \text{Tr } \Sigma_{bc} ;$$

$$E(\tilde{b}) = E(\tilde{c}) = 0 .$$

Hence,

$$\begin{aligned} E(b'c)^2 &= E(\tilde{b}'\tilde{c}\tilde{c}'\tilde{b}) + \mu_b'\Sigma_{cc}\mu_b + \mu_c'\Sigma_{bb}\mu_c + 2\mu_b'\Sigma_{cb}\mu_c + (\mu_b'\mu_c)^2 + 2\mu_b'E(\tilde{c}\tilde{c}'\tilde{b}) \\ &\quad + 2\mu_c'E(\tilde{b}\tilde{b}'\tilde{c}) + 2\mu_b'\mu_c \text{Tr } \Sigma_{bc} . \end{aligned}$$

We have

$$\mu'\Sigma^*\mu = \mu_b'\Sigma_{cc}\mu_b + \mu_c'\Sigma_{bb}\mu_c + 2\mu_b'\Sigma_{cb}\mu_c ;$$

$$\begin{aligned} V(q) &= E(\tilde{b}'\tilde{c}\tilde{c}'\tilde{b}) + \mu'\Sigma^*\mu + 2\mu_c'E(\tilde{b}\tilde{b}'\tilde{c}) + 2\mu_b'E(\tilde{c}\tilde{c}'\tilde{b}) \\ &\quad - (\text{Tr } \Sigma_{bc})^2 . \end{aligned}$$

INITIAL DISTRIBUTION LIST

	No. Copies
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	2
Dean of Research Code 023 Naval Postgraduate School Monterey, California 93940	1
Library (Code 0212) Naval Postgraduate School Monterey, California 93940	2
Library (Code 55) Naval Postgraduate School Monterey, California 93940	2