

AD-A041 444

AIR FORCE ACADEMY COLO  
THE PHOTON SCATTERING MATRIX ELEMENT FOR TRANSVERSE ELECTRIC OMODES (U)  
JUN 77 H O DOGLIANI  
USAF-TR-77-7-SUPPL

F/G 20/8

UNCLASSIFIED

NL

| of |

AD  
A041444



END

DATE  
FILMED

8 - 77

USAFA-TR-77-7  
JUNE 1977  
SUPPLEMENT

*12*  
*B.S.*

AD A 041 444

# THE PHOTON SCATTERING MATRIX ELEMENT FOR TRANSVERSE ELECTRIC OUTGOING PHOTONS

HARALD O. DOGLIANI  
DEPARTMENT OF PHYSICS ✓  
USAF ACADEMY, COLORADO 80840



*go*  
DDC  
PREPARED  
JUL 12 1977  
DISCLOSURE  
C

AD NO.  
DDC FILE COPY

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

DEAN OF THE FACULTY  
UNITED STATES AIR FORCE ACADEMY  
COLORADO 80840

**EDITORIAL REVIEW BY LT COLONEL A. G. ELSER**  
**DEPARTMENT OF ENGLISH AND FINE ARTS**  
**USAF ACADEMY, COLORADO 80840**



This research report is presented as a competent treatment of the subject, worthy of publication. The United States Air Force Academy vouches for the quality of the research, without necessarily endorsing the opinions and conclusions of the author.

This report has been cleared for open publication and/or public release by the appropriate Office of Information in accordance with AFR 190-17 and DODD 5230.9. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service.

This research report has been reviewed and is approved for publication.

ACCESSION NO.	
DATE	When Searched <input checked="" type="checkbox"/>
DDC	DDC Number <input type="checkbox"/>
ORIGINATOR	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODE	
Dist.	AVAIL. AND/OR SPECIAL
IA	

*Philip J. Erdle*

**PHILIP J. ERDLE, Colonel, USAF**  
**Vice Dean of the Faculty**

Additional copies of this document are available through the National Technical Information Service, United States Department of Commerce, 5285 Port Royal Road, Springfield, VA 22151.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 USAFA-TR-77-7-Suppl	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Photon Scattering Matrix Element for Transverse Electric Outgoing Photons	5. TYPE OF REPORT & PERIOD COVERED Supplement	
7. AUTHOR(s) Harald O. Dogliani	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics (DFP) USAF Academy, CO 80840	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 227P	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE 10 June 1977	
	13. NUMBER OF PAGES 23	
16. DISTRIBUTION STATEMENT (of this Report) This research report has been cleared for open publication and/or public release by the appropriate office of information in accordance with AFR 190-1 and DODD 5230.9. Distribution unlimited.	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Dirac Propagator, Second Order Perturbation Theory, Magnetic Multipole Scattering, K-shell scattering, Compton effect, Transverse Electric Modes, Vector Spherical Harmonics		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report is a supplement to <u>Second Order Radiative Effects Using Dirac Propagators and Vector Harmonic Photons</u> (Ph.D. dissertation by Harald O. Dogliani, University of Colorado, 1976). The matrix element for photons scattering from bound K-shell electrons is presented. It is assumed that only magnetic multipoles are important and, hence, only transverse electric mode vector harmonic, outgoing photons are assumed. Electric multipole moments are accounted for in the Ph.D. dissertation.		

011 550

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CONTENTS

	<u>Page</u>
Introduction . . . . .	1
Transverse Electric Matrix Elements . . . . .	3
Appendix . . . . .	5

## INTRODUCTION

This report is a supplement to Second Order Radiative Effects Using Dirac Propagators and Vector Harmonic Photons, henceforth referenced as SORE, by Harald O. Dogliani, Ph.D. dissertation, University of Colorado, 1976. In SORE the scattering of circularly polarized photons by K-shell atomic electrons is treated using Dirac propagators representing intermediate state electrons and using a vector harmonic expansion for the outgoing photon fields. Since electric dipole transitions are usually assumed to dominate such a scattering process, only electric multipoles were treated in SORE by considering transverse magnetic, TM, outgoing photons and by neglecting transverse electric, TE, modes.

The analytic representation of the differential scattering cross section for photons scattering off K-shell electrons with outgoing vector harmonic photons is given by

$$d\sigma = \frac{2RV}{ch} \sum_{\ell, m} |M|^2 dk. \quad 1$$

The differential scattering cross section,  $d\sigma$ , is a differential with respect to the final photon wave number  $k$ . The speed of light is  $c$  and  $V$  is the interaction volume with  $R$  being the radius of a spherical volume containing the outgoing photon of wave number  $k$  with angular momentum  $\ell$  and projection  $m$ . Planck's constant is  $h$ . The matrix element for the interaction is  $M$ .

As a supplement to SORE, the matrix element for TE mode photons is presented in this report. By combining the TE mode matrix element presented in this report with the TM mode contributions presented in SORE, one can more properly treat photon scattering events in those situations where higher order multipole transitions are significant. The relevant terms for including TE mode scattering are presented in the appendix of this report. In order to calculate the matrix element for K-shell photon scattering, Equations 2, 3 and 5 must be combined with Equations 18, 19, 21 and 22 in the appendix.

A complete discussion of the vector harmonic expansion in terms of TE and TM modes is contained in SORE. Also included in SORE is a discussion on how a typical TE or TM contribution to the matrix element is derived. Finally, it should be emphasized that scattering matrix elements were developed using Dirac propagators for the intermediate electron states (see SORE). The use of single-electron, Dirac propagators with the use of vector harmonic outgoing photons results in expansions in terms of discrete exchanges of angular momenta. As is indicated in SORE, this may be of significant utility in certain scattering problems where total cross sections are needed.

TRANSVERSE ELECTRIC MATRIX ELEMENTS

The TE matrix element, averaged over initial electron spins, is given by

$$|M|^2 = \frac{1}{2} \{ |M_1|^2 + |M_2|^2 \} \quad 2$$

where

$$M_1 = \sum_{m', \ell', p'} C(kk_0 \ell m) (I_a(1, k \ell m: TE) + I_e(1, k \ell m: TE)),$$

$$M_2 = \sum_{m', \ell', p'} C(kk_0 \ell m) (I_a(2, k \ell m: TE) + I_e(2, k \ell m: TE)),$$

$$C(kk_0 \ell m) = h e^3 c (-1)^m [(2\omega_k) / (\ell(\ell+1) R V \omega_{k_0})]^{1/2}. \quad 3$$

The following symbols are used:

$k_0$	incoming photon wave number
$k$	outgoing photon wave number
$\omega_{k_0}$	incoming photon angular frequency
$\omega_k$	outgoing photon angular frequency
$e$	charge of an electron
$\ell$	outgoing photon angular momentum quantum number
$m$	outgoing photon angular momentum projection quantum number
$m', \ell', p'$	intermediate state (propagator) quantum numbers (see SORE)
$\lambda$	incoming photon circular polarization $\lambda = 1, 2,$
$i$	$\sqrt{-1}$ . <span style="float: right;">4</span>

The  $I_a(\lambda, k\ell m: TE)$  and  $I_e(\lambda, k\ell m: TE)$  must be evaluated for the absorption first and emission first contributions to  $M_1$  and  $M_2$  for incoming photon polarization  $\lambda$ . They are evaluated as follows:

$$I_a(1, k\ell m: TE) = \sum_{i=1}^2 YATE(i) \delta_{m', 3/2} \delta_{m, -m'+3/2} WATEU(i) \\ + \sum_{i=3}^6 YATE(i) \delta_{m', 1/2} \delta_{m, -m'+1/2} WATED(i),$$

$$I_a(2, k\ell m: TE) = \sum_{i=1}^2 YATE(i) \delta_{m', -3/2} \delta_{m, -m'-3/2} WATED(i) \\ + \sum_{i=3}^6 YATE(i) \delta_{m', 1/2} \delta_{m, -m'+1/2} WATED(i),$$

$$I_e(1, k\ell m: TE) = \sum_{n=0}^{\infty} i^n (2(2n+1))^{1/2} \sum_{j=1}^4 YETE(i)$$

$$[\delta_{m', -m+1/2} \delta_{m, -m'+3/2} WETEU(i) \\ + \delta_{m', -m-1/2} \delta_{m, -m'+1/2} WETED(i)],$$

$$I_e(2, k\ell m: TE) = \sum_{n=0}^{\infty} i^n (2(2n+1))^{1/2} \sum_{j=1}^4 YETE(i)$$

$$[\delta_{m', -m+1/2} \delta_{m, -m''-1/2} WETEU(i+4) \\ + \delta_{m', -m-1/2} \delta_{m, -m''-3/2} WETED(i+4)].$$

5

The final electronic states are specified by the double primed quantum numbers  $\ell'', m''$  and  $p''$ . The integrals YETE and YATE, and the coefficients WETED, WETEU, WATED, and WATEU are provided in the appendix.

APPENDIX

Enumeration of the YETE'S, YATE'S, WETE'S and WATE'S

The functions required to evaluate the matrix elements describing photons scattering from K-shell electrons are presented in this section. Only the magnetic multipoles (TE modes) are accounted for in this scheme. These functions are inserted into Equation 5.

The following mnemonics are used in the definitions of the constants:

- 1) all constants beginning with the letter "Y" represent radial integrals;
- 2) all constants whose second letter is "A" are encountered in the "absorption first" matrix element;
- 3) similarly, all constants whose second letter is "E" are encountered in the "emission first" case;
- 4) all constants end with "TE" since this set considers TE modes only.

The  $j_n(k_0 r)$  are spherical Bessel functions of order  $n$ , whereas the lower case  $g(r)$  and  $f(r)$  are the radial wavefunctions for the K-shell electron initial states - See SORE Equations(3-54) and(3-73). The final electronic state radial wave functions are represented by  $G(r)$  and  $F(r)$  as introduced in Equation (3-58), SORE. The following functions are also used:

$$I(j,k:\ell,m:n,i) = \int Y_{jk}^*(\theta,\phi) Y_{\ell m}(\theta,\phi) Y_{ni}(\theta,\phi) d\Omega, \quad 6$$

Where the  $Y_{jk}(\theta,\phi)$  are the usual spherical harmonics.

The  $C(jk:p)$  are given by

$$C(jk:p) = C^{jp} C^{kp} / ch^p \quad 7$$

where the  $C^{jp}$  are

$$C^{1p} = [(\ell + 1/2 + pm)/(2\ell + 1)]^{1/2},$$

(6)

$$\begin{aligned}
C^{2p} &= -p[(\ell + 1/2 - pm)/(2\ell + 1)]^{1/2}, \\
C^{3p} &= -[(\ell + 1/2 + p - pm)/(2\ell + 2p + 1)]^{1/2}, \\
C^{4p} &= -p[(\ell + 1/2 + p + pm)/(2\ell + 2p + 1)]^{1/2},
\end{aligned}
\tag{8}$$

and  $y^p$  is related to the Wronskian of the radial propagators,  $S(r)$  and  $R(r)$ , where

$$y^p = S^{i*}(r) R^r(r) - S^r(r) R^{i*}(r). \tag{9}$$

The  $S^r(r)$ ,  $S^i(r)$ ,  $R^r(r)$  and  $R^i(r)$  are the regular and irregular  $S$  and  $R$  solutions, respectively, to the following coupled differential equation

$$\begin{aligned}
(mc^2 + V - E) S(r) - \hbar \left( \frac{d}{dr} + \frac{k}{r} \right) R(r) &= 0, \\
(-mc^2 + V - E) R(r) + \hbar \left( \frac{d}{dr} - \frac{k}{r} \right) S(r) &= 0.
\end{aligned}
\tag{10}$$

In Equation 10,  $k$  is a non zero, integer quantum number,  $E$  the propagator energy and  $V$  the potential. The radial propagators,  $RR(p'ri:rr')$ ,  $RS(\dots)$  and  $SS(\dots)$ , are given by

$$\begin{aligned}
RR(p'ri:rr') &= R^r(r) R^{i*}(r') / rr', \\
RS(p'ri:rr') &= R^r(r) S^{i*}(r') / rr', \\
SS(p'ri:rr) &= S^r(r) S^{i*}(r') / rr'.
\end{aligned}
\tag{11}$$

If the electron is initially in a spin up state represented by  $\psi_{\beta}(\bar{r}')$  then \*

$$\psi_{\beta_1}(\bar{r}') = \begin{bmatrix} C^{1+} Y_{00}(\theta', \phi') g(r') \\ 0 \\ C^{3+} Y_{10}(\theta', \phi') \text{ if } (r') \\ C^{4+} Y_{11}(\theta', \phi') \text{ if } (r) \end{bmatrix}
\tag{12}$$

\*Bethe, M.A. and Salpeter, E.E., Quantum Mechanics of One- and Two-Electron Atoms (Springer-Verlag, Berlin, 1957) p. 64.

For an arbitrary, spherically symmetric potential the radial wave functions,  $g(r)$  and  $f(r)$ , must usually be obtained by numerical means. They form solutions to the following coupled differential equations:\*

$$\begin{aligned} \frac{1}{\hbar c} (E - V(r) + mc^2) f - \left( \frac{dg}{dr} + (1+k) \frac{g}{r} \right) &= 0 \\ \frac{1}{\hbar c} (E - V(r) - mc^2) g + \left( \frac{df}{dr} + (1-k) \frac{f}{r} \right) &= 0 \end{aligned} \quad 13$$

The final electron state is similarly given by

$$\psi_{\gamma_1}(\vec{r}) = \frac{1}{r} \begin{bmatrix} C^{1p''} Y_{\ell'', m''-1/2}(\theta, \phi) G(r) \\ C^{2p''} Y_{\ell'', m''+1/2}(\theta, \phi) G(r) \\ C^{3p''} Y_{\ell'', m''-1/2}(\theta, \phi) iF(r) \\ C^{4p''} Y_{\ell'', m''+1/2}(\theta, \phi) iF(r) \end{bmatrix} \quad 14$$

Notice that Equation 14 has a slightly different  $r$  dependence as compared to Equation 12. The  $G(r)$  and  $F(r)$  functions satisfy the following differential equations:

$$\begin{aligned} \frac{dF}{dr} - k \frac{F}{r} &= \left[ \frac{mc}{\hbar} \left( 1 - \frac{E}{mc^2} \right) + \frac{V}{\hbar c} \right] G, \\ \frac{dG}{dr} + k \frac{G}{r} &= \left[ \frac{mc}{\hbar} \left( 1 + \frac{E}{mc^2} \right) - \frac{V}{\hbar c} \right] F \end{aligned} \quad 15$$

which can be obtained from Equation 13 by substituting  $G(r)/r$  for  $g(r)$  and  $F(r)/r$  for  $f(r)$ .\*

\*Bethe, *Ibid.*, p. 65

Similarly, for the spin down initial electron the wave function is given by\*

$$\begin{aligned}
 p &= 1 \\
 j &= 1/2, \\
 \ell &= 0, \\
 k &= 0, \\
 m &= -1/2.
 \end{aligned}
 \tag{16}$$

This then yields

$$\psi_{\beta_1} = \begin{bmatrix} 0 \\ C^{2+} Y_{00} g(r) \\ C^{3+} Y_{1-1} if(r) \\ C^{4+} Y_{10} if(r) \end{bmatrix}
 \tag{17}$$

YETE's

$$\begin{aligned}
 \text{YETE (1)} &= \int dr' r'^2 g(r') j_\ell(kr') \int dr r j_n(k_0 r) G^*(r) RR(p'ri:rr'), \\
 \text{YETE (2)} &= \int dr' r'^2 f(r') j_\ell(kr') \int dr r j_n(k_0 r) G^*(r) RS(p'ri:rr'), \\
 \text{YETE (3)} &= \int dr' r'^2 g(r') j_\ell(kr') \int dr r j_n(k_0 r) F^*(r) RS(p'ir:r'r), \\
 \text{YETE (4)} &= \int dr' r'^2 f(r') j_\ell(kr') \int dr r j_n(k_0 r) F^*(r) SS(p'ri:rr'),
 \end{aligned}
 \tag{18}$$

\*Bethe, Ibid., p. 64.

YATE's

$$\text{YATE (1)} = \int dr' r' f(r') j_{\ell'}(k_0 r') \int dr r j_{\ell}(kr) G^*(r) \text{RS}(p'ri:rr')$$

$$\text{YATE (2)} = \int dr' r' j_{\ell'}(k_0 r') f(r') \int dr r j_{\ell}(kr) F^*(r) \text{SS}(p'ri:rr'),$$

$$\text{YATE (3)} = \int dr' r'^2 j_{\ell'+p'}(k_0 r') g(r') \int dr r j_{\ell}(kr) G^*(r) \text{RR}(p'ri:rr'),$$

$$\text{YATE (4)} = \int dr' r'^2 j_{\ell'+p'}(k_0 r') g(r') \int dr r j_{\ell}(kr) F^*(r) \text{RS}(p'ir:r'r),$$

$$\text{YATE (5)} = \int dr' r'^2 j'_{\ell'}(k_0 r') f(r') \int dr r j_{\ell}(kr) G^*(r) \text{RS}(p'ri:rr'),$$

$$\text{YATE (6)} = \int dr' r'^2 j'_{\ell'}(k_0 r') f(r') \int dr r j_{\ell}(kr) F^*(r) \text{SS}(p'ri:rr'). \quad 19$$

In some of the above equations one encounters  $j'$  which should be interpreted as the first derivative of the spherical Bessel function with respect to its argument.

Define the following constants:

$$A = C^{4+} i^{\ell'+1} (3\ell'(\ell'+1)(2\ell'+1))^{\frac{1}{2}} k_0^{-1},$$

$$AB = C^{1+} i^{\ell'+p'} (4(\ell'+p')+2)^{\frac{1}{2}},$$

$$AC = C^{3+} i^{\ell'+1} (6(2\ell'+1))^{\frac{1}{2}},$$

$$B = - C^{2+} i^{\ell'+p'} (4(\ell'+p')+2)^{\frac{1}{2}},$$

$$BA = - C^{4+} i^{\ell'+1} (6(2\ell'+1))^{\frac{1}{2}},$$

$$BB = C^{3+} i^{\ell'-1} (3\ell'(\ell'+1)(2\ell'+1))^{\frac{1}{2}} k_0^{-1}.$$

WETE'S

$$\text{WETEU}(1) = C^{1p''} C^{1+} I (\ell'', m''-1/2: \ell, m''-1/2: n, 0) \delta_{\ell', \ell-p'}$$

$$((m''-3/2 C(43:p') + CC(\ell, m''-3/2) C(44:p')),$$

$$\text{WETEU}(2) = - C^{1p''} 3^{1/2} \{I(\ell'', m''-1/2: \ell+p'+1, m''-1/2: n, 0)$$

$$((2\ell+1)(2\ell+3))^{-1/2} \delta_{\ell', \ell+1} [(m''-3/2) C^{3+} C(41:p')$$

$$((\ell+m''-1/2)(\ell-m''+5/2))^{1/2} + C^{3+} CC(\ell, m''-3/2) C(42:p')$$

$$((\ell+m''+1/2)(\ell-m''+3/2))^{1/2} + C^{4+} 2^{-1/2} CC(\ell, -m''+3/2) C(41:p')$$

$$((\ell+m''-3/2)(\ell+m''-1/2)/2)^{1/2} C^{4+} 2^{-1/2} (m''-3/2) C(42:p')$$

$$((\ell+m''-3/2)(\ell+m''+1/2)/2)^{1/2} \}$$

$$+ I(\ell'', m''-1/2: \ell'+p'-1, m''-1/2: n, 0) ((2\ell+1)(2\ell-1))^{-1/2} \delta_{\ell', \ell-1}$$

$$[C^{3+} (m''-3/2) C(41:p') ((\ell+m''-3/2)(\ell-m''+3/2))^{1/2}$$

$$+ C^{3+} CC(\ell, -m) C(42:p') ((\ell+m''-1/2)(\ell-m''-5/2))^{1/2}$$

$$- C^{4+} CC(\ell, -m''+3/2) C(41:p') ((\ell-m''+5/2)(\ell-m''+3/2)/2)^{1/2}$$

$$+ C^{4+} (m''-3/2) C(42:p') ((\ell-m''+1/2)(\ell-m''+3/2)/2)^{1/2} \},$$

$$\text{WETEU}(3) = -C^{3p''} C^{1+} I (\ell''+p'', m''-1/2: \ell-p', m''-1/2: n, 0) \delta_{\ell', \ell-p'}$$

$$((m''-3/2 C(23:p') + CC(\ell, m''-3/2) C(24:p')),$$

$$\begin{aligned}
\text{WETE}(4) = & - C^{3p''} 3^{1/2} \{ I(\ell''+p'', m''-1/2:\ell+1, m''-1/2:n, 0) \delta_{\ell', \ell+1} \\
& ((2\ell+1)(2\ell+3))^{-1/2} [C^{3+}(m''-3/2)C(21:p') \\
& ((\ell+m''-1/2)(\ell-m''+5/2))^{1/2} + C^{3+} CC(\ell, m''-3/2) C(22:p') \\
& ((\ell+m''+1/2)(\ell-m''+3/2))^{1/2} + C^{4+} 2^{-1/2} CC(\ell-m''+3/2) C(21:p') \\
& ((\ell+m''-3/2)(\ell+m''-1/2)/2)^{1/2} - C^{4+} 2^{-1/2} (m''-3/2) C(22:p') \\
& ((\ell+m''-3/2)(\ell+m''+1/2)/2)^{1/2} ] \\
& + I(\ell''+p'', m''-1/2:\ell-1, m''-1/2:n, 0) \delta_{\ell', \ell-1} ((2\ell+1)(2\ell-1))^{-1/2} \\
& [C^{3+}(m''-3/2) C(21:p') ((\ell+m''-3/2)(\ell-m''+3/2))^{1/2} \\
& + C^{3+} CC(\ell, m''-3/2) C(22:p') ((\ell+m''-1/2)(\ell-m''+1/2))^{1/2} \\
& - C^{4+} 2^{-1/2} CC(\ell, -m''+3/2) C(21:p') ((\ell-m''+5/2)(\ell-m''+3/2)/2)^{1/2} \\
& + C^{4+} 2^{-1/2} (m''-3/2) C(22:p') ((\ell-m''+1/2)(\ell-m''+3/2)/2)^{1/2} ] \}, \\
\text{WETE}(5) = & C^{3p''} C^{1+} I(\ell'', m''+1/2:\ell, m''+1/2:n, 0) \delta_{\ell, \ell-p'} \\
& ((m''+1/2) C(33:p') + CC(\ell, m''+1/2) C(34:p')),
\end{aligned}$$

$$\begin{aligned}
\text{WETEU}(6) = & - C^{3p''} 3^{1/2} \{ I (\ell'', m''+1/2: \ell+p'+1, m''+1/2: n, 0) \delta_{\ell', \ell+1} \\
& ((2\ell+1)(2\ell+3))^{-1/2} [C^{3+}(m''+1/2) C(31:p') (\ell+m''+3/2) (\ell-m''+1/2)]^{1/2} \\
& + C^{3+} CC(\ell, m''+1/2) C(32:p') ((\ell+m''+5/2) (\ell-m''-1/2))^{1/2} \\
& + C^{4+} 2^{-1/2} CC(\ell, -m''-1/2) C(31:p') ((\ell+m''+1/2) (\ell+m''3/2)/2)^{1/2} \\
& - C^{4+} 2^{-1/2} (m''+1/2) C(32:p') ((\ell+m''+3/2) (\ell+m''+5/2)/2)^{1/2} ] \\
& + I (\ell'', m''+1/2: \ell+p'-1, m''+1/2: n, 0) \delta_{\ell', \ell-1} ((2\ell+1)(2\ell-1))^{-1/2} \\
& [ C^{3+}(m''+1/2) C(31:p') ((\ell+m''+1/2) (\ell-m''-1/2))^{1/2} \\
& + C^{3+} CC(\ell, m''+1/2) C(32:p') ((\ell+m''+3/2) (\ell-m''-3/2))^{1/2} \\
& - C^{4+} 2^{-1/2} CC(\ell, -m''-1/2) C(31:p') ((\ell-m''+1/2) (\ell-m''-1/2)/2)^{1/2} \\
& + C^{4+} 2^{-1/2} (m''+1/2) C(32:p') ((\ell-m''-1/2) (\ell-m''-3/2)/2)^{1/2} ] \},
\end{aligned}$$

$$\begin{aligned}
\text{WETEU}(7) = & C^{4p''} C^{1+} I (\ell''+p'', m''+1/2: \ell-p', m''+1/2: n, 0) \delta_{\ell', \ell-p'} \\
& ((m''+1/2) C(13:p') + CC(\ell, m''+1/2) C(14:p')),
\end{aligned}$$

$$\begin{aligned}
\text{WETEU}(8) = & C^{4p''} 3^{1/2} \{ I(\ell''+p'', m''+1/2: \ell+1, m''+1/2: n, 0) \delta_{\ell', \ell+1} \\
& ((2\ell+1)(2\ell+3))^{-1/2} [ C^{3+} (m''+1/2) C(11:p') ((\ell+m''+3/2)(\ell-m''+1/2))^{1/2} \\
& + C^{3+} CC(\ell, m''+1/2) C(12:p') ((\ell+m''+5/2)(\ell-m''-1/2))^{1/2} \\
& + C^{4+} 2^{-1/2} CC(\ell, -m''-1/2) C(11:p') ((\ell+m''+1/2)(\ell+m''+3/2)/2)^{1/2} \\
& - C^{4+} 2^{-1/2} (m''+1/2) C(12:p') ((\ell+m''+3/2)(\ell+m''+5/2)/2)^{1/2} ] \\
& + I(\ell''+p'', m''+1/2: \ell-1, m''+1/2: n, 0) \delta_{\ell', \ell-1} ((2\ell+1)(2\ell-1))^{-1/2} \\
& [ C^{3+} (m''+1/2) C(11:p') ((\ell+m''+1/2)(\ell-m''-1/2))^{1/2} \\
& + C^{3+} CC(\ell, m''+1/2) C(12:p') ((\ell+m''+3/2)(\ell-m''-3/2))^{1/2} \\
& - C^{4+} 2^{-1/2} CC(\ell, -m''-1/2) C(11:p') ((\ell-m''+1/2)(\ell-m''-1/2)/2)^{1/2} \\
& + C^{4+} 2^{-1/2} (m''+1/2) C(12:p') ((\ell-m''-1/2)(\ell-m''-3/2)/2)^{1/2} ] \},
\end{aligned}$$

$$\begin{aligned}
\text{WETED}(1) = & C^{1p''} C^{2+} I(\ell'', m''-1/2: \ell, m''-1/2: n, 0) \delta_{\ell', \ell-p'} \\
& (CC(\ell, -m''+1/2) C(43:p') - (m''-1/2) C(44:p')),
\end{aligned}$$

$$\text{WETED}(2) = C^{1p''} 3^{1/2} \{ I (\ell'', m''-1/2: \ell+p'+1, m''-1/2: n, 0) \quad .$$

$$((2\ell+1)2\ell-1)^{-1/2} \delta_{\ell', \ell+1} [C^{3+} (-m''+1/2) 2^{-1/2} C(41:p')$$

$$((\ell-m''+3/2)(\ell-m''+5/2))^{1/2} - C^{3+} 2^{-1/2} CC(\ell, m''-1/2) C(41:p')$$

$$((\ell-m''+1/2)(\ell-m''+3/2))^{1/2} - C^{4+} CC(\ell, -m''+1/2) C(41:p')$$

$$((\ell+m''-1/2)(\ell-m''+5/2))^{1/2} + C^{4+} (m''-1/2) C(42:p')$$

$$((\ell+m''+1/2)(\ell-m''+3/2))^{1/2} ]$$

$$+ I (\ell'', m''-1/2: \ell+p'-1, m''-1/2: n, 0) ((2\ell+1)(2\ell-1))^{-1/2} \delta_{\ell', \ell-1}$$

$$[ C^{3+} (m''-1/2) C(41:p') ((\ell+m''-1/2)(\ell+m''-3/2))^{1/2}$$

$$+ C^{3+} CC(\ell, m''-1/2) C(42:p') ((\ell+m''+1/2)(\ell+m''-1/2))^{1/2}$$

$$- C^{4+} 2^{1/2} CC(\ell, -m''+1/2) C(41:p') ((\ell+m''-3/2)(\ell-m''+3/2))^{1/2}$$

$$+ C^{4+} 2^{1/2} (m''-1/2) C(42:p') ((\ell+m''-1/2)(\ell-m''+1/2))^{1/2} ] \} ,$$

$$\text{WETED}(3) = - C^{3p''} C^{2+} I (\ell''+p'', m''-1/2: \ell-p', m''-1/2: n, 0) \delta_{\ell', \ell-p'}$$

$$(C(\ell, -m''+1/2) C(23:p') - (m''-1/2) C(24:p')),$$

$$\text{WETED}(4) = C^{3p''} 3^{1/2} \{ I (\ell''+p'', m''-1/2: \ell+1, m''-1/2: n, 0)$$

$$((2\ell+1)(2\ell+3))^{-1/2} \delta_{\ell', \ell+1} [C^{3+} 2^{-1/2} (m''-1/2) C(21:p')$$

$$((\ell-m''+3/2)(\ell-m''+5/2))^{1/2} + C^{3+} 2^{-1/2} CC(\ell, m''-1/2) C(22:p')$$

$$((\ell-m''+1/2)(\ell-m''+3/2))^{1/2} + C^{4+} CC(\ell, -m''+1/2) C(21:p')$$

$$((\ell+m''-1/2)(\ell-m''+5/2))^{1/2} - C^{4+} (m''-1/2) C(22:p')$$

$$((\ell+m''+1/2)(\ell-m''+3/2))^{1/2}]$$

$$+ I (\ell''+p, m''-1/2: \ell-1, m''-1/2: n, 0) ((2\ell+1)(2\ell-1))^{-1/2} \delta_{\ell', \ell-1}$$

$$[-C^{3+} 2^{-1/2} (m''-1/2) C(21:p') ((\ell+m''-1/2)(\ell+m''-3/2))^{1/2}$$

$$- C^{3+} 2^{-1/2} CC(\ell, m''-1/2) C(22:p') ((\ell+m''+1/2)(\ell+m''-1/2))^{1/2}$$

$$+ C^{4+} CC(\ell, -m''+1/2) C(21:p') ((\ell+m''-3/2)(\ell-m''+3/2))^{1/2}$$

$$- C^{4+} (m''-1/2) C(22:p') ((\ell+m''-1/2)(\ell-m''+1/2))^{1/2}] ,$$

$$\text{WETED}(5) = C^{2p''} C^{2+} I (\ell'', m''+1/2: \ell, m''+1/2: n, 0) \delta_{\ell', \ell-p'}$$

$$(CC(\ell, -m''-3/2) C(33:p') - (m''+3/2) C(34:p')),$$

$$\begin{aligned}
\text{WETED}(6) = & -C^{2p''} 3^{1/2} \{ I(\ell'', m''+1/2: \ell+p'+1, m''+1/2: n, o) \\
& ((2\ell+1)(2\ell+3))^{-1/2} \delta_{\ell', \ell+1} [C^{3+} 2^{-1/2} (m''+5/2) C(31:p') \\
& ((\ell-m''-1/2)(\ell-m''+1/2))^{1/2} + C^{3+} 2^{-1/2} CC(\ell, m''+3/2) C(32:p') \\
& ((\ell-m''-3/2)(\ell-m''-1/2))^{1/2} + C^{4+} CC(\ell, -m''-3/2) C(31:p') \\
& ((\ell+m''+3/2)(\ell-m''+1/2))^{1/2} - C^{4+} (m''+3/2) C(32:p') \\
& ((\ell+m''+5/2)(\ell-m''-1/2))^{1/2} ] \\
& + I(\ell'', m''+1/2: \ell+p'-1, m''+1/2: n, o) ((2\ell+1)(2\ell-1))^{-1/2} \delta_{\ell', \ell-1} \\
& [C^{3+} 2^{-1/2} (m''+3/2) C(31:p') ((\ell+m''+3/2)(\ell+m''+1/2))^{1/2} \\
& - C^{3+} 2^{-1/2} CC(\ell, m''+3/2) C(32:p') ((\ell+m''+5/2)(\ell+m''+3/2))^{1/2} \\
& + C^{4+} CC(\ell, -m''-3/2) C(31:p') ((\ell+m''+1/2)(\ell-m''-1/2))^{1/2} \\
& - C^{4+} (m''+3/2) C(32:p') ((\ell+m''+3/2)(\ell-m''-3/2))^{1/2} ] \}
\end{aligned}$$

$$\text{WETED}(7) = -C^{4p''} C^{2+} I(\ell''+p'', m''+1/2: \ell-p', m''+1/2: n, o) \delta_{\ell', \ell-p'}$$

$$(CC(\ell, -m''-3/2) C(13:p') - (m''+3/2) C(14:p')),$$

$$\begin{aligned}
\text{WETED}(8) = & C^{4p''} 3^{1/2} \{ I(\ell''+p'', m''+1/2: \ell+1, m''+1/2: n, o) \\
& ((2\ell+1) (2\ell+3))^{-1/2} \delta_{\ell', \ell+1} [C^{3+} 2^{-1/2} (m''+3/2) C(11:p')] \\
& ((\ell-m''-1/2) (\ell-m''+1/2))^{1/2} + C^{3+} 2^{-1/2} CC(\ell, m''+3/2) C(12:p') \\
& ((\ell-m''-3/2) (\ell-m''-1/2))^{1/2} + C^{4+} CC(\ell, -m''-3/2) C(11:p') \\
& ((\ell+m''+3/2) (\ell-m''+1/2))^{1/2} - C^{4+} (m''+3/2) C(12:p') \\
& ((\ell+m''+5/2) (\ell-m''-1/2))^{1/2} ] \\
& + I(\ell''+p'', m''+1/2: \ell-1, m''+1/2: n, o) ((2\ell+1) (2\ell-1))^{-1/2} \delta_{\ell', \ell-1} \\
& [ -C^{3+} 2^{-1/2} (m''+3/2) C(11:p') ((\ell+m''+3/2) (\ell+m''+1/2))^{1/2} \\
& - C^{3+} 2^{-1/2} CC(\ell, m''+3/2) C(12:p') ((\ell+m''+5/2) (\ell+m''+3/2))^{1/2} \\
& + C^{4+} CC(\ell, -m''-3/2) C(11:p') ((\ell+m''+1/2) (\ell-m''-1/2))^{1/2} \\
& - C^{4+} (m''+3/2) C(12:p') ((\ell+m''+3/2) (\ell-m''-3/2))^{1/2} ] \}.
\end{aligned}$$

21

WATE'S

$$\text{WATEU}(1) = A[C^{1p''} \{ C(31:p') (m''-3/2)$$

$$I(\ell'', m''-1/2: \ell, m''-3/2: \ell'+p', 1)$$

$$-C(41:p') ((\ell+m''-3/2) (\ell-m''+5/2))^{1/2}$$

$$I(\ell'', m''-1/2: \ell, m''-5/2: \ell'+p', 2) \}$$

$$-C^{2p''} \{ C(31:p') ((\ell-m''+3/2) (\ell+m''-1/2))^{1/2}$$

$$I(\ell'', m''+1/2: \ell, m''-1/2: \ell'+p', 1)$$

$$+C(41:p') (m''-3/2) I(p'', m''+1/2: \ell, m''-3/2: \ell'+p', 2) \}],$$

$$\text{WATEU}(2) = -A[C^{3p''} \{ C(11:p') (m''-3/2) I(\ell''+p'', m''-1/2: \ell, m''-3/2: \ell', 1)$$

$$-C(21:p') ((\ell+m''-3/2) (\ell-m''+5/2))^{1/2}$$

$$I(\ell''+p'', m''-1/2: \ell, m''-5/2: \ell', 2) \}$$

$$-C^{4p''} \{ C(11:p') ((\ell-m''+3/2) (\ell+m''-1/2))^{1/2}$$

$$I(\ell''+p'', m''+1/2: \ell, m''-1/2: \ell', 1)$$

$$+ C(21:p') (m''-3/2) I(\ell''+p'', m''+1/2: \ell, m''-3/2: \ell', 2) \}],$$

$$\text{WATEU}(3) = \text{AB}[C^{1P''} \{ C(34:p') (m''+1/2)$$

$$I(\ell'', m''-1/2: \ell, m''+1/2: \ell'+p', -1)$$

$$- C(44:p') ((\ell+m''+1/2) (\ell-m''+1/2))^{1/2}$$

$$I(\ell'', m''-1/2: \ell, m''-1/2: \ell'+p', 0) \}$$

$$- C^{2P''} \{ C(34:p') ((\ell-m''-1/2) (\ell+m''+3/2))^{1/2}$$

$$I(\ell'', m''+1/2: \ell, m''+3/2: \ell'+p', -1)$$

$$+ C(44:p') (m''+1/2) I(\ell'', m''+1/2: \ell, m''+1/2: \ell'+p', 0) \}],$$

$$\text{WATEU}(4) = \text{AB}[C^{3P''} \{ C(24:p') ((\ell+m''+1/2) (\ell-m''+1/2))^{1/2}$$

$$I(\ell''+p'', m''-1/2: \ell, m''-1/2: \ell', 0)$$

$$- C(14:p') (m''+1/2) I(\ell''+p'', m''-1/2: \ell, m''+1/2: \ell', -1) \}$$

$$+ C^{4P''} \{ C(24:p') (m''+1/2) I(\ell''+p'', m''+1/2: \ell, m''+1/2: \ell', 0)$$

$$+ C(14:p') ((\ell-m''-1/2) (\ell+m''+3/2))^{1/2}$$

$$I(\ell''+p'', m''+1/2: \ell, m''+3/2: \ell', -1) \}],$$

$$\begin{aligned}
\text{WATEU}(5) = & AC[C^{1p''} \{ C(42:p') ((\ell+m''+1/2) (\ell-m''+1/2))^{1/2} \\
& I(\ell'', m''-1/2: \ell, m''-1/2: \ell'+p', 0) \\
& - C(32:p') (m''+1/2) I(\ell'', m''-1/2: \ell, m''+1/2: \ell'+p', -1) \} \\
& + C^{2p''} \{ C(32:p') ((\ell-m''-1/2) (\ell+m''+3/2)) \\
& I(\ell'', m''+1/2: \ell, m''+3/2: \ell'+p', -1) \\
& + C(42:p') (m''+1/2) I(\ell'', m''+1/2: \ell, m''+1/2: \ell'+p', 0) \}],
\end{aligned}$$

$$\begin{aligned}
\text{WATEU}(6) = & AC[C^{3p''} \{ C(12:p') (m''+1/2) I(\ell''+p'', m''-1/2: \ell, m''+1/2: \ell', -1) \\
& - C(22:p') ((\ell+m''+1/2) (\ell-m''+1/2))^{1/2} \\
& I(\ell''+p'', m''-1/2: \ell, m''-1/2: \ell', 0) \} \\
& - C^{4p''} \{ C(12:p') ((\ell-m''-1/2) (\ell+m''+3/2))^{1/2} \\
& I(\ell''+p'', m''+1/2: \ell, m''+3/2: \ell', -1) \\
& + C(22:p') (m''+1/2) I(\ell''+p'', m''+1/2: \ell, m''+1/2: \ell', 0) \}],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(1) = & BB[C^{1p''} (- (m''+3/2) C(32:p') I(\ell'', m''-1/2: \ell, m''+3/2: \ell'+p', -2) \\
& + ((\ell+m''+3/2) (\ell-m''-1/2))^{1/2} C(42:p') \\
& I(\ell'', m''-1/2: \ell, m''+1/2: \ell, m''+1/2: \ell'+p', -1)) \\
& + C^{2p''} (((\ell-m''-3/2) (\ell+m''+5/2))^{1/2} C(32:p') \\
& I(\ell'', m''+1/2: \ell, m''+1/2: \ell, m''+5/2: \ell'+p', -2) \\
& + (m''+3/2) C(42:p') I(\ell'', m''+1/2: \ell, m''+3/2: \ell'+p', -1))],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(2) = & -BB[C^{3p''} (-(m''+3/2) C(12:p') I(\ell''+p'', m''-1/2:\ell', m''+3/2:\ell', -2) \\
& + ((\ell+m''+3/2) (\ell-m''-1/2))^{1/2} C(22:p') \\
& I(\ell''+p'', m''-1/2:\ell, m''+1/2:\ell', -1)) \\
& + C^{4p''} (((\ell-m''-3/2) (\ell+m''+5/2))^{1/2} C(12:p') \\
& I(\ell''+p'', m''+1/2:\ell, m''+5/2:\ell', -2) \\
& + (m''+3/2) C(22:p') I(\ell''+p'', m''+1/2:\ell, m''+3/2:\ell', -1)]],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(3) = & B[C^{1p''} ((-m''+1/2) C(33:p') I(\ell'', m''-1/2:\ell, m''-1/2:\ell'+p', 0) \\
& + ((\ell+m''-1/2) (\ell-m''+3/2))^{1/2} C(43:p') \\
& I(\ell'', m''-1/2:\ell, m''-3/2:\ell'+p', 1)) \\
& + C^{2p''} (((\ell-m''+1/2) (\ell+m''+1/2))^{1/2} C(33:p') \\
& I(\ell'', m''+1/2:\ell, m''+1/2:\ell'+p', 0) \\
& + (m''-1/2) C(43:p') I(\ell'', m''+1/2:\ell, m''-1/2:\ell'+p', 1)]],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(4) = & -B[C^{3p''} ((-m''+1/2) C(13:p') I(\ell''+p'', m''-1/2:\ell, m''-1/2:\ell', 0) \\
& + ((\ell+m''-1/2) (\ell-m''+3/2))^{1/2} C(23:p') \\
& I(\ell''+p'', m''-1/2:\ell, m''-3/2:\ell', 1)) \\
& + C^{4p''} (((\ell-m''+1/2) (\ell+m''+1/2))^{1/2} C(13:p') \\
& I(\ell''+p'', m''+1/2:\ell, m''+1/2:\ell', 0) \\
& + (m''-1/2) C(23:p') I(\ell''+p'', m''+1/2:\ell, m''-1/2:\ell', 1)]],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(5) = & \text{BA} [C^{1p''} ((-m''+1/2) C(31:p')) \\
& I(\ell'', m''-1/2: \ell, m''-1/2: \ell, m''-1/2: \ell'+p', 0) \\
& + ((\ell+m''-1/2) (\ell-m''+3/2))^{1/2} C(41:p') \\
& I(\ell'', m''-1/2: \ell, m''-3/2: \ell'+p', 1)) \\
& + C^{2p''} (((\ell-m''+1/2) (\ell+m''+1/2))^{1/2} C(31:p')) \\
& I(\ell'', m''+1/2: \ell, m''+1/2: \ell'+p', 0) \\
& + (m''-1/2) C(41:p') I(\ell'', m''+1/2: \ell, m''-1/2: \ell'+p', 1)]],
\end{aligned}$$

$$\begin{aligned}
\text{WATED}(6) = & -\text{BA} [C^{3p''} (-(m''-1/2) C(11:p')) I(\ell''+p'', m''-1/2: \ell, m''-1/2: \ell', 0), \\
& + ((\ell+m''-1/2) (\ell-m''+3/2))^{1/2} C(21:p') \\
& I(\ell''+p'', m''-1/2: \ell, m''-3/2: \ell', 1)) \\
& + C^{4p''} (((\ell-m''+1/2) (\ell+m''+1/2))^{1/2} C(11:p')) \\
& I(\ell+p'', m''+1/2: \ell, m''+1/2: \ell', 0) \\
& + (m''-1/2) C(21:p') I(\ell''+p'', m''+1/2: \ell, m''-1/2: \ell', 1)]].
\end{aligned}$$