

AD-A041 545

NAVAL RESEARCH LAB WASHINGTON D C  
REVISED AND ENLARGED COLLECTION OF PLASMA PHYSICS FORMULAS AND --ETC(U)  
MAY 77 D L BOOK  
NRL-MR-3332

F/G 20/5

UNCLASSIFIED

NL

1 of 2

ADA041 545



12

NRL Memorandum Report 3332

# Revised and Enlarged Collection of Plasma Physics Formulas and Data

DAVID L. BOOK

*Plasma Dynamics Branch  
Plasma Physics Division*

May 1977

AD A 041 545

AD No. \_\_\_\_\_  
DDC FILE COPY

DDC  
JUL 14 1977  
B



NAVAL RESEARCH LABORATORY  
Washington, D.C.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 3332	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Revised and Enlarged Collection of Plasma Physics Formulas and Data	5. TYPE OF REPORT & PERIOD COVERED Final report	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) D. L. Book	8. CONTRACT OR GRANT NUMBER(s) N000147TWRT0008	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D.C. 20375	10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem A03 - 16	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, Virginia 22217	12. REPORT DATE May 1977	13. NUMBER OF PAGES 106
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12-103p.	15. SECURITY CLASS. (of this report) UNCLASSIFIED	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 14 NRL-MR-3332		ADDITIONAL FOR NTR <input checked="" type="checkbox"/> White Section DDC <input type="checkbox"/> Ball Section UNANNOUNCED <input type="checkbox"/> JUSTIFICATION BY DISTRIBUTION/AVAILABILITY CODES Dist. Avail. and/or Spec.
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Plasma Physics Formulas Reference Data		A
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report is intended to serve as a kind of users' manual for the NRL Plasma Formulary, supplying explanations worked examples and references, for which space in the pocket version was insufficient. A certain amount of supplementary material has been included.		

DD FORM 1473 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
S. N. 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251950

JP

CONTENTS

Preface ..... 1

Errata ..... 3

Numerical and Algebraic ..... 4

Vector Calculus ..... 7

Dimensions and Units ..... 26

Physical Constants ..... 31

Metric Prefixes ..... 35

Electricity and Magnetism ..... 36

Maxwell's Equations ..... 37

Electromagnetic Bands ..... 38

AC Circuits ..... 40

Dimensionless Numbers of Fluid Mechanics ..... 42

Shocks ..... 44

Plasma Dispersion Function ..... 45

Fundamental Plasma Parameters ..... 47

Approximate Magnitudes in Typical Plasmas ..... 51

Collisions and Transport ..... 53

Ionospheric Parameters ..... 74

Controlled Thermonuclear Reactions (CTR) ..... 75

Relativistic Electron Beams ..... 79

Electrostatic Streaming Instabilities ..... 82

Lasers ..... 85

Atomic Physics and Radiation ..... 88

REVISED AND ENLARGED COLLECTION OF PLASMA  
PHYSICS FORMULAS AND DATA

PREFACE

The first edition of the NRL Plasma Formulary was compiled and published in 1974. By 1976 the original printing of 5000 copies had been exhausted. The time seemed ripe to prepare a revised edition incorporating corrections of errors reported by users. Extensive new material was also added (some of it already contained in the companion manual to the first edition, NRL Memorandum Report Number 2898), and a number of changes made in the interests of greater clarity and ease of use. It is hoped that the result represents a net improvement over its predecessor, and that the iterations will soon converge to a definitive version.

Two forms of the revised Formulary were prepared, a folding card like the 1974 edition, and a small booklet. With minor exceptions, they contain the same material. Their content is reproduced in the present memorandum report, which therefore serves as a less compact (but possibly more easily read) third version of the revised Formulary. It includes a list of errata for the folding card and booklet, complete as of the date of publication of the report. Where a formula in the pocket versions was incorrect, the corrected formula appears in this report marked with an asterisk. (Since the report itself was typed independently, however, it may contain errors not present in the pocket versions.)

In addition, the present report is intended to serve as a kind of users' manual for the Formulary, supplying explanations, worked examples, and references, for which space in the pocket versions was insufficient.

A certain amount of supplementary material has been included, particularly in the Vector Calculus section. The elaborate reworking of the Atomic and Molecular Physics section included in the previous edition of this report has been abridged, however. Interested readers seeking background material on the formulas of this section are referred to the version originally prepared by A. W. Ali for Memorandum Report No. 2898 or to a forthcoming report, "Summary of Plasma Spectrographic Data," by J. Davis et. al., as well as to the numerous ponderous monographs on the subject.

Once again, many of my colleagues in NRL's Plasma Physics Division helped with the Formulary by suggesting changes and corrections and by proof reading. In addition to those persons previously acknowledged, particular thanks for specific contributions are due to Dr. Verne Jacobs of Science Applications, Inc. (Atomic Physics and Radiation), and Drs. Niels Winsor and Stephen Bodner of NRL (Collisions and Transport and Lasers, respectively). My special thanks go to Ms. Jean Monnette for managing to compose and lay out the Formulary while I was making continual changes in it.

ERRATA

March 1, 1977

(Foldout)	(Booklet)	Erratum and Correction
<u>Panel</u>	<u>Page</u>	
2	4	Eq. (11): replace $\underline{B_x} \nabla \underline{x_A}$ by $\underline{B_x} (\nabla \underline{x_A})$
2	5	Eq. (17-19), (27): See p. 8
3	7	Under "Components of $(\underline{A} \cdot \nabla) \underline{B}$ ": replace $\frac{\partial B_z}{\partial z}$ by $\frac{\partial B_z}{\partial z}$
4	10	Under "Rationalized mks" replace "mho" by "siemens"
5	11	Under "Rationalized mks" replace "weber/m <sup>2</sup> " by "tesla"
5	12	Under "Rationalized mks" replace "newton/m <sup>2</sup> " by "pascal"
8	34	Ion gyrofrequency, F region: replace 250 by 300
9	21	Def. of L: "horixontal" should be "horizontal"
14	33	In <u>Weakly Ionized Plasmas</u> , Eq. for collision frequency $\nu_\alpha$ : $T_\alpha$ should be multiplied by k.
16	36	Eq. (5c): interchange 11.9 and 2.4; last Eq.: replace $A_1 \sqrt{E}$ by $A_1 / \sqrt{E}$
19	44	Eq. (15): should read $\alpha_s = 8.75 \times 10^{-27} T_e^{-+ .5}$ cm <sup>2</sup> /sec

NUMERICAL AND ALGEBRAIC

Euler-Mascheroni constant<sup>[1]</sup>  $\gamma = 0.57722$

Gamma Function  $\Gamma(x+1) = x \Gamma(x)$

$$\Gamma\left(\frac{1}{5}\right) = 5.5663$$

$$\Gamma\left(\frac{1}{5}\right) = 4.5908$$

$$\Gamma\left(\frac{1}{4}\right) = 3.6256$$

$$\Gamma\left(\frac{1}{3}\right) = 2.6789$$

$$\Gamma\left(\frac{2}{5}\right) = 2.2182$$

$$\Gamma\left(\frac{1}{2}\right) = 1.7725 = \pi^{\frac{1}{2}}$$

$$\Gamma\left(\frac{3}{5}\right) = 1.4892$$

$$\Gamma\left(\frac{2}{3}\right) = 1.3541$$

$$\Gamma\left(\frac{3}{4}\right) = 1.2254$$

$$\Gamma\left(\frac{4}{5}\right) = 1.1542$$

$$\Gamma\left(\frac{5}{3}\right) = 1.1288$$

$$\Gamma(1) = 1.0$$

Binomial Theorem (good for  $|x| < 1$  or  $\alpha =$  positive integer):

$$(1+x)^\alpha = \sum_{k=0}^{\infty} \binom{\alpha}{k} x^k \cong 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!} x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} x^3 + \dots$$

Here

$$\binom{\alpha}{k} = \frac{\Gamma(\alpha+1)}{k! \Gamma(\alpha+1-k)} = \frac{\alpha(\alpha-1)\dots(\alpha-k+1)}{k!}$$

is the binomial coefficient.

A very general relation involving the binomial coefficients is the  
 Rothe-Hagen identity <sup>[2]</sup> (good for all complex  $x, y, z$  except when singular):

$$\sum_{k=0}^n \frac{x}{x+kz} \binom{x+kz}{k} \frac{y}{y+(n-k)z} \binom{y+(n-k)z}{n-k} = \frac{x+y}{x+y+nz} \binom{x+y+nz}{n}.$$

Many formulas involving binomial coefficients are special cases of the  
 Rothe-Hagen identity. For example, with  $z = 0$ ,

$$\sum_{k=0}^n \binom{x}{k} \binom{y}{n-k} = \binom{x+y}{n};$$

with  $x = y = n$ ,  $z = 1$ ,

$$\sum_{k=0}^n \frac{(n+k-1)!}{k!} \frac{(2n-k-1)!}{(n-k)!} = \frac{2}{3} \frac{n!(3n)!}{n^2(2n)!};$$

with  $x = -y = \frac{1}{2}$ ,  $z = \frac{1}{n}$ ,  $n$  odd,

$$\sum_{k=0}^n \frac{1}{n+2k} \frac{1}{n-2k} \binom{\frac{n+2k}{2n}}{k} \binom{\frac{n-2k}{2n}}{n-k} = 0,$$

etc.

Gain in decibels of  $P_2$  relative to  $P_1$

$$G = 10 \log_{10} (P_2 / P_1) .$$

To within two percent

$$(2\pi)^{1/2} \approx 2.5; \quad e^3 \approx 20; \quad \pi^2 \approx 10; \quad 2^{10} \approx 10^3 .$$

VECTOR CALCULUS

Notation:  $f, g, \text{ etc.}$ , are scalars;  $\underline{A}, \underline{B}, \text{ etc.}$  are vectors;  $\underline{T}$  is a tensor

- (1)  $\underline{A} \cdot \underline{B} \times \underline{C} = \underline{A} \times \underline{B} \cdot \underline{C} = \underline{B} \cdot \underline{C} \times \underline{A} = \underline{B} \times \underline{C} \cdot \underline{A} = \underline{C} \cdot \underline{A} \times \underline{B} = \underline{C} \times \underline{A} \cdot \underline{B}$
- (2)  $\underline{A} \times (\underline{B} \times \underline{C}) = (\underline{A} \cdot \underline{C}) \underline{B} - (\underline{A} \cdot \underline{B}) \underline{C}$
- (3)  $\underline{A} \times (\underline{B} \times \underline{C}) + \underline{B} \times (\underline{C} \times \underline{A}) + \underline{C} \times (\underline{A} \times \underline{B}) = 0$
- (4)  $(\underline{A} \times \underline{B}) \cdot (\underline{C} \times \underline{D}) = (\underline{A} \cdot \underline{C})(\underline{B} \cdot \underline{D}) - (\underline{A} \cdot \underline{D})(\underline{B} \cdot \underline{C})$
- (5)  $(\underline{A} \times \underline{B}) \times (\underline{C} \times \underline{D}) = (\underline{A} \times \underline{B} \cdot \underline{D}) \underline{C} - (\underline{A} \times \underline{B} \cdot \underline{C}) \underline{D}$
- (6)  $\underline{\nabla} \cdot (fg) = \underline{\nabla} \cdot (gf) = f \underline{\nabla} \cdot g + g \underline{\nabla} \cdot f$
- (7)  $\underline{\nabla} \cdot (f\underline{A}) = f \underline{\nabla} \cdot \underline{A} + \underline{A} \cdot \underline{\nabla} f$
- (8)  $\underline{\nabla} \times (f\underline{A}) = f \underline{\nabla} \times \underline{A} + \underline{\nabla} f \times \underline{A}$
- (9)  $\underline{\nabla} \cdot (\underline{A} \times \underline{B}) = \underline{B} \cdot \underline{\nabla} \times \underline{A} - \underline{A} \cdot \underline{\nabla} \times \underline{B}$
- (10)  $\underline{\nabla} \times (\underline{A} \times \underline{B}) = \underline{A} (\underline{\nabla} \cdot \underline{B}) - \underline{B} (\underline{\nabla} \cdot \underline{A}) + (\underline{B} \cdot \underline{\nabla}) \underline{A} - (\underline{A} \cdot \underline{\nabla}) \underline{B}$
- \* (11)  $\underline{\nabla} (\underline{A} \cdot \underline{B}) = \underline{A} \times (\underline{\nabla} \times \underline{B}) + \underline{B} \times (\underline{\nabla} \times \underline{A}) + (\underline{B} \cdot \underline{\nabla}) \underline{A} + (\underline{A} \cdot \underline{\nabla}) \underline{B}$
- (12)  $\underline{\nabla}^2 f = \underline{\nabla} \cdot \underline{\nabla} f$
- (13)  $\underline{\nabla}^2 \underline{A} = \underline{\nabla} (\underline{\nabla} \cdot \underline{A}) - \underline{\nabla} \times \underline{\nabla} \times \underline{A}$
- (14)  $\underline{\nabla} \times \underline{\nabla} f = 0$
- (15)  $\underline{\nabla} \cdot \underline{\nabla} \times \underline{A} = 0$

If  $\underline{e}_1, \underline{e}_2, \underline{e}_3$  are orthonormal unit vectors, a second-order tensor  $\underline{T}$  can be written in the dyadic form

$$(16) \quad \underline{T} = \sum_{i,j} T_{ij} \underline{e}_i \underline{e}_j$$

In cartesian coordinates the divergence of a tensor is a vector with components given by

$$(17) \quad (\underline{\nabla} \cdot \underline{T})_i = \sum_j (\partial T_{ji} / \partial x_j)$$

In general

$$(18) \quad \underline{\nabla} \cdot (\underline{A}\underline{B}) = (\underline{\nabla} \cdot \underline{A})\underline{B} + (\underline{A} \cdot \underline{\nabla})\underline{B}$$

$$(19) \quad \underline{\nabla} \cdot (f\underline{T}) = \underline{\nabla} f \cdot \underline{T} + f \underline{\nabla} \cdot \underline{T}$$

\*Instead of (17), an alternative definition [4] was employed in the pocket versions of the Formulary:

$$(17') \quad (\nabla \cdot \underline{T})_i = \sum_j (\partial T_{ij} / \partial x_j)$$

Since we write the operator  $\nabla$  to the left, this choice, involving contraction with respect to the second index instead of the first, appears less "natural." When it is adopted, (18) and (19)

become

$$(18') \quad \nabla \cdot (\underline{AB}) = \underline{A} \nabla \cdot \underline{B} + \underline{B} \cdot \nabla \underline{A}$$

$$(19') \quad \nabla \cdot (f\underline{T}) = \underline{T} \cdot \nabla f + f \nabla \cdot \underline{T}$$

But Gauss' theorem for tensors, formula (27), becomes

$$(27') \quad \int_V \nabla \cdot \underline{T} \, dV = \int_S \underline{T} \cdot d\underline{S} \, ,$$

which to the unsuspecting eye is equally "natural" looking. For symmetric tensors, the primed and unprimed formulas are identical. In plasma physics, fortunately, most of the tensors we deal with are symmetric. Nevertheless, the choice (17') is a potential source of confusion and its use in the other editions is regrettable. Here I have abandoned it. For uniformity, all of the integral formulas (25-34) have been written with differentials at the left, although this results in a significant change only in (27). The tensor divergence  $\nabla \cdot \underline{T}$  in cylindrical and spherical coordinates (p. 12 and p. 15, respectively) now appears with indices transposed.

Let  $\underline{r} = ix + jy + kz$  be the radius vector of magnitude  $r$ , from the origin to the point  $x, y, z$ . Then

$$(20) \quad \underline{\nabla} \cdot \underline{r} = 3$$

$$(21) \quad \underline{\nabla} \times \underline{r} = 0$$

$$(22) \quad \underline{\nabla} r = \underline{r}/r$$

$$(23) \quad \underline{\nabla} (1/r) = -\underline{r}/r^3$$

$$(24) \quad \underline{\nabla} \cdot (\underline{r}/r^3) = 4\pi\delta(\underline{r})$$

If  $V$  is a volume enclosed by a surface  $S$  and  $d\underline{S} = \underline{n} dS$  where  $\underline{n}$  is the unit normal outward from  $V$ ,

$$(25) \quad \int_V dV \underline{\nabla} f = \int_S d\underline{S} f$$

$$(26) \quad \int_V dV \underline{\nabla} \cdot \underline{A} = \int_S d\underline{S} \cdot \underline{A}$$

$$(27) \quad \int_V dV \underline{\nabla} \cdot \underline{T} = \int_S d\underline{S} \cdot \underline{T}$$

$$(28) \quad \int_V dV \underline{\nabla} \times \underline{A} = \int_S d\underline{S} \times \underline{A}$$

$$(29) \quad \int_V dV (f \underline{\nabla}^2 g - g \underline{\nabla}^2 f) = \int_S d\underline{S} \cdot (f \underline{\nabla} g - g \underline{\nabla} f)$$

$$(30) \quad \int_V dV (\underline{A} \cdot \underline{\nabla} \times \underline{\nabla} \times \underline{B} - \underline{B} \cdot \underline{\nabla} \times \underline{\nabla} \times \underline{A}) = \int_S d\underline{S} \cdot (\underline{B} \times \underline{\nabla} \times \underline{A} - \underline{A} \times \underline{\nabla} \times \underline{B})$$

If  $S$  is an open surface bounded by the contour  $C$  of which the line element is  $d\underline{l}$ ,

$$(31) \int_S \underline{dS} \times \underline{\nabla} f = \oint_C \underline{dl} f$$

$$(32) \int_S \underline{dS} \cdot \underline{\nabla} \times \underline{A} = \oint_C \underline{dl} \cdot \underline{A}$$

$$(33) \int_S (\underline{dS} \times \underline{\nabla}) \times \underline{A} = \oint_C \underline{dl} \times \underline{A}$$

$$(34) \int_S \underline{dS} \cdot (\underline{\nabla} f \times \underline{\nabla} g) = \oint_C f dg = - \oint_C g df$$

### Cylindrical Coordinates

Divergence

$$\nabla \cdot \underline{A} = \frac{1}{r} \frac{\partial}{\partial r} (rA_r) + \frac{1}{r} \frac{\partial A_\theta}{\partial \theta} + \frac{\partial A_z}{\partial z}$$

Gradient

$$(\nabla f)_r = \frac{\partial f}{\partial r}; \quad (\nabla f)_\theta = \frac{1}{r} \frac{\partial f}{\partial \theta}; \quad (\nabla f)_z = \frac{\partial f}{\partial z};$$

Curl

$$(\nabla \times \underline{A})_r = \frac{1}{r} \frac{\partial A_z}{\partial \theta} - \frac{\partial A_\theta}{\partial z}$$

$$(\nabla \times \underline{A})_\theta = \frac{\partial A_r}{\partial z} - \frac{\partial A_z}{\partial r}$$

$$(\nabla \times \underline{A})_z = \frac{1}{r} \frac{\partial}{\partial r} (rA_\theta) - \frac{1}{r} \frac{\partial A_r}{\partial \theta}$$

Laplacian

$$\nabla^2 f = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial f}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 f}{\partial \theta^2} + \frac{\partial^2 f}{\partial z^2}$$

Laplacian of a vector

$$(\nabla^2 \underline{A})_r = \nabla^2 A_r - \frac{2}{r^2} \frac{\partial A_\theta}{\partial \theta} - \frac{A_r}{r^2}$$

$$(\nabla^2 \underline{A})_\theta = \nabla^2 A_\theta + \frac{2}{r^2} \frac{\partial A_r}{\partial \theta} - \frac{A_\theta}{r^2}$$

$$(\nabla^2 \underline{A})_z = \nabla^2 A_z$$

Components of  $(\underline{A} \cdot \underline{\nabla}) \underline{B}$

$$(\underline{A} \cdot \underline{\nabla})_r = A_r \frac{\partial B_r}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_r}{\partial \theta} + A_z \frac{\partial B_r}{\partial z} - \frac{A_\theta B_\theta}{r}$$

$$(\underline{A} \cdot \underline{\nabla})_\theta = A_r \frac{\partial B_\theta}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_\theta}{\partial \theta} + A_z \frac{\partial B_\theta}{\partial z} + \frac{A_\theta B_r}{r}$$

$$^*(\underline{A} \cdot \underline{\nabla})_z = A_r \frac{\partial B_z}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_z}{\partial \theta} + A_z \frac{\partial B_z}{\partial z}$$

Divergence of a tensor

$$(\underline{\nabla} \cdot \underline{T})_r = \frac{1}{r} \frac{\partial}{\partial r} (r T_{rr}) + \frac{1}{r} \frac{\partial}{\partial \theta} (T_{\theta r}) + \frac{\partial T_{zr}}{\partial z} - \frac{1}{r} T_{\theta\theta}$$

$$(\underline{\nabla} \cdot \underline{T})_\theta = \frac{1}{r} \frac{\partial}{\partial r} (r T_{r\theta}) + \frac{1}{r} \frac{\partial T_{\theta\theta}}{\partial \theta} + \frac{\partial T_{z\theta}}{\partial z} + \frac{1}{r} T_{\theta r}$$

$$(\underline{\nabla} \cdot \underline{T})_z = \frac{1}{r} \frac{\partial}{\partial r} (r T_{rz}) + \frac{1}{r} \frac{\partial T_{\theta z}}{\partial \theta} + \frac{\partial T_{zz}}{\partial z}$$

## Spherical Coordinates

### Divergence

$$\nabla \cdot \underline{A} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 A_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (A_\theta \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial A_\phi}{\partial \phi}$$

### Gradient

$$(\nabla f)_r = \frac{\partial f}{\partial r}; \quad (\nabla f)_\theta = \frac{1}{r} \frac{\partial f}{\partial \theta}; \quad (\nabla f)_\phi = \frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi}$$

### Curl

$$(\nabla \times \underline{A})_r = \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (A_\phi \sin \theta) - \frac{1}{r \sin \theta} \frac{\partial A_\theta}{\partial \phi}$$

$$(\nabla \times \underline{A})_\theta = \frac{1}{r \sin \theta} \frac{\partial A_r}{\partial \phi} - \frac{1}{r} \frac{\partial}{\partial r} (r A_\phi)$$

$$(\nabla \times \underline{A})_\phi = \frac{1}{r} \frac{\partial}{\partial r} (r A_\theta) - \frac{1}{r} \frac{\partial A_r}{\partial \theta}$$

### Laplacian

$$\nabla^2 f = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial f}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial f}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2}$$

Laplacian of a vector

$$(\nabla^2 \underline{A})_r = \nabla^2 A_r - \frac{2A_r}{r^2} - \frac{2}{r^2} \frac{\partial A_\theta}{\partial \theta} - \frac{2A_\theta \cot \theta}{r^2} - \frac{2}{r^2 \sin \theta} \frac{\partial A_\phi}{\partial \phi}$$

$$(\nabla^2 \underline{A})_\theta = \nabla^2 A_\theta + \frac{2}{r^2} \frac{\partial A_r}{\partial \theta} - \frac{A_\theta}{r^2 \sin^2 \theta} - \frac{2 \cos \theta}{r^2 \sin^2 \theta} \frac{\partial A_\phi}{\partial \phi}$$

$$(\nabla^2 \underline{A})_\phi = \nabla^2 A_\phi - \frac{A_\phi}{r^2 \sin^2 \theta} + \frac{2}{r^2 \sin \theta} \frac{\partial A_r}{\partial \phi} + \frac{2 \cos \theta}{r^2 \sin^2 \theta} \frac{\partial A_\theta}{\partial \phi}$$

Components of  $(\underline{A} \cdot \nabla) \underline{B}$

$$(\underline{A} \cdot \nabla \underline{B})_r = A_r \frac{\partial B_r}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_r}{\partial \theta} + \frac{A_\phi}{r \sin \theta} \frac{\partial B_r}{\partial \phi} - \frac{A_\theta B_\theta + A_\phi B_\phi}{r}$$

$$(\underline{A} \cdot \nabla \underline{B})_\theta = A_r \frac{\partial B_\theta}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_\theta}{\partial \theta} + \frac{A_\phi}{r \sin \theta} \frac{\partial B_\theta}{\partial \phi} + \frac{A_\theta B_r}{r} - \frac{A_\phi B_\phi \cot \theta}{r}$$

$$(\underline{A} \cdot \nabla \underline{B})_\phi = A_r \frac{\partial B_\phi}{\partial r} + \frac{A_\theta}{r} \frac{\partial B_\phi}{\partial \theta} + \frac{A_\phi}{r \sin \theta} \frac{\partial B_\phi}{\partial \phi} + \frac{A_\theta B_r}{r} + \frac{A_\phi B_\theta \cot \theta}{r}$$

Divergence of a tensor

$$\begin{aligned}
 (\nabla \cdot \underline{T})_r &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 T_{rr}) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (T_{\theta r} \sin \theta) \\
 &+ \frac{1}{r \sin \theta} \frac{\partial T_{\phi r}}{\partial \phi} - \frac{T_{\theta\theta} + T_{\phi\phi}}{r}
 \end{aligned}$$

$$\begin{aligned}
 (\nabla \cdot \underline{T})_\theta &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 T_{r\theta}) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (T_{\theta\theta} \sin \theta) \\
 &+ \frac{1}{r \sin \theta} \frac{\partial T_{\phi\theta}}{\partial \phi} + \frac{T_{\theta r}}{r} - \frac{\cot \theta}{r} T_{\phi\phi}
 \end{aligned}$$

$$\begin{aligned}
 (\nabla \cdot \underline{T})_\phi &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 T_{r\phi}) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (T_{\theta\phi} \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial T_{\phi\phi}}{\partial \phi} + \frac{T_{\phi r}}{r} \\
 &+ \frac{\cot \theta}{r} T_{\phi\theta}
 \end{aligned}$$

Note that the expressions given for  $\nabla \cdot \underline{T}$  are consistent with the definition (17).

### General Curvilinear Coordinates <sup>[3,4]</sup>

In any orthogonal curvilinear coordinate system, all of the operations involving  $\nabla$  can be defined in terms of the scale factors  $h_i$ . These are related to the differential arc length  $d\ell$  by

$$\begin{aligned} d\ell^2 &= dx^2 + dy^2 + dz^2 \\ (35) \quad &= h_1^2 d\xi_1^2 + h_2^2 d\xi_2^2 + h_3^2 d\xi_3^2, \end{aligned}$$

where  $\{\xi_1, \xi_2, \xi_3\}$  designate the curvilinear coordinates. For example, the scale factors associated with the spherical coordinates  $\{r, \theta, \phi\}$  are 1,  $r$  and  $r \sin \theta$ , respectively.

In orthogonal curvilinear coordinate systems a vector  $\underline{A}$  can be written as

$$(36) \quad \underline{A} = \underline{e}_1 A_1 + \underline{e}_2 A_2 + \underline{e}_3 A_3,$$

where the  $\{\underline{e}_i\}$  form a right-handed set of unit vectors normal to the surfaces of constant  $\xi_i$ . Then the divergence of  $\underline{A}$  is

$$(37) \quad \nabla \cdot \underline{A} = \frac{1}{H} \sum_i \frac{\partial}{\partial \xi_i} \left( \frac{H A_i}{h_i} \right),$$

where  $H = h_1 h_2 h_3$ . The gradient of a scalar  $f$  is given by

$$(38) \quad (\nabla f)_i = \frac{1}{h_i} \frac{\partial f}{\partial \xi_i}.$$

From Eqs. (12), (37) and (38), it follows that the Laplacian of  $f$  is

$$(39) \quad \nabla^2 f = \frac{1}{H} \sum_i \frac{\partial}{\partial \xi_i} \left( \frac{H}{h_i^2} \frac{\partial f}{\partial \xi_i} \right).$$

The curl of  $\underline{A}$  is

$$(40) \quad (\underline{\nabla} \times \underline{A})_i = \sum_{j,k} \epsilon_{ijk} \frac{1}{h_j h_k} \frac{\partial}{\partial \xi_j} (h_k A_k),$$

where  $\epsilon_{ijk}$  equals +1 if  $\{ijk\}$  is even (i.e., an even permutation of  $\{123\}$ ), equals -1 if  $\{ijk\}$  is odd, and vanishes if  $i, j$  and  $k$  are not all distinct.

To obtain expressions for the remaining operations, it is useful to introduce the Christoffel symbols  $\Gamma_{jk}^i$ , defined by

$$(41) \quad \Gamma_{jk}^i = \frac{1}{h_j h_k} \left[ \frac{\partial h_j}{\partial \xi_k} \delta_j^i - \frac{\partial h_k}{\partial \xi_j} \delta_k^i \right],$$

where  $\delta_j^i$  equals 1 if  $i = j$  and vanishes otherwise.  $\Gamma_{jk}^i$  vanishes if  $j$  and  $k$  both differ from  $i$  or if  $j = k$ . In addition,

$$(42) \quad \Gamma_{jk}^i + \Gamma_{kj}^i = 0;$$

$$(43) \quad \Gamma_{jk}^i + \Gamma_{ki}^j + \Gamma_{ij}^k = 0;$$

and

$$(44) \quad \sum_i \Gamma_{ik}^i = \frac{1}{H} \frac{\partial}{\partial \xi_k} \left( \frac{H}{h_k} \right).$$

There are at most six independent non-zero components  $\Gamma_{jk}^i$ , and less in most coordinate systems of physical interest. The curvature tensor,

$$R_{ijkl} = \frac{1}{h_j} \frac{\partial \Gamma_{kl}^i}{\partial \xi_j} - \frac{1}{h_i} \frac{\partial \Gamma_{kl}^j}{\partial \xi_i} +$$

(45)

$$\sum_n \left\{ \Gamma_{ij}^n \Gamma_{kl}^n - \Gamma_{nk}^j \Gamma_{nl}^i + \Gamma_{kn}^i \Gamma_{ln}^j \right\}$$

always vanishes in a flat space, i.e., one which can be described by Cartesian coordinates. (This is the only kind of space we usually deal with in plasma physics.)

In terms of the  $\Gamma$ 's, the gradient of a vector  $\underline{A}$  is a tensor  $\underline{T}$  given by

$$T_{ji} = (\underline{\nabla} \underline{A})_{ji} = \frac{\partial A_i}{h_j \partial \xi_j} + \sum_l \Gamma_{il}^j A_l .$$

(46)

Contracting this expression (i.e., multiplying by  $\delta_j^i$  and summing over  $i$  and  $j$ ) recovers Eq. (37). The directional derivative is

$$\left[ (\underline{A} \cdot \underline{\nabla}) \underline{B} \right]_i = \sum_j A_j (\underline{\nabla} \underline{B})_{ji}$$

(47)

$$= \sum_j \left\{ \frac{A_j}{h_j} \frac{\partial B_i}{\partial \xi_j} + \frac{A_i B_j}{h_i h_j} \frac{\partial h_i}{\partial \xi_j} - \frac{A_i B_j}{h_i h_j} \frac{\partial h_j}{\partial \xi_i} \right\} .$$

The gradient of a (second-order) tensor is a third-order tensor given by

$$(48) \quad (\nabla_{\underline{\underline{T}}})_{kji} = \frac{\partial T_{ji}}{h_k \partial \xi_k} + \sum_{\ell} \left\{ \Gamma_{i\ell}^k T_{j\ell} + \Gamma_{j\ell}^k T_{\ell i} \right\}.$$

The divergence of the tensor  $\underline{\underline{T}}$  is obtained by contracting Eq. (48) with respect to the first two indices:

$$\begin{aligned} (\nabla_{\underline{\underline{T}}})_i &= \sum_j \left\{ \frac{\partial T_{ji}}{h_j \partial \xi_j} + \sum_{\ell} \left[ \Gamma_{i\ell}^j T_{j\ell} + \Gamma_{j\ell}^j T_{\ell i} \right] \right\} \\ &= \sum_j \left\{ \frac{1}{H} \frac{\partial}{\partial \xi_j} \left[ \frac{HT_{ji}}{h_j} \right] + \sum_{\ell} \Gamma_{i\ell}^j T_{j\ell} \right\}. \end{aligned}$$

If  $\underline{\underline{T}}$  is defined according to Eq. (46),  $\nabla \cdot \underline{\underline{T}}$  is the Laplacian of the vector  $\underline{A}$ :

$$\begin{aligned} (\nabla^2 \underline{A})_i &= \sum_j (\nabla_{\underline{\underline{T}}})_{jji} = \sum_j \left\{ \frac{1}{H} \frac{\partial}{\partial \xi_j} \left[ \frac{H}{h_j^2} \frac{\partial A_i}{\partial \xi_j} \right] \right. \\ (50) \quad &+ \frac{h_i A_i}{H} \frac{\partial}{\partial \xi_j} \left[ \frac{H}{h_i^2 h_j^2} \frac{\partial h_i}{\partial \xi_j} \right] + \frac{2}{h_i^2 h_j^2} \frac{\partial h_i}{\partial \xi_j} \frac{\partial}{\partial \xi_i} (h_j A_j) \\ &\left. - \frac{2}{h_i h_j^3} \frac{\partial h_j}{\partial \xi_i} \frac{\partial}{\partial \xi_j} (h_j A_j) + \frac{h_j A_j}{h_i} \frac{\partial}{\partial \xi_i} \left[ \frac{1}{H} \frac{\partial}{\partial \xi_j} \left( \frac{H}{h_j^2} \right) \right] \right\}. \end{aligned}$$

Evidently the first term on the RHS is  $\nabla^2 A_i$ , the scalar Laplacian applied to  $A_i$  [Eq. (39)]. Fortunately, Eq. (50) simplifies considerably for most coordinate systems of interest.

In the following section, the results of Eqs. (35) - (50) are applied to a coordinate system occasionally encountered in plasma physics problems.

### Toroidal Coordinates

An arbitrary point in 3-space can be uniquely identified by the coordinates  $\{\rho, \phi, \psi\}$ , where  $\rho$  is the distance between the given point and a fixed toroidal axis of radius  $R$  (the major radius),  $\phi$  is the azimuthal or toroidal angle (i.e., the one sweeping around the major circumference), and  $\psi$  is the poloidal angle. (Note that this system differs from the "toroidal" coordinates in Ref. 3, p. 666, and from the flux surface coordinates commonly used in connection with toroidal devices.) The values assumed by these coordinates are restricted according to  $0 \leq \rho < \infty$ ,  $0 \leq \phi < 2\pi$ , and  $0 \leq \psi < 2\pi$ . If  $\{r, \phi, z\}$  are the cylindrical coordinates of the point in question, where  $z$  is normal to the plane of the torus, then  $\phi$  is the same in both systems and

$$\begin{aligned} R + \rho \cos \psi &= r, \\ \rho \sin \psi &= z. \end{aligned}$$

Hence the square of the differential arc length is

$$\begin{aligned} d\ell^2 &= dr^2 + r^2 d\phi^2 + dz^2 \\ &= d\rho^2 + (R + \rho \cos \psi)^2 d\phi^2 + \rho^2 d\psi^2. \end{aligned}$$

Consequently the scale factors are 1,  $R + \rho \cos \psi$ , and  $\rho$ , respectively.

From Eq. (37), the divergence of  $\underline{A}$  is

$$\nabla \cdot \underline{A} = \frac{1}{R + \rho \cos \psi} \left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \left[ \rho(R + \rho \cos \psi) A_{\rho} \right] + \frac{\partial A_{\psi}}{\partial \psi} + \frac{1}{\rho} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) A_{\psi} \right] \right\}.$$

From Eq. (38), the gradient of  $f$  has components

$$(\nabla f)_{\rho} = \frac{\partial f}{\partial \rho};$$

$$(\nabla f)_{\psi} = \frac{1}{R + \rho \cos \psi} \frac{\partial f}{\partial \psi};$$

$$(\nabla f)_{\psi} = \frac{1}{\rho} \frac{\partial f}{\partial \psi}.$$

From Eq. (39), the Laplacian of  $f$  is

$$\nabla^2 f = \frac{1}{R + \rho \cos \psi} \left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \left[ \rho(R + \rho \cos \psi) \frac{\partial f}{\partial \rho} \right] + \frac{1}{R + \rho \cos \psi} \frac{\partial^2 f}{\partial \psi^2} + \frac{1}{\rho^2} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) \frac{\partial f}{\partial \psi} \right] \right\}.$$

From Eq. (40), the curl of  $\underline{A}$  is given by

$$(\underline{\nabla} \times \underline{A})_o = \frac{1}{R + \rho \cos \psi} \left\{ \frac{\partial A_\psi}{\partial \phi} - \frac{1}{\rho} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) A_\phi \right] \right\} ;$$

$$(\underline{\nabla} \times \underline{A})_\phi = \frac{1}{\rho} \left\{ \frac{\partial A_o}{\partial \psi} - \frac{\partial}{\partial \rho} (\rho A_\psi) \right\} ;$$

$$(\underline{\nabla} \times \underline{A})_\psi = \frac{1}{R + \rho \cos \psi} \left\{ \frac{\partial}{\partial \rho} \left[ (R + \rho \cos \psi) A_\phi - \frac{\partial A_o}{\partial \phi} \right] \right\} .$$

The only nonvanishing Christoffel symbols are

$$\Gamma_{21}^2 = -\Gamma_{12}^2 = \frac{\cos \psi}{R + \rho \cos \psi} ;$$

$$\Gamma_{23}^2 = -\Gamma_{32}^2 = \frac{-\sin \psi}{R + \rho \cos \psi} ;$$

$$\Gamma_{31}^3 = -\Gamma_{13}^3 = \frac{1}{\rho} .$$

Then from Eq. (47),

$$\begin{aligned} \left[ (\underline{A} \cdot \underline{\nabla}) \underline{B} \right]_o &= A_o \frac{\partial B_o}{\partial \rho} + \frac{A_\phi}{R + \rho \cos \psi} \frac{\partial B_o}{\partial \phi} \\ &+ \frac{A_\psi}{\rho} \frac{\partial B_o}{\partial \psi} - \frac{A_\phi B_\phi \cos \psi}{R + \rho \cos \psi} - \frac{A_\psi B_\psi}{\rho} ; \end{aligned}$$

$$\left[ (\underline{A} \cdot \underline{\nabla}) \underline{B} \right]_{\phi} = A_{\phi} \frac{\partial B_{\phi}}{\partial \rho} + \frac{A_{\phi}}{R + \rho \cos \psi} \frac{\partial B_{\phi}}{\partial \phi} +$$

$$\frac{A_{\psi}}{\rho} \frac{\partial B_{\phi}}{\partial \psi} + \frac{A_{\phi} B_{\rho} \cos \psi}{R + \rho \cos \psi} - \frac{A_{\phi} B_{\psi} \sin \psi}{R + \rho \cos \psi} ;$$

$$\left[ (\underline{A} \cdot \underline{\nabla}) \underline{B} \right]_{\psi} = A_{\psi} \frac{\partial B_{\psi}}{\partial \rho} + \frac{A_{\psi}}{R + \rho \cos \psi} \frac{\partial B_{\psi}}{\partial \phi} +$$

$$\frac{A_{\psi}}{\rho} \frac{\partial B_{\psi}}{\partial \psi} + \frac{A_{\psi} B_{\rho}}{\rho} + \frac{A_{\psi} B_{\phi} \sin \psi}{R + \rho \cos \psi} .$$

From Eq. (49),

$$(\underline{\nabla} \cdot \underline{T})_{\rho} = \frac{1}{R + \rho \cos \psi} \left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \left[ \rho (R + \rho \cos \psi) T_{\rho\rho} \right] \right.$$

$$\left. + \frac{\partial T}{\partial \phi} \phi_{\rho} + \frac{1}{\rho} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) T_{\psi\rho} \right] \right.$$

$$\left. - \cos \psi T_{\phi\phi} \right\} - \frac{1}{\rho} T_{\psi\psi} ;$$

$$\begin{aligned}
(\nabla \cdot \underline{T})_{\phi} &= \frac{1}{R + \rho \cos \psi} \left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \left[ \rho(R + \rho \cos \psi) T_{\rho\phi} \right] \right. \\
&+ \frac{\partial T_{\phi\phi}}{\partial \phi} + \frac{1}{\rho} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) T_{\psi\phi} \right] \\
&\left. + \cos \psi T_{\phi\rho} - \sin \psi T_{\phi\psi} \right\} ;
\end{aligned}$$

$$\begin{aligned}
(\nabla \cdot \underline{T})_{\psi} &= \frac{1}{R + \rho \cos \psi} \left\{ \frac{1}{\rho} \frac{\partial}{\partial \rho} \left[ \rho(R + \rho \cos \psi) T_{\rho\psi} \right] \right. \\
&+ \frac{\partial T_{\phi\psi}}{\partial \phi} + \frac{1}{\rho} \frac{\partial}{\partial \psi} \left[ (R + \rho \cos \psi) T_{\psi\psi} \right] \\
&\left. + \sin \psi T_{\phi\rho} \right\} + \frac{1}{\rho} T_{\psi\rho} .
\end{aligned}$$

Finally, from Eq. (50),

$$\begin{aligned}
(\nabla^2 \underline{A})_{\rho} &= \nabla^2 A_{\rho} - A_{\rho} \left[ \frac{1}{\rho^2} + \frac{\cos^2 \psi}{(R + \rho \cos \psi)^2} \right] \\
&- \frac{2 \cos \psi}{(R + \rho \cos \psi)^2} \frac{\partial A_{\phi}}{\partial \phi} - \frac{2}{\rho^2} \frac{\partial A_{\psi}}{\partial \psi} + \\
&\frac{\sin \psi A_{\psi}}{R + \rho \cos \psi} \left[ \frac{1}{\rho} + \frac{\cos \psi}{R + \rho \cos \psi} \right] ;
\end{aligned}$$

$$\begin{aligned}
 (\nabla^2 A)_\phi &= \nabla^2 A_\phi - \frac{1}{(R + \rho \cos \psi)^2} A_\phi \\
 &+ \frac{2}{(R + \rho \cos \psi)^2} \left[ \cos \psi \frac{\partial A_\rho}{\partial \phi} - \sin \psi \frac{\partial A_\psi}{\partial \phi} \right];
 \end{aligned}$$

$$\begin{aligned}
 (\nabla^2 A)_\psi &= \nabla^2 A_\psi - A_\psi \left[ \frac{1}{\rho^2} + \frac{\sin^2 \psi}{(R + \rho \cos \psi)^2} \right] \\
 &+ \frac{2}{\rho^2} \frac{\partial A_\rho}{\partial \psi} + \frac{\sin \psi}{(R + \rho \cos \psi)^2} \left[ 2 \frac{\partial A_\phi}{\partial \psi} - \frac{R A_\rho}{\rho} \right].
 \end{aligned}$$

DIMENSIONS AND UNITS [5]

To get the value of a quantity in Gaussian units, multiply the value expressed in mks units by the conversion factor.

Physical Quantity	Symbol	Dimensions		Rationalized mks	Conversion Factor	Gaussian
		mks Units	Gaussian Units			
Capacitance	C	$\frac{t^2 q^2}{m \ell^2}$	$\ell$	farad	$9 \times 10^{11}$	cm
Charge	q	q	$\frac{1/2 m \ell^{3/2}}{t}$	coulomb	$3 \times 10^9$	statcoulomb
Charge density	$\rho$	$\frac{q}{\ell^3}$	$\frac{1/2 m}{\ell^{5/2} t}$	coulomb/m <sup>3</sup>	$3 \times 10^3$	statcoulomb/cm <sup>3</sup>
Conductance		$\frac{t q^2}{m \ell^2}$	$\frac{\ell}{t}$	siemens*	$9 \times 10^{11}$	cm/sec
Conductivity	$\sigma$	$\frac{t q^2}{m \ell^3}$	$\frac{1}{t}$	siemens/m*	$9 \times 10^9$	sec <sup>-1</sup>
Current	I	$\frac{q}{t}$	$\frac{1/2 m \ell^{3/2}}{t^2}$	ampere	$3 \times 10^9$	statampere
Current density	J	$\frac{q}{\ell^2 t}$	$\frac{1/2 m}{\ell^{3/2} t^2}$	ampere/m <sup>2</sup>	$3 \times 10^5$	statampere/cm <sup>2</sup>
Density	$\rho$	$\frac{m}{\ell^3}$	$\frac{m}{\ell^3}$	kg/m <sup>3</sup>	$10^{-3}$	g/cm <sup>3</sup>

Physical Quantity	Symbol	Dimensions		Rationalized mks	Conversion Factor	Gaussian
		mks Units	Gaussian Units			
Displacement	$\underline{D}$	$\frac{q}{\ell^2}$	$\frac{m^{1/2}}{\ell^{1/2}t}$	coulomb/m <sup>2</sup>	$4\pi \times 10^5$	statcoulomb/cm <sup>2</sup>
Electric field	$\underline{E}$	$\frac{m\ell}{t^2q}$	$\frac{m^{1/2}}{t\ell^{1/2}}$	volt/m	$\frac{1}{3} \times 10^{-4}$	statvolt/cm
Electromotance	$\underline{\mathcal{E}}, \text{Emf}$	$\frac{m\ell^2}{t^2q}$	$\frac{m^{1/2}\ell^{1/2}}{t}$	volt	$\frac{1}{3} \times 10^{-2}$	statvolt
Energy	$U, W$	$\frac{m\ell^2}{t}$	$\frac{m\ell^2}{t}$	joule	$10^7$	erg
Energy density		$\frac{m}{\ell t^2}$	$\frac{m}{\ell t^2}$	joule/m <sup>3</sup>	10	erg/cm <sup>3</sup>
Force	$\underline{F}$	$\frac{m\ell}{t^2}$	$\frac{m\ell}{t^2}$	newton	$10^5$	dyne
Frequency	$f, \nu, \omega$	$\frac{1}{t}$	$\frac{1}{t}$	hertz	1	hertz

Physical Quantity	Symbol	Dimensions		Rationalized mks	Conversion Factor	Gaussian
		mks Units	Gaussian Units			
Impedance	Z	$\frac{m\ell^2}{tq^2}$	$\frac{t}{\ell}$	ohm	$\frac{1}{9} \times 10^{-11}$	sec/cm
Inductance	L	$\frac{m\ell^2}{q^2}$	$\frac{t^2}{\ell}$	henry	$\frac{1}{9} \times 10^{-11}$	sec <sup>2</sup> /cm
Length	ℓ	ℓ	ℓ	meter (m)	10 <sup>2</sup>	centimeter (cm)
Magnetic intensity	H	$\frac{q}{\ell t}$	$\frac{m^{1/2}}{\ell^{1/2} t}$	ampere-turn/m	$\frac{4\pi}{9} \times 10^{-3}$	oersted
Magnetic flux	Φ	$\frac{m\ell^2}{tq}$	$\frac{m^{1/2}\ell^{3/2}}{t}$	weber	10 <sup>6</sup>	maxwell
Magnetic induction	B	$\frac{m}{tq}$	$\frac{m^{1/2}}{\ell^{1/2} t}$	tesla *	10 <sup>4</sup>	gauss
Magnetization	M	$\frac{q}{\ell t}$	$\frac{m^{1/2}}{\ell^{1/2} t}$	ampere-turn/m	10 <sup>-3</sup>	oersted
Magnetomotance	Mmf, N	$\frac{q}{t}$	$\frac{m^{1/2}\ell^{1/2}}{t}$	ampere-turn	$\frac{4\pi}{9}$	gilbert
Mass	m, M	m	m	kilogram (kg)	10 <sup>3</sup>	gram (g)
Momentum	p, P	$\frac{m\ell}{t}$	$\frac{m\ell}{t}$	kg-m/sec	10 <sup>3</sup>	g-cm/sec

Physical Quantity	Symbol	Dimensions		Rationalized mks	Conversion Factor	Gaussian
		mks Units	Gaussian Units			
Momentum density		$\frac{m}{\ell^2 t}$	$\frac{m}{\ell^2 t}$	kg/m <sup>2</sup> -sec	10 <sup>-1</sup>	g/cm <sup>2</sup> -sec
Permeability	$\mu$	$\frac{m\ell}{t^2 q}$	1	henry/m	$\frac{1}{4\pi} \times 10^7$	(dimensionless)
Permittivity	$\epsilon$	$\frac{t^2 q^2}{m\ell^3}$	1	farad/m	$56\pi \times 10^9$	(dimensionless)
Polarization	$\underline{P}$	$\frac{q}{\ell^2}$	$\frac{1/2}{\ell^2/t}$	coulomb/m <sup>2</sup>	$3 \times 10^5$	statcoulomb/cm <sup>2</sup>
Potential	V, $\phi$	$\frac{m\ell^2}{t^2 q}$	$\frac{m^{1/2}\ell^{1/2}}{t}$	volt	$\frac{1}{3} \times 10^{-2}$	statvolt
Power	P	$\frac{m\ell^2}{t^3}$	$\frac{m\ell^2}{t^3}$	watt	10 <sup>7</sup>	erg/sec
Power density		$\frac{m}{\ell t^3}$	$\frac{m}{\ell t^3}$	watt/m <sup>3</sup>	10	erg/cm <sup>3</sup> -sec
Pressure	P	$\frac{m}{\ell t^2}$	$\frac{m}{\ell t^2}$	pascal*	10	dyne/cm <sup>2</sup>
Reluctance	$\mathcal{R}$	$\frac{q^2}{m\ell^2}$	$\frac{1}{\ell}$	ampere-turn/weber	$4\pi \times 10^{-9}$	cm <sup>-1</sup>

Physical Quantity	Symbol	Dimensions		Rationalized mks	Conversion Factor	Gaussian
		mks Units	Gaussian Units			
Resistance	R	$\frac{m^2}{t^2 q}$	$\frac{t}{\ell}$	ohm	$\frac{1}{9} \times 10^{-11}$	sec/cm
Resistivity	$\eta, \rho$	$\frac{m^2}{t^2 q}$	t	ohm-m	$\frac{1}{9} \times 10^{-9}$	sec
Thermal conductivity	$\kappa$	$\frac{m\ell}{t^3}$	$\frac{m\ell}{t^3}$	watt/m <sup>2</sup> -deg(K)	10 <sup>5</sup>	erg/cm-sec-deg(K)
Time	t	t	t	second(sec)	1	second
Vector potential	$\vec{A}$	$\frac{m\ell}{tq}$	$\frac{m^{1/2} \ell^{1/2}}{t}$	weber/m	10 <sup>9</sup>	gauss-cm
Velocity	$\vec{v}$	$\frac{\ell}{t}$	$\frac{\ell}{t}$	m/sec	10 <sup>2</sup>	cm/sec
Viscosity	$\eta, \mu$	$\frac{m}{\ell t}$	$\frac{m}{\ell t}$	kg/m-sec	10	poise
Vorticity	$\zeta$	$\frac{1}{t}$	$\frac{1}{t}$	sec <sup>-1</sup>	1	sec <sup>-1</sup>
Work	W	$\frac{m\ell^2}{t^2}$	$\frac{m\ell^2}{t^2}$	joule	10 <sup>7</sup>	erg

PHYSICAL CONSTANTS [5, 2]

CGS

Physical Quantity	Symbol	Value	Units
Boltzmann constant	$k$	$1.3807 \times 10^{-16}$	erg/deg (K)
Elementary charge	$e$	$4.8032 \times 10^{-10}$	statcoulomb
Electron mass	$m_e$	$9.1095 \times 10^{-28}$	g
Proton mass	$m_p$	$1.6726 \times 10^{-24}$	g
Gravitational constant	$G$	$6.6720 \times 10^{-8}$	dyne-cm <sup>2</sup> /g <sup>2</sup>
Planck constant	$h$	$6.6262 \times 10^{-27}$	erg-sec
	$\hbar = h/2\pi$	$1.0546 \times 10^{-27}$	erg-sec
Speed of light in vacuum	$c$	$2.9979 \times 10^{10}$	cm/sec

Physical Quantity	Symbol	Value	Units
Proton/electron mass ratio	$m/p_e$	$1.8362 \times 10^3$	
Electron charge/mass ratio	$e/m_e$	$5.2728 \times 10^{17}$	statcoulomb/g
Rydberg constant	$R_\infty = \frac{2\pi^2 m_e^4}{ch^3}$	$1.0974 \times 10^5$	$\text{cm}^{-1}$
Bohr radius	$a_0 = \frac{h^2}{me^2}$	$5.2918 \times 10^{-9}$	cm
Atomic cross section	$\pi a_0^2$	$8.7974 \times 10^{-17}$	$\text{cm}^2$
Classical electron radius	$r_e = e^2/mc^2$	$2.8179 \times 10^{-13}$	cm
Thomson cross section	$(8\pi/3)r_e^2$	$6.6524 \times 10^{-25}$	$\text{cm}^2$
Compton wavelength of electron	$h/m_e c$	$2.4263 \times 10^{-10}$	cm
	$h/m_e c$	$3.8616 \times 10^{-11}$	cm
Fine-structure constant	$\alpha = \frac{e^2}{hc}$	$7.2974 \times 10^{-3}$	
	$\alpha^{-1}$	137.04	

Physical Quantity	Symbol	Value	Units
First radiation constant	$c_1 = 2\pi^5 hc^2$	$3.7418 \times 10^{-5}$	erg-cm <sup>2</sup> /sec
Second radiation constant	$c_2 = hc/k$	1.4388	cm-deg (K)
Stefan-Boltzmann constant	$\sigma$	$5.6703 \times 10^{-5}$	erg/cm <sup>2</sup> -sec-deg <sup>4</sup>
Wavelength associated with 1 eV	$\lambda_0$	$1.2399 \times 10^{-4}$	cm
Frequency associated with 1 eV	$\nu_0$	$2.4180 \times 10^{14}$	Hz
Wave number associated with 1 eV	$k_0$	$8.0655 \times 10^3$	cm <sup>-1</sup>
Energy associated with 1 eV		$1.602 \times 10^{-12}$	erg
Energy associated with 1 cm <sup>-1</sup>		$1.9865 \times 10^{-16}$	erg
Energy associated with 1 Rydberg		13.606	eV
Energy associated with 1° Kelvin		$8.6173 \times 10^{-5}$	eV
Temperature associated with 1 eV		$1.1605 \times 10^4$	deg(K)
Avogadro number	$N_A$	$6.0220 \times 10^{23}$	mol <sup>-1</sup>

Physical Quantity	Symbol	Value	Units
Faraday constant	$F = N_A e$	$2.8925 \times 10^{14}$	statcoulomb/mol
Gas constant	$R = N_A k$	$8.3144 \times 10^7$	erg/deg-mol
Loschmidt's number (no. density at STP)	$n_0$	$2.6868 \times 10^{19}$	$\text{cm}^{-3}$
Atomic mass unit	$m_u$	$1.6605 \times 10^{-24}$	g
Standard temperature	$T_0$	273.15	deg (K)
Atmospheric pressure	$p_0 = n k T_0$	$1.0133 \times 10^9$	dyne/cm <sup>2</sup>
Pressure of 1 mm Hg (torr)		$1.3332 \times 10^9$	dyne/cm <sup>2</sup>
Molar volume at STP	$V_0 = RT_0/p_0$	$2.2415 \times 10^4$	cm <sup>3</sup>
calorie (cal)		$4.1858 \times 10^7$	erg
Gravitational acceleration	g	980.67	cm/sec <sup>2</sup>

METRIC PREFIXES <sup>(5)</sup>

Multiple	Prefix	Symbol	Multiple	Prefix	Symbol
$10^{-1}$	deci	d	10	deca	da
$10^{-2}$	centi	c	$10^2$	hecto	h
$10^{-3}$	milli	m	$10^3$	kilo	k
$10^{-6}$	micro	$\mu$	$10^6$	mega	M
$10^{-9}$	nano	n	$10^9$	giga	G
$10^{-12}$	pico	p	$10^{12}$	tera	T
$10^{-15}$	femto	f	$10^{15}$	peta	P
$10^{-18}$	atto	a	$10^{18}$	exa	E

## ELECTRICITY AND MAGNETISM

In the following,  $\sigma$  = conductivity,  $f = \omega/2\pi$  = radiation frequency,  $\kappa_m = \mu/\mu_0$  and  $\kappa_e = \epsilon/\epsilon_0$ . Where subscripts are used, 1 denotes a conducting medium and 2 a propagating (lossless dielectric) medium. All units are mks unless otherwise specified.

Permittivity of free space  $\epsilon_0 = 8.8542 \times 10^{-12}$  farad/m.

Permeability of free space  $\mu_0 = 4\pi \times 10^{-7} = 1.2566 \times 10^{-6}$  henry/m.

Resistance of free space  $R_0 = (\mu_0/\epsilon_0)^{1/2} = 376.73$  ohm

Relaxation time in a lossy medium  $\tau = \epsilon/\sigma$  sec

Skin depth in a lossy medium  $\delta = (2/\omega\sigma)^{1/2} = (\pi f\mu\sigma)^{-1/2}$  m

Wave impedance in lossy medium  $Z = [\mu/(\epsilon + i\sigma/\omega)]^{1/2}$

Reflection coefficient at conducting surface (good only for  $R \approx 1$ )  $R = 1 - 4.22 \times 10^{-4} (f\kappa_m \kappa_e / \sigma)^{1/2}$

Field at distance  $r$  from straight wire carrying current  $I$   $B_a = \mu I / 2\pi r$  tesla  
 $= 0.2 I/r$  gauss ( $r$  in cm)

Electromagnetic energy in volume  $V$   $W = \frac{1}{2} \int_V dV (\underline{H} \cdot \underline{B} + \underline{E} \cdot \underline{D})$

Poynting's theorem  $\frac{\partial W}{\partial t} + \int_S \underline{N} \cdot d\underline{S} = \int_V dV \underline{J} \cdot \underline{E}$

where  $S$  is the closed surface bounding  $V$  and the Poynting vector (energy flux across  $S$ ) is  $\underline{N} = \underline{E} \times \underline{H}$ .

MAXWELL'S EQUATIONS

Name	Rationalized mks	Gaussian
Faraday's law	$\nabla \times \underline{E} = -\frac{\partial \underline{B}}{\partial t}$	$\nabla \times \underline{E} = -\frac{1}{c} \frac{\partial \underline{B}}{\partial t}$
Ampere's law	$\nabla \times \underline{H} = \frac{\partial \underline{D}}{\partial t} + \underline{J}$	$\nabla \times \underline{H} = \frac{1}{c} \frac{\partial \underline{D}}{\partial t} + \frac{4\pi}{c} \underline{J}$
Poisson's equation	$\nabla \cdot \underline{D} = \rho$	$\nabla \cdot \underline{D} = 4\pi\rho$
[ Absence of magnetic monopoles ]	$\nabla \cdot \underline{B} = 0$	$\nabla \cdot \underline{B} = 0$
Lorentz force on charge q	$q \left[ \underline{E} + \underline{v} \times \underline{B} \right]$	$q \left[ \underline{E} + \frac{1}{c} \underline{v} \times \underline{B} \right]$
Constitutive relations	$\underline{D} = \epsilon \underline{E}$ $\underline{B} = \mu \underline{H}$	$\underline{D} = \epsilon \underline{E}$ $\underline{B} = \mu \underline{H}$

In a plasma,  $\mu \approx \mu_0$  (Gaussian units:  $\mu \approx 1$ ). The permittivity satisfies  $\epsilon \approx \epsilon_0$  (Gaussian:  $\epsilon \approx 1$ ) provided all charge is regarded as free. Using the drift approximation  $\underline{v}_\perp = \underline{E} \times \underline{B}/B^2$  to calculate polarization charge density given rise to a dielectric constant  $K = \epsilon/\epsilon_0 = 1 + 36\pi \times 10^9 \rho/B^2$  (mks) =  $1 + 4\pi \rho c^2/B^2$  (Gaussian), where  $\rho$  is the mass density.

ELECTROMAGNETIC FREQUENCY/WAVELENGTH BANDS

Nomenclature

E = Extreme                      M = Medium  
 F = Frequency                  S = Super  
 H = High                          U = Ultra  
 L = Low                            V = Very

Designation <sup>[E]</sup>	Frequency Range		Wavelength Range	
	Lower	Upper	Lower	Upper
ULF*		10 Hz	3 Mm	
ELF*	10 Hz	3 kHz	100 km	3 Mm
VLf	3 kHz	30 kHz	10 km	100 km
LF	30 kHz	300 kHz	1 km	10 km
MF	300 kHz	3 MHz	100 m	1 km
HF	3 MHz	30 MHz	10 m	100 m
VHF	30 MHz	300 MHz	1 m	10 m
UHF	300 MHz	3 GHz	10 cm	1 m
SHF <sup>#</sup>	3 GHz	30 GHz	1 cm	10 cm
EHF	30 GHz	300 GHz	1 mm	1 cm
(Submillimeter)	300 GHz	3 THz	100 $\mu$	1 mm
(Infrared)	3 THz			100 $\mu$

\*The boundary between ULF and ELF is variously defined.

‡The SHF (microwave) band is further subdivided approximately as follows:<sup>[3]</sup>

Band Designation	Frequency Range (GHz)		Wavelength Range (cm)	
	Lower	Upper	Lower	Upper
S	2.6	3.95	7.6	11.5
G	3.95	5.85	5.1	7.6
J	5.3	8.2	3.7	5.7
H	7.05	10.0	3.0	4.25
X	8.2	12.4	2.4	3.7
M	10.0	15.0	2.0	3.0
P	12.4	18.0	1.67	2.4
K	18.0	26.5	1.1	1.67
R	26.5	40.0	.75	1.1

## AC CIRCUITS

For a resistance  $R$ , inductance  $L$ , and capacitance  $C$  in series with a voltage source  $V = V_0 \exp(j\omega t)$  (where  $j = \sqrt{-1}$ ), the current  $I$  satisfies  $I = dq/dt$ , where

$$L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + q/C = V .$$

Solutions are  $q(t) = q_s + q_t$ , where the steady state is  $I_s = j\omega q_s = V/Z$  in terms of the impedance  $Z = R + j(\omega L - 1/\omega C)$ . For initial conditions  $q(0) = q_0 = \bar{q}_0 + q_s(0)$ ,  $I(0) = I_0$ , the transients can be of three types, depending on  $\Delta = R^2 - 4L/C$ :

(a) Overdamped,  $\Delta > 0$

$$q_t = \left[ (I_0 - \gamma_- \bar{q}_0) e^{\gamma_+ t} - (I_0 - \gamma_+ \bar{q}_0) e^{\gamma_- t} \right] / (\gamma_+ - \gamma_-),$$

where

$$\gamma_{\pm} = \left( -R \pm \Delta^{\frac{1}{2}} \right) / 2L ;$$

(b) Critically damped,  $\Delta = 0$

$$q_t = \left[ \bar{q}_0 + (I_0 + \gamma_R \bar{q}_0) t \right] e^{-\gamma_R t} ,$$

where

$$\gamma_R = R/2L ;$$

(c) Underdamped,  $\Delta < 0$

$$q_t = \left[ \omega_1^{-1} (\gamma_R \bar{q}_0 + I_0) \sin \omega_1 t + \bar{q}_0 \cos \omega_1 t \right] e^{-\gamma_R t},$$

where

$$\omega_1 = \omega_0 \left( 1 - R^2 C / 4L \right)^{1/2}, \quad \omega_0 = (LC)^{-1/2}.$$

The quality of the circuit is  $Q = \omega_0 L / R$ ;  $\omega_0$  is the resonant frequency. At  $\omega = \omega_0$ ,  $Z = R$ . Instability results when  $L$ ,  $R$ ,  $C$  are not all of the same sign.

## DIMENSIONLESS NUMBERS OF FLUID MECHANICS

Hundreds of dimensionless groups have been defined, particularly in chemical engineering and fluid dynamics.<sup>[10]</sup> The present collection includes only a small subset of the latter category.

Notation is as follows:

c	speed of sound
$c_p$	specific heat at constant pressure (units $\text{cm}^2/\text{sec}^2 - \text{deg}$ )
g	gravitational acceleration
H	vertical length scale
$k = \kappa \rho c_p$	thermal conductivity (units $\text{g-cm}/\text{sec}^3\text{-deg}$ )
L	horizontal* length scale
n	frequency of solid-body oscillation
r	radius of curvature of a pipe or channel
R	radius of pipe
$\alpha$	Newton's law coefficient of heat transfer (units $\text{g}/\text{sec}^3\text{-deg}$ )
$\beta$	volumetric expansion coefficient, $dV/V = \beta dT$
$\Delta T$	applied temperature difference
$\kappa$	thermal diffusivity (units $\text{cm}^2/\text{sec}$ )
$\lambda$	collisional mean free path
$\mu = \rho \nu$	bulk viscosity
$\nu$	kinematic viscosity
$\Omega$	rotational velocity
$\rho$	mass density

Name	Symbol	Definition	Significance
Burgers	B	$(\beta g \Delta T)^{\frac{1}{2}} / 2\Omega H^{\frac{1}{2}}$	effect of stratification/effect of rotation
Dean	D	$(UR/\nu)(R/r)^{\frac{1}{2}}$	transverse circulation due to curvature/longitudinal flow speed
Eckert	E	$U^2/c_p \Delta T$	adiabatic temperature increase/applied temperature difference
Ekman	Ek	$\nu/\Omega H^2$	viscous forces/coriolis forces
Froude	Fr	$\Omega^2 L/g$	centrifugal forces/gravitational forces
Grashof	G	$g\beta\Delta T L^3/\nu^2$	effects of bouyancy/viscous effects
Knudsen	Kn	$\lambda/L$	hydrodynamic time scale/collisional time scale
Mach	M	$U/c$	directed motion/thermal motion
Nusselt	N	$hL/k$	total thermal transfer/thermal conduction
Péclet	Pe	$UL/\kappa$	thermal diffusion time scale/free streaming time scale
Prandtl	Pr	$\nu/\kappa$	momentum diffusion rate/heat diffusion rate
Rayleigh	Ra	$\beta g \Delta T H^3/\kappa$	bouyancy forces/diffusive forces
Reynolds	Re	$UL/\nu$	inertial forces/viscous forces
Richardson	Ri	$\beta g \Delta T H/U^2$	effects of bouyancy/effects of vertical shear
Rossby	Ro	$U/\Omega L$	inertial forces/coriolis forces
Stanton	$S_x$	$\alpha/\rho c_p U$	thermal convection/compressional heating
Strouhal	S	$nL/U$	oscillatory motion/directed motion

## SHOCKS

Ideal gas jump conditions for density  $\rho$ , velocity  $u$ , pressure  $p$ , temperature  $T$ :

$$\frac{\rho_2}{\rho_1} = \frac{u_1}{u_2} = \frac{(\gamma + 1) M_1^2}{(\gamma - 1) M_1^2 + 2}; \quad \frac{p_2}{p_1} = \frac{2 \gamma M_1^2 - \gamma + 1}{\gamma + 1};$$

$$\frac{T_2}{T_1} = \frac{\left[ \frac{2 \gamma M_1^2 - \gamma + 1}{\gamma + 1} \right] \left[ \frac{(\gamma - 1) M_1^2 + 2}{(\gamma + 1)^2 M_1^2} \right]}{(\gamma + 1)^2 M_1^2},$$

where 1, 2 denote quantities in front of and behind the shock respectively in the shock frame;  $M_1 = u_1 / c_{s1} =$  Mach number in region 1,  $c_s^2 = \gamma kT / \rho$ ;  $\gamma =$  ratio of specific heats  $= (f + 2) / f$ , where  $f$  is the number of degrees of freedom. Entropy change across the shock is

$$\Delta S = S_2 - S_1 = c_v \ln \left[ \frac{p_2}{p_1} \left( \frac{\rho_1}{\rho_2} \right)^\gamma \right],$$

where  $c_v$  is the specific heat at constant volume. For weak shocks,

$$\Delta S \rightarrow c_v \frac{2 \gamma (\gamma - 1)}{3 (\gamma + 1)^2} (M_1^2 - 1)^3.$$

Evidently in the limit  $M_1 \rightarrow \infty$ ,  $p_2 / \rho_1 \rightarrow (\gamma + 1) / (\gamma - 1)$  ( $= 4$  for  $\gamma = 5/3$ ), while the temperature and pressure jumps become infinite.

PLASMA DISPERSION FUNCTION [11]

Definition

$$Z(\zeta) = \pi^{-\frac{1}{2}} \int_{-\infty}^{\infty} dt \exp(-t^2)/(t - \zeta) \quad (\text{Im}\zeta > 0)$$

$$= 2i \exp(-\zeta^2) \int_{-\infty}^{i\zeta} dt \exp(-t^2) .$$

Physically  $\zeta = x + iy$  is the ratio of phase to thermal velocity.

Differential equation

$$\frac{dZ}{d\zeta} = -2 [1 + \zeta Z] , \quad Z(0) = i\pi^{\frac{1}{2}} ;$$

$$\frac{d^2Z}{d\zeta^2} + 2\zeta \frac{dZ}{d\zeta} + 2Z = 0 .$$

Real argument ( $y = 0$ )

$$Z(x) = i\pi^{\frac{1}{2}} \exp(-x^2) - 2x Y(x) ,$$

$$Y(x) = x^{-1} \exp(-x^2) \int_0^x dt \exp(t^2) .$$

Imaginary argument

$$Z(iy) = i\pi^{\frac{1}{2}} \exp(y^2) [1 - \text{erf}(y)] = i\pi^{\frac{1}{2}} \exp(y^2) \text{erfc}(y)$$

Power series (small argument)

$$Z(\zeta) = i\pi^{\frac{1}{2}} \exp(-\zeta^2) - 2\zeta \left[ 1 - 2\zeta^2/3 + 4\zeta^4/15 - 8\zeta^6/105 + \dots \right]$$

Asymptotic series (large argument,  $x > 0$ ) <sup>[12]</sup>

$$Z(\zeta) = i\pi^{\frac{1}{2}} \sigma \exp(-\zeta^2) - \zeta^{-1} \left[ 1 + 1/2 \zeta^2 + 3/4 \zeta^4 + 15/8 \zeta^6 + \dots \right]$$

$$\sigma = \begin{cases} 0 & y > 1/x \\ 1 & |y| < 1/x \\ 2 & y < -1/x \end{cases}$$

Symmetry properties ( $\zeta^* = x - iy$ )

$$Z(\zeta^*) = - \left[ Z(-\zeta) \right]^*$$

$$\text{For } y > 0, \quad Z(x - iy) = \left[ Z(x + iy) \right]^* + 2 i\pi^{\frac{1}{2}} \exp[-(x - iy)^2]$$

Two-pole approximations <sup>[13]</sup>

For  $\zeta$  in upper half plane (good except when  $y < \pi^{\frac{1}{2}} x^2 e^{-x^2}$ ,  $x \gg 1$ )

$$Z(\zeta) \approx \frac{0.50 + 0.81 i}{a - \zeta} - \frac{0.50 - 0.81 i}{a^* - \zeta} ;$$

$$Z'(\zeta) \approx \frac{0.50 + 0.86 i}{(b - \zeta)^2} + \frac{0.50 - 0.86 i}{(b^* - \zeta)^2} ;$$

$$a = 0.51 - 0.81 i ,$$

$$b = 0.48 - 0.91 i .$$

## FUNDAMENTAL PLASMA PARAMETERS

All quantities are in Gaussian units except temperature ( $T_e, T_i, T$ ) expressed in eV and ion mass ( $m_i$ ) expressed in units of proton mass,  $\mu = m_i/m_p$ ;  $Z$  is charge state;  $k$  is Boltzmann's constant;  $K$  is wavenumber.

### Frequencies

electron gyrofrequency

$$f_{ce} = \omega_{ce}/2\pi = 2.80 \times 10^9 B \quad \text{Hz}$$

$$\omega_{ce} = eB/m_e c = 1.76 \times 10^7 B \quad \text{rad/sec}$$

ion gyrofrequency

$$f_{ci} = \omega_{ci}/2\pi = 1.52 \times 10^3 Z\mu^{-1} B \quad \text{Hz}$$

$$\omega_{ci} = eB/m_i c = 9.58 \times 10^3 Z\mu^{-1} B \quad \text{rad/sec}$$

hybrid gyrofrequency

$$f_{LH} = \omega_{LH}/2\pi = 6.53 \times 10^4 Z^{\frac{1}{2}} \mu^{-\frac{1}{2}} B \quad \text{Hz}$$

$$\begin{aligned} \omega_{LH} &= (\omega_{ce} \omega_{ci})^{\frac{1}{2}} \\ &= 4.10 \times 10^5 Z^{\frac{1}{2}} \mu^{-\frac{1}{2}} B \quad \text{rad/sec} \end{aligned}$$

electron plasma frequency

$$f_{pe} = \omega_{pe}/2\pi = 8.98 \times 10^3 n_e^{\frac{1}{2}} \quad \text{Hz}$$

$$\begin{aligned} \omega_{pe} &= (4\pi n_e e^2/m_e)^{\frac{1}{2}} \\ &= 5.64 \times 10^4 n_e^{\frac{1}{2}} \quad \text{rad/sec} \end{aligned}$$

ion plasma frequency

$$f_{pi} = \omega_{pi}/2\pi = 2.10 \times 10^2 Z\mu^{-\frac{1}{2}} n_i^{\frac{1}{2}} \quad \text{Hz}$$

$$\begin{aligned} \omega_{pi} &= (4\pi n_i Z^2 e^2/m_i)^{\frac{1}{2}} \\ &= 1.32 \times 10^3 Z\mu^{-\frac{1}{2}} n_i^{\frac{1}{2}} \quad \text{rad/sec} \end{aligned}$$

electron trapping rate  $v_{Te} = \left( \frac{eKE}{m_e} \right)^{\frac{1}{2}} = 7.26 \times 10^8 K^{\frac{1}{2}} E^{\frac{1}{2}} \text{ sec}^{-1}$

ion trapping rate  $v_{Ti} = \left( \frac{eKE}{m_i} \right)^{\frac{1}{2}} = 1.69 \times 10^7 K^{\frac{1}{2}} E^{\frac{1}{2}} \mu^{-\frac{1}{2}} \text{ sec}^{-1}$

### Velocities

electron thermal velocity  $v_{Te} = (kT_e/m_e)^{\frac{1}{2}} = 4.19 \times 10^7 T_e^{\frac{1}{2}} \text{ cm/sec}$

ion thermal velocity  $v_{Ti} = (kT_i/m_i)^{\frac{1}{2}} = 9.79 \times 10^5 \mu^{-\frac{1}{2}} T_i^{\frac{1}{2}} \text{ cm/sec}$

ion sound velocity  $c_s = (kT_e/m_i)^{\frac{1}{2}} = 9.79 \times 10^5 \mu^{-\frac{1}{2}} T_e^{\frac{1}{2}} \text{ cm/sec}$

Alfvén velocity  $v_A = B/(4\pi n_i m_i)^{\frac{1}{2}}$   
 $= 2.18 \times 10^{11} \mu^{-\frac{1}{2}} n_i^{-\frac{1}{2}} B \text{ cm/sec}$

With this definition of thermal velocity, the Debye length and plasma frequency for a species  $j$  are connected by

$$\omega_{pj} \lambda_{Dj} = v_{Tj} ,$$

where  $\lambda_{Dj}$  is obtained by substituting  $T_j$  for  $T$  in the definition below.

Some individuals, however, prefer the definition

$$v'_{Tj} = (2 kT_j/m_j)^{\frac{1}{2}} ,$$

according to which the electron and ion thermal velocities and ion sound velocity become

$$v'_{Te} = (2 kT_e/m_e)^{\frac{1}{2}} = 5.93 \times 10^7 T_e^{\frac{1}{2}} \text{ cm/sec};$$

$$v'_{Ti} = (2 kT_i/m_i)^{\frac{1}{2}} = 1.38 \times 10^8 (T_i/\mu)^{\frac{1}{2}} \text{ cm/sec};$$

$$c'_s = (2 kT_e/m_i)^{\frac{1}{2}} = 1.38 \times 10^8 (T_e/\mu)^{\frac{1}{2}} \text{ cm/sec}.$$

The Debye length and gyroradii may or may not be redefined accordingly.

### Lengths

electron deBroglie length	$\lambda = h/(m_e kT_e)^{\frac{1}{2}} = 2.76 \times 10^{-8} T_e^{-\frac{1}{2}} \text{ cm}$
classical distance of minimum approach	$e^2/kT = 1.44 \times 10^{-7} T^{-1} \text{ cm}$
electron gyroradius	$r_e = v_{Te}/\omega_{ce} = 2.38 T_e^{\frac{1}{2}} B^{-1} \text{ cm}$
ion gyroradius	$r_i = v_{Ti}/\omega_{ci}$ $= 1.02 \times 10^2 \mu^{\frac{1}{2}} Z^{-\frac{1}{2}} T_i^{\frac{1}{2}} B^{-1} \text{ cm}$
plasma skin depth	$c/\omega_{pe} = 5.31 \times 10^5 n^{-\frac{1}{2}} \text{ cm}$
Debye length	$\lambda_D = (kT/4\pi n e^2)^{\frac{1}{2}} = 7.43 \times 10^2 T^{\frac{1}{2}} n^{-\frac{1}{2}} \text{ cm}$

### Dimensionless

(electron/proton mass ratio)<sup>1/2</sup>  $(m_e/m_p)^{\frac{1}{2}} = 2.33 \times 10^{-2} = 1/42.9$

number of particles in Debye  
sphere  $\frac{4\pi}{3} n \lambda_D^3 = 1.72 \times 10^9 T^{3/2} n^{-\frac{1}{2}}$

Alfvén velocity/speed of light	$v_A/c = 7.28 \mu^{-\frac{1}{2}} n_i^{-\frac{1}{2}} B$
magnetic/ion rest energy ratio	$B^2/8\pi n_i m_i c^2 = 26.5 \mu^{-1} n_i^{-1} B^2$
electron plasma/gyrofrequency ratio	$\omega_{pe}/\omega_{ce} = 3.21 \times 10^{-3} n_e^{\frac{1}{2}} B^{-1}$
ion plasma/gyrofrequency ratio	$\omega_{pi}/\omega_{ci} = 0.137 \mu^{\frac{1}{2}} n_i^{\frac{1}{2}} B^{-1}$
thermal/magnetic energy ratio	$\beta = 8\pi nT/B^2 = 4.03 \times 10^{-11} nT B^{-2}$

#### Miscellaneous

Bohm diffusion coefficient  $D_B = \frac{ckT}{16 eB} = 6.25 \times 10^{-1} TB^{-1} \text{ cm}^2 \text{ sec}^{-1}$

Spitzer resistivity  
(transverse)  $\eta_{\perp} = 1.15 \times 10^{-14} Z \ln \Lambda T^{-3/2} \text{ sec}$   
 $= 1.03 \times 10^{-2} Z \ln \Lambda T^{-3/2} \text{ ohm-cm}$

The anomalous collision rate due to low frequency ion sound turbulence is <sup>[14]</sup>

$$v^* \approx \omega_{pe} W/kT = 5.64 \times 10^4 n^{-1/2} W/kT \text{ sec}^{-1},$$

where  $W$  is the total energy of waves with  $\omega/k < v_{Ti}$ .

Magnetic pressure is given by

$$P = 3.98 \times 10^{-2} B^2 \text{ dyne/cm}^2,$$

when  $P$  and  $B$  are expressed in cgs units. Thus the pressure equivalent of 10 kG is

$$P(10 \text{ kG}) = 3.98 \times 10^2 \text{ dyne/cm}^2 = 3.93 \text{ atm.}$$

APPROXIMATE MAGNITUDES IN SOME TYPICAL PLASMAS

Plasma Type	$n \text{ cm}^{-3}$	T eV	$\omega_{pe} \text{ sec}^{-1}$	$\lambda_D \text{ cm}$	$n^3_D$	$\nu_{ei} \text{ sec}^{-1}$
Interstellar gas	1	1	$6 \times 10^4$	$7 \times 10^2$	$4 \times 10^6$	$7 \times 10^{-5}$
Gaseous nebula	$10^3$	1	$2 \times 10^3$	20	$10^7$	$6 \times 10^{-2}$
Solar corona	$10^8$	$10^2$	$6 \times 10^7$	7	$4 \times 10^5$	$6 \times 10^{-2}$
Diffuse hot plasma	$10^{12}$	$10^2$	$6 \times 10^{10}$	$7 \times 10^{-3}$	$4 \times 10^5$	40
Solar atmosphere, gas discharge	$10^{14}$	1	$6 \times 10^{11}$	$7 \times 10^{-5}$	40	$2 \times 10^2$
Warm plasma	$10^{14}$	10	$6 \times 10^{11}$	$2 \times 10^{-4}$	$10^9$	$10^7$
Hot plasma	$10^{14}$	$10^2$	$6 \times 10^{11}$	$7 \times 10^{-4}$	$4 \times 10^3$	$4 \times 10^3$
Thermonuclear plasma	$10^{15}$	$10^4$	$2 \times 10^{12}$	$2 \times 10^{-3}$	$10^7$	$5 \times 10^4$
Theta pinch	$10^{13}$	$10^2$	$6 \times 10^{12}$	$7 \times 10^{-5}$	$4 \times 10^3$	$3 \times 10^6$
Dense hot plasma	$10^{15}$	$10^2$	$6 \times 10^{13}$	$7 \times 10^{-3}$	$4 \times 10^2$	$2 \times 10^{10}$
Laser plasma	$10^{20}$	$10^2$	$6 \times 10^{14}$	$7 \times 10^{-7}$	40	$2 \times 10^{12}$




Fig. 1 - Information of a similar nature is depicted graphically.<sup>15</sup>

## COLLISIONS AND TRANSPORT

Temperatures are in eV; the corresponding value of Boltzmann's constant is  $k = 1.60 \times 10^{-12}$  erg/eV; masses  $\mu$ ,  $\mu'$  are in units of the proton mass;  $e_\alpha = Z_\alpha e$  is the charge of species  $\alpha$ . All other units are cgs except where noted.

### Relaxation rates [18]

Rates are associated with four relaxation processes arising from the interaction of test particles (labeled  $\alpha$ ) streaming through a background of field particles (labeled  $\beta$ ):

slowing down  $\frac{dv_\alpha}{dt} = -v_s^{\alpha/\beta} v_\alpha$  ;

transverse diffusion  $\frac{d}{dt} (v_\alpha - \bar{v}_\alpha)_\perp^2 = v_\perp^{\alpha/\beta} v_\alpha^2$  ;

parallel diffusion  $\frac{d}{dt} (v_\alpha - \bar{v}_\alpha)_\parallel^2 = v_\parallel^{\alpha/\beta} v_\alpha^2$  ;

energy loss  $\frac{d}{dt} v_\alpha^2 = -v_e^{\alpha/\beta} v_\alpha^2$  ,

where the averages are performed over an ensemble of test particles and a Maxwellian field particle distribution.

The exact formulas may be written

$$v_s^{\alpha/\beta} = (1 + m_\alpha/m_\beta) \psi(x^{\alpha/\beta}) v_o^{\alpha/\beta} \text{ sec}^{-1} ;$$

$$v_\perp^{\alpha/\beta} = 2 \left[ \psi(x^{\alpha/\beta}) (1 - 1/2x^{\alpha/\beta}) + \psi'(x^{\alpha/\beta}) \right] v_o^{\alpha/\beta} \text{ sec}^{-1} ;$$

$$v_{\parallel}^{\alpha/\beta} = \left[ \psi(x^{\alpha/\beta}) / x^{\alpha/\beta} \right] v_o^{\alpha/\beta} \text{ sec}^{-1} ;$$

$$v_e^{\alpha/\beta} = 2 \left[ (m_\alpha/m_\beta) \psi(x^{\alpha/\beta}) - \psi'(x^{\alpha/\beta}) \right] v_o^{\alpha/\beta} \text{ sec}^{-1} ,$$

where

$$v_o^{\alpha/\beta} = 4\pi e_\alpha^2 e_\beta^2 \lambda_{\alpha\beta} n_\beta / m_\alpha^2 v_\alpha^3 \text{ sec}^{-1} ;$$

$$x^{\alpha/\beta} = \frac{1}{2} m_\beta v_\alpha^2 / kT_\beta ;$$

$$\psi(x) = \frac{2}{\sqrt{\pi}} \int_0^x dt \sqrt{t} e^{-t} ;$$

$$\psi'(x) = d\psi/dx ,$$

and  $\lambda_{\alpha\beta} = \ln \Lambda_{\alpha\beta}$  is the Coulomb logarithm (see below). The limiting forms of  $v_s$ ,  $v_\perp$  and  $v_{\parallel}$  are given in the following table. The two expressions shown for each rate hold for very slow ( $x^{\alpha/\beta} \ll 1$ ) and very fast ( $x^{\alpha/\beta} \gg 1$ ) test particles, respectively. Test particle energy  $\epsilon$  and field particle temperature  $T$  are both in eV;  $\mu = m_i/m_p$ , where  $m_p$  is the proton mass; for electron-electron and ion-ion encounters, field particle quantities are distinguished by a prime.

### Electron-electron

$$v_s^{e/e'} / n_e \lambda_{ee'} \approx 5.8 \times 10^{-6} T^{-3/2} \rightarrow 7.7 \times 10^{-6} \epsilon^{-3/2}$$

$$v_{\perp}^{e/e'} / n_e \lambda_{ee'} \approx 5.8 \times 10^{-6} T^{-1/2} \epsilon^{-1} \rightarrow 7.7 \times 10^{-6} \epsilon^{-3/2}$$

$$v_{\parallel}^{e/e'} / n_e \lambda_{ee'} \approx 2.9 \times 10^{-6} T^{-1/2} \epsilon^{-1} \rightarrow 3.9 \times 10^{-6} T \epsilon^{-5/2}$$

### Electron-ion

$$v_s^{e/i} / n_i Z^2 \lambda_{ei} \approx 0.23 \mu^{3/2} T^{-3/2} \rightarrow 3.9 \times 10^{-6} \epsilon^{-3/2}$$

$$v_{\perp}^{e/i} / n_i Z^2 \lambda_{ie} \approx 2.5 \times 10^{-4} \mu^{1/2} T^{-1/2} \epsilon^{-1} \rightarrow 7.7 \times 10^{-6} \epsilon^{-3/2}$$

$$v_{\parallel}^{e/i} / n_i Z^2 \lambda_{ie} \approx 1.2 \times 10^{-4} \mu^{1/2} T^{-1/2} \epsilon^{-1} \rightarrow 2.1 \times 10^{-9} \mu^{-1} T \epsilon^{-5/2}$$

### Ion-electron

$$v_s^{i/e} / n_e \lambda_{ie} \approx 1.6 \times 10^{-9} \mu^{-1} T^{-3/2} \rightarrow 1.7 \times 10^{-4} \mu^{1/2} \epsilon^{-3/2}$$

$$v_{\perp}^{i/e} / n_e \lambda_{ie} \approx 3.2 \times 10^{-9} \mu^{-1} T^{-1/2} \epsilon^{-1} \rightarrow 1.8 \times 10^{-7} \mu^{-1/2} \epsilon^{-3/2}$$

$$v_{\parallel}^{i/e} / n_e \lambda_{ie} \approx 1.6 \times 10^{-9} \mu^{-1} T^{-1/2} \epsilon^{-1} \rightarrow 1.7 \times 10^{-4} \mu^{1/2} T \epsilon^{-5/2}$$

Ion-ion

$$v_s^{i/i'} / n_i Z^2 Z'^2 \lambda_{ii'} \approx 6.8 \times 10^{-8} \frac{u'^{1/2}}{u} \left(1 + \frac{u'}{u}\right) T^{-3/2}$$

$$\rightarrow 9.0 \times 10^{-8} u^{-1/2} \left(1 + \frac{u}{u'}\right) e^{-3/2}$$

$$v_{\perp}^{i/i'} / n_i Z^2 Z'^2 \lambda_{ii'} \approx 1.4 \times 10^{-7} u'^{1/2} u^{-1} T^{-1/2} e^{-1}$$

$$\rightarrow 1.8 \times 10^{-7} u^{-1/2} e^{-3/2}$$

$$v_{\parallel}^{i/i'} / n_i Z^2 Z'^2 \lambda_{ii'} \approx 6.8 \times 10^{-8} u'^{1/2} u^{-1} T^{-1/2} e^{-1}$$

$$\rightarrow 9.0 \times 10^{-8} u^{1/2} u'^{-1} T e^{-5/2}$$

In the same limits, the energy transfer rate follows from the identity

$$v_e = 2 v_s - v_{\perp} - v_{\parallel},$$

except for the case of fast electrons or fast ions scattered by ions, where the leading terms cancel. Here the appropriate forms are

$$v_e^{e/i} \rightarrow 4.2 \times 10^{-9} n_i Z^2 \lambda_{ei} \left[ e^{-3/2} u^{-1} - 8.9 \times 10^4 (u/T)^{1/2} e^{-1} \exp(-1836 u_s/T) \right] \text{ sec}^{-1}$$

and

$$v_e^{i/i'} \rightarrow 1.8 \times 10^{-7} n_i Z^2 Z'^2 \lambda_{ii'} \left[ e^{-\epsilon/\epsilon_{\alpha}^*} \epsilon^{1/2} / u' - 1.1 (u'/T)^{1/2} \epsilon^{-1} \exp(-u'\epsilon/T) \right] \text{ sec}^{-1} .$$

In general, the energy transfer rate  $v_e^{\alpha/\beta}$  is positive for  $\epsilon > \epsilon_{\alpha}^*$  and negative for  $\epsilon < \epsilon_{\alpha}^*$ , where  $x^* = (m_{\beta}/m_{\alpha}) \epsilon_{\alpha}^*/T_{\beta}$  is the solution of  $m_{\beta}/m_{\alpha} = \psi(x^*)\psi'(x^*)$ .

The ratio  $\epsilon_{\alpha}^*/T_{\beta}$  is given for a number of specific  $\alpha, \beta$  in the following table:

$\alpha/\beta$	i/e	e/e	i/i	e/p	e/D	e/T, e/He <sup>3</sup>	e/He <sup>4</sup>
$\frac{\epsilon_{\alpha}^*}{T_{\beta}}$	1.5	.98	.98	$4.8 \times 10^{-3}$	$2.6 \times 10^{-3}$	$1.8 \times 10^{-3}$	$1.4 \times 10^{-3}$

The collision rates for various types of encounter are represented graphically in Fig. 2. [17]




Fig. 2 - Test particle showing-down rates  $\nu_s^{\alpha/\beta}$ , scaled by the ion-electron rate  $\nu_s^{i/e}(0)$  in the limit of zero relative velocity  $v_t$ , as a function of  $v_t$  divided by the electron thermal velocity.

### Thermal equilibration

If the components of a plasma have different temperatures, but no relative drift, equilibration is described by

$$\frac{dT_\alpha}{dt} = \sum_{\beta} \bar{v}_e^{\alpha/\beta} (T_\beta - T_\alpha) ,$$

where

$$\bar{v}_e^{\alpha/\beta} = 1.8 \times 10^{-13} \frac{(m_\alpha m_\beta)^{1/2} Z_\alpha^2 Z_\beta^2 n_\alpha \lambda_{\alpha\beta}}{(m_\alpha T_\beta + m_\beta T_\alpha)^{3/2}} \text{ sec}^{-1} .$$

For electrons and ions with  $T_e \sim T_i = T$ , this implies

$$\frac{\bar{v}_e^{e/i}}{n_i} = \frac{\bar{v}_e^{i/e}}{n_e} = 3.2 \times 10^{-9} \frac{Z^2 \lambda}{\mu T^{3/2}} \text{ cm}^3/\text{sec} .$$

### Temperature anisotropy

Isotropization is described by

$$\frac{dT_\perp}{dt} = -\left(\frac{1}{2}\right) \frac{dT_\parallel}{dt} = -v_T^\alpha (T_\perp - T_\parallel) ,$$

where, if  $A = T_\perp/T_\parallel - 1 > 0$ ,

$$v_T^\alpha = \frac{2\sqrt{\pi} e^4 n_\alpha \lambda}{m_\alpha^{1/2} (kT_\parallel)^{3/2}} A^{-2} \left[ -3 + (A+3) \frac{\tan^{-1} A^{1/2}}{A^{1/2}} \right] \text{ sec}^{-1} ,$$

if  $A < 0$ ,  $\tan^{-1} A^{1/2}/A^{1/2}$  is replaced by  $\tanh^{-1} (-A)^{1/2}/(-A)^{1/2}$ .

For  $T_{\perp} \approx T_{\parallel} = T$ ,

$$v_T^e = 8.2 \times 10^{-7} n \lambda T^{-3/2} \quad \text{sec}^{-1} ;$$

$$v_T^i = 1.9 \times 10^{-8} n \lambda Z^2 / \omega T^{3/2} \quad \text{sec}^{-1} .$$

### Coulomb logarithm

For test particles of mass  $m_{\alpha}$ , charge  $e_{\alpha} = Z_{\alpha} e$ , scattering off field particles of mass  $m_{\beta}$ , charge  $e_{\beta} = Z_{\beta} e$ , the Coulomb logarithm is defined as  $\lambda \equiv \ln \Lambda = \ln(r_{\max}/r_{\min})$ . Here  $r_{\min}$  is the larger of  $e_{\alpha} e_{\beta} / (m_{\alpha\beta} \bar{u}^2)$  and  $\hbar / (2 m_{\alpha\beta} \bar{u})$ , averaged over both particle velocity distributions, where  $m_{\alpha\beta} = m_{\alpha} m_{\beta} / (m_{\alpha} + m_{\beta})$  and  $\bar{u} = v_{\alpha} - v_{\beta}$ ;

$r_{\max} = (4\pi \sum n_{\nu} e_{\nu}^2 / kT_{\nu})^{-1/2}$ , where the summation extends over all species  $\nu$  for which  $\bar{u}^2 < v_{T\nu}^2$ , with  $v_{T\nu} = (kT_{\nu}/m_{\nu})^{1/2}$ . If this inequality cannot be satisfied or if either  $\bar{u} \omega_{c\alpha}^{-1} < r_{\max}$  or  $\bar{u} \omega_{c\beta}^{-1} < r_{\max}$ , the theory breaks down. Typically  $\lambda \approx 10 - 20$ . Corrections to the transport coefficients are  $O(\lambda^{-1})$ , hence the theory is good only to  $\sim 10\%$  and fails when  $\lambda \sim 1$ .

The following cases are of particular interest.

(a) Thermal electron-electron collisions

$$\lambda_{ee} = 23 - \ln(n_e^{1/2} T_e^{-3/2}) \quad T_e \leq 10 \text{ eV}$$

$$= 24 - \ln(n_e^{1/2} T_e^{-1}) \quad T_e \geq 10 \text{ eV}$$

(b) Electron-ion collisions

$$\lambda_{ei} = \lambda_{ie} = 23 - \ln(n_e^{1/2} Z T_e^{-3/2}), \quad 10 \text{ eV} > T_e > T_i m_e / m_i ;$$

$$= 24 - \ln(n_e^{1/2} T_e^{-1}), \quad T_e > 10 \text{ eV} > T_i m_e / m_i ;$$

$$= 30 - \ln(n_i^{1/2} T_i^{-3/2} Z^2 u^{-1}), \quad T_i > T_e m_i / m_e .$$

(c) Mixed ion-ion collisions

$$\lambda_{ii'} = \lambda_{i'i} = 23 - \ln \left[ \frac{ZZ'(u+u')}{uT_i' + u'T_i} \left( \frac{n_i Z^2}{T_i} + \frac{n_i' Z'^2}{T_i'} \right)^{1/2} \right]$$

(d) Counterstreaming ions (relative velocity  $v_D = \beta_D c$ ) in the presence of warm electrons,  $kT_e/m_e > v_D^2 > kT_i/m_i, kT_i'/m_i',$

$$\lambda_{ii'} = \lambda_{i'i} = 35 - \ln \left[ \frac{ZZ'(u+u')}{u u' \beta_D^2} \left( \frac{n_e}{T_e} \right)^{1/2} \right] .$$

### Examples

1. Find the slowing-down and energy-transfer rates for a 100 keV deuteron beam in a plasma consisting of 1 keV electrons and 100 eV tritons, each having number density  $10^{14} \text{ cm}^{-3}$ .

Answer: The Coulomb logarithms are

$$\begin{aligned}\lambda_{De} &= 24 - \ln \left[ (10^{14})^{1/2} (10^3)^{-1} \right] \\ &= 15\end{aligned}$$

from formula (b), and

$$\begin{aligned}\lambda_{DT} &= 35 - \ln \left[ \frac{(5/6)}{(2.2 \times 10^8/3 \times 10^{10})^2} \left( \frac{10^{14}}{10^2} \right)^{1/2} \right] \\ &= 12\end{aligned}$$

from formula (d). The beam velocity  $v_D = 2.2 \times 10^8 \text{ cm/sec}$  satisfies  $\bar{v}_T \ll v_D \ll \bar{v}_e$ , where  $\bar{v}_T$  and  $\bar{v}_e$  are the respective thermal velocities. It is therefore appropriate to use the first form for  $v_s^{D/e}$  and the second for  $v_s^{D/T}$ . Hence

$$\begin{aligned}v_s^{D/e} &= (1.6 \times 10^{-9})(2)^{-1}(10^3)^{-3/2}(10^{14})(15) \\ &= 37 \text{ sec}^{-1},\end{aligned}$$

and

$$\begin{aligned}v_s^{D/T} &= (9 \times 10^{-5})(2)^{-1/2}(1 + 2/3)(10^5)^{-3/2}(10^{14})(12) \\ &= 3.9 \text{ sec}^{-1}\end{aligned}$$

The net rate of transfer of energy to the electrons is

$$\nu_e^{D/e} = 2 \nu_s^{D/e} - \nu_{\perp}^{D/e} - \nu_{\parallel}^{D/e} = \nu_s^{D/e} = 37 \text{ sec}^{-1},$$

while the corresponding rate for tritons follows from the second formula for fast ions:

$$\begin{aligned} \nu_e^{D/T} &= (1.8 \times 10^{-7})(10^{14})(12) \left[ (10^5)^{-3/2} (2)^{1/2} (3)^{-1} \right. \\ &\quad \left. - (1.1)(3/10^2)^{1/2} (10^5)^{-1} \exp(-3 \times 10^5/10^2) \right] \\ &= 3.1 \text{ sec}^{-1}. \end{aligned}$$

Evidently, for the parameters given, electron collisions are about an order of magnitude more effective in both processes.

2. A hydrogen plasma has  $T_e = 500 \text{ eV}$ ,  $T_i = 100 \text{ eV}$  and  $n = 10^{10} \text{ cm}^{-3}$ . If it contains an admixture of 1%  $\text{Li}^7 (Z = 3)$  ions with  $T_{\text{Li}} = 25 \text{ eV}$ , is the net effect of collisions to heat or to cool the  $\text{H}^+$  ions? What happens if  $T_e = 5 \text{ keV}$  is used?

Answer: The Coulomb logarithms are

$$\begin{aligned} \lambda_{\text{H}^+ \text{e}}^+ &= 24 - \ln \left[ (10^{10})^{1/2} (500)^{-1/2} \right] \\ &= 16, \end{aligned}$$

and

$$\begin{aligned} \lambda_{\text{H}^+ \text{Li}^{+++}}^+ &= 23 - \ln \left\{ \frac{(3)(8)}{25 + (7)(100)} \left[ \frac{10^{10}}{100} \right. \right. \\ &\quad \left. \left. + \frac{(10^8)(9)}{25} \right] \right\} = 8.0 \end{aligned}$$

from (b) and (c), respectively. The thermal equilibration rates are thus

$$\begin{aligned} \bar{\nu}_e^{H^+/e} &= 3.2 \times 10^{-9} n \lambda_{H^+} T_e^{-3/2} \\ &= 4.5 \times 10^{-2} \text{ sec}^{-1}, \end{aligned}$$

and

$$\begin{aligned} \bar{\nu}_e^{H^+/Li^{+++}} &= \frac{1.8 \times 10^{-19} (7)^{1/2} (3^2) (10^{-2} n) \lambda_{H^+ Li^{+++}}}{(1.7 \times 10^{-24})^{1/2} [25 + (7)(100)]^{3/2}} \\ &= .12 \text{ sec}^{-1}. \end{aligned}$$

Hence

$$\begin{aligned} \frac{dT_{H^+}}{dt} &= \bar{\nu}_e^{H^+/Li^{+++}} [25 - 100] + \bar{\nu}_e^{H^+/e} [500 - 100] \\ &= -9 + 18 = 9 \text{ eV/sec} > 0. \end{aligned}$$

Evidently as  $T_e \rightarrow \infty$  the electron contribution to  $\frac{dT_{H^+}}{dt}$  vanishes  $\sim T_e^{-1/2}$

and for sufficiently large  $T_e$ ,  $\frac{dT_{H^+}}{dt} < 0$ . Thus with  $T_e = 5 \text{ keV}$ ,

$$\bar{\nu}_e^{H^+/e} = 1.4 \times 10^{-3} \text{ sec}^{-1}$$

and

$$\frac{dT}{dt} \frac{H^+}{H^+} = (.12) [25 - 100] + (1.4 \times 10^{-3}) [5000 - 100]$$

$$= - 2 \text{ eV/sec.}$$

Hence heating the electrons cools the  $H^+$  ions!

Fokker-Planck equation

$$\frac{Df^\alpha}{Dt} = \frac{\partial f^\alpha}{\partial t} + \underline{v} \cdot \underline{\nabla} f^\alpha + \underline{F} \cdot \underline{\nabla}_v f^\alpha = \left( \frac{\partial f^\alpha}{\partial t} \right)_{\text{coll}}$$

where  $\underline{F}$  is an external force field. The general form of the collision integral is  $\left( \frac{\partial f^\alpha}{\partial t} \right)_{\text{coll}} = - \sum_\beta \underline{\nabla}_v \cdot \underline{j}^{\alpha/\beta}$  with

$$\underline{j}^{\alpha/\beta} = 2\pi\lambda_{\alpha\beta} \frac{e_\alpha^2 e_\beta^2}{m_\alpha} \int d^3v' (u^2 \underline{I} - \underline{u} \underline{u}) u^{-3}$$

$$\left\{ \frac{1}{m_\beta} f^\alpha(\underline{v}) \underline{\nabla}_v f^\beta(\underline{v}') - \frac{1}{m_\alpha} f^\beta(\underline{v}') \underline{\nabla}_v f^\alpha(\underline{v}) \right\}$$

(Landau form) where  $\underline{u} = \underline{v}' - \underline{v}$  and  $\underline{I}$  is the unit dyad, or alternatively

$$\underline{j}^{\alpha/\beta} = 4\pi\lambda_{\alpha\beta} \frac{e_\alpha^2 e_\beta^2}{m_\alpha^2} \left\{ f^\alpha(\underline{v}) \underline{\nabla}_v H(\underline{v}) - \frac{1}{2} \underline{\nabla}_v \cdot \left[ f^\alpha(\underline{v}) \underline{\nabla}_v \underline{\nabla}_v G(\underline{v}) \right] \right\}$$

where the Rosenbluth potentials are

$$G(\underline{v}) = \int f^\beta(\underline{v}') u d^3 v'$$

$$H(\underline{v}) = \left( 1 + \frac{m_\alpha}{m_\beta} \right) \int f^\beta(\underline{v}') u^{-1} d^3 v' .$$

If species  $\alpha$  is a weak beam (number and energy density small compared with background) streaming through a Maxwellian plasma, then

$$\begin{aligned} \underline{J}^{\alpha/\beta} = & - v_s^{\alpha/\beta} \underline{v} f^\alpha - \frac{1}{2} v_\perp^{\alpha/\beta} v_\parallel \nabla_\parallel f^\alpha \\ & + \frac{1}{2} (v_\perp^{\alpha/\beta} - v_\parallel^{\alpha/\beta}) \underline{v} \cdot \nabla_\parallel f^\alpha . \end{aligned}$$

### Transport coefficients <sup>[18]</sup>

Transport equations for a multispecies plasma:

$$\frac{D^\alpha n_\alpha}{Dt} + n_\alpha \nabla \cdot \underline{v}_\alpha = 0 ;$$

$$m_\alpha n_\alpha \frac{D^\alpha \underline{v}_\alpha}{Dt} = - \nabla p_\alpha - \nabla \cdot \underline{P}_\alpha + Z_\alpha e n_\alpha \left[ \underline{E} + \frac{1}{c} \underline{v}_\alpha \times \underline{B} \right] + \underline{R}_\alpha ;$$

$$\frac{3}{2} n_\alpha k \frac{D^\alpha T_\alpha}{Dt} + p_\alpha \nabla \cdot \underline{v}_\alpha = - \nabla \cdot \underline{q}_\alpha - \underline{P}_\alpha : \nabla \underline{v}_\alpha + Q_\alpha ,$$

where  $D^\alpha/Dt = \partial/\partial t + \underline{v}_\alpha \cdot \nabla$  and  $p_\alpha = (3/2) n_\alpha kT_\alpha$ .  $R_\alpha = \sum_B R_{\alpha B}$  and  $Q_\alpha = \sum_B Q_{\alpha B}$

where  $R_{\alpha B}$  and  $Q_{\alpha B}$  are respectively the momentum and energy gained by the  $\alpha^{\text{th}}$  species through collisions with the  $B^{\text{th}}$ ,  $\underline{P}_\alpha$  is the stress tensor, and  $\underline{q}_\alpha$  is the heat flow.

The transport coefficients in a simple two-component (electrons and singly charged ions) plasma are tabulated below. Here  $\parallel$  and  $\perp$  refer to the direction of the magnetic field  $\underline{B} = b\hat{B}$ ;  $\underline{u} = \underline{v}_e - \underline{v}_i$  is the relative streaming velocity;  $n_e = n_i = n$ ;  $\underline{j} = -ne\underline{u}$  is the current;  $\omega_{ce} = 1.76 \times 10^7 B$  and  $\omega_{ci} = (m_e/m_i)\omega_{ce}$  are the electron and ion gyrofrequencies, respectively; and the basic collisional times are taken to be

$$\tau_e = \frac{3\sqrt{m_e}(kT_e)^{3/2}}{4\sqrt{2\pi}n\lambda e^4} = 3.44 \times 10^5 \frac{T_e^{3/2}}{n\lambda},$$

and

$$\tau_i = \frac{3\sqrt{m_i}(kT_i)^{3/2}}{4\sqrt{\pi}n\lambda e^4} = 2.09 \times 10^7 \frac{T_i^{3/2}}{n\lambda} u^{1/2}.$$

In the limit of large fields ( $\omega_{cj}\tau_j \gg 1$ ) the transport processes may be summarized as follows:

momentum transfer  $\underline{R}_{ei} = -\underline{R}_{ie} = \underline{R} = \underline{R}_u + \underline{R}_T$  ;

frictional force  $\underline{R}_u = ne(\underline{j}_\perp/\sigma_\perp + \underline{j}_\parallel/\sigma_\parallel)$  ;

conductivities  $\sigma_\parallel = 2.0 \sigma_\perp = 2.0 \frac{ne^2\tau_e}{m_e}$  ;

thermal force  $\underline{R}_T = -0.71nkT_e \nabla_{\parallel} T_e - \frac{3}{2} \frac{nk}{\omega_{ce} \tau_e} \underline{b} \times \nabla_{\perp} T_e$  ;

ion heating  $Q_i = 3 \frac{m_e}{m_i} \frac{nk}{\tau_e} (T_e - T_i)$  ;

electron heating  $Q_e = -Q_i - \underline{R} \cdot \underline{u}$  ;

ion heat flux  $\underline{q}_i = -\kappa_{\parallel}^i \nabla_{\parallel} kT_i - \kappa_{\perp}^i \nabla_{\perp} kT_i + \kappa_{\wedge}^i \underline{b} \times \nabla_{\perp} kT_i$  ;

ion thermal conductivities  $\kappa_{\parallel}^i = 3.9 \frac{nkT_i \tau_i}{m_i}$  ;

$$\kappa_{\perp}^i = 2 \frac{nkT_i}{m_i \omega_{ci}^2 \tau_i} ;$$

$$\kappa_{\wedge}^i = -\frac{5}{2} \frac{nkT_i}{m_i \omega_{ci}} ;$$

electron heat flux  $\underline{q}^e = \underline{q}_{\perp}^e + \underline{q}_{\parallel}^e$  ;

frictional heat flux  $\underline{q}_{\perp}^e = 0.71nkT_e \nabla_{\parallel} T_e + \frac{3}{2} \frac{nkT_e}{\omega_{ce} \tau_e} \underline{b} \times \underline{u}_{\perp}$  ;

thermal gradient heat flux  $\underline{q}_{\parallel}^e = -\kappa_{\parallel}^e \nabla_{\parallel} kT_e - \kappa_{\perp}^e \nabla_{\perp} kT_e - \kappa_{\wedge}^e \underline{b} \times \nabla_{\perp} kT_e$  ;

electron thermal  
conductivities

$$\kappa_{\parallel}^e = 3.2 \frac{nkT_e \tau_e}{m_e} ;$$

$$\kappa_{\perp}^e = 4.7 \frac{nkT_e}{m_e \omega_{ce}^2 \tau_e} ;$$

$$\kappa_{\Lambda}^e = \frac{5}{2} \frac{nkT_e}{m_e \omega_{ce}} ;$$

stress tensor  
(both species)

$$P_{xx} = -\frac{\eta_0}{2} (W_{xx} + W_{yy}) - \frac{\eta_1}{2} (W_{xx} - W_{yy}) - \eta_3 W_{xy} ;$$

$$P_{yy} = -\frac{\eta_0}{2} (W_{xx} + W_{yy}) + \frac{\eta_1}{2} (W_{xx} - W_{yy}) + \eta_3 W_{xy} ;$$

$$P_{xy} = P_{yx} = -\eta_1 W_{xy} + \frac{\eta_3}{2} (W_{xx} - W_{yy}) ;$$

$$P_{xz} = P_{zx} = -\eta_2 W_{xz} - \eta_4 W_{yz} ;$$

$$P_{yz} = P_{zy} = -\eta_2 W_{yz} + \eta_4 W_{xz} ;$$

$$P_{zz} = -\eta_0 W_{zz}$$

(here the z axis is defined parallel to  $\underline{B}$ ) ;

ion viscosity  $\eta_0^i = 0.96 nkT_i \tau_i$  ;

$$\eta_1^i = \frac{3}{10} \frac{nkT_i}{\omega_{ci}^2 \tau_i} ;$$

$$\eta_2^i = \frac{6}{5} \frac{nkT_i}{\omega_{ci}^2 \tau_i} ;$$

$$\eta_3^i = -\frac{1}{2} \frac{nkT_i}{\omega_{ci}} ;$$

$$\eta_4^i = \frac{nkT_i}{\omega_{ci}} ;$$

electron viscosity  $\eta_0^e = 0.73 nkT_e \tau_e$  ;

$$\eta_1^e = 0.51 \frac{nkT_e}{\omega_{ce}^2 \tau_e} ;$$

$$\eta_2^e = 2.0 \frac{nkT_e}{\omega_{ce}^2 \tau_e} ;$$

$$\eta_3^e = -\frac{1}{2} \frac{nkT_e}{\omega_{ce}} ;$$

$$\eta_4^e = -\frac{nkT_e}{\omega_{ce}} .$$

For both species the rate-of-strain tensor is defined as

$$W_{jk} = \frac{\partial v_j}{\partial x_k} + \frac{\partial v_k}{\partial x_j} - \frac{2}{3} \delta_{jk} \nabla \cdot \underline{v} \quad .$$

When  $\underline{B} = 0$  the following simplifications occur:

$$\underline{R}_u = ne \underline{j} / \sigma_{\parallel} \quad ;$$

$$\underline{R}_T = - 0.71 n \nabla_{\parallel} kT_e \quad ;$$

$$\underline{q}_i = - \kappa_{\parallel}^i \nabla_{\parallel} kT_i \quad ;$$

$$\underline{q}_u^e = 0.71 n kT_e \underline{u} \quad ;$$

$$\underline{q}_T^e = - \kappa_{\parallel}^e \nabla_{\parallel} kT_e \quad ;$$

$$P_{jk} = - \eta_0 W_{jk} \quad ,$$

when  $\omega_{ce} \tau_e \gg 1 \gg \omega_{ci} \tau_i$ , the high-field expressions are obeyed by the electrons and the zero-field expressions by the ions.

Collisional transport theory is applicable when (1) macroscopic time rates of change satisfy  $\frac{d}{dt} \ll \frac{1}{\tau}$ , where  $\tau$  is the longest collisional time scale, and (in the absence of a magnetic field) (2) macroscopic length scales  $L$  satisfy  $L \gg \ell$ , where  $\ell = \bar{v} \tau$  is the mean

free path. In a strong field,  $\omega_{ce}\tau \gg 1$ , condition (2) is replaced by  $L_{\parallel} \gg \ell$  and  $L_{\perp} \gg \sqrt{\ell r_e}$  ( $L_{\perp} \gg r_e$  in a uniform field), where  $L_{\parallel}$  is a macroscopic scale parallel to the field  $\underline{B}$  and  $L_{\perp}$  is the smaller of  $(\nabla_{\perp} B/B)^{-1}$  and the transverse plasma dimension. In addition, the standard transport coefficients are valid only when (3) the Coulomb logarithm satisfies  $\lambda \gg 1$ ; (4) the electron gyroradius satisfies  $r_e \gg \lambda_D$ , or  $B^2 \ll 8\pi n_e m_e c^2$ ; (5) relative drifts  $\underline{u} = \underline{v}_{\alpha} - \underline{v}_{\beta}$  between two species are small compared with the thermal velocities,  $u \ll (kT_{\alpha}/m_{\alpha})^{1/2}, (kT_{\beta}/m_{\beta})^{1/2}$ , and (6) anomalous transport processes owing to microinstabilities are negligible.

#### Weakly Ionized Plasmas

Collision frequency for scattering of charged particles by neutrals

$$* \nu_{\alpha} = n_0 \sigma_{\alpha 0} (kT_{\alpha}/m_{\alpha})^{1/2} \quad \text{sec}^{-1} ,$$

where  $n_0$  is the neutral density and  $\sigma_{\alpha 0}$  is the cross-section, typically  $\sim 5 \times 10^{-15} \text{ cm}^2$  and weakly dependent on temperature.

When the system size  $L \ll \lambda_D$ , the charged particle diffusion coefficients are

$$D_{\alpha} = kT_{\alpha}/m_{\alpha} \nu_{\alpha} \quad \text{cm}^2/\text{sec} .$$

In the opposite limit, both species diffuse at the ambipolar rate

$$D_A = \frac{u_i D_e - u_e D_i}{u_i - u_e} = \frac{(T_i + T_e) D_i D_e}{T_i D_e + T_e D_i} \quad \text{cm}^2/\text{sec} ,$$

where  $\mu_\alpha = e_\alpha/m_\alpha v_\alpha$  is the mobility. The conductivity  $\sigma_\alpha$  satisfies  $\sigma_\alpha = n_\alpha e_\alpha \mu_\alpha$ .

In the presence of a magnetic field  $\underline{B}$ ,  $\underline{\mu}$  and  $\sigma$  become tensors

$$\underline{J}^\alpha = \underline{\sigma}^\alpha \cdot \underline{E} = \sigma_{\parallel}^\alpha \underline{E}_{\parallel} + \sigma_{\perp}^\alpha \underline{E}_{\perp} + \sigma_{\Lambda}^\alpha \underline{E} \times \underline{b} \quad ,$$

where  $\underline{b} = \underline{B}/B$  and

$$\sigma_{\parallel}^\alpha = n_\alpha e_\alpha^2 / m_\alpha v_\alpha \quad ;$$

$$\sigma_{\perp}^\alpha = \sigma_{\parallel}^\alpha v_\alpha^2 / (v_\alpha^2 + \omega_{c\alpha}^2) \quad ;$$

$$\sigma_{\Lambda}^\alpha = \sigma_{\parallel}^\alpha v_\alpha \omega_{c\alpha} / (v_\alpha^2 + \omega_{c\alpha}^2) \quad .$$

Here  $\sigma_{\Lambda}$  and  $\sigma_{\perp}$  are the Hall and Pedersen conductivities, respectively.

IONOSPHERIC PARAMETERS <sup>[13]</sup>

These are typical nighttime values. Where two numbers are entered, the first refers to the lower and the second to the upper portion of the layer.

Quantity	E Region	F Region
Number density ( $m^{-3}$ )	$1.5 \times 10^{10} - 3.0 \times 10^{10}$	$5 \times 10^{10} - 2 \times 10^{11}$
Height-integrated number density ( $m^{-2}$ )	$9 \times 10^{24}$	$4.5 \times 10^{25}$
Ion-neutral collision frequency ( $sec^{-1}$ )	$2 \times 10^3 - 10^2$	0.5 - .05
Ion gyro-/collision frequency ratio $\kappa_i$	.09 - 2.0	$4.6 \times 10^2 - 5.0 \times 10^3$
Ion Pedersen conductance $\kappa_i/(1 + \kappa_i^2)$	0.09 - 0.5	$2.2 \times 10^{-3} - 2 \times 10^{-4}$
Ion Hall conductance $\kappa_i^2/(1 + \kappa_i^2)$	$8 \times 10^{-4} - 0.8$	1.0
Electron-neutral collision frequency	$1.5 \times 10^4 - 9.0 \times 10^2$	80 - 10
Electron gyro-/collision frequency ratio $\kappa_e$	$4.1 \times 10^2 - 6.9 \times 10^3$	$7.8 \times 10^4 - 6.2 \times 10^5$
Electron Pedersen conductance $\kappa_e/(1 + \kappa_e^2)$	$2.7 \times 10^{-3} - 1.5 \times 10^{-4}$	$10^{-5} - 1.5 \times 10^{-3}$
Electron Hall conductance $\kappa_e^2/(1 + \kappa_e^2)$	1.0	1.0
Mean molecular weight	28 - 26	22 - 16
Ion gyrofrequency ( $sec^{-1}$ )	180 - 190	230 - 300*
Neutral diffusion coefficient ( $m^2/sec$ )	$30 - 5 \times 10^3$	$10^5$

The terrestrial magnetic field in the lower ionosphere at equatorial latitudes is approximately  $B_0 = .35 \times 10^{-4}$  tesla. The earth's radius is  $R_E = 4.0 \times 10^4/2\pi$  km  $\approx 6.4 \times 10^3$  km.

CTR<sup>[20]</sup>Natural abundance of deuterium  $n_D/n_H = 1.5 \times 10^{-4}$ 

Mass ratios











$$m_e/m_D = 2.72 \times 10^{-4} = 1/3670$$

$$(m_e/m_D)^{\frac{1}{2}} = 1.65 \times 10^{-2} = 1/60.6$$

$$m_e/m_T = 1.82 \times 10^{-4} = 1/5496$$

$$(m_e/m_T)^{\frac{1}{2}} = 1.35 \times 10^{-2} = 1/74.1$$

Fusion reactions (branching ratios are correct for energies near the cross-section peaks; a negative yield means the reaction is endothermic).<sup>[21]</sup>



The rates for the principal reactions, averaged over Maxwellian distributions, are [21, 22]

Temperature (keV)	$\overline{\sigma v}$ (cm <sup>3</sup> /sec)				
	D - D (1a) + (1b)	D - T (2)	D - He <sup>3</sup> (3)	T - T (4)	T - He <sup>3</sup> (5a) + (5b) + (5c)
1.0	$1.5 \times 10^{-22}$	$5.5 \times 10^{-21}$	$3 \times 10^{-26}$	$3.3 \times 10^{-22}$	$10^{-28}$
2.0	$5.4 \times 10^{-21}$	$2.6 \times 10^{-19}$	$1.4 \times 10^{-23}$	$7.1 \times 10^{-21}$	$10^{-25}$
5.0	$1.8 \times 10^{-19}$	$1.3 \times 10^{-17}$	$6.7 \times 10^{-21}$	$1.4 \times 10^{-19}$	$2.1 \times 10^{-22}$
10.0	$1.2 \times 10^{-18}$	$1.1 \times 10^{-16}$	$2.3 \times 10^{-19}$	$7.2 \times 10^{-19}$	$1.2 \times 10^{-20}$
20.0	$5.2 \times 10^{-18}$	$4.2 \times 10^{-16}$	$3.8 \times 10^{-18}$	$2.5 \times 10^{-18}$	$2.6 \times 10^{-19}$
50.0	$2.1 \times 10^{-17}$	$8.7 \times 10^{-16}$	$5.4 \times 10^{-17}$	$8.7 \times 10^{-18}$	$5.3 \times 10^{-18}$
100.0	$4.5 \times 10^{-17}$	$8.5 \times 10^{-16}$	$1.6 \times 10^{-16}$	$1.9 \times 10^{-17}$	$2.7 \times 10^{-17}$
200.0	$8.8 \times 10^{-17}$	$6.3 \times 10^{-16}$	$2.4 \times 10^{-16}$	$4.2 \times 10^{-17}$	$9.2 \times 10^{-17}$
500.0	$1.8 \times 10^{-16}$	$3.7 \times 10^{-16}$	$2.3 \times 10^{-16}$	$8.4 \times 10^{-17}$	$2.9 \times 10^{-16}$
1000.0	$2.2 \times 10^{-16}$	$2.7 \times 10^{-16}$	$1.8 \times 10^{-16}$	$8.0 \times 10^{-17}$	$5.2 \times 10^{-16}$

The total cross section as a function of E, the energy in keV of the incident particle [the first ion on the left side of Eqs. (1-5)], assuming the target ion at rest, can be fitted by [23]

$$\sigma_T(E) = \frac{A + \left[ \frac{(A - A_5 E)^2 + 1}{4} \right]^{-1/2}}{E \left[ \exp(A_1/\sqrt{E}) - 1 \right]}$$

where the Duane coefficients A<sub>j</sub> for the principal fusion reactions are as follows:

Coefficient (units)	Reaction (Eq. Number)					
	D - D (1a)	D - D (1b)	D - T (2)	D - He <sup>3</sup> (3)	T - T (4)	T - He <sup>3</sup> (5)
A <sub>1</sub> (keV) <sup>1/2</sup>	46.097	47.88	45.95	89.27	38.39	123.1
A <sub>2</sub> (keV-barn)	372	482	5.02 x 10 <sup>4</sup>	2.59 x 10 <sup>4</sup>	448	1.125 x 10 <sup>4</sup>
A <sub>3</sub> (keV) <sup>-1</sup>	4.35 x 10 <sup>-4</sup>	3.08 x 10 <sup>-3</sup>	1.368 x 10 <sup>-2</sup>	3.98 x 10 <sup>-3</sup>	1.02 x 10 <sup>-3</sup>	0
A <sub>4</sub>	1.220	1.177	1.076	1.297	2.09	0
A <sub>5</sub> (keV-barn)	0	0	4.09	647	0	0

For low energies ( $T \leq 25$  keV) the data may be represented by

$$(\overline{\sigma v})_{DD} = 2.33 \times 10^{-14} T^{-2/3} \exp(-18.76 T^{-1/3}) \text{ cm}^3/\text{sec}$$

$$(\overline{\sigma v})_{DT} = 3.68 \times 10^{-12} T^{-2/3} \exp(-19.94 T^{-1/3}) \text{ cm}^3/\text{sec}$$

The power density released in the form of charged particles is

$$P_{DD} = 3.3 \times 10^{-13} n_D^2 (\overline{\sigma v})_{DD} \text{ watt/cm}^3 \text{ (including subsequent D-T reaction)}$$

$$P_{DT} = 5.6 \times 10^{-13} n_D n_T (\overline{\sigma v})_{DT} \text{ watt/cm}^3$$

$$P_{DHe^3} = 2.9 \times 10^{-12} n_D n_{He^3} (\overline{\sigma v})_{DHe^3} \text{ watt/cm}^3$$

The curie (abbreviated Ci) is a measure of radioactivity: 1 curie =

$3.7 \times 10^{10}$  counts/sec. Absorbed radiation dose is measured in rads:

1 rad =  $10^2$  erg/g.

## RELATIVISTIC ELECTRON BEAMS

Here  $\gamma = (1 - \beta^2)^{-\frac{1}{2}}$  is the relativistic scaling factor; in analytic formulas units are mks or cgs, as indicated; in numerical formulas, I is in amps, B in gauss, electron density N in  $\text{cm}^{-3}$ , and temperature, voltage and energy in MeV;  $\beta_z = v_z/c$ .

Relativistic electron gyroradius  $r_e = \frac{mc^2}{eB} (\gamma^2 - 1)^{\frac{1}{2}} \quad (\text{cgs})$   
 $= 1.70 \times 10^3 (\gamma^2 - 1)^{\frac{1}{2}} B \quad \text{cm}$

Relativistic electron energy  $W = mc^2 \gamma \quad (\text{cgs}) = 0.511 \gamma \text{ MeV}$

Alfvén-Lawson limit  $I_A = \frac{mc^3}{e} \beta_z \gamma \quad (\text{cgs})$   
 $= \frac{4\pi}{c} \frac{mc^3}{e} \beta_z \gamma \quad (\text{mks})$   
 $= 1.70 \times 10^4 \beta_z \gamma \quad \text{amp}$

The ratio of net current to  $I_A$  is

$$I/I_A = v/\gamma,$$

where  $v = Nr_e$ , with  $r_e = e^2/mc^2 = 2.82 \times 10^{-13} \text{ cm}$ . Beam electron number density is

$$n = 2.08 \times 10^8 J/B \text{ cm}^{-3},$$

where J is current density in  $\text{amp/cm}^2$ . For a uniform beam of radius a (in cm),

$$n = 6.63 \times 10^7 I / Ba^2 \text{ cm}^{-3},$$

$$2 r_e / a = v / \gamma.$$

aturated parapotential current (magnetically self-limited flow along equipotentials in pinched diodes and transmission lines).<sup>[23]</sup>

$$I_p = 8.5 \times 10^3 G \gamma \ln \left[ \gamma + (\gamma^2 - 1)^{1/2} \right] \text{ amp},$$

where G is a geometrical factor depending on the diode structure:

$\frac{\gamma}{2\pi d}$  for parallel plane cathode and anode of width  $\gamma$ , separation  $d$ ;

$\left[ \ln \frac{R_2}{R_1} \right]^{-1}$  for cylinders of radii  $R_1$  (inner) and  $R_2$  (outer);

$\frac{R_c}{d_0}$  for conical cathode of radius  $R_c$ , maximum separation  $d_0$  (at  $r = R_c$ ) from plane anode, etc.

$\gamma \rightarrow 0$  ( $\gamma \rightarrow 1$ ), both  $I_A$  and  $I_p$  vanish.

pinch condition

$$I^2 = 2Nk(T_e + T_i)c^2 \quad (\text{cgs})$$

$$= 3.20 \times 10^{-4} N(T_e + T_i) \text{ amp}^2$$

Child's law (space-charge limited current density between parallel plates with voltage drop  $V$  in MeV and separation  $d$  in cm)

Nonrelativistic:  $J = 2.34 \times 10^3 V^{3/2} d^{-2} \frac{\text{amp}}{\text{cm}^2}$

Relativistic ( $\gamma \geq 2$ ):  $J = 2.71 \times 10^3 (\gamma^{1/2} - 0.8471)^2 d^{-2} \frac{\text{amp}}{\text{cm}^2}$

Voltage registered by Rogowski coil of minor cross-sectional area  $A$ ,  $n$  turns, major radius  $a$ , inductance  $L$ , external resistance  $R$  and capacitance  $C$  (all in mks):

self-integrating  $V = - \frac{R n A \mu_0 I}{L 2\pi a}$  volts

externally integrated  $V = \frac{1}{RC} \frac{n A \mu_0 I}{2\pi a}$  volts

X-ray production for target with average atomic number  $Z$  ( $V \leq 5$  MeV)

$$\eta = \frac{\text{X-ray power}}{\text{beam power}} = 7 \times 10^{-4} ZV$$

X-ray dose at 1 meter generated by an e-beam depositing total charge  $Q$  coulombs while  $V \geq .84 V_{\text{max}}$  in material with charge state  $Z$

$$D = 150 V_{\text{max}}^{2.8} Q Z^{1/2} \text{ rads}$$

## ELECTROSTATIC STREAMING INSTABILITIES [24]

In this table, subscripts e,i,d,b stand for "electron," "ion," "drift," and "beam," respectively. Thermal velocities are denoted by a bar. In addition, the following are used:

$m$	electron mass
$M$	ion mass
$V$	velocity
$T$	temperature
$n_e, n_i$	number density
$n$	harmonic number
$C_s = T_e/M$	ion sound speed
$\omega_e, \omega_i$	plasma frequency
$\lambda_D$	Debye length
$r_e, r_i$	gyroradius
$\beta$	plasma/magnetic energy density ratio
$V_A$	Alfvén speed
$\Omega_e, \Omega_i$	gyro frequency
$\Omega_H$	hybrid gyro frequency, $\Omega_H^2 = \Omega_e \Omega_i$
$U$	relative drift velocity of two ion species

Name	Conditions	Parameters of Most Unstable Mode				Saturation Mechanism
		Growth Rate	Frequency	Wave Number	Group Velocity	
Electron - electron	$v_d > v_{ej}, j = 1, 2$	$\frac{1}{2} \omega_e$	0	$\frac{\sqrt{2}}{2} \omega_e / v_d$	0	Electron trapping until $\bar{v}_e \sim v_d$
Beam-plasma	$(n_b/n_p)^{1/3} > \bar{v}_b/v_b$	$\frac{\sqrt{3}}{2} \frac{(n_b/n_p)^{1/3}}{(n_p/n_b)^{1/3}} \omega_e$	$\omega_e \left[ 1 - \frac{1}{2^{4/3}} (n_b/n_p)^{1/3} \right]$	$\omega_e / v_d$	$\frac{2}{3} v_b$	Trapping of beam electrons
Buneman	$v_d > (M/m)^{1/3} \bar{v}_i, v_d > \bar{v}_e$	$\omega_e / v_d$	$\frac{1}{2^{4/3}} \omega_e (m/M)^{1/3}$	$\omega_e / v_d$	$\frac{2}{3} v_d$	Electron trapping until $\bar{v}_e \sim v_d$
Weak beam-plasma	$\bar{v}_b/v_b > (n_b/n_p)^{1/3}$	$\frac{1}{2} \frac{n_b}{n_p} \left( \frac{v_b}{v_b} \right)^2 \omega_e$	$\omega_e$	$\omega_e / v_b$	$3\bar{v}_e^2 / v_b$	Quasilinear or non-linear
Beam-plasma (hot electron)	$\bar{v}_e > v_b > \bar{v}_b, (n_b/n_p)^{1/2} \bar{v}_e > \bar{v}_b$	$\left( \frac{n_b}{n_p} \right)^{1/2} \frac{\bar{v}_e}{v_b} \omega_e$	$\frac{v_b}{v_e} \omega_e$	$\lambda_D^{-1}$	$v_b$	Quasilinear or non-linear
Ion acoustic	$T_e \gg T_i, v_d > C_s$	$\left( \frac{m}{M} \right)^{1/2} \omega_i$	$C_s \frac{\omega_e}{v_e}$	$\lambda_D^{-1}$	$C_s$	Quasilinear; ion tail formation; non-linear scattering; or resonance broadening.

Name	Condition	Parameters of Most Unstable Mode				
		Growth Rate	Frequency	Wave Number	Group Velocity	
Anisotropic temperature (Hydro)	$\frac{T_e}{T_i} < \frac{1}{2}$	$\Omega_e$	$\omega_e \cos \theta \sim \Omega_e$	$r_e^{-1}$	$\bar{V}_{e1}$	Isotropization
Ion Cyclotron	$V_d > 20 \bar{V}_i$	$0.1 \Omega_i$	$1.2 \Omega_i$	$r_i^{-1}$	$\frac{1}{2} \bar{V}_i$	Ion heating
Beam Cyclotron (Hydro)	$V_d > C_s$	$\frac{1}{\sqrt{2}} \Omega_e$	$n \Omega_e$	$\frac{1}{\sqrt{2}} \lambda_D^{-1}$	$v_d \leq v_g \leq C_s$	Resonance broadening
Modified two stream (Hydro)	$\sqrt{1 + \beta V_A} > v_d > C_s$	$\frac{1}{2} \Omega_H$	$\frac{\sqrt{2}}{2} \Omega_H$	$\sqrt{3} \frac{\Omega_H}{v_d}$	$\frac{1}{2} v_d$	Trapping
Ion-Ion	$U < 2V_A \sqrt{1 + \beta}$	$\frac{1}{2\sqrt{2}} \Omega_H$	0	$\sqrt{3/2} \frac{\Omega_H}{U}$	0	Ion trapping
Ion-Ion	$U < 2 C_s$	$\frac{1}{2\sqrt{2}} \omega_i$	0	$\sqrt{3/2} \frac{\omega_i}{U}$	0	Ion trapping

## LASERS

### Parameters of Important Laser Systems

Efficiencies and power levels are approximately state-of-the-art (1976)

Type	Wavelength (microns)	Efficiency	Power levels available(W)	
			Pulsed	CW
CO <sub>2</sub>	10.6	.01 - .02 pulsed	> 10 <sup>11</sup>	> 10 <sup>5</sup>
CO	5	0.4	> 10 <sup>8</sup>	> 100
Iodine	1.315	3 x 10 <sup>-3</sup>	> 10 <sup>11</sup>	—
Nd-glass, YAG	1.06	10 <sup>-3</sup>	> 10 <sup>12</sup>	1 - 300
Ruby	.6943	< 10 <sup>-3</sup>	10 <sup>10</sup>	1
He - Ne	.6328	10 <sup>-4</sup>	—	1 - 50 x 10 <sup>-3</sup>
Argon ion	.45 - .60	10 <sup>-3</sup>	5 x 10 <sup>4</sup>	1 - 10
N <sub>2</sub>	.3371	10 <sup>-3</sup> - .05	10 <sup>5</sup> - 10 <sup>8</sup>	—
Xenon	.175	.02	> 10 <sup>8</sup>	—

3. Ponderomotive force

$$\underline{f} = N \nabla \langle E^2 \rangle / 8\pi N_c$$

where

$$N_c = 1.1 \times 10^{21} / \lambda_0^2 \text{ cm}^{-3} .$$

4. For uniform illumination of a lens with f-number F, the diameter d at focus (85% of the energy) and the depth of focus  $l$  (first zero in intensity) are given by

$$d \approx 2.44 F \lambda_0 \theta / \theta_{DL} \mu \text{ and } l \approx \pm 2 F^2 \lambda_0 \theta / \theta_{DL} \mu .$$

Here  $\theta$  is the beam divergence containing 85% of energy and  $\theta_{DL}$  is the diffraction limited divergence:

$$\theta_{DL} = \frac{2.44 \lambda_0}{b} \quad (\text{aperture } b \text{ in } \mu)$$

These formulas are modified for nonuniform illumination of the lens (such as gaussian) or for pathological laser profiles.

### Formulas

1. An e-m wave with  $\underline{k} \parallel \underline{B}$  has an index of refraction given by

$$n_{\pm} = \left[ 1 - \frac{\omega_{pe}^2}{\omega(\omega \pm \omega_{ce})} \right]^{1/2}$$

where  $\pm$  refers to the velocity. The rate of change of polarization angle  $\theta$  as a function of displacement  $s$  (Faraday rotation) is given by

$$\frac{d\theta}{ds} = \frac{1}{2} k(n_{-} - n_{+}) = 2.36 N B f^{-2} \text{ cm}^{-1},$$

where  $N$  is the electron number density,  $B$  is the field strength, and  $f$  is the wave frequency. For  $N = 10^{20} \text{ cm}^{-3}$ ,  $B = 10 \text{ kG}$  and  $f = 3 \times 10^{14} \text{ Hz}$  (corresponding to a wavelength  $\lambda = 1 \mu$ ),  $\frac{d\theta}{ds} = 2.6 \times 10^{-3} \text{ rad/m}$ .

2. The quiver velocity of an electron in an e-m field of angular frequency  $\omega$  is

$$v_0 = \frac{eE_{\max}}{m\omega} = 25.6 \sqrt{I} \lambda_0 \text{ cm/sec}$$

in terms of the laser flux  $I = \frac{cE_{\max}^2}{8\pi}$ , with  $I$  in watts/cm<sup>2</sup>, laser

wavelength  $\lambda_0$  in microns. The ratio of quiver energy to thermal energy

is

$$\frac{m_e v_0^2}{2 kT} = 1.81 \times 10^{-13} \frac{\lambda_0^2 I}{T},$$

$T$  in eV. E.g., if  $I = 10^{15} \text{ watts/cm}^2$ ,  $\lambda_0 = 1 \mu$ ,  $T = 2 \text{ keV}$ , then

$$W_{\text{quiver}}/W_{\text{thermal}} \approx .1$$

## ATOMIC AND RADIATION PROCESSES

The subject of atomic physics is central and indispensable for understanding the behavior of real plasmas. It is ill-suited for distillation into a few general formulas, however. Those compiled here should not be applied blindly. The inexperienced user would do well to first consult the references cited, particularly [31] and [35], which are more accessible to plasma physicists.

Energies and temperatures are in eV; all other units are cgs except where noted.  $Z$  is the charge state ( $Z = 0$  refers to a neutral atom); the subscript  $e$  labels electrons.  $N$  refers to number density,  $n$  to principal quantum number. Asterisk superscripts on level population densities denote local thermodynamic equilibrium (LTE) values. Thus  $N_n^*$  is the LTE number density of atoms (or ions) in level  $n$ .

Characteristic atomic collision cross section

$$(1) \quad \pi a_0^2 = 8.80 \times 10^{-17} \text{ cm}^2$$

Binding energy for outer electron in level labelled by quantum numbers

$n, \ell$

$$(2) \quad E_{\infty}^Z(n, \ell) = - \frac{Z^2 E_{\infty}^H}{(n - \Delta_{\ell})^2} \text{ eV},$$

where  $E_{\infty}^H = 13.6 \text{ eV}$  is the hydrogen ionization energy and  $\Delta_{\ell} = .75\ell^{-5}$ ,

$\ell \geq 5$ , is the quantum defect.

### Excitation and Decay

Cross section (Bethe approximation) for electron excitation by dipole allowed transition  $m \rightarrow n$  [25,28]

$$(3) \quad \sigma_{mn} = 2.36 \times 10^{-13} \frac{f_{nm} g(n,m)}{\epsilon \Delta E_{nm}} \text{ cm}^2,$$

where  $f_{nm}$  is the oscillator strength,  $g(n,m)$  is the Gaunt factor,  $\epsilon$  is the incident electron energy, and  $\Delta E_{nm} = E_n - E_m$ .

Electron excitation rate averaged over Maxwellian velocity distribution [27,28]

$$(4) \quad X_{mn} = N_e \langle \sigma_{mn} v \rangle = \frac{1.6 \times 10^{-5} f_{nm} \langle g(n,m) \rangle N_e}{\Delta E_{nm} \sqrt{T_e}} \exp\left(-\frac{\Delta E_{nm}}{T_e}\right) \text{ sec}^{-1},$$

where  $\langle g(n,m) \rangle$  denotes the thermal averaged Gaunt factor whose behavior as a function of energy is shown in Fig. 3 (generally  $\sim 1$  for atoms,  $\sim 0.2$  for ions).

Rate for electron collisional de-excitation

$$(5) \quad Y_{nm} = \left(\frac{N_m^*}{N_n^*}\right) X_{nm} \text{ sec}^{-1}.$$

Here  $\frac{N_m^*}{N_n^*} = \left(\frac{g_m}{g_n}\right) \exp(\Delta E_{nm}/T_e)$  is the Boltzmann relation for level population densities, where  $g_n$  is the statistical weight of level  $n$ .

Rate for spontaneous radiative decay  $n \rightarrow m$  (Einstein A coefficient) [27]

$$(6) \quad A_{nm} = 4.3 \times 10^7 \frac{g_n}{g_m} f_{nm} (\Delta E_{nm})^2 \text{ sec}^{-1}.$$




Fig. 3 - The effective Gaunt factor  $\bar{g}$  for neutral atoms and positive ions. For neutrals the curve is deduced from the experimental results for  $1s \rightarrow 2p$  in H.

Intensity emitted per unit volume from the transition  $n \rightarrow m$  in an optically thin plasma

$$(7) \quad I_{nm} = 1.6 \times 10^{-19} A_{nm} N_n \Delta E_{nm} W/cm^3.$$

Condition for steady state in a corona model

$$(8) \quad N_o N_e \langle \sigma_{on} \nu \rangle = N_n A_{no},$$

where the ground state is labelled by a subscript zero.

Hence for a transition  $n \rightarrow m$  in ions, where  $\langle g(n,m) \rangle \approx 0.2$ ,

$$(9) \quad I_{nm} = 5.1 \times 10^{-25} f_{nm} (g_o/g_m) (\Delta E_{nm}/\Delta E_{no})^3 N_e N_o T_e^{-1/2} \exp(-\Delta E_{no}/T_e) W/cm^3.$$

#### Ionization and Recombination

In a general time-dependent situation the number density of the charge state  $Z$  satisfies

$$(10) \quad \frac{dN(Z)}{dt} = N_e [ -S(Z)N(Z) - \alpha(Z)N(Z) + S(Z-1)N(Z-1) + \alpha(Z+1)N(Z+1) ].$$

Here  $S(Z)$  is the ionization rate. The recombination rate  $\alpha(Z)$  has the form  $\alpha(Z) = \alpha_r(Z) + N_e \alpha_3(Z)$ , where  $\alpha_r$  and  $\alpha_3$  are the radiative and three-body recombination rates, respectively.

Classical ionization cross-section for any atomic shell  $k$  <sup>[23]</sup>

$$(11) \quad \sigma_i = 6 \times 10^{-14} \frac{b_k}{U_k^2} g_k(x) \text{ cm}^2;$$

Here  $b_k$  is the number of shell electrons,  $U_k$  is the binding energy of the ejected electron,  $x = \epsilon/U_k$  where  $\epsilon$  is the incident electron energy, and  $g$  is a universal function with a maximum value  $\approx 0.2$  at  $x \approx 4$ .

Ionization from ion ground state, averaged over Maxwellian electron distribution [22]

$$(12) \quad S(Z) = \frac{10^{-5} (T_e/E_\infty^Z)^{1/2}}{(E_\infty^Z)^{3/2} (6 + T_e/E_\infty^Z)} \exp(-E_\infty^Z/T_e) \text{ cm}^3/\text{sec},$$

where  $E_\infty^Z$  is the ionization energy.

Electron-ion radiative recombination rate ( $e + N(Z) \rightarrow N(Z-1) + h\nu$ ) [30]

$$(13) \quad \alpha_r(Z) = 5.2 \times 10^{-14} Z \sqrt{E_\infty^Z/T_e} \left\{ 0.43 + \frac{1}{2} \ln(E_\infty^Z/T_e) + 0.469 (E_\infty^Z/T_e)^{-1/3} \right\} \text{ cm}^3/\text{sec}.$$

For  $10 \text{ keV} < T_e/Z^2 < 150 \text{ keV}$ , this becomes approximately [28]

$$(14) \quad \alpha_r(Z) = 2.7 \times 10^{-13} \frac{Z^2}{\sqrt{T_e}} \text{ cm}^3/\text{sec}.$$

Collisional (three-body) recombination rate for singly ionized plasma [31]

$$*(15) \quad \alpha_3 = 8.75 \times 10^{-27} T_e^{-3/2} \text{ cm}^3/\text{sec}$$

(limited to low temperatures,  $kT_e < E_\infty^H$ ). Photoionization cross


section for ions in level  $n$ ,  $\ell$  (short wavelength limit) <sup>[24]</sup>

$$(16) \quad \sigma_{\text{ph}}(n, \ell) = 1.64 \times 10^{-18} Z^5 / (n^3 K^7 + 2\ell),$$

where  $K$  is wavenumber in Rydberg units ( $1 \text{ Rydberg} = 1.0974 \times 10^5 \text{ cm}^{-1}$ ).

### Charge Exchange

Two kinds of charge exchange take place in a plasma, depending on  $\Delta E$  in



When  $\Delta E = 0$  (the internal energy difference vanishes) the process is called resonant. When  $A = B$  the resonance is symmetric; otherwise it is asymmetric. When  $\Delta E \neq 0$  the process (17) is just a charge transfer reaction. In this case the adiabatic principle of Massey suggests that the peak of the cross section  $\sigma$  arises when

$$v \approx \ell \frac{\Delta E}{h} .$$

Here  $\ell$  is the interaction range and  $v$  is a critical relative velocity.

For larger values of  $v$ ,

$$(18) \quad \sigma \approx k \exp \left( - \frac{\ell |\Delta E|}{4 h v} \right),$$

with  $k$  being a constant for each reaction. However, for symmetric charge exchange the cross-section as given by Fursov can be expressed as <sup>[22]</sup>

$$(19) \quad \sigma = \left[ a - b \log_{10} E \right]^2 \text{ cm}^2 .$$

AD-A041 545

NAVAL RESEARCH LAB WASHINGTON D C  
REVISED AND ENLARGED COLLECTION OF PLASMA PHYSICS FORMULAS AND --ETC(U)  
MAY 77 D L BOOK

F/G 20/5

UNCLASSIFIED

NRL-MR-3332

NL

2 OF 2  
AD-A041 545



END

DATE  
FILMED  
'8 - 77

Values of a and b are given below for some selected reactions<sup>32</sup> for the energy range  $E_{\min} < E < E_{\max}$

Process	a ( $10^{-8}$ cm)	b ( $10^{-8}$ cm)	$E_{\min}$ (eV)	$E_{\max}$ (eV)
$H^+ - H$	7.6	1.06	20	10,000
$He^+ - He$	4.7	1.1	20	6,000
$Ne^+ - Ne$	6.1	1.7	20	6,000
$H^+ - O$	4.2	0.38	25	8,000

#### Equilibrium Models

Saha equilibrium [33]

$$(20) \quad \frac{N_e N^*(Z)}{N^*(Z-1)} = 6.0 \times 10^{21} \frac{g_1^Z T_e^{3/2}}{g_n^{Z-1}} \exp\left(-\frac{E_{\infty}^Z(n,l)}{T_e}\right) \text{ cm}^{-3},$$

where  $g_n^Z$  is the statistical weight for level n of charge state Z and  $E_{\infty}^Z(n,l)$  is the ionization energy of the neutral atom initially in level (n,l), given by Eq. (2).

In a steady state at high electron density

$$(21) \quad \frac{N_e N^*(Z)}{N^*(Z-1)} = \frac{s(Z-1)}{\alpha_3} \text{ cm}^{-3},$$

a function only of T.

Conditions for LTE

(a) Collisional and radiative excitation rates for a level  $n$  must satisfy [33]

$$(22) \quad Y_{nm}/A_{nm} > 10.$$

(b) Electron density must satisfy [33]

$$(23) \quad N_e \geq 7 \times 10^{13} Z^7 n^{-17/2} (T/E_\infty^Z)^{1/2} \text{ cm}^{-3}.$$

Steady state condition in corona model

$$(24) \quad \frac{N(Z-1)}{N(Z)} = \frac{\alpha_r}{S(Z-1)}$$

Corona model is applicable if [34]

$$(25) \quad 10^{12} t_I^{-1} < N_e < 10^{16} T_e^{7/2} \text{ cm}^{-3},$$

where  $t_I^{-1}$  is the inverse ionization time.

### Radiation<sup>†</sup>

Average radiative decay rate of state with principal quantum number  $n$  is

$$(26) \quad A_n = \sum_{m < n} A_{nm} = 1.6 \times 10^{10} Z^4 n^{-9/2} \text{ sec}^{-1}$$

Natural line width ( $\Delta E$  in eV)

$$(27) \quad \Delta E \Delta t = h = 4.14 \times 10^{-15} \text{ eV-sec.}$$

where  $\Delta t$  is the lifetime of the line.

Doppler width

$$(28) \quad \frac{\Delta \lambda}{\lambda} = 7.7 \times 10^{-5} \sqrt{\frac{T}{\mu}},$$

where  $\mu$  is the emitting atom or ion mass in units of the proton mass.

Optical depth for a Doppler broadened line<sup>[33]</sup>

$$(29) \quad \tau = 1.76 \times 10^{-13} \lambda \left( \frac{Mc^2}{kT} \right)^{1/2} N\ell = 5.4 \times 10^{-9} \lambda \left( \frac{\mu}{T} \right)^{1/2} N\ell,$$

where  $\lambda$  is wavelength and  $\ell$  the physical depth of the plasma;  $M$ ,  $N$  and  $T$  are atomic mass, number density and temperature of the absorber;  $\mu$  is  $M$  divided by the proton mass. Optically thin means  $\tau < 1$ .

---

<sup>†</sup> Note that the units employed here are those summarized on p. 88.

Resonance absorption cross section at center of line

$$(30) \quad \sigma_{\lambda=\lambda_c} = 5.6 \times 10^{-13} \frac{\lambda^2}{\Delta\lambda} \text{ cm}^2 .$$

Wien displacement law: wavelength of maximum black body emission is given by

$$(31) \quad \lambda_{\text{max}} = 2.50 \times 10^{-5} T^{-1} \text{ cm} .$$

Radiation from surface of black body of temperature T

$$(32) \quad W = 1.03 \times 10^5 T^4 \text{ watts/cm}^2 .$$

Bremsstrahlung from hydrogen-like plasma <sup>[19]</sup>

$$(33) \quad P_{\text{Br}} = 1.69 \times 10^{-32} N_e T_e^{1/2} \sum [Z^2 N(Z)] \text{ watts/cm}^3 ,$$

where the sum is over all ionization states Z.

Bremsstrahlung optical depth <sup>[35]</sup>

$$(34) \quad \tau = 5.0 \times 10^{-38} N_e N_i Z^2 \bar{g} \ell T^{-7/2} ,$$

where  $\bar{g} \approx 1.2$  is an average Gaunt factor and  $\ell$  is the physical path length.

Recombination (free-bound) radiation

$$(35) \quad P_r = 1.69 \times 10^{-32} N_e T_e^{1/2} \sum \left[ Z^2 N(Z) \left( \frac{E_\infty^{Z-1}}{T_e} \right) \right] \text{watts/cm}^3.$$

Cyclotron radiation in magnetic field B<sup>[19]</sup>

$$(36) \quad P_c = 6.21 \times 10^{-21} B^2 N_e T_e \text{ watt/cm}^3$$

For  $N_e kT_e = N_i kT_i = B^2/16\pi$  ( $\beta = 1$ , isothermal plasma),<sup>[19]</sup>

$$(37) \quad P_c = 5.00 \times 10^{-31} N_e^2 T_e^2 \text{ watt/cm}^3.$$

Cyclotron radiation e-folding time for a single electron<sup>[35]</sup>

$$(38) \quad \tau \approx \frac{9.0 \times 10^8 B^{-2}}{2.5 + \gamma} \text{ sec}^{-1},$$

where  $\gamma$  is the energy divided by  $mc^2$ .

Number of cyclotron harmonics trapped in a medium of finite depth  $l$ <sup>[35]</sup>

$$(39) \quad m^* = (57 \beta B l)^{1/3},$$

where  $\beta = NkT/8\pi B^2$ .

Line radiation is given by summing Eq. 9 over all species in the plasma.

## REFERENCES

Most of the material in this report is quite well-known and for all practical purposes is in the "public domain." The books and articles cited below are intended primarily not for the purpose of giving credit to the original workers, but (i) to guide the reader to sources containing additional related material and (ii) to indicate where derivations, explanations, examples, etc., omitted from this compilation can be found.

1. See M. Abramowitz and I. A. Stegun (Editors), Handbook of Mathematical Functions (Dover, New York, 1968), pp. 1-3, for a tabulation of some mathematical constants not available on pocket calculators.
2. H. W. Gould, "Note on Some Binomial Coefficient Identities of Rosenbaum," *J. Math. Phys.* 10, 49 (1969); H. W. Gould and J. Kaucky, "Evaluation of a Class of Binomial Coefficient Summations," *J. Comb. Theory* 1, 233 (1966).
3. P. M. Morse and H. Feshbach, Methods of Theoretical Physics (McGraw-Hill, New York, 1953), vol. I, pp. 47-52 and pp. 656-666.
4. W. D. Hayes, "A Collection of Vector Formulas," Princeton University, Princeton, N.J., 1956 (unpublished), and personal communication (1977).

5. See Quantities, Units and Symbols, report of the Symbols Committee of the Royal Society, 2<sup>nd</sup> Edition (Royal Society, London, 1975) for a discussion of nomenclature in SI units.
6. E. R. Cohen and B. N. Taylor, "Least-Squares Adjustment of the Fundamental Constants," J. Phys. Chem. Ref. Data 2, 663 (1973).
7. J. Stratton, Electromagnetic Theory (McGraw-Hill Book Co., New York, 1941), p. 508.
8. ITT Reference Data for Engineers (Eams and Co., Indianapolis, Indiana, 1975), p. I-1. These definitions are International Telecommunications Union (ITU) Standards.
9. Harry E. Thomas, Handbook of Microwave Techniques and Equipment Prentice-Hall, Englewood Cliffs, N. J., 1972), p. 8. Further subdivisions are defined in Ref. 8, p. I-3.
10. John P. Catchpole and George Fulford, Ind. and Eng. Chem. 58, 47 (1966); reprinted in recent editions of the Handbook of Chemistry and Physics (Chemical Rubber Co., Cleveland, Ohio) on pp. F201-213.
11. B. D. Fried and S. D. Conte, The Plasma Dispersion Function (Academic Press, Inc., New York, 1961).
12. R. W. Landau and S. Cuperman, "Stability of Anisotropic Plasmas to Almost-Perpendicular Magnetosonic Waves," J. Plasma Phys. 6, 495 (1971).

13. B. D. Fried, C. L. Hedrick, J. McCune, "Two-Pole Approximation for the Plasma Dispersion Function," *Phys. Fluids* 11, 249 (1968).
14. K. Papadopoulos, personal communication (1976).
15. John Sheffield, Plasma Scattering of Electromagnetic Radiation (Academic, New York, 1975), p. 6 (after J. W. Paul).
16. B. A. Trubnikov, "Particle Interactions in a Fully Ionized Plasma," *Reviews of Plasma Physics*, vol. 1 (Consultants Bureau, New York, 1965), p. 105.
17. After J. G. Siambis and S. N. Stitzer, "Collisional Parameters of a Test Particle and a Streaming Plasma," *Phys. Fluids* 17, 2192 (1974).
18. S. I. Braginskii, "Transport Processes in a Plasma," Reviews of Plasma Physics, vol. 1 (Consultants Bureau, New York, 1965), p. 205.
19. K. H. Lloyd and G. Härendel, "Numerical Modeling of the Drift and Deformation of Ionospheric Plasma Clouds and of their Interaction with Other Layers of the Ionosphere," *J. Geophys. Res.*, 78, 7389 (1973).
20. S. Glasstone and R. H. Lovberg, Controlled Thermonuclear Reactions (Van Nostrand, New York, 1960), chapt. 2.
21. References to experimental measurements of branching ratios and cross sections are listed in F. K. McGowan, et al., *Nucl. Data Tables* A6, 353 (1969); and *Nucl. Data Tables* A8, 199 (1970). The yields listed in the table are calculated directly from the mass defect.

22. G. H. Miley, H. Towner and N. Ivich, Fusion Cross Section and Reactivities, Rept. COO-2218-17, University of Illinois, Urbana, Ill. 61801 (1974).
23. J. M. Creedon, "Relativistic Brillouin Flow in the High  $v/\gamma$  Limit," J. App. Phys. 46, 2946 (1975).
24. See, for example, A. B. Mikhailovskii, Theory of Plasma Instabilities (Consultants Bureau, New York, 1974), vol. I.
25. M. J. Seaton, Atomic and Molecular Processes, Ed. by D. R. Bates (New York, Academic Press 1962), chapt. 11.
26. H. Van Regemorter, "Rate of Collisional Excitation in Stellar Atmospheres," Astrophys. J. 136, 906 (1962).
27. A. C. Kolb and R. W. P. McWhirter, "Ionization Rates and Power Loss from  $\theta$ -Pinches by Impurity Radiation," Phys. Fluids 7, 519 (1964).
28. R. W. P. McWhirter, Plasma Diagnostic Techniques, Huddleston and Leonard, Eds. (Academic Press, New York, 1965).
29. M. Gryzinski, "Classical Theory of Atomic Collisions I. Theory of Inelastic Collisions," Phys. Rev. 138A, 336 (1965).
30. M. J. Seaton, "Radiative Recombination of Hydrogen Ions," Monthly Notices Roy. Astron. Soc. 119, 81 (1959).
31. Y. B. Zel'dovich and Yu. P. Raizer, Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena, (Academic Press, New York, 1966), vol. I, p. 407.
32. J. W. Bond, Jr., K. M. Watson and J. A. Welch Jr., Atomic Theory of Gas Dynamics (Addison Wesley Publishing Co., New York, 1966).

33. H. R. Griem, Plasma Spectroscopy (Academic Press, New York, 1966).
34. T. F. Stratton, Plasma Diagnostic Techniques, Huddlestone and Leonards, Eds. (Academic Press, New York, 1965).
35. G. Bekefi, Radiation Processes in Plasmas (Wiley, New York, 1966).