

12
P.S.

AD A 042375

OFFICE OF NAVAL RESEARCH

Contract Number N00014-76-WR60021

Task No. NR 205-009

BIODEGRADATION OF OIL IN SEAWATER FOR
NAVAL POLLUTION CONTROL

by

T. B. O'Neill

Civil Engineering Laboratory
Naval Construction Battalion Center
Port Hueneme, California 93043

DDC
RECEIVED
AUG 2 1977
RESOLVED

1 June 1977

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

Distribution of this report is unlimited.

AD No. —
DDC FILE COPY

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) O'Neill, Thomas B., Code L52 Civil Engineering Laboratory Naval Construction Battalion Center Port Huene, CA 93043	2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED
	2b. GROUP

3. REPORT TITLE
 ⑥ THE BIODEGRADATION OF OIL IN SEAWATER FOR NAVAL POLLUTION CONTROL, FINAL REPORT

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)
 ⑨ FINAL REPORT, 1974-1976,

5. AUTHOR(S) (First name, middle initial, last name)
 ⑩ THOMAS B./O'NEILL ⑫ 12 p.

6. REPORT DATE ⑪ 1 JULY 1977	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS 3
---------------------------------	------------------------	----------------------

8a. CONTRACT OR GRANT NO. N0001476WR60021 b. PROJECT NO. Work Unit No. NR205009 c. d.	9a. ORIGINATOR'S REPORT NUMBER(S) 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
--	--

10. DISTRIBUTION STATEMENT
 Distribution of this report is unlimited.

11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY OFFICE OF NAVAL RESEARCH, CODE 443 ARLINGTON, VA 22217
-------------------------	---

13. ABSTRACT
 The report describes the isolation and utilization of pure and mixed microbial cultures for experiments on the biodegradation of crude oil Bunker C fuel and marine diesel. Many microbial species were found that had hydrocarbonoclastic activity. When pure cultures were combined in mixtures the activity was much greater, 91% oxidation in seven days, than the activity of any one of the component species when used in a pure culture.

391 111

SB

UNCLASSIFIED

Security Classification

14 KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Biodeterioration						
Oils						
Sea Water						
Pollution						
Seepage						
Oil Spills						
Oceans						
Shores						
Marine micro-organisms						
Oxidizers						
Degradation						
Sampling						

iv

UNCLASSIFIED

Security Classification

CONTENTS

	Page
INTRODUCTION	1
PROCEDURE	1
RESULTS	2
CONCLUSIONS	7
REFERENCES	8

ACCESSION FOR	
ATED	WITH SECTION <input checked="" type="checkbox"/>
END	DATE SECTION <input type="checkbox"/>
UNHANDLED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION AVAILABILITY CODES	
USE	APPL. AND M. SPENTAL
A	

INTRODUCTION

The problem of oil spillage is one of great magnitude and concern to the Navy, particularly in this era of increasing legal restrictions and ecological awareness. Oil on the seas is a fire hazard and a menace to organisms of the marine environment as well as an offense against aesthetics to those who use the seas for recreation and inspiration.

The negative consequences of oil spillage on waters and beaches are not new. They are concomitant with the origin and development of propulsion units and other devices utilizing fossil fuels. Pertinent to the context of this report, it is imperative to maintain that natural oil seeps from the ocean floor have undoubtedly existed for millennia. Merz (1959) reported on the almost universal presence of oil, derived from natural oil seeps, on the beaches of Southern California. The amounts of oil cited are small, approximately 60cc per 50m², and generally non-persistent. Coal Oil Point, in the Santa Barbara Channel, is an extreme example of a beach having continued deposits of large amounts of crude oil and tar. Allen, et al (1970) determined this spillage to be on the order of 50 to 70 barrels (8,000 to 11,000 liters) of oil per day into the channel and on the beaches. The beach deposits are estimated to be greater than any other beach in Southern California.

As well known as the appearance of oil and tar on the beaches and adjacent waters is its disappearance after unspecified periods of time. ZoBell (1969) reported that oil-oxidizing bacteria are often abundant in marine sediments and seawater of coastal areas where natural or man-induced pollution is persistent. ZoBell further noted that oil-oxidizing bacteria, yeasts, and molds have been identified in water and mud samples in many parts of the world.

The present report is concerned with the measurable activity of micro-organisms to oxidize certain fossil fuels. The organisms utilized were isolated from beaches and inshore marine waters where oil spills are common or were secured from type culture collections.

PROCEDURE

The literature, from 1943 to the contemporary period, dealing with microbial oxidation of hydrocarbons was searched to prepare a listing of micro-organisms reported to possess an ability to degrade hydrocarbons. Subsequently, where possible, lyophilized cultures were secured from type culture collections, notably the American Type Culture Collection of Rockville, Maryland.

Other sources of micro-organisms were the local beaches and harbor areas where oil spills are common; e.g., Coal Oil Point, Carpenteria Beach, and Los Angeles-Wilmington Harbor. Aliquot samples of beach sand or harbor soil or water were placed in a mineral salts medium made with aged filtered sea water and containing 0.75 percent hydrocarbon, either crude oil, marine diesel or Bunker C fuel as a sole source of organic material. Incubation was at room temperature, 20°C-25°C, upon a rotary shaker and varied from periods of one week to ten days. During the periods of incubation microbial populations were determined daily, by plate counts, and hydrocarbon oxidizing micro-organisms were isolated.

The isolated pure cultures were maintained on optimal media and transferred as required. Prior to the utilization of the cultures they were transferred to broth and during the late exponential growth phase the cells were centrifuged and washed three times with the mineral salts medium to remove the major portion of the organic matter of the media. In the experiments approximately 3×10^{12} cells were used as an inoculum in 250ml flasks containing 150ml of mineral salts medium containing 0.75% of hydrocarbon as the sole source of organic matter. Controls were identical flasks without added micro-organisms. The flasks were incubated at room temperature upon a rotary shaker for seven days. At the end of the incubation period, the quantity of hydrocarbon remaining was determined by an extraction process.

In the extraction procedure, several washings of chloroform were used as the hydrocarbon solvent. The extract was dried over anhydrous sodium sulfate, filtered to remove the drying agent and evaporated to dryness under reduced pressure. Evaporation was continued until three consecutive weighings were identical to the nearest milligram.

Initial experiments utilized pure cultures of one microbial species and subsequent experiments involved mixtures of two or more pure cultures in single flasks.

Over the period of experimentation many media modifications were tested as were other conditions that might optimize hydrocarbon oxidizing activity.

A longevity study was also performed. In these experiments pure cultures of one or more microbial species were added to a mineral salts hydrocarbon mixture and incubated for a period of a year with occasional additions of mineral salts solution to maintain an appropriate level of liquid.

RESULTS

In the period of experimentation, the media and conditions of

experiments significantly changed. This represents an increasing knowledge of parameters that facilitate microbial oxidation of hydrocarbons. For example, the use of ammonium salts as a nitrogen source lowered the pH of sea water used in the media from a pH of 8 to a detrimental pH of 4 as the bacteria multiplied. The unfavorable pH was a principal factor in the reduction of microbial population. The use of buffer solutions alleviated this condition. The addition of sterile air to flasks increased microbial activity approximately seven-fold. The incorporation of sterile diatomaceous earth to flasks increased adsorptive surface area for hydrocarbon and microbes and increased activity three-fold.

Fifty-five species of hydrocarbon oxidizing species of microorganisms were secured from type culture collections and 62 species were isolated from local habitats. The maintenance of this comparatively large collection of microorganisms is time consuming for they must be transferred monthly to fresh media.

Many of the organisms previously cited in the literature as having hydrocarbon-oxidizing ability and secured from culture collections failed to demonstrate activity under the conditions of the present experiments. That all media were prepared using salt water (75%) and fresh water (25%) rather than fresh water alone might well explain lack of microbial activity. Possibly the microbes could not oxidize the specific hydrocarbons used in these experiments. The species not demonstrating any hydrocarbonoclastic activity are:

Achromobacter delicatulus
Arthrobacter ureafaciens
Brevibacterium pusillum
Candida parapsilosis
Flavobacterium arborescens
Hydrogenomonas facilis
Hyphomicrobium indicum
Micrococcus varians
Mycobacterium rubrum
Mycobacterium smegmatis
Pseudomonas cruciviae
Pseudomonas dacunhae
Pseudomonas stutzeri
Serratia marinorubra
Vibrio cycloides

The pure cultures of microorganisms that demonstrated a significant hydrocarbonoclastic activity are:

Organism	Hydrocarbon Oxidized (% in a 7-day period)		
	Crude Oil	Bunker C	Marine Diesel
<u>Achromobacter cycloclastes</u>	23	12	8
<u>Aeromonas hydrophila</u> subsp <u>formicans</u>	8	11	4
<u>Arthrobacter arthrobotryoides</u>	44	22	12
<u>Arthrobacter citreus</u>	9	3	0
<u>Arthrobacter simplex</u>	31	27	19
<u>Azotobacter chroococcum</u>	28	12	8
<u>Azotomonas insoluta</u>	13	9	3
<u>Bacillus insolitus</u>	20	9	7
<u>Candida lipolytica</u>	41	33	17
<u>Candida petrophilum</u>	37	22	19
<u>Cellulomonas fimi</u>	12	7	3
<u>Cephalosporium acremonium</u>	18	11	8
<u>Clasdosporium resinae</u>	18	4	9
<u>Corynebacterium hydrocarboclastus</u>	32	27	18
<u>Cunninghamella elegans</u>	67	32	29
<u>Flavobacterium ferrungineum</u>	17	9	5
<u>Flavobacterium marinotypicum</u>	51	26	16
<u>Hyphomicrobium neptunium</u>	39	15	11
<u>Micrococcus roseus</u>	23	7	12
<u>Micrococcus parrafinolyticus</u>	32	11	9
<u>Mycobacterium hyalinum</u>	17	4	4
<u>Mycobacterium phlei</u>	17	9	8
<u>Mycobacterium rhodochrous</u>	26	11	12
<u>Mycoplana bullata</u>	30	19	5

<u>Mycoplana dimorpha</u>	26	13	8
<u>Nitrobacter agilis</u>	12	7	4
<u>Nocardia corralina</u>	31	22	8
<u>Nocardia hydrocarbonoxydans</u>	36	19	17
<u>Nocardia opaca</u>	29	11	10
<u>Nocardia parrafinica</u>	32	10	8
<u>Nocardia petreoleophila</u>	30	17	6
<u>Penicillium zonatum</u>	42	26	8
<u>Pseudomonas boreopolis</u>	24	12	7
<u>Pseudomonas creosotensis</u>	27	10	6
<u>Pseudomonas desmolytica</u>	29	11	9
<u>Pseudomonas facilis</u>	22	8	7
<u>Pseudomonas oleovorans</u>	18	14	11
<u>Pseudomonas resinovorans</u>	20	12	7
<u>Pseudomonas rubescens</u>	12	4	6
<u>Pseudomonas putida</u>	19	11	8
<u>Sarcina aurantica</u>	6	0	0
<u>Sarcina flava</u>	12	7	4
<u>Vibrio sp</u>	16	4	5
<u>Vibrio neocistis</u>	18	9	3

The selection of microbial species for experiments involving more than one species was based on performance in the above described exercises utilizing but one species. The components of the mixed cultures were individually grown as pure cultures in broth, centrifuged and washed, and then added to the flasks containing mineral salts medium and hydrocarbon. The combined total of all microorganisms added was approximately 3×10^{12} cells.

In some experiments the total of hydrocarbons oxidized did not exceed that of either component when used alone. In such a category are the mixed cultures of Arthrobacter simplex and Bacillus insolitus, Corynebacterium hydrocarboclastus and Candida petrophilum, and Micrococcus roseus with Nocardia corralina.

More encouraging are those combinations of cultures whose activity is greater than either of the component cultures individually, or greater than both collectively. Some examples of these results are as follows:

Organism	Hydrocarbon Oxidized (% in a 7-day period)		
	Crude Oil	Bunker C	Marine Diesel
<u>Corynebacterium hydrocarbo-</u> <u>clastus</u> and <u>Candida petrophilum</u>	63	31	24
<u>Flavobacterium marinotypicum</u> and <u>Hyphomicrobium neptunium</u>	57	29	25
<u>Arthrobacter arthrobotryoides</u> and <u>Candida lipolytica</u>	61	34	31
<u>Arthrobacter simplex</u> and <u>Arthrobacter arthrobotryoides</u>	44	22	17
<u>Candida petrophilum</u> and <u>Flavobacterium marinotypicum</u>	54	29	21
<u>Hyphomicrobium neptunium</u> and <u>Arthrobacter simplex</u>	51	18	11
<u>Candida petrophilum</u> and <u>Arthrobacter arthrobotryoides</u>	64	39	18
<u>Micrococcus paraffinolyticus</u> and <u>Arthrobacter arthrobotryoides</u>	67	41	27
<u>Mycobacterium rhodochrous</u> and <u>Arthrobacter arthrobotryoides</u>	85	47	32
<u>Mycoplana bullata</u> and <u>Pseudomonas desmolytica</u>	43	27	34

<u>Cunninghamella elegans</u> and <u>Arthrobacter simplex</u>	79	53	21
<u>Cunninghamella elegans</u> and <u>Arthrobacter arthrobotryoides</u>	87	42	38
<u>Nocardia paraffinica</u> and <u>Penicillium zonatum</u>	54	17	28
<u>Vibrio sp</u> and <u>Cunninghamella elegans</u>	62	40	19
<u>Nocardia hydrocarbonoxydans</u> and <u>Arthrobacter arthrobotryoides</u>	58	15	21
<u>Candida lipolytica</u> and <u>Cunninghamella elegans</u>	61	52	18
<u>Corynebacterium hydrocarboclastus</u> and <u>Pseudomonas desmolytica</u>	82	39	17

Several experiments utilized combinations of three or four cultures. The procedure was similar to that employing two cultures. With a single exception, the mixtures yielded results no better than mixtures involving but two species. The single exception was a mixture of Cunninghamella elegans, Arthrobacter arthrobotryoides, Nocardia hydrocarbonoxydans, and Pseudomonas desmolytica, which in seven days oxidized 91%, 73%, and 50% of crude oil, Bunker C, and marine diesel, respectively.

In those experiments allowed to incubate for a year at room temperature, the bacteria reached a peak activity after from 10 to 14 days and subsequently the activity significantly dropped, but nevertheless did continue at a slow rate for the entire year.

CONCLUSIONS

There appears to be no dearth of microbes, including bacteria, fungi, and yeasts that have the capacity to oxidize crude oil, Bunker C fuel, and marine diesel. In most experiments the percentage of crude oil oxidized

exceeded the percentage of Bunker C oxidized in the same period. In general, Bunker C was oxidized at a faster rate than marine diesel.

Many organisms that demonstrated an ability to survive in media with a hydrocarbon as a sole source of organic matter lacked a perceptible ability to degrade measurable amounts of hydrocarbon. Commonly, the combination of different species was more active than the activity of either of the component species alone. This effect appears to be synergistic. On the contrary, an antagonistic effect also occurs when the sum total of hydrocarbon degeneration is less than that achieved by any component of a mixed culture when used alone as a pure culture.

The author concludes that the continued endeavor of putting together different combinations of cultures will ultimately yield a mixture that, with appropriate nutrients, can be added in a lyophilized form to an oil spill, after physical methods of clean-up have been completed, to eliminate the last vestiges of oil.

REFERENCES

- Allen, A.A., Schlueter, R.S., and Mikolaj, P.G.. "Natural Oil Seepage at Coal Oil Point". Science 170, No. 3961, 1970, pp. 974-976.
- Merz, R.C.. Determination of the Quality of Oily Substances on Beaches and in Nearshore Waters. California State Water Pollution Control Board, 1959, Publication No. 21.
- O'Neill, T.B.. The Biodegradation of Oil in Seawater for Naval Pollution Control. Naval Civil Engineering Laboratory, Technical Note N-1195, Jan 1972.
- ZoBell, C.E.. Microbial Modification of Crude Oil in the Sea. Presented at the Joint API/FWPCA Conference on Prevention and Control of Oil Spills, Dec 1969.

OFFICE OF NAVAL RESEARCH
NAVAL BIOLOGY PROJECT
STANDARD DISTRIBUTION LIST

Number of copies:

- (12) Administrator, Defense Documentation Center
Cameron Station
Alexandria, VA 22314
- (6) Director, Naval Research Laboratory
Attention: Technical Information Division
Code 2627
Washington, D.C. 20375
- (6) Office of Naval Research
Code 102IP (ONRL DOC)
800 N. Quincy Street
Arlington, VA 22217
- (3) Office of Naval Research
Naval Biology Project
Code 443
Arlington, VA 22217
- (1) Office of Naval Research
Code 200
Arlington, VA 22217
- (1) Office of Naval Research Branch Office
495 Summer Street
Boston, MA 02100
- (1) Office of Naval Research Branch Office
536 South Clark Street
Chicago, IL 60605
- (1) Office of Naval Research Branch Office
1030 East Green Street
Pasadena, CA 91101
- (1) Director, Oceanic Biology Program (Code 214)
Naval Ocean Research & Development Activity
National Space Technology Laboratory
Bay St. Louis, MS 39520
- (1) Assistant Commander for Research & Development (Code 03)
Naval Facilities Engineering Command
200 Stovall Street
Alexandria, VA 22332

OFFICE OF NAVAL RESEARCH
NAVAL BIOLOGY PROJECT
STANDARD DISTRIBUTION LIST (Cont'd)

Number of copies:

- (1) Biological Sciences Staff (Code 101B)
Naval Facilities Engineering Command
200 Stovall Street
Alexandria, VA 22332
- (1) Scientific Library
Naval Biosciences Laboratory
Naval Supply Center
Oakland, California 94625
- (1) Technical Library
U. S. Army Natick Laboratories
Natick, MA 01760
- (1) Commander
Attention: Dr. Morthland
U. S. Army Research Office, Durham
Box CM, Duke Station
Durham, NC 27706
- (1) National Environmental Research Center
Edison Water Quality Research Division
Edison, NJ 08817
- (1) Agricultural & Marine Pollution Control Branch
Environmental Protection Agency
1901 Fort Myers Drive
Arlington, VA 22209
- (1) Technical Advisory Division
National Marine Fisheries Service
Department of Commerce
Washington, D.C. 20235
- (1) Director
Gulf Breeze Laboratory
Environmental Protection Agency
Sabine Island
Gulf Breeze, FL 32561
- (1) Matthew Stevenson
National Academy of Sciences
Room JH 538
2101 Constitution Avenue
Washington, D.C. 20418

OFFICE OF NAVAL RESEARCH
NAVAL BIOLOGY PROJECT
STANDARD DISTRIBUTION LIST

Number of copies:

- () Commandant, DAT
U. S. Coast Guard
400 Seventh Street, SW
Washington, DC 20511
- () Commandant, DAS
U. S. Coast Guard Research & Development Center
Avery Point
Groton, CT 06340
- () Office of the Oceanographer of the Navy
Code N5
732 North Washington Street
Alexandria, VA 22314
- () Director
U. S. Army Cold Regions Research and
Engineering Laboratory
Hanover, NH 03755
- () Commanding Officer
Attention: Dr. Hoogstraal
U. S. Naval Medical Research Unit #3
Camp Lejeune, NC 24582
- () Officer in Charge
Naval Disease Vector Control Center
Naval Air Station
Alameda, CA 94501
- () Chief, Entomology Research Branch
Preventive Medicine Division
U. S. Army Medical R & D Command
Forrestal Building
Washington, D.C. 20314
- () Office of Naval Research Resident Representative
2110 G. Street, NW
Washington, D.C. 20037