

ADA042715

12^{B.S.}

AD

COPY NO. 51


TECHNICAL REPORT ARLCD-TR-77039

PHOTOCHEMICAL STUDIES OF
SECONDARY NITRAMINES.

PART II. ULTRAVIOLET PHOTOLYSIS AND
OZONOLYSIS OF RDX IN AQUEOUS SOLUTIONS

ANDREW F. SMETANA
SURYANARAYANA BULUSU

JUNE 1977



US ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND
LARGE CALIBER
WEAPON SYSTEMS LABORATORY
DOVER, NEW JERSEY

APPROVED FOR PUBLIC RELEASE, DISIRIBUTION UNLIMITED.

AD No. _____
DDC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report/ARLCD-TR-77039 - Pt-2	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) PHOTOCHEMICAL STUDIES OF SECONDARY NITRAMINES. PART II. ULTRAVIOLET PHOTOLYSIS AND OZONOLYSIS OF RDX IN AQUEOUS SOLUTIONS.	5. TYPE OF REPORT & PERIOD COVERED Technical Report. Apr 75 - Dec 76,	
7. AUTHOR(s) Andrew F./Smetana Suryanarayana/Bulusu	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Large Caliber Weapons Systems Laboratory (DRDAR-LCE) US Army Armament Research and Development Command Dover, NJ 07801	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS 23 p.	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS OCMS 4932.05.4114.0	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE June 1977	
	13. NUMBER OF PAGES 29	
	15. SECURITY CLASS. (of this report) Unclassified	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Photolysis "Pink Water" Ozonolysis Ultraviolet photolysis Nitramines RDX		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of this study was to determine the feasibility of chemical degradation of RDX by ultraviolet-ozone process which had been previously developed for eliminating TNT from pink water. A study of UV photolysis at 254, 300 and 350 nm wavelengths, ozonolysis alone and a combination of O ₃ and UV (350 nm) on aqueous solutions of RDX was undertaken. The results indicate that O ₃ + UV in combination efficiently degrade RDX to gaseous products. These products are further compared with those obtained by photolysis and thermal decomposition of samples of crystalline RDX.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

The citation in this report of the names of commercial firms or commercially available products or services does not constitute official endorsement or approval of such commercial firms, products, or services by the US Government.

ACCESSION for	
NTIS	White Section ☒
DOC	B.I. Section ☐
CONTRACTS	☐
EXHIBITS	
FY	
EXHIBITION/AVAILABILITY NOTES	
A	

FOREWORD

This report details the work accomplished under the technical sponsorship (OCMS #4932.05.4114.0) of the Modernization and Special Technology Branch of the Manufacturing Technology Division, Large Caliber Weapons Systems Laboratory, ARRADCOM, Dover, NJ.

The report includes work on UV/Ozone process for chemical degradation of RDX as a part of the broader "pink water" pollution abatement problem at the Army ammunition loading and packaging plants.

TABLE OF CONTENTS

	Page No.
Introduction	1
Experimental and Results	2
Solution of RDX	2
Photolysis of Aqueous Solutions of RDX at 254 nm	2
Analysis of RDX	2
Relative Photolysis Rates at 254, 300 and 350 nm	3
Ozonolysis of Aqueous Solutions of RDX	3
Concurrent Photolysis and Ozonolysis	3
Analysis of Gaseous Products	4
GC/MS Analysis	5
Photolysis of Polycrystalline RDX	6
Results and Discussion	8
Photolysis of RDX in Solution	8
Ozonolysis	8
Degradation Products of RDX	9
Photolysis of Pure RDX	9
Thermal Decomposition of RDX	9
Conclusions	11
References	11
Distribution List	19

Tables

1	Ozonolysis, photolysis and their combined application for the chemical degradation of RDX	4
2	Mass spectra of the gaseous products of photolysis of RDX	5
3	GC/MS Analysis of photolyzed RDX solutions	6
4	Photolysis products of RDX at $\lambda = 254$ nm (ccmm)	7
5	Photolysis products of RDX at $\lambda = 310-330$ nm (ccmm)	7
6	Moles of product/mole of RDX normalized to 100% decomposition	10

Figures

1	UV spectra of RDX before and after photolysis	13
2	Absorption spectra of RDX solutions with photolysis	14
3	Spectra of aqueous RDX solutions by the direct and indirect methods	15
4	Change in absorbance of aqueous RDX solutions with photolysis/ozonolysis time	16
5	Spectra of (I) 1-nitroso - 3:5-dinitro-1:3:5-triazacyclohexane and of (II) 1:3:5-trinitroso-1:3:5-triazacyclohexane, both in methanol	17

INTRODUCTION

Since 1970 increased emphasis has been placed on eliminating pollution arising from munitions manufacturing and processing plants. One serious pollution problem is the removal of the nitro bodies from the so-called pink water which is the wash water from loading, assembling and packaging operations involving TNT. This water dissolves up to 150 ppm of TNT, about 80 ppm RDX and much smaller amounts of other nitro compounds. On exposure to sunlight, the photochemical reactions of TNT generate several colored species which impart the characteristic pink color to the water. Under a contract from the Volunteer Army Ammunition Plant (VAAP), Chatanooga, Tenn, IIT Research Institute (IITRI) studied ozonolysis (Ref 1) as a means to destroy the nitro compounds and showed that pink water at its normal pH of 5.6 was rendered colorless and more acidic (pH = 4-5) on treatment with ozone (O_3). However, neutralization to pH 7 returned new colored species which were presumed different from those formed photochemically. If the ozonolysis was carried out at an alkaline pH (~ 8 to 10.5) the color could be removed permanently but this process was inefficient in the use of ozone.

The IITRI project then experimented (Ref 2) with an ozone plus ultraviolet irradiation process to decontaminate the pink water, this combination process having been found previously (Ref 3) to be effective in oxidation of refractory organic materials in municipal waste waters. For these experiments synthetic pink water, made by dissolving TNT (160 ppm) in water, was used and the efficiency of the process was measured by the determination of the total organic carbon (TOC) as a function of O_3 + UV dosage. A germicidal lamp was used in these experiments as the source of ultraviolet light. A dramatic UV effect was observed, relative to the use of O_3 alone, and the initial total organic carbon levels of about 60 mg/l dropped to 17 mg/l in two hours. In addition, it was found that more than 85% of the lost organic carbon appeared as CO_2 , thus obviating the problem of secondary pollution from intermediate degradation products of TNT. The chemistry and engineering aspects of the UV + O_3 process were further investigated in the feasibility studies conducted by Westgate Research Corporation (Ref 4) and Walden Research Division (Ref 5). While further details of the above studies of the O_3 -UV process will not be included here, it is important to emphasize that this process was demonstrated to be effective in degrading the TNT and related nitro-aromatic compounds in pink water to innocuous end products. RDX, which is a common constituent of pink water (80 ppm) was not included in the above studies and the reactions of RDX under O_3 + UV oxidative treatment have not been investigated. Since it would be desirable to have a method common to both TNT and RDX, a study of the latter was initiated in this laboratory and the results are reported here.

The general approach for this RDX study consisted of an investigation of its gross photochemical degradation in aqueous solutions and, independently, its oxidative degradation by ozone. In each case, a determination of the products of degradation was attempted together with the time element for each process. It was hoped that this approach would lead to a better understanding of the chemistry involved. The combined effect of O_3 -UV was then studied. During this work a report (Ref 6) appeared in the literature describing the photolysis of tap water-dissolved RDX under laminar flow conditions. One other report of a brief study of RDX photolysis dealt with photolysis in organic solvents (Ref 7).

EXPERIMENTAL AND RESULTS

Solution of RDX

A saturated (50 mg/l) stock solution of pure RDX (> 99.8%) in distilled water was prepared. This solution was used, after appropriate dilution, for all the experiments except the irradiations of polycrystalline RDX. Absorption spectra were always measured after the solutions were diluted to give optimum absorption.

Photolysis of Aqueous Solutions of RDX at 254 nm

In the first set of experiments to photolyze RDX, a Rayonet photochemical reactor (Ref 8), Model RPR-100, consisting of a circular array of 16 low pressure Hg lamps (254 nm) with a total output of 35 watts was used. It was however, found that in this reactor even saturated solutions of RDX (approximately 50 mg/l) were fully depleted of RDX in 10 minutes (Fig 1). The subsequent experiments were carried out with a very low intensity spectroline quartz pencil lamp (UV) placed 4" from a 1 cm Beckmann quartz spectrophotometer cell containing the RDX solution. Figure 2 shows the disappearance of RDX absorption with increasing time of UV-irradiation. Under these conditions the degradation of RDX appears to be complete in less than one hour. Exact correlations of the UV energy with RDX disappearance were not attempted.

Analysis of RDX

Two methods, to be called A&B, were employed for the analysis of RDX remaining in the solutions. In method A the concentration was obtained directly from a calibration plot of concentration versus optical density (O.D.). In method B, illustrated in Figure 3 (curves 3 & 5) the absorbance of the experimental solution was measured at all times against the standard strating solution to compensate the

overlapping absorption edge near the λ_{max} of RDX. At the same time solvent and solution compartments of the spectrophotometer were interchanged for convenience in the recorded output. Thus while curves 1 and 2 in Figure 3 represent normal absorption of RDX solution before and after 30 minutes photolysis, respectively, curve 3 was obtained by the compensation method and its peak height is a direct measure of RDX consumed in the photolysis.

Relative Photolysis Rates at 254, 300 and 350 nm

The photolysis experiments were also extended to long wave UV (300,350 nm) radiation to more nearly simulate solar radiation. A comparative study of three wavelengths was accomplished using a Rayonet reactor with four lamps each of RPR-2537, RPR-3000 and RPR-3500 type with rated outputs of 9, 5 and 24 watts, respectively. Aqueous solutions of RDX at a concentration of 20 mg/l were irradiated in spectrophotometer cells placed at the center of the reactor. The results of this series of runs are summarized in Figure 4, curves 1 to 3.

Ozonolysis of Aqueous Solutions of RDX

A Welsbach, Model T-816 ozonizer (Ref 9) was used as the source of ozone. The ozone was generated by the passage of an electric discharge (70 volts, 68 watts) through oxygen flowing at a rate of 500 ml/min. Under these conditions, the ozone concentration in the exit flow was 45 mg/min as determined by iodometric analysis (Ref 9). The ozonolysis was typically conducted by bubbling the O_2-O_3 mixture through a gas wash bottle containing 250 ml of RDX solution at a concentration of 20 mg/l. Five ml portions taken at different times were analyzed for RDX content by UV spectrophotometry. The data from one such run are included in Figure 4, curve 4.

Concurrent Photolysis and Ozonolysis

For these experiments the 350 nm UV source which caused the lowest photolysis rate was used in conjunction with a $O_3 + O_2$ flow rate of 0.5 l/min (45 mg O_3 /min). The concentration of the RDX solution was 20 mg/l and, as the reaction progressed, the optical density of the solution at 238 nm was measured versus time. Table 1 summarizes these results and shows the relative rates of photolysis, ozonolysis and the combined application of both UV and O_3 . The numbers represent the progressive decrease in the optical density at 238 nm (λ_{max} of RDX).

Table 1

Ozonolysis, photolysis and their combined
application for the chemical degradation of RDX

Reaction time, hrs	Changes in optical density		
	Ozonolysis	Photolysis	Ozonolysis & photolysis
0	0	0	0
1	0.12	0.07	0.64
2	0.19	0.19	0.91
3	0.22	0.30	0.93
4	0.29	0.37	0.94
5	0.33	0.41	0.94
6	0.36	0.52	
7	0.38	0.57	
8	0.43	0.60	
9	0.46	0.64	
10	0.49	0.56	
11	0.52	0.66	
12		0.67	
13		0.70	
14		0.75	
15		0.75	

Analysis of Gaseous Products

Since UV-absorption spectroscopy indicated a drastic degradation of RDX molecule at 254 nm without giving rise to new absorption in the UV or visible region, it was decided to analyze the photolysis solution for gaseous products. A special cell was designed in which the solution could be outgassed, irradiated with UV and then outgassed a second time to collect the low molecular weight gaseous products. The gaseous products were analyzed using a DuPont 21-492 mass spectrometer. The significant portions of the mass spectrum in two runs, corrected for background are shown in Table 1.

Table 2

Mass spectra of the gaseous products of photolysis of RDX



<u>m/e*</u>	<u>Run 1</u> <u>Relative</u> <u>abundance</u>	<u>Run 2</u> <u>Relative</u> <u>abundance</u>
22	-	-
28	1230	960
29	32	50
30	840	785
31	40	14
44	1410	1300
45	20	19

These mass spectra indicate the principal products to be N_2O (m/e 44, 30), NO (m/e 30) and N_2/CO (m/e 28) with trace amounts of formaldehyde (m/e 30,29).

* Mass-to-charge ratio

GC/MS Analysis

Previous studies of thermal decomposition (Ref 10,11) of RDX identified as gaseous products N_2O , CH_2O , N_2 , CO, CO_2 , HCN and H_2O with a polymeric residue (~1%). One report (Ref 7) indicated the presence of trace amounts of two solid products, 1-nitroso-3,5-dinitro-1,3,5-triaza cyclohexane (I) and 1,3,5-trinitroso-1,3,5-triazacyclohexane (II). In the hope of detecting the same or closely related products under UV + O_3 degradation, GC and GC/MS analyses were attempted. Photolyzed aqueous solutions of RDX, as well as those subjected to both



O_3 and UV, were concentrated and analyzed by GC using a SE-30 column. The observed peaks, when admitted to a mass spectrometer, gave evidence

of only low molecular weight ($< m/e$ 44) species (Table 3). One sample partially decomposed by UV + O_3 was extracted with ethylene dichloride and the latter solution was concentrated and analyzed directly by mass spectrometry. The only species found in this sample was unreacted RDX with no evidence of impurities above the background level.

Table 3

GC/MS Analysis of photolyzed RDX solutions

Fragments m/e	Relative abundance, chart divisions		
	GC peak #1	GC peak #2	GC peak #3
28	1300	100	100
29	850	250	50
30	2100	800	170
42			30
44	2900	1000	190
54		60	100
74			130
81		90	130

Photolysis of Polycrystalline RDX

Several irradiations of polycrystalline samples of RDX were attempted in which the product analysis was expected to be more manageable. The irradiations were carried out in a quartz spectrometer cell (1 cm x 1 cm x 5 cm) attached to a vacuum manifold through a ground joint. The vacuum manifold was connected to the inlet system of the mass spectrometer (duPont 21-492). In each experiment 100 mg of RDX were placed in the cell and the latter evacuated. One sample was irradiated with 254 nm UV light (low pressure Hg lamp fitted with appropriate filters). The gas liberated by photolysis was analyzed by bleeding samples into the mass spectrometer at intervals of time up to 3 1/2 hours. The constituents of the gas were determined from the known cracking patterns and initial calibrations. Tables 4 and 5 show the detailed analysis.

Table 4

Photolysis products of RDX at $\lambda = 254$ nm (ccmm)

	Time (minutes)						
	30	60	90	120	150	180	210
N ₂	22.0	35.0	46.0	54.0	60.0	65.0	70
NO	18.0	27.8	34.0	39.0	42.0	46.0	49
N ₂ O	4.6	7.8	10.4	12.8	14.1	15.5	17.1
CO	3.9	8.8	11.1	14.1	18.0	19.5	20.5
CO ₂	2.4	5.5	8.1	9.8	11.5	13.6	15.2
HCN	0.5	0.9	1.2	1.4	1.6	1.7	1.9
HCHO	1.3	2.6	3.1	3.2	3.3	3.3	3.4
H ₂ O	0.2	0.3	0.5	0.6	0.8	1.1	1.5
H ₂	0.3	0.4	0.5	0.5	0.6	0.7	0.7

Table 5

Photolysis products of RDX at $\lambda = 310-330$ nm (ccmm)

	Time (minutes)						
	30	60	90	120	150	180	210
N ₂	10.0	30.5	53	72.1	9.12	107	121
NO	3.6	9.5	16.0	23.0	31.0	38	47
N ₂ O	2.6	8.1	13.6	19.3	24.5	28.4	33.3
CO	0.8	3.4	5.9	7.8	10.2	11.8	14.2
CO ₂	1.6	4.3	7.2	10.0	12.9	15.0	17.4
H ₂ O	0.1	0.5	1.0	1.5	2.2	2.9	3.9
H ₂	-	-	0.1	0.1	0.1	0.1	0.1

RESULTS AND DISCUSSION

Photolysis of RDX in Solution

The absorption spectra of RDX and other nitramines, as well as the photolysis of dimethylnitramine, were the subjects of earlier publications (Ref 12, 13) from this Division. Aqueous solutions of RDX showed a broad absorption extending to 300 nm in the UV with a λ_{max} at 238 nm and an extinction of 11,000. It was decided, therefore, to use UV sources in this wavelength range for the irradiation, e.g., 254, 300 and 350 nm. Since the earlier study of dimethylnitramine (DMNA) showed that a major photolytic product was the corresponding nitrosamine which absorbed in the UV at 235 and 370 nm, a preliminary experiment was run with RDX to determine whether a similar process occurred. There was no evidence of such species and it was possible to use the absorbance of RDX for analytical purposes.

Figures 1 and 2 clearly demonstrate the rapid degradation of RDX in aqueous solutions under the influence of 254 nm UV radiation. Indeed, with the use of as little as 35 watts output in these experiments most of the RDX was destroyed in about 10 minutes. This is seen more strikingly in Figure 3, curve 3 obtained by the method B of analysis.

The relative efficiencies of different wavelengths of UV light can readily be seen in Figure 4. It is apparent that the 350 nm radiation, even with 3 times the intensity of the 254 nm source, was much less effective in degrading RDX. This is understandable since the absorption edge for RDX lies in this wavelength range. It was found independently that solid RDX was very stable to 366 nm UV and evolved no gas in over 10 hours.

Ozonolysis

This study consisted of conducting photolysis and ozonolysis under identical conditions and then comparing these experiments with the combined reaction in order to determine the possible enhancement of the rate of degradation. It can be seen in Figure 4 that O_3 at a relatively high concentration was much slower in degradation of RDX when used alone. From curve 4 one can estimate that about 27 g. O_3 passed through the solution containing 5 mg RDX and yet, only ~ 50% of the RDX was consumed. This inefficiency was probably due to excessively high flow rate of O_3 which allowed little contact time with RDX.

Interestingly, an unmistakable synergistic effect was observed when O_3 and 350 nm UV were used together. It can be seen in Table 1 that each of the two reagents destroys approximately half of the

RDX in the solution in ~ 10 hours, but when used in combination, $\sim 95\%$ of the RDX was degraded in 5 hours. There has been some speculation about the role of singlet oxygen in this reaction (Ref 4) but the present work could not shed any light on this speculation.

Degradation Products of RDX

Curve 6 in Figure 2 shows that most of the degradation products of RDX were volatile because they could be driven out by sparging with N_2 . These spectra (Fig 2) also indicate that condensible products (I) and (II), the spectra of which are shown in Figure 5, were not formed in these experiments. Typical compositions of the gaseous products are shown in Tables 2 and 3. As indicated in the experimental section, they consist of low molecular weight species, the largest being N_2O . However, these analyses are not strictly quantitative.

Photolysis of Pure RDX

The aqueous solution of RDX used for this study was very dilute due to the low solubility of RDX in water. This, combined with the fact that RDX essentially degrades to gaseous products presented severe analytical problems. To avoid these problems, experiments were designed in which polycrystalline samples of RDX were directly photolyzed and the products determined quantitatively. It was possible by this means to establish the general nature of the chemical degradation products. Typical analyses are shown in Tables 4 and 5 and these products appear to be similar to those which gave rise to the mass spectral fragments shown in Tables 2 and 3.

Thermal Decomposition of RDX

It is noteworthy that the thermal decomposition products of RDX also bear a resemblance to those obtained in the photolysis and/or ozonolysis. Table 6 summarizes the thermal decomposition products found in an earlier study (Ref 7).

Table 6

Moles of product/mole of RDX normalized to 100% decomposition

<u>% RDX decomposed</u>	<u>CO₂</u>	<u>N₂O</u>	<u>N₂</u>	<u>NO</u>	<u>CO</u>	<u>HCN</u>	<u>CH₂O</u>	<u>H₂O</u>
19	.54	1.22	1.08	.23	.41	.23	--	--
30	.66	1.47	1.04	.49	.54	.23	--	1.67
69	.62	1.25	1.16	.49	.56	.16	.72	1.10
78	.50	1.26	.83	.44	.48	.09	--	--
93	.61	1.36	.99	.45	.52	.03	.97	.97

It is evident that in the photodegradation of RDX, in both aqueous solution and in the solid state, much smaller amounts of formaldehyde were formed than in the thermal decomposition. This may be an experimental artifact in that the HCHO produced in thermal decomposition runs could diffuse away from the heated region of the apparatus while in photolysis it could undergo secondary photo-reactions. Also, HCN was not observed in the present experiments in aqueous solution, whereas it was clearly detectable in the other experiments. Unfortunately, the analysis of products described in this report are not sufficiently quantitative to give exact yields. The limited scope of this work did not permit designing the large scale experiments needed for quantitative determinations. In view of the complexity of such experiments, it would be desirable to combine them with the UV + O₃ process for TNT. Past work on TNT (Ref 1) showed that most products are gaseous and experiments with TNT and RDX together would have the advantage of revealing secondary reactions among the products from the two compounds. Thus, while undesirable products like HCHO, N₂O, CO and HCN may be formed in these reactions, their relative importance can not be judged precisely until actual "pink water" samples are subjected to the UV + O₃ treatment.

CONCLUSIONS

The work reported here establishes that: (1) aqueous RDX solutions undergo rapid photochemical degradation with 254 nm UV light leading to low molecular weight gaseous products (2) ozone degrades RDX much less efficiently than UV (3) the combined use of O_3 and UV exhibits a marked synergistic effect in that the combined degradation rate is greater than the sum of the two individual degradation rates (4) photolysis at two wavelengths and thermal decomposition of crystalline RDX give products generally similar to those of the photolysis in solution, and finally (5) the appearance of HCHO and HCN is subject to variation, possibly due to secondary reactions. Condensable products do not seem to be formed in the reactions studied. A quantitative determination of gaseous products can only be attempted in large scale experiments. It thus appears that the O_3 and UV process in pink water will be effective in destroying both RDX and TNT.

REFERENCES

1. Ozonolysis of Pink Water, Phase I, Final Report, by Robert I. Brabets and Gayle E. Marks, IIT Research Institute Report IITRI-C6275, Chicago, IL 60616, 1974
2. Treatment of TNT-containing Waste Water, Proposal IITRI-75-67c, by Edward Fochtman and James E. Huff, IIT Research Institute, Chicago, IL 60616, 1974
3. Oxidation of Refractory Organic Materials by Ozone and UV Light, Final Report 7184, by Cecil G. Hewes, et al., Houston Research Inc., 7705 South Loop Street, Houston, TX 77012, 1974
4. Ultraviolet Catalyzed Oxidation of Munitions Waste Waters (UV-OX process), Proposal 2501, Westgate Research Corp, 13478 Beach Avenue, Marina Del Rey, CA 90291, February 1975. Submitted to: US Army, ARRADCOM, Dover, NJ 07801
5. Treatment of TNT Manufacturing Waste by O_3 , Proposal by Walden Research Division, Abcor Inc, Cambridge, MA, 1975. Submitted to US Army, ARRADCOM, Dover, NJ 07801
6. Pollution Potential of Explosives in Water, by E.F. Seymour, Report NSWC/WOL/X-155, Naval Surface Weapons Center, Silver Spring, MD, 1976
7. Studies in Composition B, Francis C. Rauch, et al, American Cyanamid Co, Stamford, CT 06904. Annual Reports 1969, 1970 and 1971

8. Rayonet Photochemical Reactor, Model #RPR-100, Southern New England Ultraviolet Company, 954 New Field Street, Middle Town, CT 06457
9. The Welsbach Laboratory Ozonator, Model T-816, The Welsbach Corp, Ozone Process Division, 3340 Stokley Street, Philadelphia, PA 19129. Manual of Applications
10. Bulusu Suryanarayana, Joseph R. Autera and Richard J. Graybush, Proceedings of the 1968 Army Science Conference, West Point, NY, Vol II, 423, June 1968
11. Bulusu Suryanarayana and R.J. Graybush, Proceedings of the 39th Congress on Industrial Chemistry, Brussels, Belgium, 1966
12. Photochemical Studies of Secondary Nitramines. 1., by K. Suryanarayanan and Suryanarayana Bulusu, PATR 4068, September 1970
13. Photolysis of Solid Dimethylnitramine: Nitrogen-15 Study and Evidence of Nitrosamine Rearrangement, K. Suryanarayanan and Suryanarayana Bulusu, J. Phys. Chem., 76, 496, 1972




Fig 1 UV spectra of RDX before and after photolysis




Fig 2 Absorption spectra of RDX solutions with photolysis




Fig 3 Spectra of aqueous RDX solutions by the direct and indirect methods




Fig 4 Change in absorbance of aqueous RDX solutions with photolysis/ozonolysis time




Fig. 5 Spectra of (I) 1-nitroso - 3:5-dinitro-1:3:5-triazacyclohexane and of (II) 1:3:5-trinitroso-1:3:5-triazacyclohexane, both in methanol

DISTRIBUTION LIST

Copy No.

Commander
US Army Armament Research and Development
Command
ATTN: DRDAR-TSS
DRDAR-CG
DRDAR-LCM
DRDAR-LCM-S
Dover, NJ 07801

1-5
6
7
8-28

Commander
US Army Materiel Development & Readiness
Command
ATTN: DARCOM-RD
DARCOM-RD-ES
DARCOM-IS-MD
DARCOM-PA-E
DARCOM-RP-I
5001 Eisenhower Avenue
Alexandria, VA 22333

29
30
31
32
33

Commander
USAMC Installations & Services Agency
ATTN: DARCOM-IS-RI-IU
Rock Island, IL 61201

34

Commander
US Army Armament Materiel Readiness Command
ATTN: DRSAR-NT
DRSAR-RDN
DRSAR-ISE
DRSAR-SC
DRSAR-EN
DRSAR-PPW
Rock Island, IL 61201

35
36
37-38
39
40
41

Project Manager for Munition Production
Base Modernization and Expansion
US Army Materiel Development & Readiness Command
ATTN: DARCOM-PM-PEN-EC
Dover, NJ 07801

42-43

Commander DA-DCSLOG ATTN: DALO-AMP Washington, DC 20310	44
Commander US Army Procurement Equipment Agency ATTN: DRXPE-MT Rock Island, IL 61201	45
Department of the Army ATTN: DAEN-ZCE Washington, DC 20310	46
Commander ARRADCOM, Chemical Systems Laboratory ATTN: DRDAR-TD Aberdeen Proving Ground, MD 21010	47-49
Commander Frankford Arsenal ATTN: SARFA-T-4200 Philadelphia, PA 19137	50
Defense Documentation Center Cameron Station Alexandria, VA 22314	51-62
Office Chief of Engineers ATTN: DAEN-MCZ-E Washington, DC 20314	63-64
District Engineer US Army Eng Dist, New York Corps of Engineers 26 Federal Plaza New York, NY 10007	65
District Engineer US Army Eng Dist, Baltimore Corps of Engineers P.O. Box 1715 Baltimore, MD 21203	66

District Engineer US Army Eng Dist, Norfolk Corps of Engineers 803 Front Street Norfolk, VA 23510	67
Commander US Army Environmental Hygiene Agency ATTN: USAEHA-E Aberdeen Proving Ground, MD 21010	68-69
Commander Holston Army Ammunition Plant ATTN: SARHO-E Kingsport, TN 37662	70
Commander Indiana Army Ammunition Plant ATTN: SARIN-OR Charlestown, IN 47111	71
Commander Iowa Army Ammunition Plant Middletown, IA 52638	72
Commander Joliet Army Ammunition Plant ATTN: SARJO-SS-E Joliet, IL 60436	73
Commander Lone Star Army Ammunition Plant ATTN: SARLS-IE Texarkana, TX 75501	74
Commander Louisiana Army Ammunition Plant ATTN: SARLA-S Shreveport, LA 71102	75
Commander Newport Army Ammunition Plant ATTN: SARNE-S Newport, IN 47966	76

Commander Radford Army Ammunition Plant ATTN: SARRA-IE Radford, VA 24141	77
Commander Volunteer Army Ammunition Plant ATTN: SARVO-T Chattanooga, TN 37401	78
Mr. Harry Ike 26 Federal Plaza - Room 837 New York, NY 10007	79
Mr. Ben Lacey Curtis Building, 2nd floor 6th and Walnut Streets Philadelphia, PA 19106	80
Commander Badger Army Ammunition Plant ATTN: SARBA Baraboo, WI 53913	81
Commander Cornhusker Army Ammunition Plant Grand Island, NE 68801	82
Commander Kansas Army Ammunition Plant Parsons, KS 63757	83
Commander Lake City Army Ammunition Plant Independence, MO 64056	84
Commander Louisiana Army Ammunition Plant Shreveport, LA 71102	85
Commander Milan Army Ammunition Plant ATTN: SARMI Milan, TN 38358	86
Commander Riverbank Army Ammunition Plant Riverbank, CA 95367	87

Commander Scranton Army Ammunition Plant Scranton, PA 18501	88
Commander Twin Cities Army Ammunition Plant Minneapolis, MN 55440	89
Commander Pine Bluff Arsenal Pine Bluff, AR 71601	90
Commander Watervliet Arsenal Watervliet, NY 12189	91
Commander Longhorn Army Ammunition Plant Marshall, TX 75670	92
Commander US Naval Weapons Center China Lake, CA 93555	93
Commander Naval Ammunition Depot Earle Colts Neck, NJ 07722	94
Commander Naval Ammunition Depot Hawthorn Hawthorn, NV 89415	95
Commander Newport Army Ammunition Plant Newport, IN 47966	96
Commander Naval Surface Weapons Center ATTN: Mr. John Hoffsomer Silver Springs, MD 20910	97-98
Commander US Army Natick Laboratories ATTN: Dr. A. Kaplan Natick, MA 01760	99-100

Commander US Army Natick Laboratories ATTN: Dr. N. McCormick Natick, MA 01760	101
US Army Corps of Engineers Construction Engineering Research Laboratory Environmental Engineering Group ATTN: Mr. Dan Nelson Champaign, IL 61820	102
US Army Corps of Engineers Construction Engineering Research Laboratory Environmental Engineering Group ATTN: Mr. Walter J. Mikucki Champaign, IL 61820	103
Commander Northern Division Naval Facilities Engineering Command US Naval Base ATTN: Mr. Vince Gasbaro Philadelphia, PA 19112	104
Commander Naval Ammunition Depot ATTN: Mr. Ellis Pardee McAlester, OK 74501	105
US Army Engineer District Mobile District Office ATTN: Mr. D. Shaw Mobile, AL 36600	106
Commander Naval Sea Systems Command Department of the Navy ATTN: SEA-9912E, B. Cocimano SEA-033, A. B. Amster Washington, DC 20360	107 108
Commander Naval Facilities Engineering Command ATTN: 03-A, Mr. Rockefeller PC-4, CDR J.B. Graff Washington, DC 20360	109 110

Commander Naval Civil Engineering Laboratories Director, Material Sciences Division Port Hueneme, CA 93043	111
Industrial Waste Technology Branch Edison Water Quality Research Laboratory National Environmental Research Center US Environmental Protection Agency Edison, NJ 08817	112-115
Office of Solid Waste Management Programs US Environmental Protection Agency Washington, DC 20460	116-119
Effluent Guidelines Division US Environmental Protection Agency Washington, DC 20460	120-121
Commander US Army Armament Research and Development Command ATTN: DRDAR-LCE Dover, NJ 07801	122-150