

AD-A044 435

WASHINGTON UNIV SEATTLE DEPT OF CHEMISTRY
VIBRATIONAL-TRANSLATIONAL ENERGY TRANSFER IN ATOM-POLYATOMIC MO--ETC(U)
AUG 77 I OREF, B S RABINOVITCH
NR092-549-TR05



F/6 20/8

N00014-75-C-0690


NL

UNCLASSIFIED

1 OF 1
AD
A044435



END
DATE
FILMED
10-77
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 044435

12

6
Vibrational-Translational Energy Transfer in Atom-Polyatomic
Molecule Collisions in Thermal Reaction Systems

10 by I. Oref and B. S. Rabinovitch

Prepared for publication in Chemical Physics

9 14
Technical Report No. NR092-549-TR05

Contract N00014-75-C-0690, NR092-549

15

11 15 August 1977

13 27 p.

Department of Chemistry, BG-10
University of Washington
Seattle, WA 98195

OFFICE OF NAVAL RESEARCH
Department of the Navy
Code 473
800 N. Quincy
Arlington, VA 22217

DDC
RECORDED
SEP 22 1977
B

AD No. —
DDC FILE COPY

Reproduction in whole or in part is permitted for any purpose of the
United States Government. This document has been approved for public release;
its distribution is unlimited.

370 274 —

mt

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NR092-549-TR05 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Vibrational-Translational Energy Transfer in Atom-Polyatomic Molecule Collisions in Thermal Reaction Systems	5. TYPE OF REPORT & PERIOD COVERED Technical	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) I. Oref and B. S. Rabinovitch	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0690 ✓ NR 092-549	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Professor B. S. Rabinovitch Department of Chemistry BG-10 University of Washington Seattle, WA 98195	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 473 Department of the Navy 800 N. Quincy Arlington, VA 22217	12. REPORT DATE 15 August 77	
	13. NUMBER OF PAGES 20	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted to Chemical Physics		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Angular momentum conservation Vibration-translation Transfer Energy Transfer Polyatomic molecules Statistical model Thermal systems		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Vibrational-translational energy transfer probabilities and collisional efficiencies are calculated for atom-polyatomic molecule collisions. It is assumed that a collision complex is formed and that the total internal vibra- tional energy is statistically distributed among all the modes of the complex. An attractive potential is assumed and account is taken of the centrifugal barrier. Conservation of system angular momentum is imposed. Convolution of the several thermal distribution functions is carried out and completeness and detailed balance are observed. Comparison of calculated quantum statistical		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014 6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

quantities with experiment is made for the thermal isomerization of methyl and ethyl isocyanide in the presence of heavy atomic bath gases, such as Xe or Ar, and semiquantitative agreement is found.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
	P. R. Section <input type="checkbox"/>
	<input type="checkbox"/>
DISTRIBUTION PRIORITY CODES	
	SPECIAL
A	

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

DISTRIBUTION LIST

Contractor University of Washington

NR 092-549

Contract Number N00014-75-C-0690

Date August 15, 1977

	<u>No. Copies</u>
Office of Naval Research Code 473 Arlington, VA 22217 Attn: Dr. Miller	10
Office of Naval Research Branch Office 1030 East Green Street Pasadena, CA 91106 Attn: Dr. R. Marcus	1
Office of Naval Research Branch Office 536 S. Clark Street Chicago, IL 60605 Attn: Dr. J. Smith	1
Defense Documentation Center Bldg. 5 Cameron Station Alexandria, VA 22314	12
Office of Naval Research Branch Office 495 Summer Street Boston, MA 02210 Attn: Dr. L. H. Peebles	1
Office of Naval Research Resident Representative University of Washington 3710 Brooklyn Avenue NE, Unit 2 Seattle, WA 98195	1
U. S. Naval Research Laboratory Code 2627 Washington, D.C. 20375	6
U. S. Naval Research Laboratory Code 2629 Washington, D.C. 20375	6
Naval Research Laboratory Code 6100 Washington, D.C. 20375	1
Naval Air Systems Command Code 440 Washington, D.C. 20360 Attn: Dr. H. Rosenwasser	1

No. Copies

Naval Sea Systems Command SEA-0331 Washington, D.C. 20362 Attn: Mr. J. Murrin	1
Naval Sea Systems Command SEA-0332 Washington, D.C. 20362 Attn: Dr. A. Amster	1
Naval Surface Weapons Center Research and Technology Dept.-WR Silver Spring, MD 20910	1
Naval Weapons Center Research Department Code 60 China Lake, CA 93555	1
Naval Weapons Center Code 608 China Lake, CA 93555 Attn: Ronald L. Derr	3
Air Force Office of Scientific Research Directorate of Aerospace Sciences Bolling Air Force Base Washington, D.C. 20332	1
Air Force Office of Scientific Research Directorate of Chemical Sciences Bolling Air Force Base Washington, D.C. 20332	1
Air Force Office of Scientific Research Directorate of Physics Bolling Air Force Base Washington, D.C. 20332	1
U.S. Army Research Office Chemistry Division P.O. Box 12211 Research Triangle Park, N.C. 27709	1
U.S. Army Research Office Physics Division P.O. Box 12211 Research Triangle Park, N.C. 27709	1
Professor S.N.B. Murthy Technical Director, Project SQID Purdue University Lafayette, IN 47907	5

Vibrational-Translational Energy Transfer in Atom - Polyatomic Molecule
Collisions in Thermal Reaction Systems¹

by I. Oref² and B. S. Rabinovitch

Department of Chemistry
University of Washington
Seattle, Washington 98195

Abstract

Vibrational-translational energy transfer probabilities and collisional efficiencies are calculated for atom-polyatomic molecule collisions. It is assumed that a collision complex is formed and that the total internal vibrational energy is statistically distributed among all the modes of the complex. An attractive potential is assumed and account is taken of the centrifugal barrier. Conservation of system angular momentum is imposed. Convolution of the several thermal distribution functions is carried out and completeness and detailed balance are observed. Comparison of calculated quantum statistical quantities with experiment is made for the thermal isomerization of methyl and ethyl isocyanide in the presence of heavy atomic bath gases, such as Xe or Ar, and semiquantitative agreement is found.

Introduction

Intermolecular vibrational-translational energy transfer involving polyatomic molecules is of importance in both reactive and non-reactive gas-phase chemical systems. A few models and methods which treat the details of this process have been developed in recent years and are briefly recalled.

In one theory which has been applied to the results of inelastic scattering in molecular beam experiments,³⁻⁵ it is assumed that the dominating interactions are long range r^{-5} forces and that a collision complex is formed in which available energy is statistically distributed. Decomposition of the complex occurs when the translational energy in the reaction coordinate exceeds the height of the centrifugal barrier. Extension of a statistical model to include rotations has recently been made for atom-molecule exchange.⁶

The inelastic scattering of vibrationally excited KBr by small non-polar partners⁴ has been explained by assuming near-collinear impulsive collisions with an anharmonic oscillator; the effect of rotation was not taken explicitly into consideration. In this model, certain collisional configurations were excluded. The effect was to limit the size of the energy step transferred per collision. The theory does not always agree with experiment as it is predicted to be more efficient than Ar, contrary to experimental findings. However, collision of vibrationally excited KBr with polyatomic bath molecules^{7,8} could be explained by an extension of Lin and Rabinovitch's model⁹ of complex formation with partial participation of vibrational modes of the polyatomic bath molecules. It was found by Fisk, et al.,^{7,8} that more than one model was necessary to explain all of their experimental results.

A statistical phase space theory¹⁰ for a bimolecular reaction of a spherical top with spherical top or linear species has been applied to ion-molecule reaction with use of an r^{-6} potential. It was found that the reaction cross section for $C_2H_4^+ + C_2H_4$ agrees only qualitatively with theory inasmuch as the

experimental energy dependence of the cross section decreases much faster than predicted by the theory. Better agreement of theory and experiment was obtained for the product kinetic energy distribution; however, improvements in the results in polyatomic systems requires a detailed knowledge of the potential hypersurface at the reaction zone.

Another model¹¹ for the lifetime distribution of intermediate complexes formed in ion-molecule reactions involves an r^{-4} potential and takes into consideration the centrifugal barrier. No account of the initial internal energy distribution was reported and a step-ladder type deactivation was assumed.

Quack and Troe¹² have developed a so-called statistical adiabatic channel model in which a strong complex is formed between a highly interacting atom-diatom pair. With use of an extrapolation procedure of a Morse potential between molecular complex and the separate product fragments, they obtained the energy distributions of the products.

Another approach which utilized a Morse or truncated harmonic oscillator potential function for atom-diatom collisions was developed by Johnston and Birks.¹³ It was concluded that only levels far below the threshold energy for decomposition contribute significantly to reaction.

Still another treatment for collisional energy transfer has been given by Kohmaler, Nowak and Sire¹⁴ in which the atom-polyatomic molecule interaction is approximated by a Morse interaction potential for an atom and a damped harmonic oscillator.

Several empirical distribution functions have been applied to describe collisional energy transfer probabilities, of which the step-ladder and exponential forms are the most commonly used.^{15,16} Penner and Forst¹⁷ developed a theory for vibrational-rotational energy transfer in the thermal dissociation of diatomic molecules. The transition probabilities are empirical exponential or gaussian forms. It is found that rotational, in addition to vibrational

energy transfer leads to enhancement of energy transfer. A comprehensive review of the subject has been given recently.^{18a}

In the quantum statistical model which is developed here, and which uses the potential of refs 1 and 3, collisional energy transfer for thermal systems is calculated for atom-molecule interactions under certain simplifying assumptions. A thermal system in which one of the colliding pair is a polyatomic molecule, with many vibrational-rotational internal modes, is difficult to handle. The internal energy distribution of the molecule must be convoluted with that for relative translational energy. The calculations are carried beyond the computation of energy transfer probability distribution functions, for each given initial condition, to the evaluation of average energy transfer efficiency quantities. Finally, the requirements of completeness and of detailed balance, which are usually omitted in calculations in non-thermal experimental systems, must be adhered to as a very stringent criterion on otherwise discretionary aspects of the calculation. In this circumstance, some simplifying features must be imposed on the calculation.

Not many suitable and reliable experimental systems are known in thermal studies. Application of the model is made to the isocyanide system. Collisional energy transfer probabilities and collisional efficiencies have been calculated for methyl and ethyl isocyanide molecules with Xe bath gas, and results are compared with experiment. The agreement is found to be satisfactory from a qualitative and semi-quantitative point of view. The results help to delineate some details of the energy transfer process.

Since this manuscript was written, a paper has just appeared by Holmlid and Rynefors^{18b} who have also imposed the condition of completeness on the model of refs 3,4 and have given a re-interpretation of molecular beam experiments. They used a classical density of states treatment to facilitate their calculations.

Description of the model

Consider a collision between a bath atom and a reactant molecule having translational energy, E , in the center of mass coordinate system. The bath atoms are in excess so that thermal equilibrium is established after each collision.

p. 4

If the collision trajectory is such that the impact parameter is below a given value, b_{\max} , a short-lived complex is formed. Calculations are made for the case that molecular angular momentum is small relative to system angular momentum. The energy content of the complex (exclusive of its overall rotational energy and of translational energy in the reaction coordinate) is randomized. The complex decomposes with translational energy in the reaction coordinate, E' , different from the energy in the incoming channel. The distribution in E' ($= B + \epsilon'$), the sum of rotational and translational contributions to total relative translational energy in the outgoing channel (indicated always by a prime), is calculated.

The system potential energy is

$$V(r) = -\frac{C}{R^6} + \frac{J^2}{2\mu r^2} \quad (1)$$

where C is a constant and μ is the reduced mass. The top of the centrifugal barrier is at $r = r_{\max}'$ corresponding to $dV/dr = 0$. The rotational energy of the collision complex, B , is

$$B = J^2 / (54c\mu)^{3/2} \quad (2)$$

The collision complex has maximum angular momentum corresponding to

$$J_{\max} = (54c\mu^3 E^2)^{1/6} \quad (3)$$

The rate coefficient for the dissociation of the complex in the region at r_{\max}' is given, according to RRKM theory, by

$$k(E_T, B, \epsilon') = \frac{f(\epsilon') N^+(\epsilon') N^+(\epsilon_T - B - \epsilon')}{N^+(\epsilon_T - B)} \quad (4)$$

where, $N(E)$ = density of states at energy E ; $E_T = W + E$; ϵ' = translational energy in reaction coordinate; W = internal energy of the polyatomic molecule;

and superscripts \dagger and $*$ denote complex and excited molecules, respectively. The product $f(\epsilon^{\dagger})N^*(\epsilon^{\dagger})$ may be written¹⁶ as $1/h$. The total rate coefficient for each E_T and B is given by summing eq. 4 over all possible ϵ^{\dagger} ,

$$k(E_T, B) = \frac{1}{h} \frac{\sum_{\epsilon^{\dagger}=0}^{E_T-B} N^*(E_T-B-\epsilon^{\dagger})}{N^*(E_T-B)} \quad (5)$$

The probability of the separating pair having translational energy E^{\dagger} , $p(E_T, E^{\dagger})$, is calculated under the assumption $B = B^{\dagger}$. Then $E^{\dagger} = B + \epsilon^{\dagger}$, and $P(E_T, B)$ is

$$P(E_T, B) = \frac{k(E_T, B, \epsilon^{\dagger}) P(B)}{k(E_T, B)} \quad (6)$$

where the probability of a given B, $P(B)$, is

$$P(B) = 2\alpha_0 \alpha / J^2_{\max} \quad (7)$$

Incorporating eqs. 2-5 and 7 into 6 yields the quantum statistical expression,

$$P(E_T, B) = \frac{N^*(E_T-B-\epsilon^{\dagger})}{\sum_{\epsilon^{\dagger}=0} N^*(E_T-B-\epsilon^{\dagger})} \frac{2}{3} \left(\frac{B}{E} \right)^{2/3} \frac{dB}{B} \quad (8)$$

and

$$P(E_T, E^{\dagger}) = \int_{ij} P_{ij}(E_T, B); \quad (9)$$

the summation runs over all j values of B and i values of ϵ^{\dagger} for all combinations such that $B_j + \epsilon_i^{\dagger} = E^{\dagger}$.

For a thermal system, averages of the thermal distributions must be evaluated. The average of $P(E_T, E^{\dagger})$ must be taken over all values of the incoming translational energy, in the center of mass coordinate system, and is given by

$$P(W, E^{\dagger}) = \frac{\int_0^{\infty} P(E_T, E^{\dagger}) P(E) dE}{\int_0^{\infty} P(W, E^{\dagger}) dE^{\dagger}} \quad (10)$$

$$\text{where } P(E) dE = E e^{-E/RT} dE / (RT)^2 \quad (11)$$

Calculation of Energy Transfer Probabilities

The outcome of an energy transfer event is evaluated by comparing the input probability $P(E)$ with the output $P(W, E^{\dagger})$ such that for any ΔE (where $\Delta E = E^{\dagger} - E$), activation occurs when $P(E) > P(W, E^{\dagger})$ at the same energy, and deactivation occurs when the reverse relation holds. The probability of obtaining a given ΔE for all combinations of E and E^{\dagger} is

$$P(W, \Delta E) = \int P(W, E^{\dagger}) \cdot P(E); \quad (12)$$

where 1 is a running index over all combinations of E^{\dagger} and E, i.e., E_j^{\dagger} and E_k , such that $\Delta E = E_j^{\dagger} - E_k$; $P(W, \Delta E)$ is properly normalized by the conservation relation, $\int_{-\infty}^{\infty} P(W, \Delta E) d\Delta E = 1$, and transitions with $\Delta E = 0$ are included.

The Fraction of Molecules above E_0

The fraction of molecules at each W which after collision are above the threshold energy for decomposition can be found by

$$Q(W) = \int_{E_0-W}^{\infty} P(W, \Delta E) d(\Delta E) \quad (13)$$

Equation 13 is weighted by the internal energy population and integrated over all internal energies to yield the total number of molecules with energy above E_0 which, in the low pressure limit of a thermal reaction, is proportional to the rate coefficient for decomposition. The collisional efficiency is then defined as:

$$B = \int_0^{E_0} Q(W) B_{SS}(W) dW / \int_{E_0}^{\infty} B(W) dW \quad (14)$$

$$\text{where, } B(W) = N(W) e^{-W/RT} \quad (15)$$

and the integral in the denominator is simply the strong collision expression divided by ω . $B_{SS}(W)$ is the steady state population which differs from the equilibrium population because of the depletion of levels below E_0 due to reaction. $B_{SS}(W)$ may be found¹³ by iteration of the expression

$$B_{SS}(W_i) = \frac{\int_0^{E_0} P(W_j, \Delta W) B(W_j) \delta' dW_j}{\int_0^{E_0} P(W_i, \Delta E) \delta' d(\Delta E) - \int_0^{E_0} Q(W_i) B(W_i) dW + Q(W_i)} ; \delta' = 0, \text{ for } j = i; \quad (16)$$

$$= 1, \text{ for } j \neq i;$$

where $\Delta W = |W_j - W_i|$, and subscripts j and i signify different states of the internal energy.

Exploitation of the above relations is subject to economic realities. To obtain $P(E_i, E')$ for each allowed combination of W, E, B, e^+ , mentioned before, far exceeded our computational resources and some judicious approximations had to be made. We have taken the approach which was suggested in ref. 1 and used the integral form of $P(B)$ to obtain

$$P(B) = \left(\frac{E'}{E} \right)^{2/3}, \text{ for } E' < E ; \quad (17)$$

$$= 1, \text{ for } E' > 1$$

Implicit in this approach is, of course, the fact that the number of states in the denominator of eq. 8 is assumed to be independent of B . To abide by the physics of the system, detailed balance was checked independently and made to agree by introducing a correction function for the approximation which is

introduced by the use of eq. 17. The final expression, which was used instead of eq. 9, is

$$P(E_i, E') = N(E_i - E') \left(\frac{E'}{E} \right)^{2/3} ; E' < E \quad (18)$$

$$= N(E_i - E') (E' - E)^{\alpha} ; E' > E$$

The value of α was determined by the required detailed balance of the system.

Results and Discussion

As examples, we consider the thermal isomerization of CH_3NC and $\text{C}_2\text{H}_5\text{NC}$, with heavy atoms such as Ar or Xe as bath molecules.

The thermal population distribution of CH_3NC and $\text{C}_2\text{H}_5\text{NC}$.

At low and moderate temperatures, the equilibrium population distribution function $B(E)$ for $\text{C}_2\text{H}_5\text{NC}$, and especially for CH_3NC , is discrete at the low energy end. This behavior of polyatomic molecules is unimportant if the energy region of interest is close to the value of E_0 . However, if collisions with large ΔE are important,^{13,17,19} the quantum effect will appear as fluctuations in the function $P(W, \Delta E)$.

The normalized distribution function $B(E)$ was independently checked by comparing the average thermal energy $\langle E \rangle$ computed from eq 19, evaluated by

$$\langle E \rangle = \int_0^{\infty} EB(W) dW \quad (19)$$

numerical integration, with the value obtained from the statistical - thermodynamic relation,

$$\langle E \rangle = - R d(\ln Q_V) / d(T^{-1}) \quad (20)$$

where Q_V is the vibration partition function. Agreement within 10% was required and was obtained by varying the grain size of W .

The steady state populations were evaluated from eq. 16 and were used to find B in eq. 14. It was found that there is a modest (from a few percent up to 10-15%) depletion of the populations in the levels just below E_0 ; these levels are still the major contributors to reaction, as discussed later.

Detailed balance and the value of α

The present model uses the parameter, α (eq. 18), in the probability function $P(W, E')$. It was introduced to account for the fact that the integral over all values of B (eq. 9) is replaced by an approximate expression. An independent check of the appropriate value of α is given by detailed balance,

$$B(E_j)P_{ji} = B(E_i)P_{ij} \quad (21)$$

It was required that detailed balance should be obeyed to within an average of 10% over the whole interesting range of energies. This stringent requirement on the value of α is an additional condition usually not imposed in the treatment of non-thermal systems. (If α were treated merely as an adjustable parameter, exact concordance between theory and experiment could be obtained.) The values of α used are 1.30 ± 0.05 and 1.20 ± 0.05 for CH_3NC at 550°K and 850°K, respectively, and 1.20 ± 0.05 and 1.00 ± 0.05 for $\text{C}_2\text{H}_5\text{NC}$ at 550°K and 850°K, respectively.

Translational energy distribution in the outgoing channel.

Sample curves for the translational energy distribution in the exit channel are given in Figures 1-3. For comparison, the thermal translational energy in the incoming channel is shown as well. The value of α used at each temperature was chosen independently to give the best agreement with detailed balance. The "cooling" effect of collisions can be seen clearly when the higher energy curve at 38 kcal mole⁻¹ internal energy is compared with the one at 9.2 kcal mole⁻¹.

The former has a large high energy tail because the leaving atom carries with it a relatively larger amount of translational kinetic energy than for the case of collision with a low internal energy polyatomic molecule. Large values of ΔE are not completely negligible for collisions of an atom with high internal energy molecules. This is a consequence of a mechanism in which statistical distribution prevails.

Collision probabilities $P(W, \Delta E)$

Two illustrations of the collision probability function $P(W, \Delta E)$ are given in Figure 4 for two cases: a low internal energy case (9.2 kcal mole⁻¹) and a high internal energy case (38 kcal mole⁻¹). The shapes of the curves vary with the energy content of the molecule, from almost symmetric for the low case to the skewed high energy one. The collision probability for the removal of large ΔE is larger, the higher the internal energy.

Fraction of molecules at each energy brought above E_0 by collision.

The total fraction of molecules which decompose at the low pressure limit is the sum of the fraction of molecules, at each internal energy, which are transported to energies above E_0 , as given by $Q(W)$ (eq. 13). For the isocyanides, E_0 is 38 kcal mole⁻¹. Figure 5 refers to CH_3NC and Figure 6 to $\text{C}_2\text{H}_5\text{NC}$. It can be seen that there is some contribution to reaction from levels which are far below E_0 . The extent of the contribution depends on the reacting molecules and the temperature at which the reaction is taking place.

The collision probabilities which were calculated in this work are intermediate in character between models which are used currently¹⁵ or were developed before for simple atom-diatom dissociation systems.^{13,17,19} An example of the latter case is the work of Johnston and Birks¹³ who considered three models for energy transfer: ladder climbing with a truncated harmonic oscillator, and truncated and Morse oscillators with all transitions allowed. Their conclusion was

suggests that in this model the "small" molecule behavior of CH_3NC may affect the results of energy transfer through the value of α in eq. 18, which in turn influences the collisional probabilities. Moreover, the simplifying assumption regarding conservation of angular momentum cannot be supported in general. One source of the difference between the efficiencies calculated here and the lower values calculated by Lin and Rabinovitch²¹ is the fact that a semi-empirical cut-off in the maximum amount of energy transferred to the reaction coordinate was applied by the latter authors. The cut-off was introduced as a restriction on a purely statistical model.

Dependence of β on the temperature

With increase of temperature, the substrate population shifts to higher energies; $P(k, E')$ has a larger high energy tail and the incoming translational energy distribution also shifts to higher energy (see Figure 2) to an extent that makes deactivation less probable at energies below E_0 . As a result, the value of β decreases with increase in temperature — from values of $\beta = 0.84$ and 0.92 at 550°K to values of $\beta = 0.66$ and 0.70 at 850°K , for CH_3NC and $\text{C}_2\text{H}_5\text{NC}$, respectively. Very little quantitative or qualitative information on the temperature dependence of β in any thermal system is available. For the isocyanides, a decrease in β from 0.29 to 0.21 was measured²² over the range from 483°K to 600°K in the $\text{CH}_3\text{NC-He}$ system so that the ratios are in semi-quantitative agreement with the calculation.

Conclusions

A model for energy transfer is presented which essentially contains no adjustable parameters. All probability functions were evaluated quantum statistically; completeness and detailed balance are adhered to strictly. It is found that collisional efficiencies, β , agree qualitatively and semi-quantitatively, in size and in trend, with experimental values in the $\text{CH}_3\text{NC-C}_2\text{H}_5\text{NC}$ systems and with the effect of temperature increases; it is found that small

that a truncated harmonic oscillator model with all transitions allowed provides the best agreement with the experimental results; only levels far below E_0 were found to contribute significantly to reaction. According to their model, the populations of levels just below E_0 are completely depleted and do not contribute to steady state reaction. Similar arguments were put forward by Kiefer, et al.¹⁹ Penner and Forst¹⁷ maintained that the depletion of the higher state is due not to the probability of vibrational energy transfer but to the effect of rotational energy transfer.

As mentioned earlier, popular mathematical expressions in use for collisional transfer probabilities are the empirical step-ladder and exponential models. As is clear from Figures 5,6, in the present model there is really no single group of levels which contribute to reaction; there is a distribution of ΔE values which is a function of the temperature and the individual molecule-atom pair which are involved in the collision process. The present results can be considered as intermediate between the two cases.

The collisional efficiency, β

As an independent check of the calculations, the average β was calculated for each collision pair: for $\text{C}_2\text{H}_5\text{NC}$, $\beta = 0.92 \pm 0.05$ at 550°K , as compared with the experimental value²⁰ $0.39-0.44$ for Ar and Xe at 504°K ; for CH_3NC , $\beta = 0.84 \pm 0.05$ at the same temperature, compared to the experimental results⁸ $0.31-0.25$.

If the density of states were the only factor which entered into the collisional efficiency, one would expect $\beta_{\text{CH}_3\text{NC}} > \beta_{\text{C}_2\text{H}_5\text{NC}}$; CH_3NC has fewer degrees of freedom than $\text{C}_2\text{H}_5\text{NC}$ and therefore the average energy per oscillator is larger and leads to higher probabilities for decomposition of the collision complex. That the experimental and calculational results agree, i.e., that $\beta_{\text{CH}_3\text{NC}} < \beta_{\text{C}_2\text{H}_5\text{NC}}$,¹¹

energy jumps are more important than large ones, as in the exponential or poisson models, but the latter cannot be neglected altogether.

This is the general form expected for energy transfer by weak/atomic bath colliders.¹⁸ The relative contribution to reaction from each initial energy level depends on the specifics of the colliding pair and the temperature. The higher the temperature, the less important is the contribution to reaction of the low-lying vibration states. A population depletion of levels below E_0 is found but their relative contribution is still high.

Acknowledgement. One of us (I.D.) wishes to thank the Department of Chemistry, University of Washington for its hospitality and the U.S.-Israel Binational Science Foundation for their support.

References

1. Work supported by the Office of Naval Research and in part by the National Science Foundation.
2. On sabbatical leave from the Department of Chemistry, Technion, Haifa, Israel.
3. a) S. A. Safran, N. D. Weinstein, D. R. Herschbach and J. C. Tully, *Chem. Phys. Lett.*, **12**, 564 (1972); S. S. Riley and D. R. Herschbach, *J. Chem. Phys.*, **58**, 27 (1973).
4. T. Donohue, M. S. Chow and G. A. Fisk, *Chem. Phys.*, **2**, 271 (1973).
5. J. C. Tully, *J. Chem. Phys.*, **61**, 61 (1974); **62**, 1893 (1975).
6. D. A. Case and D. R. Herschbach, *J. Chem. Phys.*, **64**, 4212 (1975).
7. F. F. Crim, M. S. Chou and G. A. Fisk, *Chem. Phys.*, **2**, 283 (1973).
8. F. F. Crim, H. B. Bente and G. A. Fisk, *J. Phys. Chem.*, **78**, 2438 (1974).
9. Y. N. Lin and B. S. Rabinovitch, *J. Phys. Chem.*, **74**, 3151 (1970).
10. W. J. Chesnovich and M. T. Bowers, *J. Am. Chem. Soc.*, **98**, 8301 (1976) and references cited therein.
11. G. G. Meisels, F. Botz and E. F. Hecke, submitted.
12. M. Quack and T. Troe, *Ber. Bunsenges. Physik. Chem.*, **78**, 240 (1974); **79**, 170 (1975).
13. H. S. Johnston and J. Birks, *Acc. Chem. Res.*, **5**, 327 (1972).
14. a) E. O. Sire and G. H. Kohlmaier, *Ber. Bunsenges. Physik. Chem.*, **80**, 504 (1976).
b) G. H. Kohlmaier, E. Nowak and E. O. Sire, *ibid.*, **80**, 515 (1976).
15. D. C. Tardy and B. S. Rabinovitch, *J. Chem. Phys.*, **45**, 3720 (1966); **48**, 1282 (1968).
16. P. J. Robinson and K. A. Holbrook, *Unimolecular Reactions*, Wiley-Interscience, London, (1972).
17. A. P. Penner and W. Forst, *Chem. Phys.*, **11**, 243 (1975); **13**, 51 (1976).
18. a) D. C. Tardy and B. S. Rabinovitch, *Chem. Revs.*, **77**, xxx (1977).
b) L. Holmlid and K. Rynefors, *Chem. Phys.*, **19**, 261 (1977).
19. J. H. Kiefer, H. P. C. Joosten and W. D. Bräsears, *Chem. Phys. Lett.*, **30**, 424 (1975).
20. S. P. Pavlow and B. S. Rabinovitch, *J. Phys. Chem.*, **75**, 1366 (1971).
21. S. C. Chan, J. T. Bryant and B. S. Rabinovitch, *J. Phys. Chem.*, **74**, 2055 (1970).

Figure 1. CH_3NC molecule at 550°K : 1) Thermal translational distribution in center of mass coordinate system; 2) $P(W, E')$ at $W = 9.2 \text{ kcal mole}^{-1}$, $\alpha = 1.3$; 3) $P(W, E')$ at $W = 38 \text{ kcal mole}^{-1}$, $\alpha = 1.3$.

Figure 2. CH_3NC molecule at 850°K : 1) Thermal translational distribution in center of mass coordinate system; 2) $P(W, E')$ at $9.2 \text{ kcal mole}^{-1}$, $\alpha = 1.2$; 3) $P(W, E')$ at $38 \text{ kcal mole}^{-1}$, $\alpha = 1.2$.

Figure 3. $\text{C}_2\text{H}_5\text{NC}$ molecule at 550°K : 1) Thermal translational distribution in center of mass coordinate system; 2) $P(W, E')$ at $9.2 \text{ kcal mole}^{-1}$, $\alpha = 1.2$; 3) $P(W, E')$ at $38 \text{ kcal mole}^{-1}$, $\alpha = 1.2$.

Figure 4. Energy transfer probabilities $P(W, \Delta E)$ for $\text{C}_2\text{H}_5\text{NC}$ at 550 and $\alpha = 1.2$: 1) $W = 9.2 \text{ kcal mole}^{-1}$; 2) $W = 38 \text{ kcal mole}^{-1}$.

Figure 5. Fraction of CH_3NC molecules of above E_0 , $Q(W)$, for a given internal energy W : 1) left hand scale, $T = 850^\circ\text{K}$, $\alpha = 1.2$; 2) right hand scale, $T = 550^\circ\text{K}$, $\alpha = 1.2$.

Figure 6. Fraction of $\text{C}_2\text{H}_5\text{NC}$ molecules above E_0 , $Q(W)$, for a given internal energy W : 1) right hand scale, $T = 550^\circ\text{K}$, $\alpha = 1.2$; 2) $T = 850^\circ\text{K}$, $\alpha = 1.0$.




Fig 3





Fig 2






Fig 4. *Original data*

