

AD-A047 728

PENNSYLVANIA STATE UNIV UNIVERSITY PARK COLL OF BUSIN--ETC F/G 12/1
NON-COOPERATIVE STOCHASTIC DOMINANCE GAMES.(U)

NOV 77 P C FISHBURN
TR-27

N00014-75-C-0857
NL

UNCLASSIFIED

| OF |
AD
A047 728



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 27	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NON-COOPERATIVE STOCHASTIC DOMINANCE GAMES		5. TYPE OF REPORT & PERIOD Interim
7. AUTHOR(s) Peter C. Fishburn		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Pennsylvania State University University Park, Pennsylvania 16802		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0857
11. CONTROLLING OFFICE NAME AND ADDRESS (code 613B: SJW) Office of Naval Research, Dept. of the Navy 800 N. Quincy Street Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 121407
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1977
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Distribution of this document is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-cooperative games Stochastic dominance Utility theory		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A non-cooperative stochastic dominance game is a non-cooperative game in which the only knowledge about the players' preferences and risk attitudes is presumed to be their preference orders on the set of n-tuples of pure strategies. Stochastic dominance equilibria are defined in terms of mixed strategies for the players that are efficient in the stochastic dominance sense against the strategies of the other players. It is shown that the set of SD equilibria equals all Nash equilibria that can be obtained from combinations of utility functions that are consistent with the		

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A 047728

AD NO. DDC FILE COPY

DDC
RECEIVED
DEC 19 1977
RECEIVED
C Dnext
Page

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. players' known preference orders. The latter part of the report looks at antagonistic stochastic dominance games in which some combination of consistent utility functions is zero-sum over the n-tuples of pure strategies.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ACCESSION NO.	
NTIS	Write Section ☒
DDC	Data Section ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

14
TR-27

9
TECHNICAL REPORT NO. 27

6
NON-COOPERATIVE STOCHASTIC DOMINANCE GAMES

10
Peter C. Fishburn
College of Business Administration
The Pennsylvania State University
University Park, Pennsylvania 16802

11
Nov 77

12
21p.

15
This research was sponsored by the Office of Naval Research, ONR
Contract Number N00014-75-C-0857, ONR Contract Authority identification
number NRO47-112/11-24-76.

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

407 732

DDC
RECEIVED
DEC 19 1977
RECEIVED
D

1. Introduction

This paper investigates equilibrium theory for non-cooperative games in which the information about the players' preferences is confined to their preference orders on the set of pure strategy combinations. Players' risk attitudes, as represented by von Neumann-Morgenstern utility functions [von Neumann and Morgenstern, 1947; Fishburn, 1976], are not presumed to be known in the analysis. We shall assume a finite number $n \geq 2$ of players, indexed by $i = 1, 2, \dots, n$, and use the following notations and assumptions throughout:

X_i , a nonempty finite set of pure strategies for i ;

P_i , the set of all mixed strategies (probability distributions on X_i) for i ;

$>_i$, an asymmetric weak order for i on $X = \prod_{i=1}^n X_i$, so that $>_i$ is asymmetric ($x >_i y \Rightarrow \text{not } (y >_i x)$) and negatively transitive ($x >_i y$ and $z \in X \Rightarrow x >_i z$ or $z >_i y$), with $x >_i y$ signifying that i prefers x to y ;

U_i , the set of all real valued functions on X that preserve $>_i$, so that $u_i \in U_i$ if and only if, for all $x, y \in X$, $x >_i y$ iff $u_i(x) > u_i(y)$.

In addition, $P = \prod_{i=1}^n P_i$ is the set of all n -tuples of mixed strategies $p = (p_1, \dots, p_n)$, and u_i is extended from X to P multilinearly as $u_i(p) = \sum_{x_1, \dots, x_n} u_i(x_1, \dots, x_n) p_1(x_1) \dots p_n(x_n)$. When $u_i \in U_i$ is individual i 's von Neumann-Morgenstern utility function on X , his preferences on P are such that he prefers p to q iff $u_i(p) > u_i(q)$.

The usual non-cooperative theory, as presented by Nash [1951] and Luce and Raiffa [1957], among others, assumes the knowledge of all players' risk attitudes. That is, in addition to $(\succ_1, \dots, \succ_n)$, it assumes that each player's von Neumann-Morgenstern utility function $u_i^* \in U_i$ (unique up to positive affine transformations) is known. Although weakening or deletion of this strong and often unrealistic assumption raises many interesting possibilities for analysis, I shall focus here on the case in which $(\succ_1, \dots, \succ_n)$ represents the state of relevant information on the preferences and risk attitudes of the players. Because absence of knowledge about an individual's risk attitudes in the risky decision context corresponds to analysis by first degree stochastic dominance [Whitmore and Findlay, 1977] from an expected-utility viewpoint, I shall refer to a non-cooperative game based on $(\succ_1, \dots, \succ_n)$ as a stochastic dominance game. The relationship between stochastic dominance and equilibria based on $(\succ_1, \dots, \succ_n)$ will become clear as we proceed.

Our focus on $(\succ_1, \dots, \succ_n)$ or on the set $U_1 \times \dots \times U_n$ of all utility functions consistent with $(\succ_1, \dots, \succ_n)$ can be interpreted in several ways. First, if the only available public information on the players' preferences and risk attitudes consists of $(\succ_1, \dots, \succ_n)$, then it seems reasonable to look at the game from this viewpoint so far as a "public" analysis is concerned. Alternatively, if the purpose of the analysis is to examine the collection of all non-cooperative games whose utility-function n-tuples are consistent with $(\succ_1, \dots, \succ_n)$, then XU_1 provides the relevant information base for utilities. Another way to look at our setting is to suppose that every player knows every other player's preferences on X but is ignorant of their risk attitudes and knows that every other player is ignorant of his risk attitudes. Even if every player knows his own risk attitudes, it can be argued that each should use XU_1 as his primary

basis for analysis. For example, when $n = 2$, player 1 may presume that player 2's actual information about utilities is $U_1 \times \{u_2^*\}$, but since 1 does not know which $u_2^* \in U_2$ obtains and since 2 knows that 1 does not know which $u_2^* \in U_2$ obtains, both should proceed as if any $(u_1, u_2) \in U_1 \times U_2$ might obtain. This of course need not preclude either player from looking at the $U_1 \times U_2$ -based analysis from the perspective of his own true u_1^* , but in doing this he should bear in mind that the other player does not know his u_1^* .

The paper is organized as follows. The next section presents the basic stochastic dominance relations, defines the set $SD(>_1, \dots, >_n)$ of SD equilibria for the stochastic dominance game as the set of all $p \in P$ such that p_1 is SD-efficient against $p_{(-1)} = (p_1, \dots, p_{i-1}, p_{i+1}, \dots, p_n)$, and then proves that the set of SD equilibria equals the union of all Nash equilibria for the different $(u_1, \dots, u_n) \in \prod U_i$. The section concludes with comments on uniform equilibria, which are Nash equilibria for every possible choice of $(u_1, \dots, u_n)$.

The latter part of the paper looks at antagonistic stochastic dominance games, which are stochastic dominance games in which some $(u_1, \dots, u_n) \in \prod U_i$ has $\sum_i u_i(x) = 0$ for all $x \in X$. Section 3 presents a characterization of antagonism in terms of stochastic dominance relations on the set P^+ of all probability distributions on X . It then identifies some aspects of two-person antagonistic games that are similar to aspects of two-person zero-sum games. The final section presents two examples of two-person antagonistic games that illustrate preceding results and definitions.

2. Stochastic Dominance and Equilibria

This section justifies the phrase "stochastic dominance games" by showing that a straightforward application of stochastic dominance concepts to a game

whose analysis is based on $(\succ_1, \dots, \succ_n)$ is tantamount to a Nash-equilibrium analysis of all non-cooperative games whose utility functions are congruent with $(\succ_1, \dots, \succ_n)$. The latter part of the section discusses SD equilibria that are Nash equilibria for all possible risk attitudes of the players.

The basic results for stochastic dominance that we shall use will be developed first. Given probability distributions α and β on a nonempty finite set A , and an asymmetric weak order $\succ_0$ on A , we say that α stochastically dominates β with respect to $\succ_0$, and write this as $\alpha D(\succ_0) \beta$, if and only if for every $y \in A$ the probability that α yields an $x \in A$ worse than y ($y \succ_0 x$) is no greater than the probability that β yields an $x \in A$ that is worse than y . Thus

$$\alpha D(\succ_0) \beta \text{ iff } \sum_{\{x: y \succ_0 x\}} \alpha(x) \leq \sum_{\{x: y \succ_0 x\}} \beta(x) \text{ for all } y \in A.$$

Strict stochastic dominance is denoted by $SD(\succ_0)$, with $\alpha SD(\succ_0) \beta$ iff $\alpha D(\succ_0) \beta$ and not $[\beta D(\succ_0) \alpha]$.

Two main lemmas on stochastic dominance will be needed in this section. The first lemma identifies a primary reason for interest in the stochastic dominance idea in connection with expected utility theory. It has been proved by Lehmann [1955], Quirk and Saposnik [1962], and Fishburn [1964], among others. The second lemma is tantamount to Proposition 2 in Fishburn [1974b]. In the lemmas, $\alpha, \beta, \alpha_k, \beta_k$ are probability distributions on finite A , $\succ_0$ is an asymmetric weak order on A , $U_0 = \{\text{all real valued functions } u_0 \text{ on } A \text{ that have } u_0(x) > u_0(y) \text{ iff } x \succ_0 y, \text{ for all } x, y \in A\}$, and $u_0(\alpha) = \sum u_0(x)\alpha(x)$.

Lemma 1: $\alpha D(\succ_0) \beta$ iff $u_0(\alpha) \geq u_0(\beta)$ for all $u_0 \in U_0$; $\alpha SD(\succ_0) \beta$ iff $u_0(\alpha) > u_0(\beta)$ for all $u_0 \in U_0$.

Lemma 2: If $\sum_{k=1}^m \lambda_k \alpha_k \text{ SD}(>_0) \sum_{k=1}^m \lambda_k \beta_k$ for no $(\lambda_1, \dots, \lambda_m)$ for which $\sum \lambda_k = 1$ and $\lambda_k \geq 0$ for $k = 1, \dots, m$, then there is a $u_0 \in U_0$ such that $u_0(\beta_k) \geq u_0(\alpha_k)$ for $k = 1, \dots, m$.

In the non-cooperative game context of the present paper, each $p \in P$ induces a probability distribution on X in the natural way, so for all $p, q \in P$ and for all $i \in \{1, \dots, n\}$ we define $D(>_i)$ by analogy as

$$p D(>_i) q \text{ iff } \sum_{\{x: y >_i x\}} p_1(x_1) \dots p_n(x_n) \leq \sum_{\{x: y >_i x\}} q_1(x_1) \dots q_n(x_n)$$

for all $y \in X$,

with $p \text{ SD}(>_i) q$ iff $p D(>_i) q$ and not $[q D(>_i) p]$. For each $p \in P$ and $i \in \{1, \dots, n\}$ let $p_{(i)} = (p_1, \dots, p_{i-1}, p_{i+1}, \dots, p_n)$ and write p as $(p_i, p_{(i)})$. With $P_{(i)}$ the set of all such $p_{(i)}$, we shall let $S_i(p_{(i)})$ be the set of mixed strategies for player i that are $\text{SD}(>_i)$ efficient against $p_{(i)} \in P_{(i)}$, so that

$$p_i \in S_i(p_{(i)}) \text{ iff } (q_i, p_{(i)}) \text{ SD}(>_i) (p_i, p_{(i)}) \text{ for no } q_i \in P_i.$$

It is easily seen that $S_i(p_{(i)})$ must contain a pure strategy in P_i and that the correspondence $S_i: P_{(i)} \rightarrow P_i$ is not generally continuous.

On the basis of the foregoing we shall say that $p \in P$ is an SD equilibrium if and only if each p_i is $\text{SD}(>_i)$ efficient against $p_{(i)}$. The set $\text{SD}(>_1, \dots, >_n)$ of all SD equilibria for $(>_1, \dots, >_n)$ is

$$\text{SD}(>_1, \dots, >_n) = \{p \in P: p_i \in S_i(p_{(i)}) \text{ for } i = 1, \dots, n\}.$$

Juxtaposed against this definition, let $S(u_1, \dots, u_n)$ be the set of Nash equilibria in P for $(u_1, \dots, u_n)$ so that

$$S(u_1, \dots, u_n) = \{p \in P: u_i(p_i, p_{(i)}) \geq u_i(q_i, p_{(i)}) \text{ for all } q_i \in P_i$$

and all $i \in \{1, \dots, n\}\}$.

We know from Nash [1951] that $S(u_1, \dots, u_n)$ is never empty. We know also from Nash [1951], Gale [1953], and Luce and Raiffa [1957], among others, that if $S(u_1, \dots, u_n)$ contains more than one equilibrium, then the elements in $S(u_1, \dots, u_n)$ might not be interchangeable or equivalent, and in fact it is possible to have $u_i(p) > u_i(q)$ for all i when $p, q \in S(u_1, \dots, u_n)$. The difficulties that these phenomena pose for the analysis of non-cooperative games are discussed at length by Luce and Raiffa [Chapter 5, Section 7.8], and I shall not dwell on them here.

The basic relationship between SD equilibria and Nash equilibria is given by the following theorem. This theorem shows that $SD(>_1, \dots, >_n)$ is never empty, and it provides a way of computing all SD equilibria from the Nash equilibria.

Theorem 1: $SD(>_1, \dots, >_n) = U\{S(u_1, \dots, u_n): (u_1, \dots, u_n) \in \prod_{i=1}^n U_i\}$.

Proof: Suppose first that $p \notin SD(>_1, \dots, >_n)$, so that $p_i \notin S_i(p_{(i)})$ for some i . Given $p_i \notin S_i(p_{(i)})$, Lemma 1 implies that $u_i(q_i, p_{(i)}) > u_i(p)$ for some $q_i \in P_i$ and for all $u_i \in U_i$, so that $p \notin US(u_1, \dots, u_n)$. Therefore $p \in US(u_1, \dots, u_n) \Rightarrow p \in SD(>_1, \dots, >_n)$. Suppose next that $p \in SD(>_1, \dots, >_n)$, so that $p_i \in S_i(p_{(i)})$ for all i , and consider $i = 1$. Let $X_1 = \{x_{11}, x_{12}, \dots, x_{1K}\}$, let $p^k = (x_{1k}, p_{(1)})$ and $q^k = p$ for $k = 1, \dots, K$ --where $(x_{1k}, p_{(1)})$ stands for

$(r_1, p_{(1)})$ with $r_1(x_{1k}) = 1$ —and let $\lambda_k = q_1(x_{1k})$ for all k . Since $p_1 \in S_1(p_{(1)})$, there is no $q_1 \in P_1$ for which $(q_1, p_{(1)})$ SD($>_1$) p , which is to say that there is no $(\lambda_1, \dots, \lambda_K) = (q_1(x_{11}), \dots, q_1(x_{1K}))$ for which $\sum \lambda_k p^k$ SD($>_1$) $\sum \lambda_k q^k$. Hence, by Lemma 2, there is a $u_1 \in U_1$ for which $u_1(p) \geq u_1(x_1, p_{(1)})$ for all $x_1 \in X_1$, which by linearity implies that $u_1(p) \geq u_1(q_1, p_{(1)})$ for all $q_1 \in P_1$. A similar application of Lemma 2 to each $i > 1$ shows that there is a $u_i \in U_i$ such that $u_i(p) \geq u_i(q_i, p_{(i)})$ for all $q_i \in P_i$. Thus $p \in \text{SD}(>_1, \dots, >_n) \Rightarrow p \in S(u_1, \dots, u_n)$ for some $(u_1, \dots, u_n) \in \prod U_i$, and the proof is complete.

To illustrate further the connection between stochastic dominance and Nash equilibria we define $p \in P$ as a uniform equilibrium for $(>_1, \dots, >_n)$ if and only if $p \in S(u_1, \dots, u_n)$ for all $(u_1, \dots, u_n) \in \prod U_i$. Uniform equilibria are attractive from the stochastic dominance viewpoint since they are Nash equilibria for all possible risk attitudes of the players that are consistent with $(>_1, \dots, >_n)$. However, uniform equilibria may not exist for an SD game, and even if they do exist then some of them may not be desirable from the players' viewpoints. For example, in Nash's sixth example [1951, p. 292], the two-person game with $X_i = \{x_i, y_i\}$ for $i = 1, 2$ with $(x_1, x_2) >_1 (x_1, y_2) \sim_1 (y_1, x_2) \sim_1 (y_1, y_2)$ for $i = 1, 2$ [where $\sim_i$ denotes indifference for i] has uniform pure equilibria (x_1, x_2) and (y_1, y_2) , the latter of which is obviously inferior.

The definition just given says that p is a uniform equilibrium iff $u_i(p) \geq u_i(x_i, p_{(i)})$ for all $u_i \in U_i$, all $x_i \in X_i$ and all $i \in \{1, \dots, n\}$. Lemma 1 then implies that p is a uniform equilibrium iff $p D(>_i)(x_i, p_{(i)})$ for all x_i and all i . The definition of $D(>_i)$ then says that p is a uniform equilibrium iff

$$\sum_{\{z: y >_i z\}} p_1(z_1) \dots p_n(z_n) \leq \min_{X_i} \sum_{\{z_{(i)}: y >_i (x_i, z_{(i)})\}} p_1(z_1) \dots p_{i-1}(z_{i-1}) p_{i+1}(z_{i+1}) \dots p_n(z_n)$$

for all $y \in X$ and all i , where $z_{(i)}$ is z with z_i omitted and $(x_i, z_{(i)}) = (z_1, \dots, z_{i-1}, x_i, z_{i+1}, \dots, z_n)$. The preceding inequality must in fact be an equality, and by looking only at the x_i for which $p_i(x_i) > 0$, it is easily seen that a necessary but not sufficient condition for p to be a uniform equilibrium is that it satisfy the following system of equalities:

$$\begin{aligned} \sum_{\{z_{(i)} : y >_1 (x_i, z_{(i)})\}} p_1(z_1) \dots p_{i-1}(z_{i-1}) p_{i+1}(z_{i+1}) \dots p_n(z_n) \\ = \sum_{\{z_{(i)} : y >_1 (x'_i, z_{(i)})\}} p_1(z_1) \dots p_{i-1}(z_{i-1}) p_{i+1}(z_{i+1}) \dots p_n(z_n) \end{aligned}$$

for all $y \in X$, all $x_i, x'_i \in X_i$ for which $p_i(x_i)p_i(x'_i) > 0$, and all $i \in \{1, \dots, n\}$.

In the two-person game with $X_i = \{x_i, y_i\}$, $(x_1, x_2) \sim_1 (y_1, y_2) >_1 (x_1, y_2) \sim_1 (y_1, x_2)$ and $(x_1, y_2) \sim_2 (y_1, x_2) >_2 (x_1, x_2) \sim_2 (y_1, y_2)$, there is a unique uniform equilibrium that has $p_i(x_i) = p_i(y_i) = 1/2$ for each i . This equilibrium is the only Nash equilibrium for each $(u_1, u_2) \in U_1 \times U_2$ in what amounts to a zero-sum game.

3. Antagonistic Games

The preceding example illustrates a two-person antagonistic game, which is a two-person game in which the players' preference orders on X are the duals (converses, inverses) of each other so that $x >_1 y$ iff $y >_2 x$, for all $x, y \in X$. Since $>_1$ is the dual of $>_2$ if and only if there are $u_1 \in U_1$ and $u_2 \in U_2$ such that $u_1(x) + u_2(x) = 0$ for all $x \in X$, two-person antagonistic games in the $(>_1, >_2)$ setting correspond in a natural way to two-person zero-sum games in the traditional setting. However, since non-constant-sum (u_1, u_2) pairs will exist for any two-person antagonistic game that has $x >_1 y >_1 z$ for some

$x, y, z \in X$, the analysis of such games from the stochastic dominance viewpoint is more involved than the zero-sum analysis. We return to this shortly.

In general I shall use the zero-sum basis to define antagonism and say that $(\succ_1, \dots, \succ_n)$ is antagonistic if and only if there exists $(u_1, \dots, u_n) \in \prod U_i$ for which $\sum_i u_i(x) = 0$ for all $x \in X$. For $n \geq 3$ this definition is not tantamount to the requirement that, for all $x, y \in X$, if $x \succ_i y$ for some i then $y \succ_j x$ for some $j \neq i$ although every antagonistic $(\succ_1, \dots, \succ_n)$ must satisfy this requirement. For example, if $n = 3$, if $\{a, b, c, x, y, z\} \subseteq X$, and if the three players have preference orders whose restrictions on $\{a, \dots, z\}$ by decreasing preference are

1. axbycz
2. bzcxya
3. cyazbx,

then $(\succ_1, \succ_2, \succ_3)$ cannot be antagonistic since it must be true that $\sum_i [u_i(a) + u_i(b) + u_i(c)] > \sum_i [u_i(x) + u_i(y) + u_i(z)]$, but for all $r, s \in \{a, \dots, z\}$ with $r \neq s$ there are i and j such that $r \succ_i s$ and $s \succ_j r$. If α and β are distributions on X with $\alpha(a) = \alpha(b) = \alpha(c) = 1/3$ and $\beta(x) = \beta(y) = \beta(z) = 1/3$, then $\alpha \text{ SD}(\succ_i) \beta$ for $i = 1, 2, 3$. On the other hand, if $(\succ_1, \dots, \succ_n)$ is antagonistic, then this type of uniform SD result is impossible. This is stated precisely in the following theorem, which characterizes antagonism in terms of stochastic dominance. In the theorem, P^+ denotes the set of all probability distributions on X , and the $D(\succ_i)$ and $SD(\succ_i)$ are to be viewed in terms of the definitions of $D(\succ_0)$ and $SD(\succ_0)$ near the beginning of section 2.

In general, P^+ will contain many distributions that do not correspond to any $p \in P$. For example, the distribution α on $X = \{x_1, y_1\} \times \{x_2, y_2\}$ that has $\alpha(x_1, x_2) = \alpha(y_1, y_2) = 1/2$ cannot be realized by any $(p_1, p_2) \in P$.

Theorem 2: $(>_1, \dots, >_n)$ is antagonistic if and only if there do not exist $\alpha, \beta \in P^+$ such that $\alpha D(>_i) \beta$ for all i along with $\alpha SD(>_i) \beta$ for some i .

The proof of this theorem uses a generalization of Lemma 2 whose proof is similar to the proof of Lemma 5 in Fishburn [1974a] and will not be redrawn here. The new lemma is as follows.

Lemma 3: Suppose that U is a nonempty convex cone of real valued functions on X , so that $au + bv \in U$ whenever $u, v \in U$ and $a, b > 0$, and that $>^+$ on P^+ is defined by $\alpha >^+ \beta$ iff $u(\alpha) > u(\beta)$ for all $u \in U$, where u on P^+ is the linear extension of u on X . Then for each $m \in \{1, 2, \dots\}$ there exists $(\lambda_1, \dots, \lambda_m)$ with $\lambda_k \geq 0$ for all k and $\sum \lambda_k = 1$ such that $\sum_1^m \lambda_k \alpha^k >^+ \sum_1^m \lambda_k \beta^k$, where the α^k and β^k are in P^+ , if and only if for every $u \in U$ there is a $k \in \{1, \dots, m\}$ such that $u(\alpha^k) > u(\beta^k)$.

Proof of Theorem 2: Suppose first that $(>_1, \dots, >_n)$ is antagonistic and $\alpha D(>_i) \beta$ for all i . Then, by Lemma 1, $u_i(\alpha) \geq u_i(\beta)$ for all $u_i \in U_i$ and all i , so that $\sum_i u_i(\alpha) \geq \sum_i u_i(\beta)$. By the definition of antagonism let the $u_i^0 \in U_i$ satisfy $\sum_i u_i^0(x) = 0$ for all $x \in X$. Then $\sum_i u_i^0(\alpha) = 0 = \sum_i u_i^0(\beta)$, which along with $u_i^0(\alpha) \geq u_i^0(\beta)$ for all i implies that $u_i^0(\alpha) = u_i^0(\beta)$ for all i . The second part of Lemma 1 then says that not $[\alpha SD(>_i) \beta]$ for every i , and in fact we get both $\alpha D(>_i) \beta$ and $\beta D(>_i) \alpha$ for all i .

For the converse proof let $U = \{\sum_i u_i : u_i \in U_i \text{ for all } i\}$, which is a nonempty convex cone of real valued functions on X , and let $>^+$ on P^+ be as

defined in Lemma 3. Now if there exist $\alpha, \beta \in P^+$ such that $u(\alpha) > u(\beta)$ for all $u \in U$, or $\alpha >^+ \beta$, then $u_i(\alpha) \geq u_i(\beta)$ for all $u_i \in U_i$ and all i --hence $\alpha D(>_i) \beta$ for all i --and, in addition, $u_i(\alpha) > u_i(\beta)$ for some u_i , which by Lemma 1 says that $\alpha SD(>_i) \beta$ for this i . Hence suppose that there do not exist $\alpha, \beta \in P^+$ such that $\alpha D(>_i) \beta$ for all i and $\alpha SD(>_i) \beta$ for some i . Then, as just proved, there do not exist $\alpha, \beta \in P^+$ such that $\alpha >^+ \beta$. Consequently, with $X = \{x^1, \dots, x^K\}$ and with $[x^k]$ the distribution in P^+ that assigns probability 1 to x^k , there does not exist a K -by- K ρ matrix with $\rho_{jk} \geq 0$ for all $j, k \in \{1, \dots, K\}$ and $\sum_j \sum_k \rho_{jk} = 1$ such that $\sum_k \sum_j \rho_{jk} [x^k] >^+ \sum_j \sum_k \rho_{jk} [x^j]$. Therefore, by Lemma 3 with $m = K^2$, there exists a $u \in U$ such that $u([x^k]) \leq u([x^j])$ for all j and k . (In other words, with $x^{jk} = x^k$ and $y^{jk} = x^j$, Lemma 3 implies that $u(x^{jk}) \leq u(y^{jk})$ for all (j, k) pairs, so that $u(x^k) \leq u(x^j)$ for all j and k .) Hence $u(x^k) = u(x^j)$ for all j and k so that u is constant on X . Since $u = \sum u_i$ for some $u_i \in U_i$, it follows that $(>_1, \dots, >_n)$ must be antagonistic. This completes the proof.

Just as traditional zero-sum games have features that are not shared by general games, antagonistic games exhibit aspects that are not shared by general non-cooperative games whose analyses are based on $(>_1, \dots, >_n)$. A case in point is seen from Theorem 2, which says in part that if $(>_1, \dots, >_n)$ is antagonistic then it cannot be true for $p, q \in P$ that $p D(>_i) q$ for all i and $p SD(>_i) q$ for some i . In particular, if $(>_1, \dots, >_n)$ is antagonistic and $p, q \in SD(>_1, \dots, >_n)$ then it must be false that $p D(>_i) q$ for all i and $p SD(>_i) q$ for some i . On the other hand it is easy to construct games with non-antagonistic $(>_1, \dots, >_n)$ that have $p, q \in SD(>_1, \dots, >_n)$ along with $p SD(>_i) q$ for all i .

There are also important differences between two-person antagonistic games and n -person antagonistic games for $n \geq 3$ that reflect some of the differences

between two-person zero-sum games and n -person zero-sum games for $n \geq 3$.

We note first some results for two-person antagonistic games.

Theorem 3: Suppose $n = 2$ and $(\succ_1, \succ_2)$ is antagonistic. If $e = (e_1, e_2)$ is a pure-strategies equilibrium in $SD(\succ_1, \succ_2)$, and if $p \in SD(\succ_1, \succ_2)$, then both $p \succ_1 e$ and $e \succ_1 p$ must be false for $i = 1$ and for $i = 2$. In addition, all pure-strategies equilibria are interchangeable and equivalent. That is, if $e = (e_1, e_2)$ and $e' = (e'_1, e'_2)$ are pure-strategies equilibria in $SD(\succ_1, \succ_2)$, then so are (e_1, e'_2) and (e'_1, e_2) , and $(e_1, e_2) \sim_i (e'_1, e'_2)$ for $i = 1, 2$, where $x \sim_i y$ iff neither $x \succ_i y$ nor $y \succ_i x$.

Proof: Suppose with $(\succ_1, \succ_2)$ antagonistic that $e \in X$ and $p \in P$ are such that $p \succ_1 e$, and that $e, p \in SD(\succ_1, \succ_2)$. (By a slight abuse of notation we consider $e \in X$ to be in P also, where $e \in P$ signifies the $(q_1, q_2) \in P$ that has $q_1(e_1) = q_2(e_2) = 1$.) Then Lemma 1 and $p \succ_1 e$ imply that $u_1(p) > u_1(e)$ for all $u_1 \in U_1$. Also, since $e \in SD(\succ_1, \succ_2)$ and Theorem 1 imply that $u_1(e) \geq u_1(x_1, e_2)$ for all $x_1 \in X_1$ for some (hence for all) $u_1 \in U_1$, we get $u_1(e) \geq u_1(p_1, e_1)$ for all $u_1 \in U_1$. In addition, since $p \in SD(\succ_1, \succ_2)$, there exists $u_2 \in U_2$ such that $u_2(p_1, p_2) \geq u_2(p_1, e_2)$ and hence there is a $u_1 \in U_1$, e.g. $u_1 = -u_2$, such that $u_1(p_1, e_2) \geq u_1(p)$. Altogether then the hypotheses of this paragraph imply that there is a $u_1 \in U_1$ for which $u_1(e) \geq u_1(p_1, e_2) \geq u_1(p) > u_1(e)$, which is absurd. Hence $e, p \in SD(\succ_1, \succ_2)$ and $(\succ_1, \succ_2)$ antagonistic imply not $[p \succ_1 e]$. Under the same hypotheses, similar proofs show that none of $p \succ_2 e$, $e \succ_2 p$ and $e \succ_1 p$ can be true. The proof of the second part of Theorem 3 is similar to the simple footnote proof on page 66 in Luce and Raiffa [1957] and will be omitted.

The two-person results of Theorem 3 do not generalize in any straightforward way to antagonistic games with $n \geq 3$. For example, it is easy to construct such games for $n = 3$ that have exactly two pure-strategies equilibria in $SD(>_1, >_2, >_3)$, say (e_1, e_2, e_3) and (e'_1, e'_2, e'_3) , with $e_i \neq e'_i$ for $i = 1, 2, 3$ and with $e >_1 e'$, $e >_2 e'$ and $e' >_3 e$. The latter imply of course that $e SD(>_1) e'$ for $i = 1, 2$ and that $e' SD(>_3) e$.

Hence the difficulties encountered with n -person antagonistic games for $n \geq 3$ are not unlike those faced by n -person non-cooperative zero-sum games for $n \geq 3$. But this still leaves open the possibility that two-person antagonistic games have certain nice features beyond those already noted in Theorem 3. As shown in the next section, this possibility is largely unfulfilled.

4. Two-Person Antagonistic Examples

This section presents two examples for two-person antagonistic games that illustrate the theory developed in preceding sections and show that such games lack certain properties that hold for two-person zero-sum games. The first example is a prototypical 2-by-2 game with no saddle point in pure strategies; the other has a pure-strategies equilibrium. In each case $>_1$ is identified by an order-preserving u_1 function on $X = X_1 \times X_2$ that is not intended to reflect the risk attitudes of the two players. For convenience we shall let $a_{ij} = u_1(x)$ when x is composed of the i th pure strategy for player 1 and the j th pure strategy of player 2. Hence each matrix is an a_{ij} matrix. Similarly, $b_{ij} = u_2(x)$ when x is composed of the i th pure strategy for player 1 and the j th pure strategy of player 2. Finally, we shall let λ_i and μ_j be respectively the probability in a mixed strategy for player 1 for his i th pure strategy and the probability in a mixed strategy for player 2 for his j th pure strategy.

Example 1: To illustrate Theorem 1 we consider the following 2-by-2 a_{ij} matrix:

$$\begin{array}{c|cc} & \mu_1 & 1 - \mu_1 \\ \hline \lambda_1 & 2 & 0 \\ 1 - \lambda_1 & 1 & 3 \end{array}$$

For convenience we shall denote a mixed strategy for player 1 as λ_1 instead of $(\lambda_1, 1 - \lambda_1)$, and similarly for the second player. By the stochastic dominance definitions of section 2, we have $\lambda_1 \in S_1(\mu_1)$ iff $(\lambda_1', \mu_1) \text{ SD}(>) (\lambda_1, \mu_1)$ for no $\lambda_1' \in [0, 1]$. Using the definition immediately following Lemma 2, $\lambda_1 \in S_1(\mu_1)$ iff, for each $\lambda_1' \in [0, 1]$,

- either $(1 - \mu_1)(\lambda_1' - \lambda_1) > 0$, [y has $a_{ij} = 1$]
- or $(1 - \mu_1)(\lambda_1' - \lambda_1) + \mu_1(\lambda_1 - \lambda_1') > 0$, [y has $a_{ij} = 2$]
- or $(1 - \mu_1)(\lambda_1' - \lambda_1) + \mu_1(\lambda_1 - \lambda_1') + \mu_1(\lambda_1' - \lambda_1) > 0$, [y has $a_{ij} = 3$]
- or equality holds in place of $>$ in the preceding three lines.

Inspection of this system then shows that

$$\begin{aligned} S_1(\mu_1) &= \{0\} & \text{if } \mu_1 \leq 1/2 \\ S_1(\mu_1) &= [0, 1] & \text{if } 1/2 < \mu_1 < 1 \\ S_1(\mu_1) &= \{1\} & \text{if } \mu_1 = 1, \end{aligned}$$

and a similar analysis for the efficient sets for player 2 gives

$$\begin{aligned} S_2(\lambda_1) &= \{1\} & \text{if } \lambda_1 = 0 \\ S_2(\lambda_1) &= [0, 1] & \text{if } 0 < \lambda_1 < 1 \\ S_2(\lambda_1) &= \{0\} & \text{if } \lambda_1 = 1. \end{aligned}$$

Therefore $\lambda_1 \in S_1(\mu_1)$ and $\mu_1 \in S_2(\lambda_1)$ iff $0 < \lambda_1 < 1$ and $1/2 < \mu_1 < 1$, so that $SD(>_1, >_2) = (0,1) \times (1/2,1)$. In other words, $p \in SD(>_1, >_2)$ if and only if player 1 uses some probability strictly between 0 and 1 for his first pure strategy and player 2 uses some probability strictly between 1/2 and 1 for his first pure strategy.

Consider next the determination of Nash equilibria for various (u_1, u_2) pairs. For convenience we normalize the u_i functions between 0 and 1 with $a_{12} = b_{22} = 0$, $a_{22} = b_{12} = 1$, $1 > a_{11} > a_{21} > 0$ and $1 > b_{21} > b_{11} > 0$. The expected utilities are $u_1(\lambda_1, \mu_1) = \mu_1 \lambda_1 a_{11} + \mu_1 (1 - \lambda_1) a_{21} + (1 - \mu_1)(1 - \lambda_1)$ and $u_2(\lambda_1, \mu_1) = \mu_1 \lambda_1 b_{11} + \mu_1 (1 - \lambda_1) b_{21} + (1 - \mu_1) \lambda_1$. For a particular choice of the a_{ij} and b_{ij} , the pair (λ_1, μ_1) is a Nash equilibrium iff $u_1(\lambda_1, \mu_1) \geq u_1(\lambda'_1, \mu_1)$ for all $\lambda'_1 \in [0,1]$ and $u_2(\lambda_1, \mu_1) \geq u_2(\lambda_1, \mu'_1)$ for all $\mu'_1 \in [0,1]$. Since it is clear that there are no such equilibria for λ_1 or μ_1 in $\{0,1\}$, assume that $0 < \lambda_1 < 1$ and $0 < \mu_1 < 1$. Reduction of the preceding inequalities then shows that $(\lambda_1, \mu_1) \in US(u_1, u_2)$ if and only if the a_{ij} and b_{ij} can be chosen such that

$$\begin{aligned}\mu_1 &= 1/(1 + a_{11} - a_{21}) \\ \lambda_1 &= b_{21}/(1 + b_{21} - b_{11}),\end{aligned}$$

which under the restrictions $1 > a_{11} > a_{21} > 0$ and $1 > b_{21} > b_{11} > 0$ gives $US(u_1, u_2) = (0,1) \times (1/2,1)$. This is of course the same result arrived at for $SD(>_1, >_2)$ in the preceding paragraph, as required by Theorem 1.

One further point for this example is in order, and that is that the set of all zero-sum equilibria in $US(u_1, -u_1)$ is a proper subset of $SD(>_1, >_2)$. In the present case $US(u_1, -u_1)$ is all (λ_1, μ_1) of the form $((1 - a_{21})/(1 + a_{11} - a_{21}))$,

$1/(1 + a_{11} - a_{21})$ with $1 > a_{11} > a_{21} > 0$, which is the union of $(t/(1+t), 1/(1+t)) \times \{1/(1+t)\}$ over all t strictly between 0 and 1. Thus there will often be $p \in SD(>_1, >_2)$ that are Nash equilibria for (u_1, u_2) only when (u_1, u_2) is not constant or zero-sum.

Example 2: The a_{ij} matrix for our second two-person antagonistic example is

	μ_1	μ_2	μ_3	μ_4	μ_5
λ_1	6	1	3	3	2
λ_2	1	6	3	3	2
λ_3	3	3	5	0	2
λ_4	3	3	0	5	2
λ_5	4	4	4	4	3

Normalizing again so that $\max a_{ij} = 1$, $\min a_{ij} = 0$, and similarly for the b_{ij} , we require

$$1 = a_{11} > a_{33} > a_{51} > a_{55} > a_{15} > a_{21} > a_{34} = 0,$$

$$1 = b_{34} > b_{21} > b_{15} > b_{55} > b_{51} > b_{33} > b_{11} = 0.$$

The equilibria in $SD(>_1, >_2)$ include p , q and e where

$$p \text{ has } \lambda_1 = \lambda_2 = \mu_1 = \mu_2 = 1/2,$$

$$q \text{ has } \lambda_3 = \lambda_4 = \mu_3 = \mu_4 = 1/2,$$

$$e \text{ has } \lambda_5 = \mu_5 = 1.$$

The pure-strategies equilibrium e is a uniform equilibrium, but p is in $S(u_1, u_2)$

iff $1 + a_{21} \geq 2a_{51}$ and $b_{21} \geq 2b_{15}$, and q is in $S(u_1, u_2)$ iff $a_{33} \geq 2a_{51}$
and $1 + b_{33} \geq 2b_{15}$.

According to Theorem 3, $SD(>_1)$ cannot hold between e and p or between e and q in either direction. However, since $a_{11} > a_{33}$ and $a_{21} > a_{34}$, p $SD(>_1)$ q . Similarly q $SD(>_2)$ p . Hence even when a two-person antagonistic game has a saddle point, stochastic dominance relationships can hold between SD equilibria of the game.

References

- Fishburn, P. C.: Decision and Value Theory, New York 1964.
- _____: Convex Stochastic Dominance with Continuous Distribution Functions, Journal of Economic Theory 7, 1974, 143-158.a.
- _____: Convex Stochastic Dominance with Finite Consequence Sets, Theory and Decision 5, 1974, 119-137.b.
- _____: Axioms for Expected Utility in n-Person Games, International Journal of Game Theory 5, 1976, 137-149.
- Gale, D.: A Theory of N-Person Games with Perfect Information, Proceedings of the National Academy of Sciences (U.S.) 39, 1953, 496-501.
- Lehmann, E.: Ordered Families of Distributions, Annals of Mathematical Statistics 26, 1955, 399-419.
- Luce, R. D., and H. Raiffa: Games and Decisions, New York 1957.
- Nash, J. F.: Non-cooperative Games, Annals of Mathematics 54, 1951, 286-295.
- Quirk, J. P., and R. Saposnik: Admissibility and Measurable Utility Functions, Review of Economic Studies 29, 1962, 140-146.
- von Neumann, J., and O. Morgenstern: Theory of Games and Economic Behavior, Princeton 1947.
- Whitmore, G. A., and M. C. Findlay (eds.): Stochastic Dominance: An Approach to Decision Making Under Risk, Lexington, Massachusetts 1977.

1 Oct. 77

BASIC DISTRIBUTION LIST FOR
UNCLASSIFIED TECHNICAL REPORTS
Operations Research Program

Code 434

Operations Research Office of Naval Research (Code 434) Arlington, Virginia 22217	3	Library Naval Electronics Laboratory Center San Diego, California 92152	1
Director, Naval Research Laboratory Attn: Library, Code 2029 (ONRL) Washington, D.C. 20390	2	Dr. A. L. Slafkosky Scientific Advisor Commandant Marine Corps (Code AX) Washington, D.C. 20380	1
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	12	Naval Undersea Warfare Center 3202 E. Foothill Boulevard Pasadena, California 91107 Attn: Technical Library	1
Defense Logistics Studies Information Exchange Army Logistics Management Center Fort Lee, Virginia 23801	1	Stanford University Department of Statistics Stanford, California 94305 Attn: Prof. G. J. Lieberman	1
Technical Information Center Naval Research Laboratory Washington, D.C. 20390	6	Stanford University Department of Statistics Stanford, California 94305 Attn: Prof. T. W. Anderson	1
Director, Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. Lau	1	University of California Department of Engineering Los Angeles, California 90024 Attn: Prof. R. R. O'Neill	1
Mr. William Dejka Naval Electronics Laboratory Center, Code 360 San Diego, California 92132	1	Maritime Transportation Research Board National Academy of Sciences 2101 Constitution Avenue Washington, D.C. 20418 Attn: RADM J. B. Oren, USCG (RET)	1
Director Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. A. L. Powell	1	Director Office of Naval Research Branch Office 536 South Clark Street Chicago, Illinois 60605 Attn: Dr. R. Buchal	1
Library Naval Postgraduate School Code 0212 Monterey, California 93940	1	Stanford University Department of Operations Research Stanford, California 94305 Attn: Prof. A. F. Veinott, Jr.	1
Naval Ordnance Station Louisville, Kentucky 40214	1		

Texas A&M Foundation
College Station, Texas 77843
Attn: Prof. H. O. Hartley

U.S. Naval Postgraduate School
Department of Operations Research
and Administrative Services
Monterey, California 93940

Captain Robert Bush
AFOSR (SRMA)
1400 Wilson Blvd.
Arlington, Virginia 22209

Professor Richard L. Fox
School of Engineering
Case Western Reserve University
Cleveland, Ohio 44106

Director, National Security Agency
Ft. George G. Meade, Maryland 20755
Attn: Dr. Joseph Blum, R44

Logistics Research Project
The George Washington University
707 22nd Street, N.W.
Washington, D.C. 20037
Attn: Dr. W. H. Marlow

Harvard University
Department of Economics
Cambridge, Massachusetts 02138
Attn: Prof. K. J. Arrow

Cowles Commission for Research
in Economics
Yale University
New Haven, Connecticut 06520
Attn: Prof. Martin Shubik

University of California
Operations Research Center
Institute of Engineering Research
Berkeley, California 94720
Attn: Prof. D. Gale

The University of Michigan
Department of Mathematics, W.E.
Ann Arbor, Michigan 48104
Attn: Prof. R. M. Thrall

1 University of Texas
Department of Mathematics
Austin, Texas 78712
Attn: Dr. A. Charnes 1

1 Stanford University
Department of Operations Research
Stanford, California 94305
Attn: Prof. F. S. Hillier 1

1 Stanford University
Department of Operations Research
Stanford, California 94305
Attn: Prof. G. B. Dantzig 1

1 Cornell University
Department of Industrial Engineering
Ithaca, New York 14850
Attn: Prof. W. Lucas 1

1 Dr. Jay Lewallen
E-D-13 Theory and Analysis Division
NASA Manned Spacecraft Center
Houston, Texas 77058 1

1 Applied Mathematics Laboratory
Naval Ships Research Development
Center
Washington, D.C. 20007 1

1 Mr. A. S. Rhode
Office of Chief of Naval Operations
Code 964
Arlington, Virginia 22203 2

1 Dr. Del Cilmer
Naval Weapons Center (Code 607)
China Lake, California 93555 1

1 Carnegie-Mellon University
Graduate School of Industrial Adm.
Pittsburgh, Pennsylvania 15213
Attn: Prof. G. Thompson 1

1 Dr. Thom J. Hodgson
Department of Industrial and
Systems Engineering
University of Florida
Gainesville, Florida 32601 1