

AD-A049 443

NAVAL RESEARCH LAB WASHINGTON D C
TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELL--ETC(U)
DEC 77 H HUANG

F/G 20/11

JNCLASSIFIED

NRL-8185

SBIE-AD-E000 099

NL

1 OF 1
AD
A049443



END
DATE
FILMED
3 - 78

DOC

AD A 049443

ade 000099
NRL Report 8185

12
£

Transient Response of Two Fluid-Coupled Spherical Elastic Shells to an Incident Pressure Pulse

H. HUANG

Ocean Technology Division

December 23, 1977

AD No. _____
JDC FILE COPY

This research was sponsored by the Defense Nuclear Agency under
Subtask 99QAXSF502, work unit 10, work unit title Double Hull Response Evaluation



D D C
RECEIVED
FEB 6 1978
B

NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release: distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER NRL 8185 8185 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9	
4. TITLE (and Subtitle) TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELLS TO AN INCIDENT PRESSURE PULSE.		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL Problem	
7. AUTHOR(s) H. Huang		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D. C. 20375 ✓		8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS Defense Nuclear Agency Washington, D.C. 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem M02-12B DNA Subtask 99QAXSF502, Work Unit 10	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 15p.		12. REPORT DATE 23 December 1977	
16. DISTRIBUTION STATEMENT (of this Report) 18 SBIE 19 AD-E000 099 Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 28	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
18. SUPPLEMENTARY NOTES This research was sponsored by the Defense Nuclear Agency under Subtask 99QAXSF502, work unit 10, work unit title Double Hull Response Evaluation		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Transient analysis Shock wave loading Fluid-structure interaction Double hull		DDC RECEIVED FEB 6 1978 REGULATED B	
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The transient response of a system of two concentric spherical elastic shells coupled by an ideal fluid and impinged by an incident plan pressure pulse is analyzed. The classical techniques of separation of variables and Laplace transforms are employed for solving the wave equations governing the fluid motions and the shell equations of motion. A scheme of iterative convolution was devised for the inversion of the Laplace transforms that facilitates the calculation of accurate transient solutions of the response of the shells. A sample calculation of shell responses was performed and results are compared to the case in which the outer shell is absent. This set of results demonstrates that a thin outer shell tends to be transparent to the incident pulse.			

DD FORM 1 JAN 73 1473


EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251950

LB

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



CONTENTS

INTRODUCTION 1

 Description of the Problem 2

 Solutions in the Laplace Transform Domain 4

 The Inversion of the Transformed Solutions 6

 Results and Discussions 8

ACKNOWLEDGMENT 11

REFERENCES 12

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

TRANSIENT RESPONSE OF TWO FLUID-COUPLED SPHERICAL ELASTIC SHELLS TO AN INCIDENT PRESSURE PULSE

INTRODUCTION

An underwater weak shock wave sufficiently far away from its generating explosion source is often treated as an acoustic pulse. There is prolific literature on the studies of the transient interaction among such pressure waves and single elastic shells of simple shapes [1,2]. The results not only reveal many of the essential physical phenomena involved in the interaction problem but also are quite useful for the verification of approximation methods for predicting the underwater explosion response of submerged structures surrounded by an exterior fluid medium of infinite extent [2-5]. In the present endeavor, the transient response of a system of two fluid coupled concentric spherical elastic shells impinged by an external incident plane shock wave is analyzed. This purports to gain physical insight in the response as well as to provide a data base for the development of general numerical methods for predicting the underwater explosion response of fluid coupled shell systems such as the double hull section of a submarine. The problem with the spherical geometry permits the separation of variables in the wave equation governing the fluid motion and the shell equations of motion. The use of Laplace transforms then facilitates the calculation of satisfactory transient solutions of the response of the shells.

A paper appears in the Russian literature [6] independently dealing with exactly the same problem. The results obtained, however, only pertain to the point-symmetric and the translational motions of the shells and these two terms of the series solution are insufficient for the description of the complete shell response. It would also seem that the scheme of dual Volterra integral equations for the calculation of the inverse Laplace transform used in Ref. 6 is unnecessarily complex and numerically inefficient. It can be readily shown that only a single and simpler integral equation is needed if this scheme is used.

In this report, a simple, straightforward method is devised for the inverse Laplace transform, and the asymptotic behaviors of the shells are analytically discussed. An example calculation is also carried out using eight terms of the series solution for adequate convergence of shell displacements, velocities, and stresses. Time histories of the transient response of the interior shell are presented. This example demonstrates that a thin exterior shell tends to be transparent to the incident pulse.

Description of the Problem

Figure 1 sketches the fluid-coupled spherical shell system and the incident plane pressure wave. The fluid surrounding the outer shell and that between the two shells are considered to be ideal compressible fluids in linear wave motions and can be characterized by their unperturbed mass densities and sound speeds, i.e., by (ρ^e, c^e) and (ρ, c) , respectively. The shells are initially concentric. In this study, the strength of the incident wave is sufficiently weak such that the shell deflections are elastic and small and the deviation from the concentricity remains negligible for the time duration of interest. The mass densities, Young's moduli and Poisson's ratios of the outer and inner shells are (ρ_s^e, E^e, ν^e) and (ρ_s, E, ν) , respectively. The middle surface radii and thicknesses of the outer and inner shells are (a^e, h^e) and (a, h) , respectively. The Φ -coordinate of the spherical system (r, θ, Φ) is not shown in Fig. 1 since it is not needed due to the symmetry of the problem. The origin O coincides with the unperturbed center of the shells.




Fig. 1 - Geometry of the problem

The deflections of the inner shell in the r - and θ -direction, normalized with respect to the outer shell radius a^e , are denoted by w and u respectively and those of the outer shell by w^e and u^e respectively. The total pressure field exterior to the outer shell is denoted by $p^e(r, \theta, t)$ and that between the shells by $p(r, \theta, t)$ where t designates time. The following dimensionless parameters will be used in the mathematical formulation:

$$\begin{aligned}
 R &= r/a^e, T = c^e t/a^e, \zeta = a/a^e, \eta = c_r(1 - \zeta), \\
 c_r &= c^e/c, \rho_r = \rho/\rho^e, M = \rho^e a/(\rho_s h), M^e = \rho^e a^e/(\rho_s^e h^e), \\
 \Pi &= p/[\rho^e (c^e)^2], \Pi^e = p^e/[\rho^e (c^e)^2], I = \frac{1}{12} (h/a)^2, I^e = \frac{1}{12} (h^e/a^e)^2, \\
 C^2 &= E/[\rho_s (1 - \nu) (c^e)^2], C_e^2 = E^e/[\rho_s^e (1 - \nu^e) (c^e)^2], \\
 \alpha_0 &= \alpha_0^e = \mu_0 = \mu_0^e = 0, \lambda_0 = 2C^2, \lambda_0^e = 2C_e^2, \\
 \alpha_m &= (1 + I)C^2 [m(m+1) - (1 - \nu)]/(1 + \nu), \\
 \alpha_m^e &= (1 + I^e)C_e^2 [m(m+1) - (1 - \nu^e)]/(1 + \nu^e), \\
 \lambda_m &= C^2 [2 + [1 + (m^2 + m + 1)I] [m(m+1) - (1 - \nu)]/(1 + \nu)], \\
 \lambda_m^e &= C_e^2 [2 + [1 + (m^2 + m + 1)I^e] [m(m+1) - (1 - \nu^e)]/(1 + \nu^e)], \\
 \mu_m &= C^4 (m^2 + m - 2) [1 - \nu + [m(m+1) - (1 - \nu)] \times \\
 &\quad [m(m+1) - (1 + \nu)] / (1 - \nu)] / (1 - \nu) \\
 \mu_m^e &= C_e^4 (m^2 + m - 2) [1 - \nu^e + [m(m+1) - (1 - \nu^e)] \times \\
 &\quad [m(m+1) - (1 + \nu_e)] / (1 - \nu_e)] / (1 - \nu_e) \\
 m &= 1, 2, 3, \dots
 \end{aligned} \tag{1}$$

Π^e and Π satisfy the wave equations

$$\nabla^2 \Pi^e = \frac{\partial^2 \Pi^e}{\partial T^2} \quad (2)$$

and

$$\nabla^2 \Pi = c_r^2 \frac{\partial^2 \Pi}{\partial T^2} \quad (3)$$

respectively, where ∇^2 is the Laplacian operator. The boundary conditions of the problem are that Π^e satisfies the radiation condition at far field and that

$$\frac{\partial \Pi}{\partial R} = \frac{\partial \Pi^e}{\partial R} = -\frac{\partial^2 w^e}{\partial T^2} \text{ at } R = 1 \quad (4)$$

and

$$\frac{\partial \Pi}{\partial R} = -\rho_r \frac{\partial^2 w}{\partial T^2} \text{ at } R = \zeta. \quad (5)$$

All quantities except the incident pressure field have quiescent initial conditions.

A Laplace transform pair is defined as

$$\begin{aligned} \bar{w}(\theta, s) &= \int_0^{\infty} w(\theta, T) e^{-sT} dT \\ w(\theta, T) &= \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \bar{w}(\theta, s) e^{sT} ds \end{aligned} \quad (6)$$

where γ lies to the right of all singularities of \bar{w} in the complex s -plane and $i = (-1)^{1/2}$.

Due to the spherical geometry of the problem, the solutions can be expanded in terms of series of Legendre polynomials as the following.

$$\begin{aligned} \Pi(R, \theta, T) &= \sum_{m=0}^{\infty} \Pi_m(R, T) P_m(\cos \theta) \\ \Pi^e(R, \theta, T) &= \sum_{m=0}^{\infty} \Pi_m^e(R, T) P_m(\cos \theta) \\ w(\theta, T) &= \sum_{m=0}^{\infty} w_m(T) P_m(\cos \theta) \\ w^e(\theta, T) &= \sum_{m=0}^{\infty} w_m^e(T) P_m(\cos \theta) \\ u(\theta, T) &= \sum_{m=1}^{\infty} u_m(T) \frac{dP_m(\cos \theta)}{d\theta} \\ u^e(\theta, T) &= \sum_{m=1}^{\infty} u_m^e(T) \frac{dP_m(\cos \theta)}{d\theta} \end{aligned} \quad (7)$$

where P_m is the Legendre polynomial of the first kind and m th degree.

In the Laplace transform domain, the equations of motion of the elastic shells are [7]

$$\begin{aligned} \bar{w}_m &= \frac{-M\zeta\Pi_m(\zeta, s)(\zeta^2s^2 + \alpha_m)}{\zeta^4s^4 + \lambda_m\zeta^2s^2 + \mu_m} \\ \bar{w}_m^e &= \frac{-M^e[\Pi_m^e(1, s) - \Pi_m(1, s)](s^2 + \alpha_m^e)}{s^4 + \lambda_m^es^2 + \mu_m^e} \\ m &= 0, 1, 2, \dots \end{aligned} \quad (8)$$

$$\begin{aligned} (\zeta^2s^2 + \alpha_m)\bar{u}_m &= \left[C^2 + \frac{I}{1+I}\alpha_m \right] \bar{w}_m \\ (s^2 + \alpha_m^e)\bar{u}_m^e &= \left[C_e^2 + \frac{I^e}{1+I^e}\alpha_m^e \right] \bar{w}_m^e \\ m &= 1, 2, 3, \dots \end{aligned} \quad (9)$$

It should be noted that the solution method developed here is applicable for any linear elastic theory for the spherical shells. The choice of the version in Eqs. (8) and (9) is for the comparison of results previously obtained in Ref. 7.

The total pressure field exterior to the outer shell Π^e consists of the pressure due to the incident wave and those due to scattering and radiation by the outer shell. An arbitrary incident plane pressure wave impinging the vertex of the outer shell ($R = 1, \theta = 0$) at $T = 0$ can be expressed by the following series [7]

$$\Pi^i(R, \theta, s) = f(s)e^{-s} \sum_{m=0}^{\infty} (2m+1)i_m(Rs)P_m(\cos\theta), \quad (10)$$

where $f(s)$ is the Laplace transform of the time characteristics of the incident wave and $i_m(Rs)$ is the abbreviated notation for the modified spherical Bessel function of the first kind $[\pi/(2Rs)]^{1/2}I_{m+1/2}(Rs)$ [8].

Solutions in the Laplace Transform Domain

It can be shown that the solutions to the system of Eqs. (2) through (10) are:

$$\begin{aligned} \bar{\Pi}_m^e(R, s) &= (2m+1) \frac{f(s)e^{-s}}{k_m'(s)} [i_m(Rs)k_m'(s) - k_m(Rs)i_m'(s)] \\ &\quad - \frac{s\bar{w}_m^ek_m(Rs)}{k_m'(s)} \end{aligned} \quad (11)$$

$$\begin{aligned} \bar{\Pi}_m(R, s) &= \frac{\rho_r s}{c_r [i_m'(c_r\zeta s)k_m'(c_r s) - i_m'(c_r s)k_m'(c_r\zeta s)]} \\ &\quad \times \{ [\bar{w}_m^ek_m'(c_r\zeta s) - \bar{w}_mk_m'(c_r s)]i_m(c_rRs) \\ &\quad - [\bar{w}_m^ei_m'(c_r\zeta s) - \bar{w}_mi_m'(c_r s)]k_m(c_rRs) \} \end{aligned} \quad (12)$$

$$\begin{aligned} \bar{w}_m^e = & -\frac{(2m+1)}{\Delta_m(s)} \frac{\pi e^{-s}}{2s^2} f(s) M^e (s^2 + \alpha_m^e) \{ (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ & \times [k_m'(c_r s) i_m'(c_r \zeta s) - i_m'(c_r \zeta s) k_m'(c_r s)] + \rho_r M \frac{\zeta s}{c_r} (\zeta^2 s^2 + \alpha_m) \\ & \times [i_m'(c_r s) k_m(c_r \zeta s) - i_m(c_r \zeta s) k_m'(c_r s)] \} \end{aligned} \quad (13)$$

$$\bar{w}_m = -\frac{(2m+1)}{\Delta_m(s)} \frac{\pi e^{-s}}{2s^2} f(s) \frac{\pi}{2c_r^3 \zeta s} \rho_r M M^e (\zeta^2 s^2 + \alpha_m) (s^2 + \alpha_m^e). \quad (14)$$

In the above equations $k_m(s)$ is the abbreviated notation for the modified spherical Bessel function of the third kind, $[\pi/2s]^{1/2} K_{m+1/2}(s)$. The prime denotes differentiation of the Bessel functions with respect to their arguments, and

$$\begin{aligned} \Delta_m(s) = & \left\{ (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) [k_m'(c_r s) i_m'(c_r \zeta s) - k_m'(c_r \zeta s) i_m'(c_r s)] \right. \\ & \left. + \rho_r M \frac{\zeta s}{c_r} (\zeta^2 s^2 + \alpha_m) [i_m'(c_r s) k_m(c_r \zeta s) - k_m'(c_r s) i_m(c_r \zeta s)] \right\} \\ & \times [k_m'(s) (s^4 + \lambda_m^e s^2 + \mu_m^e) - M^e s k_m(s) (s^2 + \alpha_m^e)] \\ & - \frac{\rho_r}{c_r} M^e s (s^2 + \alpha_m^e) (\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) k_m'(s) \\ & \times [k_m'(c_r \zeta s) i_m'(c_r s) - i_m'(c_r \zeta s) k_m(c_r s)] \\ & + \left(\frac{\rho_r}{c_r} \right)^2 M M^e \zeta s^2 (s^2 + \alpha_m^e) (\zeta^2 s^2 + \alpha_m) k_m'(s) \\ & \times [i_m'(c_r s) k_m(c_r \zeta s) - i_m'(c_r \zeta s) k_m(c_r s)] \end{aligned} \quad (15)$$

With use of the Tauber's theorem of Laplace transforms [9], some of the asymptotic behaviors of the shell responses at late time can readily be revealed from Eqs. (13) and (14). Specifically, for the case where the incident wave Π^i is a unit step wave, i.e., $f(s) = 1/s$,

$$w_0^e(T) \underset{T \rightarrow \infty}{=} \frac{-2C^2 c_r^2 (1 - \zeta^3) M^e + 3\zeta^3 \rho_r M M^e}{4C^2 C_e^2 c_r^2 (1 - \zeta^3) + 6\rho_r M \zeta^3 C_e^2 + 6\rho_r M^e C^2} \quad (16)$$

and

$$w_0(T) \underset{T \rightarrow \infty}{=} \frac{-3\zeta \rho_r M M^3}{4C^2 C_e^2 c_r^2 (1 - \zeta^3) + 6\rho_r M \zeta^3 C_e^2 + 6\rho_r M^e C^2}. \quad (17)$$

These shell deflections occur long after the incident wave has engulfed the outer shell and can also be found by static analysis.

Again, for the unit step incidence case,

$$\begin{aligned} \dot{w}_1^e(T) \underset{T \rightarrow \infty}{=} & \frac{-3M^e [(1 + 2\zeta^2) \rho_r M + 6(1 - \zeta^3)]}{6\rho_r M (1 + 2\zeta^3) + 6M^e [(1 - \zeta^3) + (2 + \zeta^3) \rho_r] + \rho_r M M^e [1 + 2\zeta^3 + 2\rho_r (1 - \zeta^3)] + 36(1 - \zeta^3)} \end{aligned} \quad (18)$$

and

$$\dot{w}_1(T) = \frac{-9\rho_r MM^e}{6\rho_r M(1+2\zeta^3) + 6M^e[(1-\zeta^3) + (2+\zeta^3)\rho_r] + \rho_r MM^e[1+2\zeta^3+2\rho_r(1-\zeta^3)] + 36(1-\zeta^3)} \quad (19)$$

where the dot denotes differentiation with respect to T . These are late time translational velocities of the shells in the direction of propagation of the incident wave. It can be seen that the only condition under which $\dot{w}_1 = \dot{w}_1^e$ at late time is $\rho_r M = 3$, i.e., when the inner shell is neutrally buoyant in the interior fluid. Otherwise, they are not equal to each other. Therefore, when interpreting the results of the present problem for large values of T , the relative positions of the shell must also be examined. If, at some time, the deviation from concentricity becomes excessive, the results of analysis thereafter are no longer physically meaningful.

Formulae (16) through (19) are convenient for parametric studies of the effects of various shell properties and fluid properties on the symmetric and translational responses of the shells, and they are also quite useful for providing asymptotic checks for the numerical calculations.

The Inversion of the Transformed Solutions

The spherical Bessel functions can be expressed in finite series of elementary functions [8]. If this property is used, the transformed solutions, e.g., Eq. (14), can be rearranged in the following form:

$$\bar{w}_m(s) = \frac{\Gamma_m(s)}{A_m(s)e^{\eta s} - B_m(s)e^{-\eta s}}, \quad (20)$$

where

$$\begin{aligned} \Gamma_m(s) &= -2(-1)^{m+1}(2m+1)(\rho_r/c_r)f(s)MM^e(\zeta s)^{m+1}(c_r s)^{2m+2}(s^2 + \alpha_m^e)(\zeta^2 s^2 + \alpha_m) \\ A_m(s) &= \{X_m(s)X_m(-c_r s)(s^4 + \alpha_m^e s^2 + \mu_m^e) + M^e s^2(s^2 + \alpha_m^e)[Y_m(s)X_m(-c_r s) \\ &\quad - \rho_r Y_m(-c_r s)X_m(s)]\}[X_m(c_r \zeta s)(\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ &\quad + \rho_r M \zeta^2 s^2(\zeta^2 s^2 + \alpha_m)Y_m(c_r \zeta s)] \\ B_m(s) &= \{X_m(s)X_m(c_r s)(s^4 + \lambda_m^e s^2 + \mu_m^e) + M^e s^2(s^2 + \alpha_m^e)[Y_m(s)X_m(c_r s) \\ &\quad - \rho_r Y_m(c_r s)X_m(s)]\}[X_m(-c_r \zeta s)(\zeta^4 s^4 + \lambda_m \zeta^2 s^2 + \mu_m) \\ &\quad + \rho_r M \zeta^2 s^2(\zeta^2 s^2 + \alpha_m)Y_m(-c_r \zeta s)] \\ X_m(s) &= \sum_{k=0}^{m+1} (m+3/2, k)2^{-k} s^{m+1-k} - m \sum_{k=0}^m (m+1/2, k)2^{-k} s^{m-k} \\ Y_m(s) &= \sum_{k=0}^m (m+1/2, k)2^{-k} s^{m-k} \\ (m+1/2, k) &= \frac{(m+k)!}{k!(m-k)!}. \end{aligned} \quad (21)$$

Both A_m and B_m are $(3m + 11)$ th order real polynomials of s and, for the plane incidence case, Γ_m is a $(3m + 6)$ th order real polynomial.

Equations of the form of Eq. (20) appear in many applied problems such as wave propagation in layered media, transmission lines, etc. The exponential factors in the denominator signify the reflections of waves, in the present case, to and from $R = \zeta$ and $R = 1$. The dimensionless time for a disturbance to traverse the distance $(1 - \zeta)$ is η . There is a variety of schemes for calculating the inverse Laplace transform of Eq. (20). One of the standardized techniques is the D'Alembert expansion [10] by which Eq. (20) can be rewritten as

$$\bar{w}_m(s) = \frac{\Gamma_m(s)e^{-\eta s}}{A_m(s)} \left\{ 1 + \frac{B_m}{A_m} e^{-2\eta s} + \left(\frac{B_m}{A_m}\right)^2 e^{-4\eta s} + \left(\frac{B_m}{A_m}\right)^3 e^{-6\eta s} + \dots \right. \\ \left. + \left(\frac{B_m}{A_m}\right)^n e^{-2n\eta s} + \dots \right\} \\ n = 1, 2, 3, \dots \quad (22)$$

Since A_m is a real polynomial of s , its roots can be accurately computed by well-established numerical procedures and the inversion of every individual term on the right-hand side of Eq. (22) can then be obtained by the method of residue. The calculation of residues, however, would be rather clumsy for terms with A_m of high power. The following scheme utilizing the convolution theorem is devised to circumvent this. From Eq. (22),

$$w_m(T) = w_m^0(T - \eta)H(T - \eta) + w_m^1(T - 3\eta) + w_m^2(T - 5\eta)H(T - 5\eta) \\ + \dots + w_m^n[T - (2n + 1)\eta]H[T - (2n + 1)\eta] + \dots \quad (23)$$

where H is the Heaviside step function and

$$w_m^1(T) = w_m^0(T) - \int_0^T G_m(T - \tau)w_m^0(\tau) d\tau, \\ w_m^2(T) = w_m^1(T) - \int_0^T G_m(T - \tau)w_m^1(\tau) d\tau, \\ \dots \\ w_m^n(T) = w_m^{n-1}(T) - \int_0^T G_m(T - \tau)w_m^{n-1}(\tau) d\tau. \quad (24)$$

In Eq. (24), $w_m^0(T)$ and $G_m(T)$ are the inverse transforms of $\Gamma_m(s)/A_m(s)$ and $[A_m(s) - B_m(s)]/A_m(s)$ respectively. They are to be first accurately obtained by the method of residue. This is quite practical with the current computing technology for up to moderate values of m . For large m asymptotic expansions for the spherical Bessel functions can be used and A_m and B_m will assume simpler forms. For the calculations of shell responses in the present problem, large m terms are not needed. The successive convolutions required in Eq. (24) can be conveniently programmed and carried out in a modern digital computer. Since $w_m^0(T)$ and $G_m(T)$ are composed of terms formed by an exponential multiplied by a trigonometrical function, Trauboth's fast convolution integration algorithm [11], which only requires about the same number of computation steps as for ordinary integration, is used here.

It can be seen from Eqs. (23) and (24) that each successive integration advances the solution time by 2η and the number of successive integrations required depends on T and η . The control of numerical accuracy lies in finding the zeros of A_m and the subsequent numerical integrations, and both are well established numerical techniques. Other quantities such as $w_m^e(T)$ can be found by the same procedure or calculated from their relationships with w_m .

Equation (20) can also be transformed into the following Volterra integral equation:

$$w_m(T) - w_m(T - \eta)H(T - \eta) + \int_0^T G_m(\tau)w_m(T - 2\eta - \tau)H(T - 2\eta - \tau)d\tau = w_m^0(T - \eta)H(T - \eta). \quad (25)$$

On close examination, however, the solution of Eq. (25) requires repeating exactly the same convolutions as in Eq. (24). Therefore it is quite unnecessary to use the numerical techniques of integral equations, even if only one single integral equation is involved, for obtaining the inverse Laplace transform of Eq. (20). On the contrary, the use of two simultaneous Volterra integral equations for obtaining the inverse Laplace transforms of \bar{w}_m and \bar{w}_m^e , as in Ref. 6 would have unduly increased the numerical difficulty and the computation effort.

Results and Discussions

Numerical results are obtained for a case in which both shells are made of steel and both fluids are water. The material properties and dimensions used are such that

$$\begin{aligned} c_r = \rho_r = 1, \quad C^2 = C_e^2 = 17.79133 \\ M^e = 32.09875, \quad M = 6.41975 \\ h^e/a^e = 1/250, \quad h/a = 1/50 \\ \zeta = 0.8, \quad \eta = 0.2. \end{aligned} \quad (26)$$

The incidence is a step wave with $f(s) = 1/s$. The transient responses of the shells are calculated for four transit times of the incident wavefront, i.e., $T = 0$ to 8. For this duration of time, it requires twenty successive convolution integrations in Eq. (24). The parabolic rule is used for the numerical integration in conjunction with Trauboth's fast convolution scheme. Since numerical roundoff and truncation errors accumulate at each successive integration, the integration intervals $\Delta\tau$ are kept sufficiently small to minimize these errors. The roots of $A_m(s)$, and from these, $w_m^0(T)$, and $G_m(T)$, are evaluated with high accuracy before starting the convolution integration. It is also found that, similar to the case of a single shell [7], eight terms ($m = 0 - 7$) in the series solution are sufficient for the representations of the shell deflections, velocities, and stresses. The integration intervals used are: $\Delta\tau = 0.01$ for w_0 and w_1 , $\Delta\tau = 0.0025$ for w_2 , and $\Delta\tau = 0.00125$ for the higher terms. If the solution must be carried out for shorter time duration, say up to $T = 4$, a much coarser $\Delta\tau$ will suffice.

In all subsequent figures, the present solutions are plotted in dotted lines and compared to the solutions obtained by the method of Ref. 7 and plotted in solid lines for the case in which the outer shell is absent.

Figure 2 shows the results for the w_0 's and \dot{w}_1 's. Similar to the single-shell case, the present results approach their respective asymptotic values after about two transit times, and




Fig. 2 - Time histories of w_0 and w_1

the numerical asymptotes of w_0 and \dot{w}_1 agree with formulae (17) and (19) respectively within three digits. These results are indicative of the accuracy of the present solution computation scheme. For this particular example, the outer shell is rather thin and its effect on the response of the inner shell is observable in the w_0 , w_1 , and w_2 terms. Its effect on the higher modes of the inner shell, i.e., w_m with $m > 2$, is quite undiscernible. Sample results for higher w_m 's are exhibited in Fig. 3. It can be seen there that the present results for w_3 and w_4 coalesce with the respective results for the outer shell absent case. The same can be said about w_5 , w_6 , and w_7 , and they are not replotted here.




Fig. 3 - Time histories of w_2 , w_3 , and w_4

Eight terms of w_m 's are summed in Eq. (7) for representing the total response of the inner shell. Figure 4 plots the time history of the relative radial deflection between the two apexes of the inner shell. The presence of the outer shell causes a downward shift of the curve representing the present result. This is due to the diminished w_0 term.

Figure 5 shows the time histories of the radial velocities of the same two apexes. The presence of the outer shell causes a slight reduction of the velocities with little alteration of their profiles. Again this can also be expected by observing the slight reduction of the \dot{w}_1 term.




Fig. 4 - Relative radial deflection between the apexes of the inner shell




Fig. 5 - Time histories of radial velocities at the apexes of the inner shell

It is reiterated here that the series solution in the form of Eq. (7) is not effective for obtaining the early time acceleration and pressure if the incident wave has a steep front. The customary remedy is to apply a modified Watson's transform to Eqs. (11) through (14) and calculate the asymptotic results for large s [1]. This is, however, much more complex than the single shell case studied in Ref. 12, and is outside the scope of the present report.

The polar membrane stress (force per unit length) N_θ and the hoop membrane stress N_ϕ are calculated by

$$N_\theta = \frac{aC^2}{(1 + \nu)M\zeta} \left[(1 + \nu)w + \frac{\partial u}{\partial \theta} + \nu u \cot \theta \right] \quad (27)$$

and

$$N_\phi = \frac{aC^2}{(1 + \nu)M\zeta} \left[(1 + \nu)w + \nu \frac{\partial u}{\partial \theta} + u \cot \theta \right]. \quad (28)$$

The time histories of the polar and/or hoop stress at $\theta = \pi/2$ and at $\theta = \pi$ are plotted in Fig. 6 and 7 respectively. The shielding of the outer shell reduces the stresses slightly. The mean stresses also decrease in proportion to w_0 . The maximum stresses occur after the incident wave has completely engulfed the outer shell and are about twice the corresponding static values.




Fig. 6 - Time histories of the polar stress at $\theta = \pi/2$




Fig. 7 - Time histories of the polar and hoop stresses at $\theta = \pi$

All results of the present example infer that a thin outer shell tends to be transparent to the incident pulse.

In conclusion, it can be said that an accurate solution method for this problem has been developed and can be used for parametric studies. Moreover, a set of numerical results has been meticulously obtained to serve as a data base for the development of numerical methods for the analysis of the transient response of double hulls of practical configurations. One immediate scheme is to apply the currently available Doubly Asymptotic Technique [2] in the outer fluid field and the fluid finite element technique for the entrained fluid.

ACKNOWLEDGMENT

The work was supported by Defense Nuclear Agency under DNA subtask 99QAXSF502, Work Unit 10, Title Double Hull Response Evaluation, with SEA 03511 of Naval Sea Systems Command as the secondary manager. The author also wishes to express his appreciation to

Mr. G.J. O'Hara, Lt. R. Wade, Dr. M. Baron, and Dr. T.L. Geers for their helpful discussions. Specifically, it was Dr. Geers who brought the author's attention to and made available the English translation of Ref. 6.

REFERENCES

1. Überall, H., and Huang, H., "Acoustical Response of Submerged Elastic Structures Obtained Through Integral Transform," *Physical Acoustics*, Mason, W.P., and Thurston, R.N., eds., Academic Press, New York, 1977, pp. 217-275.
2. Geers, T.L. "Transient Responses Analysis of Submerged Structures," *Finite Element Analysis of Nonlinear Structure Behavior*, Belytachko, T., Osias, J.R. and Marcel. P.V., eds., ASME, AMD-14, 15-84 (1975).
3. Geers, T.L., "Excitation of an Elastic Cylindrical Shell by A Transient Acoustic Wave," *J. Appl. Mech.*, **35**, 459-469 (1969).
4. Huang, H., "A Qualitative Appraisal of the Double Asymptotic Approximation for Transient Analysis of Submerged Structures Excited by Weak Shock Waves," NRL Memorandum Report 3135, September 1975.
5. Huang, H., Everstine, G.C. and Wang, Y.F., "Retarded Potential Techniques for the Analysis of Submerged Structures Impinged by Weak Shock Waves," *Computational Methods for Fluid-Structure Interaction Problems*, Belytschko, T. and Geers, T.L., eds., ASME, AMD-26, 83-93 (1977).
6. Babaev, A.E. and Kubenko, V.D., "Interaction of a Nonstationary Pressure Wave with a System of two Spherical Shells," *Mekh. Tverdogo Tela*, **2**, 135-141 (1977).
7. Huang, H., "Transient Interaction of Plane Acoustic Waves with a Spherical Elastic Shell," *J. Acoust. Soc. Amer.*, **45**, 661-670 (1969).
8. Abramowitz, M. and Stegun, I.A., eds., *Handbook of Mathematical Functions*, AMS 55, National Bureau of Standard, Washington, D.C., 1964, Chap. 10, pp. 437-445.
9. Van der Pol, B. and Bremmer, H., *Operational Calculus*, Cambridge University Press, New York, 1964, Chap. VII, pp. 122.
10. Ibid, Chap. IX, p. 189.
11. Trauboth, H.H., "Recursive Formulae for the Evaluation of the Convolution Integral," *J. Assoc. Comput. Mach.* **16**, (No. 1), 63-72 (1969).
12. Tang, S. and Yen, D., "Interaction of a Plane Acoustic Wave with an Elastic Spherical Shell," *J. Acoust. Soc. Amer.* **47**, 1325-1333 (1970).

DISTRIBUTION LIST

**ASSISTANT TO THE SECRETARY DEFENSE
ATOMIC ENERGY
WASHINGTON , DC 20301
ATTN DONALD COTTER**

**DIRECTOR
DEFENSE ADVANCED RSCH PROJ AGENCY
ARCHITECT BUILDING
1400 WILSON BLVD
ARLINGTON VA. 22209
ATTN A TACHMINDJI
ATTN STO KENT KRESA
ATTN TECHNICAL LIBRARY
ATTN R CHAPMAN**

**DIRECTOR
DEFENSE INTELLIGENCE AGENCY
WASHINGTON D.C. 20301
ATTN DI-7D E. OFARRELL
ATTN DI-7E
ATTN DT-1C J. VERONA
ATTN DT-2 WPNS + SYS DIV
ATTN TECHNICAL LIBRARY**

**DIRECTOR
DEFENSE NUCLEAR AGENCY
WASHINGTON D.C. 20305
ATTN STTL TECH LIBRARY
ATTN STST ARCHIVES
ATTN DDST
ATTN SPSS**

(2 COPIES)

(2 COPIES)

**CHAIRMAN
DEPT OF DEFENSE EXPLO SAFETY BOARD
RM-GB270, FORRESTAL BUILDING
WASHINGTON D.C. 20301
ATTN DD/S+SS**

DIR OF DEFENSE RSCH + ENGINEERING
WASHINGTON D.C. 20301
ATTN AD/SW
ATTN DD/TWP
ATTN DD/S+SS
ATTN AD/NP

COMMANDER
FIELD COMMAND
DEFENSE NUCLEAR AGENCY
KIRTLAND AFB, NM 87115
ATTN FCTA
ATTN FCTA-D

INTERSERVICE NUCLEAR WEAPONS SCHOOL
KIRTLAND AFB, NM 87115
ATTN TECH LIB

DIRECTOR
JOINT STRAT TGT PLANNING STAFF JCS
OFFUTT AFB
OMAHA, NB 68113
ATTN STINFO LIBRARY

WEAPONS SYSTEMS EVALUATION GROUP
400 ARMY NAVY DRIVE
ARLINGTON VA 22202
ATTN DOC CON

CHEIF OF RES, DEV + ACQUISITION
DEPARTMENT OF THE ARMY
WASHINGTON D.C. 20310
ATTN TECHNICAL LIBRARY
ATTN DAMA-CSM-N LTC E. DEBOESER JR

COMMANDER
HARRY DIAMOND LABORATORIES
WASHINGTON D.C. 20438
ATTN AMXDO-NP
ATTN AMXDO-TI TECH LIB

DIRECTOR
U S ARMY BALLISTIC RESEARCH LABS
ABERDEEN PROVING GROUND, MD 21005
ATTN TECH LIB-E. BAICY

COMMANDER
U S ARMY COMM COMMAND
FORT HUACHUCA, AZ 85613
ATTN TECHNICAL LIBRARY

COMMANDER
U S ARMY MAT + MECHANICS RSCH CTR
WATERTOWN, MA 02172
ATTN R SHEA

COMMANDER
U S ARMY NUCLEAR AGENCY
FORT BLISS, TX 79916
ATTN TECH LIB

COMMANDER
U S ARMY WEAPONS COMMAND
ROCK ISLAND ARSENAL
ROCK ISLAND, IL 61201
ATTN TECHNICAL LIBRARY

CHIEF OF NAVAL MATERIAL
NAVY DEPT
WASHINGTON D.C. 20360
ATTN MAT 0323

CHIEF OF NAVAL OPERATIONS
NAVY DEPARTMENT
WASHINGTON D.C. 20350
ATTN OP 03FG
ATTN OP 985F

CHIEF OF NAVAL RESEARCH
DEPARTMENT OF THE NAVY
ARLINGTON VA 22217
ATTN N PERRONE CD 474
ATTN TECHNICAL LIBRARY

OFFICER IN CHARGE
CIVIL ENGR LAB
NAVAL CONST. BATTALION CTR
PORT HUENEME CA 93041
ATTN R ODELLO
ATTN TECH LIB

COMMANDER
NAVAL ELECTRONIC SYSTEMS COMMAND
NAVAL ELECTRONIC SYSTEMS COMMAND HQS
WASHINGTON D.C. 20360
ATTN PME 117-21A

COMMANDER
NAVAL FACILITIES ENGINEERING COMMAND
HEADQUARTERS
WASHINGTON D.C. 20390
ATTN TECHNICAL LIBRARY

SUPERINTENDENT
NAVAL POSTGRADUATE SCHOOL
MONTEREY CA 93940
ATTN CODE 2124 TECH RPTS LIBRARIAN

DIRECTOR
NAVAL RESEARCH LABRATORY
WASHINGTON D.C. 20375
ATTN CODE 2027 TECHNICAL LIB
ATTN CODE 840 J GREGORY
ATTN CODE 8440 F ROSENTHAL
ATTN CODE 8403A G OHARA
ATTN CODE 8442 H HUANG

(3 COPIES)

(4 COPIES)

COMMANDER
NAVAL SEA SYSTEMS COMMAND
NAVY DEPARTMENT
WASHINGTON D.C. 20362
ATTN ORD - 91313 LIB
ATTN CODE 03511 C POHLER

COMMANDER
NAVAL SHIP RSCH AND DEVELOPMENT CTR
UNDERWATER EXPLOSIONS RSCH DIVISION
PORTSMOUTH V.A. 23709
ATTN E PALMER
ATTN TECHNICAL LIBRARY

COMMANDER
NAVAL SHIP ENGINEERING CENTER
CENTER BUILDING
HYATTSVILLE MD 20782
ATTN NSEC 6120D
ATTN NSEC 6110.01
ATTN NSEC 6105G
ATTN NSEC 6105
ATTN 6105C1
ATTN TECHNICAL LIBRARY

COMMANDER
NAVAL SHIP RESEARCH AND DEV CENTER
BETHESDA MD 20034

ATTN CODE 17 WW MURRAY
ATTN CODE 142-3 LIBRARY
ATTN CODE 174
ATTN CODE 11
ATTN CODE 2740 Y WANG
ATTN CODE 1962
ATTN CODE 1903
ATTN CODE 1731C
ATTN CODE 1171
ATTN CODE 19

COMMANDER
NAVAL SURFACE WEAPONS CENTER
WHITE OAK
SILVER SPRING MD 20910

ATTN CODE 241 J PETES
ATTN CODE 1224 NAVY NUC PRGMS OFF
ATTN CODE 730 TECH LIB
ATTN CODE 240 H SNAY
ATTN CODE 243 G YOUNG

COMMANDER
NAVAL SURFACE WEAPONS CENTER
DAHLGREN LABORATORY
DAHLGREN VA 22448
ATTN TECHNICAL LIBRARY

COMMANDER
NAVAL UNDERSEA CENTER
SAN DIEGO, CA 92152
ATTN TECHNICAL LIBRARY

COMMANDER
NAVAL WEAPONS CENTER
CHINA LAKE CA 93555
ATTN CODE 533 TECH LIB

COMMANDING OFFICER
NAVAL WEAPONS EVALUATION FACILITY
KIRTLAND AIR FORCE BASE
ALBUQUERQUE NM 87117
ATTN TECHNICAL LIBRARY

DIRECTOR
STRATEGIC SYSTEMS PROJECTS OFFICE
NAVY DEPARTMENT
WASHINGTON D.C. 20376
ATTN NSP-272
ATTN NSP-43 TECH LIB

AF CAMBRIDGE RSCH LABS, AFSC
L.G. HANSCOM FIELD
BEDFORD MA 01730
ATTN SUOL AFCRL RSCH LIB

HEADQUARTERS
AIR FORCE SYSTEMS COMMAND
ANDREWS AFB
WASHINGTON D C 20331
ATTN TECHNICAL LIBRARY

COMMANDER
ARMAMENT DEVELOPMENT & TEST CENTER
ELGIN AFB FL 32542
ATTN TECHNICAL LIBRARY

LOS ALAMOS SCIENTIFIC LAB
P O BOX 1663
LOS ALAMOS NM 87544
ATTN DOC CONTROL FOR REPORTS LIB

SANDIA LABS
LIVERMORE LAB
P O BOX 969
LIVERMORE CA 94550
ATTN DOC CON FOR TECH LIB

SANDIA LABORATORIES
P.O. BOX 5800
ALBUQUERQUE NM 87115
ATTN DOC CON FOR 3141 SANDIA RPT COLL

U S ENERGY RSCH & DEV ADMIN
DIVISION OF HEADQUARTERS SERVICES
LIBRARY BRANCH G-043
WASHINGTON D C 20545
ATTN DOC CONTROL FOR CLASS TECH LIB

UNIV OF CALIFORNIA
LAWRENCE LIVERMORE LAB
P.O. BOX 808
LIVERMORE CA 94550
ATTN TECHNICAL LIBRARY

AGBABIAN ASSOCIATES
250 NORTH NASH STREET
EL SUGONDO CA 90245
ATTN M AGBABIAN

BATTELLE MEMORIAL INSTITUTE
505 KING AVENUE
COLUMBUS OH 43201
ATTN TECHNICAL LIBRARY

BELL TELEPHONE LABORATORIES INC.
MOUNTAIN AVE
MURRAY HILL NJ 07974
ATTN TECH RPT CTR

BOEING COMPANY
P.O. BOX 3707
SEATTLE WA 98124
ATTN AEROSPACE LAB

CAMBRIDGE ACOUSTICAL ASSOC
1033 MASSACHUSETTS AVE
CAMBRIDGE MA 02138
ATTN M JUNGER

CIVIL / NUCLEAR SYSTEMS CORP
1200 UNIVERSITY N. F.
ALBUQUERQUE NM 87102
ATTN T DUFFY

ELECTRIC BOAT DIV
GENERAL DYNAMICS CORP.
GROTON CN 06340
ATTN L. CHEN

GENERAL ELECTRIC CO.
TEMPO-CENTER FOR ADVANCED STUDIES
816 STATE STREET (P.O. DRAWER 00)
SANTA BARBARA CA 93102
ATTN DASIAE

ITT RESEARCH INST
10 WEST 35TH ST
CHICAGO IL 60616
ATTN TECHNICAL LIBRARY

INST FOR DEFENSE ANALYSIS
400 ARMY NAVY DRIVE
ARLINGTON VA 22202
ATTN IDA LIBRARIAN R SMITH

J.L. MERRITT
CONSULTING + SPECIAL ENGR SVS INC
P.O. BOX 1206
REDLANDS CA 92373
ATTN TECHNICAL LIBRARY

KAMAN AVIDYNE
DIV OF KAMAN SCEINCES CORP
83 SECOND AVE
NW INDUSTRIAL PARK
BURLINGTON MA 01803
ATTN E CRISCIONE
ATTN TECHNICAL LIBRARY
ATTN G ZARTARIAN

KAMAN SCIENCES CORP.
P.O. BOX 7463
COLORADO SPRINGS CO 80933
ATTN TECHNICAL LIBRARY

LOCKHEED MISSILES AND SPACE CO.
3251 HANOVER ST
PALO ALTO CA 94304
ATTN TECH INFO CTR D/COLL
ATTN T GEERS D/52-33 BLDG 205

NATHAN M. NEWMARK
CONSULTING ENGINEERING SERVICES
1114 CIVIL ENGINEERING BLDG
URBANA IL 61801
ATTN N NEWMARK

POLYTECHNIC INST OF NEW YORK
DEPT OF APPLIED MECH
333 JAY STREET
BROOKLYN NY 11201
ATTN J KLOSNER

R+D ASSOCIATES
P.O. BOX 3580
SANTA MONICA CA 90403
ATTN TECHNICAL LIBRARY

STANFORD RESEARCH INST
333 RAVENSWOOD AVE
MENLO PARK CA 94025
ATTN SRT LIB ROOM G021
ATTN B GASTEN
ATTN G ABRAHAMSON

TETRA TECH INC.
630 N ROSEMEAD BLVD
PASEDNA CA 91107
ATTN LI-SAN HWANG
ATTN TECH LIB

THE BDM CORP
1920 ALINE AVE
VIENNA VA 22180
ATTN TECH LIB

UNIV OF MARYLAND
DEPT OF CIVIL ENGR
COLLEGE PARK MD 20742
ATTN B BERGER

URS RESEARCH CO.
155 BOVET RD.
SAN MATEO CA 94402
ATTN TECH LIB

WEIDLINGER ASSOCIATES
3000 SAND HILL ROAD
BUILDING 4 SUITE 245
MENLO PARK CA 94025
ATTN J. ISENBERG

WEIDLINGER ASSOCIATES
110 EAST 59TH STREET
NEW YORK, NY 10022
ATTN DR. M. BARON

ASSISTANT CHIEF FOR TECHNOLOGY
OFFICE OF NAVAL RESEARCH
ARLINGTON, VIRGINIA 22217
ATTN CODE 200

TELEDYNE BROWN ENG.
MAIL STOP 44
300 SPARKMAN DRIVE
RESEARCH PARK
HUNTSVILLE, ALABAMA 35807
ATTN DR. MANU PATEL

DIRECTOR
U S ARMY WATERWAYS EXPERIMENT STN
P.O. BOX 631
VICKSBURG MS 39180
ATTN J STRANGE
ATTN W FLATHAU
ATTN TECH LIB (UNCL ONLY)

AF INSTITUTE OF TECHNOLOGY, AU
WRIGHT PATTERSON AFB, OH 45433
ATTN LIB AFIT BLDG 640 AREA B (UNCL ONLY)

NKF ENGINEERING ASSOCIATES, INC.
8720 GEORGIA AVENUE
SILVER SPRING, MD. 20910
ATTN DR. R.O. BELSHEIM

M&T CO.
2130 ARCH STREET
PHILADELPHIA, PA. 19103
ATTN B. McNAIGHT

CATHOLIC UNIV. OF AMERICA
DEPT OF PHYSICS AND INSTITUTE
OF OCEAN SCIENCE AND ENGINEERING
WASHINGTON, D.C.
ATTN H. ÜBERALL

CORNELL UNIV.
COLLEGE OF ENGINEERING
DEPT OF THEORETICAL AND
APPLIED MECHANICS
ITHACA, N.Y. 14850
ATTN Y.H. PAO

SANDIA LABORATORIES
SANDIA CORPORATION
DIVISION 9324
ALBUQUERQUE, NEW MEXICO 87115
ATTN M.J. FORRESTAL