

AD-A049 783

STANFORD UNIV CALIF DEPT OF STATISTICS

F/G 12/1

A NON-LINEAR RENEWAL THEORY WITH APPLICATIONS TO SEQUENTIAL ANA--ETC(U)

MAY 77 T L LAI, D SIEGMUND

N00014-77-C-0306

UNCLASSIFIED

TR-1

NL

AD
A049783

END
DATE
FILMED

3-78

DDC

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

AD A 049783

AD No.
JDC FILE COPY

A NON-LINEAR RENEWAL THEORY WITH APPLICATIONS
TO SEQUENTIAL ANALYSIS II

by

T. L. Lai* and D. Siegmund**

Technical Report No. 1

May 3, 1977

Prepared under Contract

N00014-77-C-0306 (NR-042-373)
for the Office of Naval Research

D. Siegmund, Project Director

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*Research supported by National Science Foundation Grant
NSF-MCS 76-09179 at Columbia University

**Also partially supported by ONR Grant N00014-75C-0560 at
Columbia University

Department of Statistics
Stanford University
Stanford, California

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

12

See back page
for 1473

DDC
RECEIVED
FEB 9 1978
F

A NON-LINEAR RENEWAL THEORY WITH APPLICATIONS
TO SEQUENTIAL ANALYSIS II

by

T. L. Lai and D. Siegmund

SUMMARY

This paper continues earlier work of the authors. An analogue of Blackwell's renewal theorem is obtained for processes $Z_n = S_n + \xi_n$, where S_n is the n^{th} partial sum of a sequence $X_1, X_2, \dots$ of independent identically distributed random variables with finite positive mean and ξ_n is independent of $X_{n+1}, X_{n+2}, \dots$ and has sample paths which are slowly changing in a sense made precise below. As a consequence, asymptotic expansions up to terms tending to 0 are obtained for the expected value of certain first passage times. Applications to sequential analysis are given.

1. INTRODUCTION

Let $X_1, X_2, \dots$ be independent identically distributed random variables with positive mean μ and finite variance σ^2 . Let $S_n = X_1 + \dots + X_n$ and $Z_n = S_n + \xi_n$, where for each n ξ_n is independent of $X_{n+1}, X_{n+2}, \dots$. This paper continues the program begun by Lai and Siegmund (1977) of developing a renewal theory for Z_n under conditions which guarantee that the sample paths of the ξ_n

process are slowly changing in a suitable sense made precise below. In order to facilitate comparison of these conditions for different theorems and to provide a convenient reference, the main result of Lai and Siegmund (1977) is stated as Theorem 1. The interested reader may find the informal discussion contained in that paper helpful in motivating the decomposition of Z_n and the conditions imposed on ξ_n .

For $b \geq 0$ define

$$(1) \quad T = T(b) = \inf\{n : Z_n > b\}$$

and

$$(2) \quad \tau = \tau(b) = \inf\{n : S_n > b\}, \quad \tau_+ = \tau(0) .$$

Theorem 1. (Lai and Siegmund, 1977). Let $1/2 < \alpha \leq 1$ and assume that

$$(3) \quad b^{-\alpha}(T - b\mu^{-1}) \rightarrow 0$$

in probability. Suppose that for each $\eta > 0$ there exist n' and $\rho > 0$ such that for all $n \geq n'$

$$(4) \quad P\left\{ \max_{n \leq j \leq n+\rho n} \alpha |\xi_j - \xi_n| \geq \eta \right\} < \eta .$$

If X_1 is non-lattice, then

$$(5) \quad \lim_{b \rightarrow \infty} P\{Z_T - b \leq x\} = (E S_{\tau_+})^{-1} \int_{(0,x]} P\{S_{\tau_+} > y\} dy .$$

The first results of this paper are an analogue of Blackwell's renewal theorem and a corollary.

Theorem 2. Suppose there exists $1/2 < \alpha \leq 1$ such that the following three conditions hold:

$$(6) \quad E|X_1|^{2/\alpha} < \infty ,$$

$$(7) \quad \text{for each } \varepsilon > 0 \quad \sum_1^\infty P\{|\xi_n| > n^\alpha \varepsilon\} < \infty ,$$

and for each $\eta > 0$ there exist n' and $\rho > 0$ such that

$$(8) \quad \sum_{n \leq j \leq n+\rho n}^\infty P\{|\xi_j - \xi_n| \geq \eta\} < \eta \quad (n \geq n') .$$

If X_1 is non-lattice, then

$$(9) \quad \sum_1^\infty P\{b < Z_n \leq b+h\} \rightarrow h/\mu \quad (b \rightarrow \infty) .$$

ACCESSION for	
NTIS	Write Section ☒
DDC	Buff Section ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Date / /	
A	

Corollary. Suppose there exists $1/2 < \alpha \leq 1$ such that (6), (7), and (8) hold. Let $p > 0$ and assume $E(X_1^+)^{p+1} < \infty$. If $\{A_b, b \geq b_0\}$ is a family of events and $\{((\xi_T - \xi_{T-1})^+)^p I_{A_b}, b \geq b_0\}$ is uniformly integrable, then so is $\{(Z_T - b)^p I_{A_b}, b \geq b_0\}$. In particular, if $E(\sup_n (\xi_n - \xi_{n-1})^+)^p < \infty$, then $\{(Z_T - b)^p, b \geq b_0\}$ is uniformly integrable.

Theorem 2 and its corollary are proved in Section 2.

Theorem 3 of Section 2 is a somewhat different renewal theorem required by some applications (cf. Section 3). Theorems 1 and 2 together imply the main result of this paper, Theorem 4, which

contains an asymptotic expansion for $ET(b)$ up to terms which vanish as $b \rightarrow \infty$.

In many applications the behavior of ξ_n is governed by a term involving $(S_n - np)^2/n$; and to the extent this is so, diverse conditions on ξ_n may be replaced by moment conditions on X_1 . For technical reasons required by different applications the statement of Theorem 4 is quite complicated. Proposition 1 is designed to facilitate applications in those cases in which a single moment condition suffices to replace several of the more cumbersome conditions on ξ_n . For motivation of this formulation and method of proof see Pollak and Siegmund (1975). The proofs of Theorem 4 and Proposition 1 are given in Section 3, which also contains some information on $\text{Var } T$ as $b \rightarrow \infty$. Some applications are discussed in Section 4, and Section 5 contains a comparison of the results of this paper with those of Woodroffe (1976a, 1977).

Let $\mathcal{F}_n = \mathcal{B}((X_1, \xi_1), \dots, (X_n, \xi_n))$, $n = 1, 2, \dots$.

Theorem 4. Assume that for some $\delta > 0$

$$(10) \quad P\{T \leq \delta b\} = o(b^{-1}) \quad (b \rightarrow \infty).$$

Assume that there exists a sequence of events $A_n \in \mathcal{F}_n$ such that

$$(11) \quad \sum_{k=n}^{\infty} P\left(\bigcup_{k=n}^{\infty} \tilde{A}_k\right) < \infty \quad (\tilde{A} = \text{complement of } A),$$

and on A_n

$$(12) \quad \xi_n = f(n) + v_n,$$

where for some $1/2 < \alpha \leq 1$ the following conditions hold:

$$(13) \quad f: [0, \infty) \rightarrow \mathbb{R} \text{ satisfies } |x^{-\alpha} f(x)| + \sup_{x \leq y \leq x+x^\alpha} |f(y) - f(x)| \rightarrow 0$$

as $x \rightarrow \infty$; V_n is $\mathcal{F}_n$ -measurable and satisfies

$$(14) \quad \sum_n P\{\sup_{k \geq n} k^{-\alpha} |V_k| > \varepsilon\} < \infty \quad (\varepsilon > 0) ;$$

$$(15) \quad V_n \text{ converges in distribution to a random variable } V ;$$

$$(16) \quad \text{the sequence } V_n^* = \max_{n \leq j \leq n+n^\alpha} |V_j| \text{ is uniformly integrable ;}$$

and for each $\eta > 0$ there exist n' and $\rho > 0$ such that

$$(17) \quad \sum_{n \leq j \leq n+\rho n} \alpha P\{|V_j - V_n| \geq \eta\} < \eta \quad (n \geq n') .$$

Suppose X_1 is non-lattice and (6) holds. Then

$$(18) \quad \mu ET = b - f(\mu^{-1}b) - EV + ES_{\tau_+}^2 / 2ES_{\tau_+} + o(1)$$

as $b \rightarrow \infty$.

Proposition 1. Let $Y_1, Y_2, \dots$ be independent and identically distributed with mean 0 and finite variance $\tilde{\sigma}^2$. Let u_n and w_n be random variables such that for some positive constants $c, c_n \rightarrow 0$, and β

$$(19) \quad |u_n - c| < n^{-\beta} \quad \text{and} \quad |w_n| \leq c_n .$$

Assume that $E|Y_1|^p < \infty$ for some $p > 2$. Let $V_n = u_n (\sum_{k=1}^n Y_k)^2 / n + w_n$.

Then $V_n \xrightarrow{L} V$, where V has the distribution of $c \tilde{\sigma}^2 \chi_1^2$. Also V_n

satisfies (14) for any $\alpha \geq 4/p - 1$ and (16) for any $0 < \alpha \leq 1$. In addition for

$$0 < \alpha \leq \min(p\beta/2, p/(p+2)) ,$$

given any $\eta > 0$ there exist n' and $\rho > 0$ such that

$$(20) \quad \sum_{n \leq j \leq n+\rho n} \alpha P\{\max_{n \leq i \leq j} |V_i - V_n| \geq \eta\} < \eta \quad (n \geq n') .$$

In particular (17) holds.

2. NON-LINEAR BLACKWELL'S THEOREMS

Proof of Theorem 2. The notation below is chosen to facilitate comparisons with the proof of Theorem 1 of Lai and Siegmund (1977), which contains similar basic ideas although their technical implementation is different.

Let α , η , and ρ be as in the statement of the Theorem. Set

$$(21) \quad n_0 = \mu^{-1}(b+h), \quad n_1 = [n_0 - \rho n_0^\alpha/4], \quad n_2 = [n_0 + \rho n_0^\alpha/4] .$$

By Lemma 1 below for m sufficiently large and fixed, for all sufficiently large b

$$(22) \quad \sum_{m \leq n \leq n_1} P\{b \leq Z_n \leq b+h\} < \eta ,$$

and also

$$(23) \quad \sum_{n \geq n_2} P\{b \leq Z_n \leq b+h\} \rightarrow 0 \quad \text{as } b \rightarrow \infty .$$

Obviously,

$$(24) \quad \sum_{n < m} P\{b \leq Z_n \leq b+h\} \rightarrow 0 \quad \text{as } b \rightarrow \infty.$$

It remains to estimate the series of terms $P\{b \leq Z_n \leq b+h\}$ for $n_1 < n < n_2$. For each $n_1 < n < n_2$

$$(25) \quad \begin{aligned} P\{b \leq Z_n \leq b+h\} &\leq P\{|\xi_n - \xi_{n_1}| \geq \eta\} \\ &+ P\{b - \eta \leq Z_{n_1} + (S_n - S_{n_1}) \leq b+h+\eta\}. \end{aligned}$$

By (8) and (21) for all large b

$$(26) \quad \sum_{n_1 < n < n_2} P\{|\xi_n - \xi_{n_1}| \geq \eta\} < \eta.$$

Furthermore,

$$(27) \quad \sum_{n_1 < n < n_2} P\{b - \eta \leq Z_{n_1} + (S_n - S_{n_1}) \leq b+h+\eta\} = E g(b - Z_{n_1}),$$

where

$$(28) \quad g(t) = \sum_{j < n_2 - n_1} P\{t - \eta \leq S_j \leq (t - \eta) + h + 2\eta\}.$$

It will be shown in Lemma 2 below that as a consequence of Blackwell's Theorem

$$(29) \quad E g(b - Z_{n_1}) \rightarrow (h + 2\eta)/\mu.$$

Then by (22), (23), (24), (25), (26), and (29)

$$\limsup_{b \rightarrow \infty} \sum_1^\infty P\{b \leq Z_n < b+h\} \leq 2\eta + (h + 2\eta)/\mu.$$

Letting $\eta \rightarrow 0$ gives one inequality. The inequality in the other direction follows by a similar but easier argument, which completes the proof.

Lemma 1. Under conditions (6) and (7) for m sufficiently large and fixed, for all large b (22) holds; also (23) holds.

Proof. Let $0 < \epsilon < \rho\mu/9$. Note that for all large b and $n \geq n_2$, if $S_n - n\mu \geq -\epsilon n^\alpha$ and $\xi_n \geq -\epsilon n^\alpha$, then by (21)

$$S_n + \xi_n \geq n\mu - 2\epsilon n^\alpha \geq n_2\mu - 2\epsilon n_2^\alpha > n_0\mu = b + h.$$

From (6) it follows that $\sum P\{|S_n - n\mu| > n^\alpha \epsilon\} < \infty$ (cf. Baum and Katz, 1965, Theorem 3) and hence by (7) as $b \rightarrow \infty$

$$\sum_{n \geq n_2} P\{b \leq Z_n \leq b + h\} \leq \sum_{n \geq n_2} (P\{|S_n - n\mu| > \epsilon n^\alpha\} + P\{|\xi_n| > \epsilon n^\alpha\}) \rightarrow 0.$$

This proves (23), and (22) follows by a similar argument if m is chosen so large that

$$\sum_{n \geq m} (P\{|S_n - n\mu| > \epsilon n^\alpha\} + P\{|\xi_n| > \epsilon n^\alpha\}) < \eta.$$

Lemma 2. Under conditions (7) and (8), for g defined by (28), the limit (29) holds.

Proof. It suffices to show

$$(30) \quad g(b - Z_{n_1}) \rightarrow (h + 2\eta)/\mu \quad \text{a.s.} \quad (b \rightarrow \infty)$$

and that g is bounded, for then (29) follows by dominated convergence. Let $v(b) = n_2 - n_1$. By (7) and the strong law of numbers $Z_{n_1} = \mu n_1 + o(n_1^\alpha) = b - \rho\mu^{1-\alpha} b^\alpha/4 + o(b^\alpha)$. Hence by (28), to prove (30) it suffices to show that for arbitrary real numbers $z(b) = \rho\mu^{1-\alpha} b^\alpha/4 + o(b^\alpha)$

$$(31) \quad \sum_{j < v(b)} P\{z(b) \leq S_j \leq z(b) + h + 2\eta\} \rightarrow (h + 2\eta)/\mu.$$

But if $j \geq v(b)$ and $S_j \geq 2j\mu/3$; then by (21) $S_j \geq \rho\mu^{1-\alpha} b^\alpha/3 + o(1) > z(b) + h + 2\eta$ for all large b , and it follows that

$$\sum_{j \geq v(b)} P\{S_j \leq z(b) + h + 2\eta\} \leq \sum_{j \geq v(b)} P\{S_j < 2j\mu/3\} \rightarrow 0.$$

Thus (31) and with it (30) follow from Blackwell's Theorem. That g is bounded is a consequence of

$$(32) \quad g(t) \leq 1 + \sum_{n=0}^{\infty} P\{-h - 2\eta \leq S_n \leq h + 2\eta\} < \infty.$$

The series in (32) converges because the random walk $\{S_n\}$ is transient (cf. Feller, 1966, pp. 199 ff.).

Proof of Corollary to Theorem 2. Assume $E(\sup_{n \geq 1} (\xi_n - \xi_{n-1})^+) < \infty$.

For $x > 0$ and all large b

$$(33) \quad \begin{aligned} P\{Z_T - b \geq 2x\} &\leq \sum_{n=0}^{\infty} P\{Z_n \leq b, Z_n + X_{n+1} \geq b + x\} \\ &\quad + P\{\sup_n (\xi_n - \xi_{n-1})^+ \geq x\}. \end{aligned}$$

Also

$$(34) \quad \begin{aligned} \sum_{n=0}^{\infty} P\{Z_n \leq b, Z_n + X_{n+1} \geq b + x\} &= \int_{(-\infty, b]} P\{X_1 \geq b + x - y\} \sum_{n=0}^{\infty} P\{Z_n \in dy\} \\ &\leq \sum_{k=-\infty}^{[b]+1} P\{X_1 \geq b + x - k\} \sum_{n=0}^{\infty} P\{k-1 < Z_n \leq k\} \\ &\leq \text{const.} \left(\int_x^{\infty} P\{X_1 \geq y\} dy + P\{X_1 \geq x-1\} \right). \end{aligned}$$

To see the last inequality in (34), note that by Theorem 2 there

exists a k_0 such that for all $k \geq k_0$ $\sum_{n=0}^{\infty} P\{k < Z_n \leq k+1\} \leq 2/\mu$, while (6) and (7) imply $\sum P\{Z_n \leq k_0 + 1\} < \infty$ (cf. Baum and Katz, 1965, Theorem 3). The uniform integrability of $(Z_T - b)^p$ follows from (33) and (34). If only $\{((\xi_T - \xi_{T-1})^+)^p I_{A_b}\}$ is assumed to be uniformly integrable, (33) may be replaced by $P(\{Z_T - b \geq 2x\} \cap A_b)$

$$\leq \sum_0^{\infty} P\{Z_n < b, Z_n + X_{n+1} \geq b+x\} + P(\{\xi_T - \xi_{T-1} \geq x\} \cap A_b) ,$$

and the rest of the proof follows as above.

The following theorem is equivalent to one of the main results of Woodroffe (1976a) in a number of special cases, although its abstract formulation and proof are different. (See Section 5 for a more systematic comparison of the results of this paper with those of Woodroffe, 1976a.)

Theorem 3. Suppose there exists $1/2 < \alpha \leq 1$ such that conditions (6) and (7) hold, and for each $\eta > 0$ there exist n' and $\rho > 0$ such that

$$(35) \quad \sum_{n \leq j \leq n+\rho n^\alpha} P\{\max_{n \leq i \leq j} |\xi_i - \xi_n| \geq \eta\} < \eta \quad (n \geq n') .$$

If X_1 is non-lattice, then for all $y > 0$

$$(36) \quad \sum_{n=0}^{\infty} P\{T > n, Z_n > b-y\} \rightarrow \mu^{-1} \int_{-y}^0 P\{S_n > t \text{ for all } n \geq 0\} dt .$$

Proof. Like Theorem 2 the proof of Theorem 3 consists of reducing the general case to the case $\xi_n \equiv 0$. This reduction is similar to the proof of Theorem 2 and is omitted. However, unlike Theorem 2, which reduces to Blackwell's Theorem in the case $\xi_n \equiv 0$, the

corresponding version of Theorem 3 does not seem to have appeared in the literature (although under stronger assumptions it is implied by Theorem 3.1 of Woodroffe, 1976a).

Assume then that $\xi_n \equiv 0$ so that $Z_n = S_n$ and $T = \tau$. Let $M_n = \max(0, S_1, \dots, S_n)$ and $M^* = \min(0, S_1, S_2, \dots)$. Then (36) becomes

$$(37) \quad \sum_{n=0}^{\infty} P\{M_n \leq b, S_n > b-y\} + \mu^{-1} \int_{(-y, 0]} P\{M^* > t\} dt.$$

Let $\sigma(n)$ denote the n^{th} (strict) ascending ladder time, i.e., $\sigma(0) = 0$ and for $n \geq 1$ $\sigma(n) = \inf\{n: n > \sigma(n-1), S_n > S_{\sigma(n-1)}\}$ ($\sigma(1) = \tau_+$). Let $\tau_- = \inf\{n: n \geq 1, S_n \leq 0\}$. By considering the (uniquely defined) smallest index $k \leq n$ for which $S_k = M_n$ one obtains for $0 \leq y < b$

$$P\{M_n \leq b, S_n > b-y\} = \sum_{k=0}^{\infty} \int_{(b-y, b]} P\{S_i < S_k \ \forall i < k, S_k \in dx, \\ S_j \leq S_k \ \forall k \leq j \leq n, S_n - S_k > b-y-x\}.$$

Summing these terms for $n = 0, \dots$, and reversing the order of summation yields

$$(38) \quad \sum_{n=0}^{\infty} P\{M_n \leq b, S_n > b-y\} \\ = \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} \int_{(b-y, b]} P\{S_i < S_k \ \forall i < k, S_k \in dx\} P\{\tau_+ > n-k, S_{n-k} > b-y-x\} \\ = \int_{[0, y)} \sum_{n=0}^{\infty} P\{\tau_+ > n, S_n > x-y\} \sum_{k=0}^{\infty} P\{S_i < S_k \ \forall i < k, S_k \in b-dx\}.$$

Now

$$\begin{aligned} \sum_{k=0}^{\infty} P\{S_i < S_k \forall i < k, S_k \in b - dx\} &= \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} P\{\sigma(n) = k, S_k \in b - dx\} \\ (39) \quad &= \sum_{n=0}^{\infty} P\{S_{\sigma(n)} \in b - dx\} . \end{aligned}$$

Since $P\{\tau_+ > n, S_n > x - y\} = P\{S_n - S_i \leq 0 \forall 0 \leq i \leq n, S_n > x - y\}$, it follows that

$$\begin{aligned} \sum_{n=0}^{\infty} P\{\tau_+ > n, S_n > x - y\} &= \sum_{n=0}^{\infty} P\{S_i \geq S_n \forall 0 \leq i \leq n, S_n > x - y, \\ (40) \quad &S_j > S_n \forall j > n\} / P\{\tau_- = \infty\} \\ &= E\tau_+ P\{M^* > x - y\} \quad (0 \leq x < y) . \end{aligned}$$

The last equality uses the well known fact that $E\tau_+ = 1/P\{\tau_- = \infty\}$ (cf. Feller, 1966, p. 379). Since $P\{M^* > t\}$ is decreasing in t , applying Blackwell's renewal theorem to the right hand side of (39) and taking into account (38), (39), and (40) yields

$$\begin{aligned} \sum_{n=0}^{\infty} P\{M_n \leq b, S_n > b - y\} &\rightarrow E\tau_+ \int_{[0, y)} P\{M^* > x - y\} dx / ES_{\tau_+} \\ &= \mu^{-1} \int_{(-y, 0]} P\{M^* > t\} dt . \end{aligned}$$

Remark. The condition that $\sigma^2 = \text{Var } X_1 < \infty$ was not used in the preceding proof for the special case $\xi_n \equiv 0$.

3. EXPANSIONS OF $ET(b)$ AND $\text{Var } T(b)$

Intuitively the random variables Z_n and $T(b)$ are the same variables in Theorem 4 as in Theorems 1 and 2. For technical

reasons, in the proof that follows new random variables Z'_n and $T'(b)$ will be defined in terms of the original Z_n and $T(b)$, and Theorems 1 and 2 will be applied to these new variables.

Proof of Theorem 4. Let $\{\varepsilon_n\}$ be a sequence of positive numbers tending to 0 to be further specified below. Since $f(x) = o(x^\alpha)$, for appropriately chosen $\{\varepsilon_n\}$ there exists an integer n_0 such that $|f(x)| < \varepsilon_n n^\alpha$ for all $x \geq n \geq n_0$. Let $L_1 = \sup\{n : \tilde{A}_n \text{ occurs}\}$, $L_2 = \sup\{n : |V_n| \geq \varepsilon_n n^\alpha\}$, $L_3 = \sup\{n : |S_n - n\mu| \geq \varepsilon_n n^\alpha\}$, and $L = 1 + \max(n_0, L_1, L_2, L_3)$. By (11) $EL_1 < \infty$. By (6) (cf. Baum and Katz, 1965, Theorem 3) and (14) for appropriately chosen $\{\varepsilon_n\}$ $EL_3 < \infty$ and $EL_2 < \infty$. Hence, $EL < \infty$.

Let $\xi'_n = 0$ for $n \leq n_0$ and for $n > n_0$ set $\xi'_n = \varepsilon_n I_{A_n}\{|V_n| < \varepsilon_n n^\alpha\}$. Let $Z'_n = S_n + \xi'_n$, $T'(b) = \inf\{n : Z'_n > b\}$, and $B = B_b = \{L < \min(T, T')\}$. Note that $|\xi'_n| \leq 2\varepsilon_n n^\alpha$ for all n . For all $\varepsilon > 0$ for sufficiently large b on B

$$(41) \quad T_b = T'_b, \quad Z_{T_b} = Z'_{T_b}, \quad \text{and} \quad b\mu^{-1} - \varepsilon b^\alpha < T' < b\mu^{-1} + \varepsilon b^\alpha.$$

The two equalities in (41) are obvious. The inequality in (41) follows from

$$b < Z'_{T'} \leq \mu T' + 3\varepsilon_{T'}(T')^\alpha$$

and the corresponding inequality for $T' - 1$, which hold on B . Hence by Wald's lemma

$$(42) \quad \mu ET - \int_{\tilde{B}} S_T dP = \int_B S_{T'} dP = bP(B) + \int_B (Z_{T'}', -b) dP - \int_B \xi_{T'}', dP .$$

It will be shown in Lemma 3 below that

$$(43) \quad b P(\tilde{B}) \rightarrow 0$$

and in Lemma 4 that

$$(44) \quad \int_{\tilde{B}} S_T dP \rightarrow 0 .$$

as $b \rightarrow \infty$. It is easy to see that condition (3) is satisfied for the random variables T' and hence with the help of (15), (16), (17), (41), and (43)

$$(45) \quad \int_B \xi_{T'}', dP = f(b/\mu) + EV + o(1) .$$

Finally, it will be shown in Lemma 5 as an application of Theorems 1 and 2 that

$$(46) \quad \int_B (Z_{T'}', -b) dP \rightarrow ES_{\tau_+}^2 / 2 ES_{\tau_+} .$$

Relations (42)-(46) yield the theorem.

Lemma 3. Under the conditions (6), (11), and (14) the relation (43) holds.

Proof. The conditions of the lemma imply that $EL < \infty$, where L is defined as in the proof of Theorem 4. To prove (43) it suffices to show that

$$(47) \quad b P\{T \leq L\} \rightarrow 0$$

and

$$(48) \quad b P\{T' \leq L\} \rightarrow 0.$$

Now for arbitrary $\delta > 0$

$$\begin{aligned} P\{T \leq L\} &\leq P\{L \geq \delta b\} + P\{T \leq \delta b\} \\ &\leq (\delta b)^{-1} \int_{\{L \geq \delta b\}} L dP + P\{T \leq \delta b\}. \end{aligned}$$

Hence (47) follows from (10) and the finiteness of EL. Since by definition $|\xi'_n| < 2 \epsilon_n n^\alpha$, it may be shown that $b P\{T' \leq \delta b\} \rightarrow 0$, and then (48) follows from a similar argument.

Lemma 4. Under the conditions (6), (11), and (14) the relation (44) holds

Proof. By the Schwarz inequality and Wald's lemma for squared sums

$$\begin{aligned} \left| \int_{\tilde{B}} S_T dP \right| &= \left| \int_{\tilde{B}} (S_T - \mu T) dP + \mu \int_{\tilde{B}} T dP \right| \\ &\leq \{E(S_T' - \mu T)^2 P(\tilde{B})\}^{1/2} + \mu \int_{\tilde{B}} T dP \\ &= \{\sigma^2 E T P(\tilde{B})\}^{1/2} + \mu \int_{\{T \leq L\}} L dP + \mu \int_{\{T' \leq L < T\}} T dP. \end{aligned}$$

It is easy to see that $ET = O(b)$ and hence by Lemma 3

$$E(T)P(\tilde{B}) \rightarrow 0 \quad (b \rightarrow \infty).$$

The conditions of the lemma imply $EL < \infty$ and hence

$$\int_{\{T \leq L\}} L dP \rightarrow 0.$$

To complete the proof it remains to show

$$\int_{\{T' \leq L < T\}} T dP \rightarrow 0 .$$

If $T > L$, then $|S_{T-1} - \mu(T-1)| < \varepsilon_{T-1} T^\alpha$ and $|\xi_{T-1}| < \varepsilon_T T^\alpha$, so $b > S_{T-1} + \xi_{T-1} \geq \mu(T-1) - 3T^\alpha$. Thus for all large b $\{T > L\} \subset \{T < 2b/\mu\}$ and it follows that

$$\int_{\{T' \leq L < T\}} T dP \leq 2b/\mu P\{T' \leq L\} \rightarrow 0$$

by Lemma 3. This completes the proof.

Lemma 5. Under the conditions of Theorem 4, the relation (46) holds.

Proof. Lemma 3 and (41) show that condition (3) holds for the stopping times T' . Also the conditions of the corollary to Theorem 2 are satisfied with the events A of the corollary being the events B defined in the proof of Theorem 4 (recall especially (13), (16), and (41)). It follows from Theorem 1 that $Z'_T - b$ converges in distribution and from the corollary to Theorem 2 that the $(Z'_T - b)I_{B_b}$ are uniformly integrable. The lemma now follows by simple computation.

Proof of Proposition 1. The convergence in law of V_n is immediate from the central limit theorem. That (14) is satisfied provided $p \geq 4/(1+\alpha)$ follows from Theorem 3 of Baum and Katz (1965).

Let $U_n = \sum_{k=1}^n Y_k$. The calculations given below prove (16) and (20). Several applications are made of Kolmogorov's inequality for submartingales (cf. Chow, Robbins, and Siegmund, 1971, p. 24) and the inequality

$$(50) \quad E|U_n|^p \leq C n^{p/2}$$

(cf. Doob, 1954, p. 225). Here and in what follows C denotes constants which may differ from one appearance to the next. The proof of (16) is an immediate consequence of the inequalities

$$\begin{aligned} P\left\{\max_{n \leq j \leq n+n^\alpha} |U_j^2/j| > x\right\} &\leq P\left\{\max_{n \leq j \leq n+n^\alpha} U_j^2 > nx\right\} \\ &\leq (nx)^{-p/2} E|U_{[n+n^\alpha]}|^p \leq C x^{-p/2}. \end{aligned}$$

To prove (20) note that for $i \geq n$

$$\begin{aligned} u_i U_i^2/i - u_n^2 U_n/n &= n^{-1} U_n^2 (u_i - u_n) \\ &\quad + u_i \{i^{-1} (U_i - U_n)^2 + 2i^{-1} U_n (U_i - U_n) - (ni)^{-1} (i-n) U_n^2\}. \end{aligned}$$

Hence (20) is a consequence of the following inequalities.

$$\begin{aligned} \sum_{j=n}^{n+\rho n^\alpha} P\left\{\max_{n \leq i \leq j} i^{-1} (U_i - U_n)^2 > \eta\right\} &\leq \sum_{j=n}^{n+\rho n^\alpha} P\left\{\max_{n \leq i \leq j} |U_i - U_n|^2 > \eta n\right\} \\ &\leq (\eta n)^{-p/2} \sum_{j=n}^{n+\rho n^\alpha} E|U_j - U_n|^p \\ &\leq C(\eta n)^{-p/2} \sum_{j=n}^{n+\rho n^\alpha} (j-n)^{p/2} \\ &\leq C(\eta n)^{-p/2} (\rho n^\alpha)^{\frac{p}{2}+1}. \end{aligned}$$

$$\begin{aligned} \sum_{j=n}^{n+\rho n^\alpha} P\left\{\max_{n \leq i \leq j} i^{-1} |U_n (U_i - U_n)| > \eta\right\} \\ \leq \sum_{j=n}^{n+\rho n^\alpha} E\left\{P\left\{\max_{n \leq i \leq j} |U_i - U_n| > \eta n / |U_n|\right\} \middle| U_n\right\} \\ (\eta n)^{-p} \sum_{j=n}^{n+\rho n^\alpha} E[|U_n|^p |U_j - U_n|^p] \leq C \eta^{-p} n^{-p/2} (\rho n^\alpha)^{\frac{p}{2}+1}. \end{aligned}$$

$$\sum_{j=n}^{n+\rho n^\alpha} P\left\{ \max_{n \leq i \leq j} (1 - n/j) U_n^2/n > \eta \right\} \leq \rho n^\alpha P\{n^{-1} U_n^2 \geq \eta n^{1-\alpha}/\rho\}$$

$$\leq C \rho (\rho/\eta)^{p/2} n^{-\frac{p}{2} + \alpha(1+p/2)}.$$

$$\sum_{j=n}^{n+\rho n^\alpha} P\{n^{-1} U_n^2 \max_{n \leq i \leq j} |u_i - u_n| > \eta\} \leq \rho n^\alpha P\{n^{-1} U_n^2 (2n^{-\beta}) > \eta\}$$

$$\leq C \rho \eta^{-p/2} n^{\alpha-\beta p/2}.$$

The preceding results show that to a first order approximation the behavior of $\{Z_n\}$ and T is asymptotically the same as that of $\{S_n\}$ and τ , although differences appear with higher order asymptotic expansions. According to Chow, Robbins, and Siegmund (1971, p. 31) $\text{Var}(\tau) \sim \sigma^2 b/\mu^3$ as $b \rightarrow \infty$; and it should come as no surprise that under conditions similar to those of Theorem 4, one may show that $\text{Var } T \sim \sigma^2 b/\mu^3$ also. Indeed, such a result has been proved by a different method and applied by Woodroffe (1976a, 1977) in several special cases. The details of such an analysis seem sufficiently similar to the proof of Theorem 4 that they have been omitted.

It would be more in the spirit of the present paper to obtain an expansion for $\text{Var } T$ up to terms which vanish as $b \rightarrow \infty$. Unfortunately, the authors have been unable to produce such a result even in the simplest special (non-linear) cases. For the linear case, in which $\xi_n \equiv 0$, $Z_n = S_n$, and $\tau = T$, it is possible to obtain an expansion of $\text{Var } T = \text{Var } \tau$ up to terms which vanish as $b \rightarrow \infty$ as an application of Theorem 3. This result seems to be new except under the further assumption that $X_1 \geq 0$ --see Smith (1959).

Theorem 5. Assume that $E(X_1^+)^3 < \infty$ and X_1 is strongly non-lattice in the sense of Stone (1965). Then as $b \rightarrow \infty$

$$(51) \quad \text{Var } \tau = \mu^{-3} \sigma^2 b + \mu^{-2} K + o(1) ,$$

where K is given by

$$(52) \quad K = \sigma^2 ES_{\tau+}^2 / 2\mu ES_{\tau+} + \frac{3}{4} \{ES_{\tau+}^2 / ES_{\tau+}\}^2 - \frac{2}{3} ES_{\tau+}^3 / ES_{\tau+} \\ - (ES_{\tau+}^2 / ES_{\tau+}) E\{\min_{n \geq 0} S_n\} - 2 \int_0^\infty E\{S_{\tau+}(x) - x\} P\{\min_{n \geq 0} S_n \leq -x\} dx .$$

Proof. It is well-known (and is the linear case of Theorem 4) that as $b \rightarrow \infty$

$$(53) \quad \mu E\tau = b + ES_{\tau+}^2 / 2 ES_{\tau+} + o(1) .$$

Similarly, for $i = 1$ or 2

$$(54) \quad E(S_{\tau+} - b)^i + (ES_{\tau+})^{-1} \int_{(0, \infty)} x^i P\{S_{\tau+} > x\} dx .$$

Also $E\tau^2 < \infty$, so by Wald's lemma for second moments (cf. Chow, Robbins, and Siegmund, 1971, p. 23) and elementary algebra

$$(55) \quad \begin{aligned} \mu^2 \text{Var } \tau &= E(\mu\tau - S_{\tau+} + S_{\tau+} - \mu E\tau)^2 \\ &= E(S_{\tau+} - \mu\tau)^2 + E(S_{\tau+} - \mu E\tau)^2 - 2E[(S_{\tau+} - \mu\tau)(S_{\tau+} - \mu E\tau)] \\ &= \sigma^2 E\tau + E(S_{\tau+} - b + b - \mu E\tau)^2 - 2E[(S_{\tau+} - b + b - \mu\tau)(S_{\tau+} - b + b - \mu E\tau)] \\ &= \sigma^2 E\tau - (\mu E\tau - b)^2 - E(S_{\tau+} - b)^2 + 2E\{(\mu\tau - b)(S_{\tau+} - b)\} . \end{aligned}$$

Hence by (53) and (54)

$$\begin{aligned}
 \mu^2 \text{Var } \tau &= \sigma^2 b / \mu + \sigma^2 ES_{\tau+}^2 / 2\mu ES_{\tau+} + \frac{1}{4} \{ES_{\tau+}^2 / ES_{\tau+}\}^2 \\
 (56) \quad &- ES_{\tau+}^3 / 3 ES_{\tau+} + o(1) + 2\mu E\{(\tau - E\tau)(S_{\tau} - b)\} .
 \end{aligned}$$

It remains to evaluate the last term on the right hand side of (56).

By an easy renewal argument

$$(57) \quad E\{(\tau - E\tau)(S_{\tau} - b)\} = \sum_{n=0}^{\infty} \int_{[0, \infty)} \{E(S_{\tau(x)} - x) - E(S_{\tau(b)} - b)\} P\{\tau > n, S_n \in b - dx\} .$$

It follows from standard fluctuation identities (especially Feller,

1966, p. 570, equation (3.6)) that $S_{\tau+}$ has a distribution which is

strongly non-lattice in the sense of Stone (1965). Also $E(X_1^+)^3 < \infty$ implies $ES_{\tau+}^3 < \infty$. Hence by Theorem 3 of Stone (1965) applied to the

renewal process determined by $S_{\tau+}$, equation (54) for $i = 1$ may be sharpened to read

$$(58) \quad E(S_{\tau(b)} - b) = ES_{\tau+}^2 / 2 ES_{\tau+} - H(b) + o(b^{-2} \log b) ,$$

where

$$(59) \quad H(b) = \int_b^{\infty} \int_x^{\infty} P\{S_{\tau+} > y\} dy$$

is integrable at $+\infty$. Hence by (58)

$$|E(S_{\tau(x)} - x) - ES_{\tau+}^2 / 2 ES_{\tau+}|$$

is a directly Riemann integrable function of x . It follows from

(53), (57), (58), and Theorem 3 that

$$(60) \quad E\{(\tau - E\tau)(S_{\tau} - b)\} \rightarrow \mu^{-1} \int_{[0, \infty)} \{E(S_{\tau}(x) - x) - ES_{\tau+}^2 / 2ES_{\tau+}\} P\{\min_{n \geq 0} S_n \geq -x\} dx.$$

Since $Z(x) = E(S_{\tau}(x) - x)$ satisfies the renewal equation $Z = z + F * Z$ with $F(y) = P\{S_{\tau+} \leq y\}$, it may be shown by taking Laplace transforms and making a Taylor series expansion that

$$(61) \quad \int_{[0, \infty)} \{E(S_{\tau}(x) - x) - ES_{\tau+}^2 / 2ES_{\tau+}\} dx = 1/4 \{ES_{\tau+}^2 / ES_{\tau+}\}^2 - 1/6 ES_{\tau+}^3 / ES_{\tau+}^2,$$

and obviously

$$(62) \quad \int_{[0, \infty)} P\{\min_{n \geq 0} S_n \leq -x\} dx = -E\{\min(0, S_1, S_2, \dots)\}.$$

The theorem follows by substituting (60)-(62) into (56).

Remark. Even in those cases where the moments of $S_{\tau+}$ can be computed, the authors know of no general way to compute the integral appearing in (52). However, for numerical purposes the last two terms in (52) are "almost equal" and opposite in sign and can probably be neglected. To see that they are "almost equal" observe that $E\{S_{\tau}(x) - x\} \rightarrow ES_{\tau+}^2 / 2ES_{\tau+}$ as $x \rightarrow \infty$, and if this were actually an equality rather than just a limit relation, the two terms would be equal.

4. APPLICATIONS TO SEQUENTIAL ANALYSIS

Theorem 4 may be applied to yield asymptotic expansions for the expected sample size of a variety of sequential tests, including the classical sequential χ^2 , t , and F tests. Many of these

applications are conceptually similar, and for brevity only two have been included here. The first example was studied by Pollak and Siegmund (1975), who ignored the problem of the excess over the boundary in their analysis but otherwise provided a concrete model from which Theorem 4 has been abstracted.

For θ in some open interval J containing 0 assume that $\exp(\theta x - \psi(\theta))$ is a probability density function with respect to a probability distribution H and that $\psi(0) = \psi'(0) = 0$ and $\psi''(\theta) > 0$ for all θ . Let F be a probability on J and define

$$f(x, t) = \int_J \exp(yx - t\psi(y)) dF(y) .$$

Assume that $x_1, x_2, \dots$ are independent, identically distributed random variables such that for some $\theta \in J - \{0\}$

$$(63) \quad Ex_1 = \psi'(\theta) ,$$

and that F' exists in some neighborhood of θ where it is continuous and positive. Let $s_n = \sum_{k=1}^n x_k$ and $Z_n = \log f(s_n, n)$. Take

$0 < \gamma_1 < \gamma_2 < \frac{1}{2}$ and $A_n = \{|s_n - n\psi'(\theta)| < n^{\frac{1}{2} + \gamma_1}\}$. A straightforward modification of the proof of Theorem 1 of Pollak and Siegmund (1975) shows that on A_n

$$\begin{aligned} Z_n &= \theta s_n - n\psi(\theta) + \log \int_J \exp[(y - \theta)s_n - n(\psi(y) - \psi(\theta))] dF(y) \\ &= S_n - \frac{1}{2} \log n + \frac{1}{2} \log 2\pi(F'(\theta))^2 / \psi''(\theta) + u_n (s_n - n\psi'(\theta))^2 / 2\psi''(\theta)n + w_n . \end{aligned}$$

Here $S_n = \theta s_n - n\psi(\theta)$ and u_n and w_n are random variables for which

$|u_n - 1| \leq n^{-\frac{1}{2} + \gamma_2}$ and $|w_n| \leq c_n$ with c_n non-random and converging to 0. Now assume that for some $p \geq 4$

$$(64) \quad E|x_1|^p < \infty.$$

Choose $\gamma_1 + 2\gamma_2 \leq \frac{1}{2}$ and set $\alpha = \frac{1}{2} + \gamma_1$ and $\beta = \frac{1}{2} - \gamma_1$, so $\alpha \leq \min(2\beta, 2/3)$. Let $V_n = u_n(s_n - n\psi'(\theta))^2 / 2\psi''(\theta)n + w_n$ on A_n and 0 elsewhere. It follows from (64) and Theorem 3 of Baum and Katz (1965) that (11) holds, and by Proposition 1 that (14), (16), and (17) are satisfied. It is easy to see that (10) need not hold without further assumptions, but it does hold if either $\exp(\theta x - \psi(\theta))$ is the true density function of x_1 , i.e.,

$$(65) \quad P\{x_1 \in dx\} = \exp(\theta x - \psi(\theta)) dH(x),$$

or (64) holds for some $p > 4$ and for some $\underline{\sigma}^2 > 0$

$$(66) \quad \psi''(\theta) \geq \underline{\sigma}^2 \quad \text{for all } \theta \in J.$$

That (65) implies (10) follows from Lemma 3 of Pollak and Siegmund (1975). A simple application of the Hájek-Rényi-Chow inequality to modify the proof of their Lemma 7 shows that (64) with $p > 4$ and (66) imply (10). Hence by Theorem 4 as $b \rightarrow \infty$

$$(67) \quad \begin{aligned} I(\theta)E(T) &= b + \frac{1}{2}[\log(b/I(\theta)) - \log\{2\pi[F'(\theta)]^2/\psi''(\theta)\} - \tilde{\sigma}^2/\psi''(\theta)] \\ &\quad + ES_{\tau_+}^2 / 2ES_{\tau_+} + o(1), \end{aligned}$$

where $I(\theta) = \theta\psi'(\theta) - \psi(\theta)$ and $\tilde{\sigma}^2 = \text{Var } x_1$.

The approximation (67) without the term involving S_{τ_+} was given by Pollak and Siegmund (1975). Classical random walk theory leads to an evaluation of $ES_{\tau_+}^2 / 2 ES_{\tau_+}$ in terms of

$$(68) \quad \sum_1^\infty n^{-1} \int_{(S_n < 0)} S_n dP ,$$

which in general is very difficult to compute. For the special case in which the x_i are $N(\theta, 1)$,

$$(69) \quad ES_{\tau_+}^2 / 2 ES_{\tau_+} = 2 + \theta^2/2 - 2\theta B(\theta/2) ,$$

where

$$B(\theta) = \sum_1^\infty \{ n^{-\frac{1}{2}} \phi(\theta n^{\frac{1}{2}}) - \theta \phi(-\theta n^{\frac{1}{2}}) \} .$$

A brief table of values for B is given by Siegmund (1975). A simple useful approximation is given by

$$ES_{\tau_+}^2 / 2 ES_{\tau_+} = \theta(.584 + \theta/8) + o(\theta^2) \quad (\theta \rightarrow 0) .$$

For the numerical values considered in Tables 1 and 2 of Pollak and Siegmund (1975), use of the term $ES_{\tau_+}^2 / 2 ES_{\tau_+}$ reduces an error of about 10% by a factor of roughly one half. Except for special cases computation of this term is quite difficult and perhaps not worth the necessary effort. This is in marked contrast to the approximation of error probabilities, where analyzing the excess over the boundary can lead to dramatic improvement in the accuracy of the approximation (cf. Siegmund, 1975, or part I of this paper).

The second example involves a stopping rule suggested by Siegmund (1977) for testing whether a normal mean is 0 when the variance is unknown. Let $x_1, x_2, \dots$ be independent and normally distributed with mean $\tilde{\mu}$ and variance $\tilde{\sigma}^2$. Put $s_n = x_1 + \dots + x_n$, $\bar{x}_n = n^{-1} s_n$, $s_n^* = (s_n - n\tilde{\mu})/\tilde{\sigma}$, $v_n^2 = n^{-1} \sum_{k=1}^n (x_k - \bar{x}_n)^2$, $t_n^* = \sum_{k=1}^n (x_k - \tilde{\mu})^2/\tilde{\sigma}^2 - n$, and $\theta = \tilde{\mu}/\tilde{\sigma}$. Let $Z_n = n/2 \log\{1 + \bar{x}_n^2/v_n^2\}$, and for $m \geq 3$ define $T = T(b) = \inf\{n : n \geq m, Z_n > b\}$. Obviously $Z_n = n/2 \log(\bar{x}_n^2, n^{-1} \sum_{k=1}^n x_k^2)$, where $g(x, y) = -\log(1 - x^2/y)$. Expanding g in a Taylor series about $(\tilde{\mu}, \tilde{\sigma}^2 + \tilde{\mu}^2)$ and collecting terms yields

$$\begin{aligned} Z_n &= \frac{n}{2} \log(1 + \theta^2) + \theta(1 + \theta^2)^{-1} s_n^* - \frac{1}{2} \theta^2 (1 + \theta^2)^{-1} t_n^* \\ &\quad + \frac{1}{2} (1 + \theta^2)^{-2} (1 + \theta^4) s_n^{*2}/n + \frac{1}{2} (1 + \theta^2)^{-2} \theta^2 (\theta^2 + 2) t_n^{*2}/n \\ &\quad - (1 + \theta^2)^{-2} \theta s_n^* t_n^*/n + w_n, \end{aligned}$$

where $|w_n| \leq W(n^{-2} |s_n^*|^3 + n^{-2} |t_n^*|^3)$ for some function W which equals 0 at 0 and is continuous there.

Let $A_n = \{n^{-2} |s_n^*|^3 < \epsilon_n, n^{-2} |t_n^*|^3 < \epsilon_n\}$, where $\epsilon_n \rightarrow 0$ sufficiently slowly that (11) holds. Let $S_n = n/2 \log(1 + \theta^2) + (1 + \theta^2)^{-1} \theta s_n^* - 1/2 (1 + \theta^2)^{-1} \theta^2 t_n^*$, $v_n^{(1)} = s_n^{*2}/n$, $v_n^{(2)} = t_n^{*2}/n$, and $v_n^{(3)} = s_n^* t_n^*/n$. Also let

$$v_n = \frac{1}{2} (1 + \theta^2)^{-2} \{ (1 + \theta^4) v_n^{(1)} + \theta^2 (\theta^2 + 2) v_n^{(2)} - 2\theta v_n^{(3)} \} + w_n$$

on A_n and 0 elsewhere. With the aid of the identity

$2v_n^{(3)} = (s_n^* + t_n^*)^2/n - v_n^{(1)} - v_n^{(2)}$ and Proposition 1 it is easy to check that $v_n^{(i)}$ for $i = 1, 2$, and 3 and hence v_n satisfy (14)-(17).

As in the preceding example (10) requires a special argument. Let P_0 denote the probability under which the x 's have expectation $\tilde{\mu} = 0$. From the trivial inequality $P_0\{T \leq n\} \leq \sum_{k=m}^n P_0\{Z_k > b\}$ and an analysis of the tail of the t -distribution, it may be shown for $x > 0$ that

$$(68) \quad P_0\{T \leq bx\} = O(b^{\frac{1}{2}} \exp[-b(1 - \frac{2}{m})])$$

as $b \rightarrow \infty$. By Lemma 3 of Pollak and Siegmund (1975), for arbitrary $\theta \neq 0$ and $y > 0$

$$(69) \quad P\{T \leq n\} \leq 1 - \Phi(y) + \exp(\theta^2 n/2 + y\theta n^{\frac{1}{2}}) P_0\{T \leq n\}.$$

Putting $y = b^{1/4}$ in (69) and appealing to (68) yields (10) for any $\delta < 2\theta^{-2}(1 - 2/m)$. Hence by Theorem 4, for $\theta \neq 0$ and $m \geq 3$

$$(70) \quad \log(1 + \theta^2)ET = 2b - 1 + \theta^4/(1 + \theta^2)^2 + ES_{\tau_+}^2/2ES_{\tau_+} + o(1),$$

as $b \rightarrow \infty$.

5. COMPARISON WITH WOODROOFE'S RESULTS

The purpose of this section is to discuss briefly the relation of the results of this paper to similar results obtained recently by Woodroffe (1976a, 1977) by completely different methods. In general terms the methods of this paper and its companion develop renewal theory for non-linear functions of a random walk S_n by expanding the function and applying classical renewal theory to the dominant linear term. In contrast Woodroffe considers the first

passage of a random walk S_n to a non-linear boundary which he analyzes by expanding the boundary around an appropriate point. One consequence of this difference in formulation is that in this paper and its companion a fairly small number of theorems provide a unified theory, whereas Woodroffe is required to reapply his method with its fairly elaborate computations to deal with different stopping rules. A technical difference is that Woodroffe requires a blanket smoothness condition on the distribution of his random variables, which has no counterpart in the present development. Other technical differences are described below.

Let $x_1, x_2, \dots$ be independent identically distributed random variables with positive expectation $\tilde{\mu}$ and finite variance $\tilde{\sigma}^2$. Let $s_n = x_1 + \dots + x_n$. Woodroffe (1976a) studies the behavior of the stopping rule

$$(71) \quad T_1 = \inf\{n : s_n > cn^\gamma\} \quad (c > 0, 0 \leq \gamma < 1)$$

as $c \rightarrow \infty$. (Actually for some results a slightly more general class of stopping rules is considered, but since Woodroffe gives no application for these more general rules, and since their introduction would complicate this discussion, they have been omitted.)

Statistical applications of the stopping rule (71) have been described by Woodroffe (1976b) and Siegmund (1977). Under the additional restriction

$$(72) \quad P\{x_1 \leq 0\} = 0 \quad .$$

Woodroffe (1977) studies the behavior of

$$(73) \quad T_2 = \inf\{n: n \geq m, s_n < cn^\gamma L(n)\} \quad (c > 0, \gamma > 1, m = 1, 2, \dots)$$

as $c \rightarrow 0$, where $L(n) = 1 + \text{const.}/n + o(n^{-1})$ as $n \rightarrow \infty$. Both (71) and (73) may be written in the form

$$(74) \quad T = \inf\{n: n \geq m, (n + \delta + \delta_n)g(n^{-1}s_n) > b\},$$

where $g(x) = (x^+)^{1/(1-\gamma)}$, $b = g(c)$, and $\delta_n \rightarrow 0$. Suppose more generally that $g: (-\infty, \infty) \rightarrow [0, \infty)$ is three times continuously differentiable in a neighborhood of $\tilde{\mu}$ and that $g'(\tilde{\mu}) > 0$. Let $\varepsilon_n \rightarrow 0$ and $A_n = \{n^{-2}|s_n - n\tilde{\mu}|^3 < \varepsilon_n\}$. By Taylor's theorem, on A_n

$$ng(n^{-1}s_n) = ng(\tilde{\mu}) + (s_n - n\tilde{\mu})g'(\tilde{\mu}) + \frac{(s_n - n\tilde{\mu})^2}{2n}g''(\tilde{\mu}) + w_n,$$

where $|w_n| \leq W(n^{-2}|s_n - n\tilde{\mu}|^3)$ for some function W which vanishes at 0 and is continuous there. Let $S_n = ng(\tilde{\mu}) + (s_n - n\tilde{\mu})g'(\tilde{\mu})$ and $V_n = (s_n - n\tilde{\mu})^2/2n + (\delta + \delta_n)g(n^{-1}s_n)$ on A_n and 0 otherwise. It may be shown as in the second example of Section 4 with the aid of Proposition 1 that for suitable ε_n (11) and (14)-(17) are satisfied with $\alpha = 2/3$, provided $E|x_1|^4 < \infty$. Thus Theorem 4 applies to give an asymptotic expansion for $E(T)$ provided that (10) holds, and as always a special argument is required here.

For the stopping rules (71) and (73) proofs of (10) under appropriate conditions follow from Woodroffe (1976a, Lemma 7.1 and 1977, Lemma 2.3). Use of the Hájek-Rényi-Chow inequality (cf. Chow, Robbins, and Siegmund, 1971, p. 25) together with (50) would simplify these arguments.

Hence Woodroffe's expansions of ET_1 and ET_2 follow from Theorem 4.

In Woodroffe's work a central role is played by results resembling Theorem 3, which form the basis for subsequent calculations. One example is Theorem 3.1 of Woodroffe (1976a), which says that if $E|x_1|^3 < \infty$ and Woodroffe's blanket smoothness condition is satisfied, then for T_1 defined by (71)

$$(75) \quad \sum_1^\infty P\{T_1 > n, s_n > cn^\gamma - y\} \rightarrow [(1-\gamma)\tilde{\mu}]^{-1} \int_{-y}^0 P\{s_n \geq n\gamma\tilde{\mu} - x \text{ for all } n \geq 1\} dx.$$

Deriving (75) from Theorem 3 requires the slightly stronger moment condition $E|x_1|^p < \infty$ for some $p \geq 1 + 5^{1/2}$. With (71) rewritten in the form of (74) (with $\delta = \delta_n = 0$) the inequality $s_n > cn^\gamma - y$ becomes $Z_n = ng(n^{-1}s_n) > (b^{1-\gamma} - yn^{-\gamma})^{1/(1-\gamma)}$. As in the proof of Theorem 2, it is easily shown that only the terms with $n_1 \leq n \leq n_2$, where n_1 and n_2 are defined in (21), are non-negligible in evaluating the limit of the left hand sides of (75). For these values of n , which are $\sim b/\mu$, a simple expansion gives $(b^{1-\gamma} - yn^{-\gamma})^{1/(1-\gamma)} = b - \mu^\gamma y/(1-\gamma) + o(1)$. Now Theorem 3 and a simple change of variable yield (75). A similar argument applies to the stopping rule T_2 of (73).

REFERENCES

- Baum, L.E. and Katz, M. (1965). Convergence rates in the law of large numbers. Trans. Amer. Math. Soc. 120, 108-123.
- Chow, Y.S., Robbins, H., and Siegmund, D. (1971). Great Expectations: the Theory of Optimal Stopping, Houghton Mifflin, Boston.
- Doob, J.L. (1954). Stochastic Processes, John Wiley and Sons, New York.
- Feller, W. (1966). An Introduction to Probability Theory and Its Applications, Vol. II, John Wiley and Sons, New York.
- Lai, T.L. and Siegmund, D. (1977). A non-linear renewal theory with applications to sequential analysis I. Ann. Statist. 5.
- Pollak, M. and Siegmund, D. (1975). Approximations to the expected sample size of certain sequential tests. Ann. Statist. 3, 1267-1282.
- Siegmund, D. (1975). Error probabilities and average sample number of the sequential probability ratio test. J. Roy. Statist. Soc. Ser. B 37, 394-401.
- Siegmund, D. (1977). Repeated significance tests for a normal mean. Biometrika 64.
- Smith, W.L. (1959). Cumulants of renewal processes. Biometrika 46, 1-29.
- Stone, C. (1965). On characteristic functions and renewal theory. Trans. Amer. Math. Soc. 120, 327-342.
- Woodroffe, M. (1976a). A renewal theorem for curved boundaries and moments of first passage times. Ann. Prob. 4, 67-80.
- Woodroffe, M. (1976b). Frequentist properties of Bayesian sequential tests. Biometrika 63, 101-110.
- Woodroffe, M. (1977). Second order approximations for sequential point and interval estimation. Ann. Statist. 5.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #1 14 TR-1	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A NON-LINEAR RENEWAL THEORY WITH APPLICATIONS TO SEQUENTIAL ANALYSIS II.		5. TYPE OF REPORT & PERIOD COVERED 9 Technical Report
7. AUTHOR(s) 10 T. L./Lai D./Siegmund		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics Stanford University Stanford, California		8. CONTRACT OR GRANT NUMBER(s) 15 N00014-77-C-0306, N00014-75-C-0564
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 436 Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 11 (NR-042-373) 3 May 77
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 3, 1977
		13. NUMBER OF PAGES 30 (12) 33 p.
		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale; distribution unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Renewal theorem Sequential tests Expected sample size		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See reverse side		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601UNCLASSIFIED 332 580
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

All

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

xi This paper continues earlier work of the authors. An
sub n analogue of Blackwell's renewal theorem is obtained for processes
 $Z_n = S_n + \xi_n$, where S_n is the n^{th} partial sum of a sequence
 $X_1^n, X_2^n, \dots$ of independent identically distributed random variables
with finite positive mean and ξ_n *xi sub n* is independent of $X_{n+1}, X_{n+2}, \dots$ and
has sample paths which are slowly changing in a sense made precise *sub n+2*
below. As a consequence, asymptotic expansions up to terms tending *sub n+1*
to 0 are obtained for the expected value of certain first passage
times. Applications to sequential analysis are given.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)