

AD-A052 755

COLORADO STATE UNIV FORT COLLINS DEPT OF CHEMISTRY
QUADRATIC JAHN-TELLER COUPLING IN OCTAHEDRAL SYSTEMS.(U)
MAR 78 E R BERNSTEIN, J D WEBB

F/G 20/10

N00014-75-C-1179

UNCLASSIFIED

TR-20

NL

1 of 1
AD
A052755



END
DATE
FILMED
5 -78

DDC

AD A 052755

AD No. 1
DDC FILE COPY

OFFICE OF NAVAL RESEARCH
Contract N00014-75-C-1179
Task No. NR 056-607
TECHNICAL REPORT NO. 20

12

4

"QUADRATIC JAHN-TELLER COUPLING IN OCTAHEDRAL SYSTEMS"

by

E. R. BERNSTEIN AND J. D. WEBB

Prepared for Publication in
Molecular Physics

Department of Chemistry
Colorado State University
Fort Collins, Colorado 80523

March 1978

DDC
RECEIVED
APR 17 1978
A

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

Approved for Public Release; Distribution Unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(12) 18p.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER (14) <u>TR-28</u>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (6) <u>Quadratic Jahn-Teller Coupling in Octahedral Systems.</u>	5. TYPE OF REPORT & PERIOD COVERED <u>Technical Report.</u>	
7. AUTHOR(s) (10) <u>E. R. Bernstein J. D. Webb</u>	8. CONTRACT OR GRANT NUMBER(s) (15) <u>NO0014-75-C-1179</u>	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Colorado State University Fort Collins, Colorado 80523	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-607	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217	12. REPORT DATE (11) <u>Mar 1978</u> NUMBER OF PAGES 19	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Jahn-Teller Effect Quadratic Jahn-Teller Effect Γ_8 Vibronic Coupling e_g Vibrations		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper Jahn-Teller vibronic coupling calculations are presented for the general case $[\Gamma_{(8g)} \times e_g]$ in O_h symmetry. The Hamiltonian employed is complete up to terms quadratic in vibrational coordinates q_α and q_ϵ which (epsilon) (cepilon)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

404 992

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. (continued...)

→ span the $\sqrt{2}(e_g)$ mode of MF_6 systems. Computational details are discussed.

It is concluded that, in agreement with previously reported spectroscopic data for ReF_6 and IrF_6 , quadratic terms in the vibronic interaction are essential (both qualitatively and quantitatively) to the complete understanding of intrastate vibronic coupling in transition metal hexafluorides.

↑

ACCESSION FOR	
NTIS	WHITE SERVICE ☒
ODG	BUTL SERVICE ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
BWL	AVAIL. CODE OR SPECIAL
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

I. INTRODUCTION

Recent experimental studies of transition metal hexafluorides¹⁻³ have indicated that the spectroscopically observed dynamic Jahn-Teller (JT) effects are not adequately explained by the usual linear JT theory⁴; quadratic terms, at least, are required in the vibronic Hamiltonian. In order to treat hexafluoride data quantitatively, it would be desirable to solve the problem of two JT-active vibrations (ν_2 (e_g) and ν_5 (t_{2g})) vibronically coupled to a Γ_{8g} electronic state for a Hamiltonian complete up to quadratic terms $[\Gamma_{8g} \times (e_g + t_{2g})]_{\text{QUAD}}$. Unfortunately, however, the requisite numerical methods are not feasible given currently available computers. In fact, even the $[\Gamma_{8g} \times t_{2g}]_{\text{QUAD}}$ problem alone presents an arduous task. The $[\Gamma_{8g} \times e_g]_{\text{QUAD}}$ coupling problem is, however, readily soluble. Its solution is presented here and is found to generate a qualitative insight into the nature of the quadratic portion of the JT interaction.

Previous work on $[\Gamma_{8g} \times e_g]_{\text{QUAD}}$ (or the very closely related problem: $[E_g \times e_g]_{\text{QUAD}}$)⁴ has focused on the weak coupling limit⁵ or on the case for which linear coupling is large.⁶ In the present work matrix elements are given which allow full secular matrices for $[\Gamma_{8g} \times e_g]_{\text{QUAD}}$ to be set up and diagonalized numerically. Given a sufficiently large enough truncated secular matrix, accurate eigenvalues and eigenvectors can be obtained for any values of the linear and quadratic parameters. These secular matrices are also applicable to the $[E_g \times e_g]_{\text{QUAD}}$ situation.

II. THEORY

The first step toward obtaining the necessary matrix elements for $[r_{8g} \times e_g]_{\text{QUAD}}$ is to find the vibronic Hamiltonian. Englman⁴ has discussed in detail the group theoretical techniques which allow a parametric Hamiltonian to be obtained; it is given here with only minor modifications:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}^{(0)} + \hat{\mathcal{H}}^{(1)} + \hat{\mathcal{H}}^{(2)}_{(a_1g)} + \hat{\mathcal{H}}^{(2)}_{(e_g)} + \dots \quad (1)$$

in which

$$\hat{\mathcal{H}}^{(0)} = \left\{ \frac{1}{2} (\hat{p}_\theta^2 + \hat{p}_\epsilon^2) + \frac{k_\epsilon}{2} (q_\theta^2 + q_\epsilon^2) \right\} \underline{I}, \quad (2)$$

$$\hat{\mathcal{H}}^{(1)} = l_\epsilon (q_\theta \underline{\rho}_1 + q_\epsilon \underline{\rho}_2), \quad (3)$$

$$\hat{\mathcal{H}}^{(2)}_{(a_1g)} = \frac{C_a}{2} (q_\theta^2 + q_\epsilon^2) \underline{I}, \quad (4)$$

and

$$\hat{\mathcal{H}}^{(2)}_{(e_g)} = C_\epsilon \left\{ (q_\theta^2 - q_\epsilon^2) \underline{\rho}_1 - 2q_\theta q_\epsilon \underline{\rho}_2 \right\}. \quad (5)$$

(q_θ, q_ϵ) are the mass-weighted normal coordinates of the e_g vibration, $(\hat{p}_\theta, \hat{p}_\epsilon)$ are the conjugate mass-weighted momenta, k_ϵ is the vibrational force constant, $\underline{I}$ is the 4×4 identity matrix, $(l_\epsilon, C_a, C_\epsilon)$ are the linear and quadratic coupling parameters, and $(\underline{\rho}_1, \underline{\rho}_2, \underline{\rho}_3)$ are Dirac matrices.⁴ In the following discussion we will also make use of $\underline{\sigma}_1, \underline{\sigma}_2, \underline{\sigma}_3$, which are additional Dirac matrices as given in reference 4.

Choice of vibronic basis set is important since many matrix elements need to be calculated. In any case, the basis will involve two-dimensional harmonic

oscillator functions, $\{X(n_2, \ell_2)\}$ and vectors which represent the four electronic components $\{\vec{V}(\rho_3, \sigma_3)\}$. The $\{\vec{V}(\rho_3, \sigma_3)\}$ are eigenvectors of both ρ_3 and σ_3 with eigenvalues $\rho_3 = \pm 1$ and $\sigma_3 = \pm 1$. The question of how to combine these factors into a vibronic basis set that is most convenient is now addressed.

One approach, which will only be outlined, involves utilizing molecular point group theory. The vibrational factors, however, do not transform as standard irreducible representations of O_h^* but appropriate linear combinations of these functions can be generated which do so transform. The symmetrized vibrational factors and the electronic factors, which transform as Γ_{8g} , can be combined using Clebsch-Gordon coefficients to give vibronic functions which transform as Γ_{6g} , Γ_{7g} , or Γ_{8g} . Separate secular matrices can then be formed in the usual fashion. This method will not be used, however, because the calculation of matrix elements within this basis set is more cumbersome than need be.

An alternate approach is to use the same vibronic basis that is used in the linear problem. These are written as follows:

$$\vec{b}(n_2, \ell_2; \rho_3, \sigma_3; J_2) = X(n_2, \ell_2) \vec{V}(\rho_3, \sigma_3) \quad (6)$$

in which

$$J_2 = \ell_2 + \frac{\rho_3}{2} = \pm 1/2, \pm 3/2, \dots$$

$$n_2 = 0, 1, 2, \dots; \ell_2 = -n_2, -n_2 + 2, \dots, n_2$$

$$\rho_3 = \pm 1, \sigma_3 = \pm 1.$$

In linear coupling, J_2 is a good quantum number and separate secular matrices are generated for each value of J_2 . Upon introduction of quadratic terms ($\hat{\mathcal{H}}_0^{(2)}(e_g)$ in particular), J_2 ceases to be a good quantum number. One can determine, by examining a general matrix element, that $\{2J_2 \bmod 3\}$ remains a good quantum number. Therefore, the following three sets of basis vectors, labeled by J_2 , are coupled by the quadratic terms:

$$\begin{aligned} & \{\pm 3/2, \pm 9/2, \pm 15/2, \dots\} (0 \bmod 3), \\ & \{\dots, -5/2, +1/2, +7/2, \dots\} (1 \bmod 3), \\ & \{\dots, -7/2, -1/2, +5/2, \dots\} (2 \bmod 3). \end{aligned}$$

It is of interest to correlate the $\{2 J_2 \bmod 3\}$ quantum number with irreducible representations of O_h^* . It can be shown that $\{0 \bmod 3\}$ correlates with Γ_{6g} and Γ_{7g} while both $\{1 \bmod 3\}$ and $\{2 \bmod 3\}$ correlate with Γ_{8g} . Thus, one disadvantage of this basis set is that the $\{0 \bmod 3\}$ block is not factored as much as possible (separate Γ_{6g} and Γ_{7g} sets); nonetheless, with modern computers and diagonalization routines, this is not a serious drawback.

The main task remaining is to find integrals of powers of vibrational coordinates over harmonic oscillator functions; these are often referred to as the primitive matrix elements. General formulae exist⁷ which allow these matrix elements to be evaluated simply.

Primitive matrix elements relevant to the case at hand are:

$$\int X^*(n_2, \ell_2) q_{\pm} X(n_2 + 1, \ell_2 \mp 1) d\tau = \left\{ \frac{1}{2\alpha} (n_2 \mp \ell_2 + 2) \right\}^{1/2}, \quad (7)$$

$$\int X^*(n_2, \ell_2) q_{\pm}^2 X(n_2 + 2, \ell_2 \pm 2) d\tau = \frac{1}{2\alpha} \left\{ (n_2 \pm \ell_2 + 2)(n_2 \pm \ell_2 + 4) \right\}^{1/2}, \quad (8)$$

$$\int X^*(n_2, \ell_2) q_{\mp}^2 X(n_2, \ell_2 \pm 2) d\tau = \frac{1}{2\alpha} \left\{ 4(n_2 \mp \ell_2)(n_2 \pm \ell_2 + 2) \right\}^{1/2}, \quad (9)$$

$$\int X^*(n_2, \ell_2) q_{+} q_{-} X(n_2 + 2, \ell_2) d\tau = \frac{1}{2\alpha} \left\{ (n_2 - \ell_2 + 2)(n_2 + \ell_2 + 2) \right\}^{1/2}, \quad (10)$$

$$\int X^*(n_2, \ell_2) q_{+} q_{-} X(n_2, \ell_2) d\tau = \frac{1}{2\alpha} \{2n_2 + 2\}, \quad (11)$$

in which

$$q_{\pm} = q_{\theta} \pm i q_{\epsilon}$$

$$\alpha = \frac{\omega_{\epsilon}}{\hbar} = \frac{k_{\epsilon}^{1/2}}{\hbar}.$$

The desired vibronic matrix elements are then readily found:

$$\int \vec{b}^{\dagger}(n_2, \ell_2; \rho_3, \sigma_3; J_2) \frac{\hat{\mathcal{H}}^{(1)}}{\hbar \omega_E} \vec{b}(n_2 + 1, \ell_2 + \rho_3; -\rho_3, \sigma_3; J_2) d\tau =$$

$$\{D_2(n_2 + \rho_3 \ell_2 + 2)\}^{1/2}, \quad (12)$$

$$\int \vec{b}^{\dagger}(n_2, \ell_2; \rho_3, \sigma_3; J_2) \frac{\hat{\mathcal{H}}^{(2)}}{\hbar \omega_E} (a_{1g}) \vec{b}(n_2, \ell_2; \rho_3, \sigma_3; J_2) d\tau = \frac{Q_2[a_{1g}]}{2} \{n_2 + 1\}, \quad (13)$$

$$\int \vec{b}^{\dagger}(n_2, \ell_2; \rho_3, \sigma_3; J_2) \frac{\hat{\mathcal{H}}^{(2)}}{\hbar \omega_E} (a_{1g}) \vec{b}(n_2 + 2, \ell_2; \rho_3, \sigma_3; J_2) d\tau =$$

$$\frac{Q_2[a_{1g}]}{4} \{(n_2 + \ell_2 + 2)(n_2 - \ell_2 + 2)\}^{1/2}, \quad (14)$$

$$\int \vec{b}^{\dagger}(n_2, \ell_2; \rho_3, \sigma_3; J_2) \frac{\hat{\mathcal{H}}^{(2)}}{\hbar \omega_E} (e_g) \vec{b}(n_2, \ell_2 - 2\rho_3; -\rho_3, \sigma_3; J_2 - 3\rho_3) d\tau =$$

$$Q_2[e_g] \{4(n_2 + \rho_3 \ell_2)(n_2 - \rho_3 \ell_2 + 2)\}^{1/2}, \quad (15)$$

$$\int \vec{b}^{\dagger}(n_2, \ell_2; \rho_3, \sigma_3; J_2) \frac{\hat{\mathcal{H}}^{(2)}}{\hbar \omega_E} (e_g) \vec{b}(n_2 + 2, \ell_2 - 2\rho_3; -\rho_3, \sigma_3; J_2 - 3\rho_3) d\tau =$$

$$Q_2[e_g] \{(n_2 - \rho_3 \ell_2 + 2)(n_2 - \rho_3 \ell_2 + 4)\}^{1/2}, \quad (16)$$

in which the dimensionless coupling parameters are defined as: $D_2 = \frac{\ell_E^2}{2\hbar \omega_E^3}$,

$$Q_2[a_{1g}] = \frac{C_a}{k_E}, \quad Q_2[e_g] = \frac{C_g}{2k_E}.$$

Note that σ_3 does not appear on the right hand side of any of these equations and thus two identical secular matrices are generated, one for each value (± 1) of the quantum number σ_3 . It will prove to be interesting, for later considerations (see Section III), to use perturbation theory to find the splitting of the $n_2 = 1$ level due to $\hat{\mathcal{H}}^{(2)}(e_g)$ in the unrealistic case of $D_2 = 0$ (no linear coupling). The secular determinant which results from a first order degenerate perturbation theory treatment of the $n_2 = 1$ level, employing equation 15, is

$$\begin{vmatrix} (2 - \lambda) & 4 Q_2[e_g] \\ 4 Q_2[e_g] & (2 - \lambda) \end{vmatrix} = 0 \quad (17)$$

with solutions

$$\lambda = 2 \pm 4 Q_2[e_g].$$

These quantities are measured in units of vibrational energy, $\hbar \omega_e$. It is possible to obtain a first order expression for $\hat{\mathcal{H}}^{(2)}(a_{1g})$ also; however, this is not necessary since an exact expression easily obtains:

$$\lambda = (n_e + 1) \sqrt{1 + Q_2[a_{1g}]} = (n_e + 1) \left(1 + \frac{Q_2[a_{1g}]}{2} + \dots \right) \quad (18)$$

III. DISCUSSION

Examples of the secular matrix calculation of energy levels of $[\Gamma_8 \times e_g]_{\text{QUAD}}$ are given in Figures 1 and 2. The secular matrix used in these calculations was truncated after $n_2 = 10$; a basis of this size results in 44×44 matrices for both the $\{0 \bmod 3\}$ and $\{1 \bmod 3\}$ blocks (the $\{2 \bmod 3\}$ block gives the same eigenvalues as $\{1 \bmod 3\}$). Under these circumstances it is required that D_2 be less than 1 for an accurate description of the levels. The upper limit of $Q_2[e_g]$ was not numerically determined, but is probably about 0.5. The behavior of the levels at these large values of D_2 and $Q_2[e_g]$ is quite complicated due to specific level repulsions.

D_2 may be taken as positive since the relevant matrix elements are all off-diagonal (Eqn. 12). The situation with respect to $Q_2[e_g]$ is more complicated even though it too has all off-diagonal matrix elements in the employed linear JT basis (Eqn. 15, 16). $\hat{\mathcal{H}}^{(2)}(e_g)$ will have diagonal matrix elements in a symmetry adapted basis set. The eigenvalues of $\hat{\mathcal{H}}^{(2)}(e_g)$ are independent of the sign of $Q_2[e_g]$, however, the concomitant changes in the eigenvectors upon changing the sign of $Q_2[e_g]$ may alter their symmetry transformation properties. For $Q_2[e_g] \rightarrow -Q_2[e_g]$, $\Gamma_6 \rightarrow \Gamma_7$ and $\Gamma_7 \rightarrow \Gamma_6$.

Several general observations can be made based on these calculations:

i) For the case $D_2 = 0$ (Figure 1), the perturbation expression Eq. 17 is found to be useful over a large range of $Q_2[e_g]$ values ($Q_2[e_g] < 0.1$) by direct comparison with results of the truncated secular equation calculation. This situation arises because $\hat{\mathcal{H}}^{(2)}(e_g)$ couples n_2 and $n_2 \pm 2$ levels, but not n_2 and $n_2 \pm 1$, as in the linear case.

ii) In the regime ($D_2 < 0.2$, $Q_2[e_g] \leq 0.05$), for the $n_2 = 1$ levels, effects of linear and quadratic terms are found to be approximately independent; that is, the ($n_2 = 1$, $J_2 = 1/2$) level is not shifted appreciably as $Q_2[e_g]$ increases, center-

of-gravity of the ($n_2 = 1, J_2 = 3/2$) levels is preserved, and splitting of ($n_2 = 1, J_2 = 3/2$) is approximately the same as in the $D_2 = 0$ case.

iii) Behavior of the $n_2 = 2$ levels are, however, qualitatively different from those of $n_2 = 1$ (see Figures 1 and 2); one difference is in the behavior of the ($n_2 = 2, J_2 = 3/2$) levels (Figure 1). For $D_2 = 0$, these levels do not split under $\hat{\mathcal{H}}^{(2)}(e_g)$, whereas $D_2 \neq 0$ allows admixture of $n_2 = 1, 3, \dots$ levels, thus causing ($n_2 = 2, J_2 = 3/2$) to split. It should be noted that existing perturbative treatments of $\hat{\mathcal{H}}$ are not able to generate this splitting,⁵ although a more complete perturbation calculation would, of course, reproduce these results.

Based on the form of the perturbation energy expressions for the linear and quadratic JT terms, it can be seen that certain n_2 levels are split by $\hat{\mathcal{H}}^{(2)}(e_g)$ in the first order of perturbation theory (e.g., Eq. 17) while the same n_2 levels are split by $\hat{\mathcal{H}}^{(1)}$ only in the second order.⁸ It is thus possible for quadratic JT terms to be more effective at splitting vibronic levels than are linear JT terms. Examination of the perturbation expressions for the $n_2 = 1$ levels verifies this idea:

$$\begin{aligned} \lambda (\hat{\mathcal{H}}^{(1)}) &= 2 \pm 2(D_2^{1/2})^2 \\ \lambda (\hat{\mathcal{H}}^{(2)}(e_g)) &= 2 \pm 4Q_2[e_g]. \end{aligned} \quad (19)$$

$D_2^{1/2}$ is used for comparison with $Q_2[e_g]$ since it is proportional to λ_e just as $Q_2[e_g]$ is proportional to C_e . Thus, for $D_2^{1/2} = Q_2[e_g] = 0.1$, Eqs. 19 show that the splitting induced by $\hat{\mathcal{H}}^{(2)}(e_g)$ is an order of magnitude more than that induced by $\hat{\mathcal{H}}^{(1)}$. Similar results obtain for $\hat{\mathcal{H}}^{(2)}(a_{1g})$ (see Eq. 18).

Available hexafluoride data¹⁻³ show that the splitting induced by the linear terms is of the same order of magnitude as the quadratically-induced splittings and shifts. Although these data indicate the expansion of $\hat{\mathcal{H}}$ in powers of q_θ and q_ϵ is not converging as rapidly as might be desired, the entire effect cannot be attributed to non-linearity of $\hat{\mathcal{H}}$. Greater effectiveness of $\hat{\mathcal{H}}^{(2)}(e_g)$ and $\hat{\mathcal{H}}^{(2)}(a_{1g})$ in causing shifts and splittings is also an important consideration.

IV. CONCLUSION

An accurate numerical method is given for treating vibronic coupling of $r_{8g}(0_h^*)$ electronic state to an e_g vibration for a vibronic Hamiltonian with linear and quadratic terms. Examples of the calculation are given and general comments on the behavior of the levels under the influence of quadratic terms are made. It is also found that the quadratic coupling term is a more effective perturbation than the linear coupling term with respect to the observed spectroscopic splitting.

REFERENCES

1. G. R. Meredith, J. D. Webb and E. R. Bernstein, Mol. Phys. 34, 995 (1977).
2. E. R. Bernstein and J. D. Webb, Mol. Phys. __, (1978).
3. E. R. Bernstein and J. D. Webb, Mol. Phys. __, (1978).
4. R. Englman, "The Jahn-Teller Effect in Molecules and Crystals," (Wiley-Interscience, New York, 1972).
5. B. Weinstock and G. L. Goodman, Adv. Chem. Phys. 9, 169 (1965).
6. M. C. M. O'Brien, Proc. Roy. Soc. A 281, 323 (1964).
7. C. DiLauro, J. Mol. Spec. 41, 598 (1972).
8. W. Moffitt and W. Thorson, Phys. Rev. 108, 1251 (1957).

Figure 1.

Quadratic Jahn-Teller calculation for coupling of a Γ_8 electronic state with a $\nu_2(e_g)$ vibration, $(\Gamma_8 \times e_g)_{\text{QUAD}}$. The fixed parameter values are $D_2 = 0$ and $\nu_2^0 = 670 \text{ cm}^{-1}$. D_2 is the linear Jahn-Teller coupling parameter and ν_2^0 is the unperturbed ν_2 harmonic oscillator frequency. $Q_2 [e_g]$ is the quadratic coupling parameter. Note the large range over which the splitting is linear in $Q_2 [e_g]$. In the method employed here, Γ_6 and Γ_7 levels both arise from the same secular matrix (the $\{0 \bmod 3\}$ block, see text). The symmetry labels of the eigenvectors are generated from their transformation properties. Changing the sign of $Q_2 [e_g]$ will not alter the energies, but will interchange Γ_6 and Γ_7 symmetry labels.





Figure 2.

Similar to Figure 1 but D_2 has been set to 0.125. Note that the $n_2 = 1$ for this value of D_2 levels behave similarly to those for which $D_2 = 0$ in that one pair of levels splits symmetrically and the other level remains largely unaffected as $Q_2 [e_g]$ varies. This situation does not hold, however, for the $n_2 = 2$ levels. The J_2 quantum number, which is good at $Q_2[e_g] = 0$, is indicated.



TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP 1	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1		
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1		

TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. A. El-Sayed University of California Department of Chemistry Los Angeles, California 90024	1	Dr. G. B. Schuster University of Illinois Chemistry Department Urbana, Illinois 61801	1
Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163	1	Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1
Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1
Dr. M. H. Chisholm Princeton University Department of Chemistry Princeton, New Jersey 08540	1	Dr. M. Rauhut American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805	1
Dr. J. R. MacDonald Naval Research Laboratory Chemistry Division Code 6110 Washington, D.C. 20375	1		