

AD-A054 546

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER  
UNIQUENESS OF SOLUTIONS TO  $-\Delta u = 0$ . (U)  
FEB 78 R JENSEN

F/G 12/1



DAAG29-75-C-0024  
NL

UNCLASSIFIED


MRC-TSR-1827

| of |

AD  
2054546



END  
DATE  
FILMED  
6 -78  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

FOR FURTHER TRAN

17

AD A 054546

MRC Technical Summary Report #1827

UNIQUENESS OF SOLUTIONS TO

$-\Delta u - qu = 0$

Robert Jensen

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

February 1978

(Received December 1, 1977)

DDC FILE COPY

DDC  
RECEIVED  
JUN 2 1978  
D

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

National Science Foundation  
Washington, D. C. 20550

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

UNIQUENESS OF SOLUTIONS TO  $-\Delta u - qu = 0$

Robert Jensen

Technical Summary Report #1827  
February 1978

ABSTRACT

We give conditions for the function  $q(x)$  which guarantee uniqueness of solutions to  $-\Delta u(x) + \lambda_0 u(x) - q(x)u(x) = f$ , where  $\lambda_0 \geq 0$  is a constant. We consider separately the cases for solutions in  $H^1$  and  $L^2$ . In fact we even prove existence of  $H^1$  solutions under the additional assumptions  $q(x) \geq 0$  and  $f \in L^2$ . We apply our results to the problem of essential selfadjointness for a class of Schrödinger operators and we are able to prove an interesting generalization of a result due to Kato.

Gaveau has proved a result similar to ours on  $L^2$  uniqueness. His proof, however, depends on a probabilistic argument, while in contrast we do not use any probability theory in our proofs.

AMS (MOS) Subject Classifications: 35A05, 35J10

Key Words: Bessel potential, Distribution, Measure, Schrödinger operator

Work Unit Number 1 (Applied Analysis)

ACQUISITION NO.	
DTIC	White Section <input checked="" type="checkbox"/>
DDI	Diff Section
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024 and the National Science Foundation under Grant No. MCS75-17385 A01.

## SIGNIFICANCE AND EXPLANATION

The Schrödinger wave equation is of fundamental importance in quantum mechanics. In the early 1930's von Neumann made use of the theory of self-adjoint operators on a Hilbert space to place quantum mechanics on a firm mathematical foundation. Thus it is important to know under what conditions a Schrödinger operator with singular potential is selfadjoint on a Hilbert space. The obvious Hilbert space is  $L^2$ , the space of square-integrable functions.

We consider Schrödinger operators of the form

$$T \equiv \sum_{j=1}^n \left( \frac{\partial}{\partial x_j} - ib_j(x) \right)^2,$$

with a singular potential  $q(x)$ . The operator  $T + q$  is difficult to interpret as an operator on  $L^2$  since  $L^2$  functions are not necessarily differentiable. This difficulty is circumvented by the following device: define

$$\overline{(T + q)}(u) \equiv w \quad \text{for } u, w \in L^2$$

if there is a sequence of infinitely differentiable functions  $\phi_i \in C_0^\infty$  such that  $\phi_i \rightarrow u$  in  $L^2$  and  $(T + q)(\phi_i) \rightarrow w$  in  $L^2$ . The problem now is that  $\overline{T + q}$  may not be well defined. If it is then we say  $T + q$  is essentially selfadjoint and  $\overline{T + q}$  is a selfadjoint operator on the Hilbert space  $L^2$ . Our paper generalizes results of T. Kato on conditions which guarantee  $T + q$  is essentially selfadjoint.

Robert Jensen

Mathematics Research Center  
University of Wisconsin-Madison

We give conditions on  $q(x)$  which guarantee that if

$u(x) \in L^2(\mathbb{R}^n)$  and

$$(0.1) \quad -\Delta u(x) + \lambda_0 u(x) - q(x)u(x) = 0, \quad \lambda_0 \geq 0 \text{ a constant,}$$

then  $u(x) \equiv 0$ . In fact, we show that if there exists a constant  $\theta_0$  such that

$$(0.2) \quad (-\Delta + \lambda I)^{-1} (q^+)(x) \leq \theta_0 < 1, \text{ for all } \lambda > \lambda_0,$$

then  $u(x) \in L^2(\mathbb{R}^n)$  satisfies (0.1) if and only if  $u(x) \equiv 0$ .

This result is contained in our Theorem 2.2. (Gaveau has proved this result by probabilistic means. Our techniques are entirely different from his and use no probability theory.)

Our results can also be applied to a larger class of second order elliptic operators. Indeed, remark 2 of section 1 gives a class of operators to which our results can be extended.

Furthermore, by using a beautiful lemma due to T. Kato (the lemma

Sponsored by the United States Army under Contract No. DAMG29-75-C-0024 and the National Science Foundation under Grant No. MCS75-17385 A01.

and proof are found in [2]) it is possible to apply uniqueness results about  $-\Delta + \lambda_0 - q$  to a larger class of Schrödinger operators. A statement of Kato's lemma is found in section 2 of this paper.

The paper proceeds as follows. In section 1 we obtain existence and uniqueness results for

$$(0.3) \quad -\Delta u + \lambda_0 u - \mu u = f$$

where  $\mu$  is a non-negative measure. We obtain these results for  $f \in L^2$  and  $u \in H^1$ . Section 2 contains the  $L^2$  uniqueness result for (0.2). The proof depends heavily on the material developed in section 1. Finally, section 3 contains an application of the results of section 2. We prove a strong generalization of the results found in [2] on essential self-adjointness.

## 1. Existence and uniqueness of solutions in $H^1$

We use the following notation throughout this paper.

$$C_0^\infty \equiv C_0^\infty(\mathbb{R}^n), \quad C^\infty \equiv C^\infty(\mathbb{R}^n),$$

$$L^p \equiv L^p(\mathbb{R}^n), \quad L_{loc}^p \equiv L_{loc}^p(\mathbb{R}^n),$$

$$H^s \equiv H^s(\mathbb{R}^n), \quad H_{loc}^s \equiv H_{loc}^s(\mathbb{R}^n),$$

$$L^p(\mu) \equiv L^p(\mu; \mathbb{R}^n) \text{ for } \mu \text{ a measure on } \mathbb{R}^n,$$

$$(\psi_1, \psi_2) \equiv \int_{\mathbb{R}^n} \psi_1 \cdot \bar{\psi}_2(x) dx \text{ for } \psi_i \in L^2, \quad i = 1, 2,$$

$$(\psi_1, \psi_2)_\mu \equiv \int_{\mathbb{R}^n} \psi_1 \cdot \bar{\psi}_2(x) d\mu(x) \text{ for } \psi_i \in L^2(\mu), \quad i = 1, 2.$$

Until further notice (near the end of section 2) all functions and measures are real valued. We shall also use the Bessel potentials,

$$J_\lambda \equiv F^{-1}((4\pi^2|\xi|^2 + \lambda)^{-\lambda/2}), \quad \lambda > 0,$$

where  $F^{-1}$  is the inverse Fourier transform. For each measurable function  $\phi: \mathbb{R}^n \rightarrow \mathbb{R}$  define

$$(-\Delta + \lambda I)^{-1} \phi(x) \equiv \int_{\mathbb{R}^n} J_\lambda(x-y) \phi(y) dy,$$

whenever  $J_\lambda(x-\cdot) \phi(\cdot) \in L^1$  for a.e.  $x \in \mathbb{R}^n$ . The functions  $J_\lambda$  have the property,

$$(1.1) \quad \text{if } \lambda > \lambda' \text{ then } 0 \leq J_\lambda \leq J_{\lambda'}.$$

(More information about  $J_\lambda$  can be found in [3].)

The following assumption will be used.

$$(1.2) \quad \left[ \begin{array}{l} \mu \text{ is a non-negative measure on } \mathbb{R}^n; \text{ there is} \\ \text{a constant } \lambda_0 \geq 0 \text{ such that for all } \lambda > \lambda_0 \\ (-\Delta + \lambda I)^{-1}(\mu)(x) \leq \theta_0 \leq 1 \text{ for some constant } \theta_0 \geq 0 \end{array} \right.$$

and whenever  $\lambda_0$  or  $\theta_0$  appear in this paper they are the constants in (1.2). (We define

$$(-\Delta + \lambda I)^{-1}(\mu)(x) \equiv \int_{\mathbb{R}^n} J_\lambda(x-y) d\mu(y).)$$

Although we shall delay the proof until later we now state the main theorem of this section:

Theorem 1.1. Let (1.2) hold and let  $\phi \in H^1$ . Then  $\phi \in L^2(\mu)$  and

$$(1.3) \quad (\nabla\phi, \nabla\phi) + \lambda_0(\phi, \phi) - (\phi, \phi)_\mu \geq (1 - \theta_0) (\nabla\phi, \nabla\phi),$$

$$\text{where } \nabla\phi \equiv \left( \frac{\partial\phi}{\partial x_1}, \dots, \frac{\partial\phi}{\partial x_n} \right).$$

Corollary 1.2. Let (1.2) hold, let  $\lambda > \lambda_0$ ,  $f \in L^2$  and  $\theta_0 < 1$ . Then there is a unique function  $w \in H^1$  such that

$$(1.4) \quad (-\Delta w + \lambda w, \phi) - (w, \phi)_\mu = (f, \phi),$$

for all  $\phi \in C_0^\infty$ . Furthermore, there is a constant  $C(\theta_0, \lambda)$  such that

$$(1.5) \quad \|w\|_{H^1} \leq C(\theta_0, \lambda) \|f\|_{L^2}.$$

Proof. Using Theorem 1.1 we see that for any  $\phi \in H^1$

$$(1.6) \quad (\nabla\phi, \nabla\phi) + \lambda(\phi, \phi) - (\phi, \phi)_\mu \geq (1 - \theta_0) (\nabla\phi, \nabla\phi) + (\lambda - \lambda_0) (\phi, \phi).$$

The existence and uniqueness of  $w$  and the estimate (1.5) all follow from the Riesz theorem in the usual way.

The following remarks (verifications of which are of varying degrees of difficulty) represent several directions in which generalizations of our results can be made.

Remark 1. Suppose we replace  $\mathbb{R}^n$  by a domain  $\Omega$  and change (1.2) to

$$(1.7) \quad \left[ \begin{array}{l} \mu \text{ is a non-negative measure on } \Omega, \text{ there are} \\ \text{constants } \lambda_0 \geq 0, 1 \geq \theta_0 \geq 0 \text{ and a measurable} \\ \text{function } v(x), 0 \leq v(x) \leq \theta_0 \text{ such that} \\ -\Delta v + \lambda_0 v = \mu \text{ (in the sense of distributions)}. \end{array} \right.$$

Then all our assertions remain valid with  $H^1$  replaced by  $H_0^1(\Omega)$  and  $L^2$  replaced by  $L^2(\Omega)$ . Although it may not be apparent, if  $\Omega = \mathbb{R}^n$  then (1.2) and (1.7) are equivalent.

Remark 2. Let  $A$  be the partial differential operator

$$(1.8) \quad A\phi(x) \equiv \sum_{i,j=1}^n -\frac{\partial}{\partial x_i} \left( a_{ij}(x) \frac{\partial \phi}{\partial x_j}(x) \right) \quad \text{for } \phi \in C_0^\infty,$$

where  $(a_{ij}(x))$  is uniformly positive definite and symmetric and the  $a_{ij}(x)$  are real valued, smooth and bounded. If we now replace  $-\Delta$  by  $A$  and (1.2) by (1.7) then analogous results are valid.

We return again to the business at hand, namely, the proof of Theorem 1.1. We remind the reader of a definition:

$$(1.9) \quad \left[ \begin{array}{l} \{b_j(x)\}_{j=1}^\infty \subset L_{loc}^1 \text{ converges in } D'(\mathbb{R}^n) \text{ to a} \\ \text{measure } \mu \text{ if for any } \phi \in C_0^\infty, (\phi, b_j) \rightarrow (\phi, \mu) \\ \text{as } j \rightarrow \infty. \end{array} \right.$$

Lemma 1.3. Let (1.2) hold and let  $\lambda$  be a constant. Then:

(i) There exists a sequence of functions  $\{a_j\}_{j=1}^\infty \subset C_0^\infty \cap L^\infty$  such that  $a_j(x) \geq 0 \forall j$ ,  $\{a_j\}_{j=1}^\infty$  converges in  $D'(\mathbb{R}^n)$  to  $\mu$  and if  $\lambda > \lambda_0$  then

$$(1.10) \quad (-\Delta + \lambda I)^{-1}(a_j)(x) \leq \theta_j < \theta_0.$$

(ii) Let  $\bar{a}(x) \in L^\infty$ ,  $\bar{a}(x) \geq 0$  and  $(-\Delta + \lambda I)^{-1}(\bar{a})(x) \leq \bar{\theta} < 1$ . Then for any function  $g \in L^2$ ,  $g \geq 0$  which satisfies

$$(1.11) \quad (-\Delta + \lambda I)^{-1}(g)(x) \in L^\infty,$$

there is a function  $G \in H_{loc}^2 \cap L^\infty$  such that

$$(1.12) \quad \begin{cases} -\Delta G + \lambda G - \bar{a}G = g \\ 0 \leq G(x) \leq (1 - \bar{\theta})^{-1} \|(-\Delta + \lambda I)^{-1}(g)\|_{L^\infty}. \end{cases}$$

Furthermore, if in addition  $\bar{a}(x)$ ,  $g(x) \in C^\infty$  then  $G \in C^\infty$ .

Proof of (i). Simply define  $a_j(x)$  by

$$a_j(x) \equiv \left\{ \left( \frac{j-1}{j} \right) \int_{\mathbb{R}^n} K_j e^{-j|x-y|} d\mu(y) \right\} \xi_j(x)$$

where  $K_j$  is the normalizing constant

$$(K_j)^{-1} \equiv \int_{\mathbb{R}^n} e^{-j|x|} dx,$$

and  $\xi_j \in C_0^\infty$ ,  $\xi_j \rightarrow 1$  uniformly on compact sets. It amounts to an exercise to verify that the sequence  $\{a_j\}_{j=1}^\infty$  just defined has the required properties.

Proof of (ii). We shall obtain  $G$  as the limit of a sequence of functions  $\{G_i\}_{i=1}^\infty$ . We define  $G_i$  inductively by

$$(1.13) \quad \begin{cases} G_0(x) \equiv 0, \\ G_{i+1}(x) \equiv (-\Delta + \lambda I)^{-1}(\bar{a}G_i + g)(x). \end{cases}$$

First we establish that  $G_i$  is monotone increasing. Indeed, setting

$$\bar{G}_i = G_i - G_{i-1}$$

we see  $\bar{G}_i$  satisfies the equation

$$(1.14) \quad \bar{G}_{i+1} = (-\Delta + \lambda I)^{-1}(\bar{a}\bar{G}_i).$$

Since  $\bar{G}_1 = G_1 - G_0 = G_1 \geq 0$ , we conclude by induction that

(1.15)  $\bar{G}_i \geq 0$  for all  $i$ ,

which proves the monotonicity.

Next, let us estimate  $\|\bar{G}_i\|_{L^\infty}$ . From (1.14) we see that

since  $\bar{a}(x) \geq 0$

$$\begin{aligned} \|\bar{G}_{i+1}\|_{L^\infty} &\leq \|(-\Delta + \lambda I)^{-1} (\|\bar{G}_i\|_{L^\infty} \bar{a})\|_{L^\infty} \\ &\leq \|\bar{G}_i\|_{L^\infty} \|(-\Delta + \lambda I)^{-1} \bar{a}\|_{L^\infty}. \end{aligned}$$

By the above and the hypothesis on  $\bar{a}(x)$  we have the inequality

$$\|\bar{G}_{i+1}\|_{L^\infty} \leq \|\bar{G}_i\|_{L^\infty} \bar{\theta}.$$

Using induction we conclude

$$(1.16) \quad \|\bar{G}_{i+1}\|_{L^\infty} \leq \|(-\Delta + \lambda I)^{-1}(g)\|_{L^\infty} (\bar{\theta})^i.$$

From the inequality above we see that  $\{\bar{G}_i\}_{i=1}^\infty$  is a

Cauchy sequence in the  $L^\infty$  norm. Therefore the sequence has

a limit which we denote by  $G(x)$ . By (1.15) and (1.16) we have

$$0 \leq G(x) \leq (1 - \bar{\theta})^{-1} \|(-\Delta + \lambda I)^{-1}(g)\|_{L^\infty}.$$

So it remains only to show  $G \in H_{loc}^2$ . This fact can be

proved by noting that for all  $i$ ,  $G_i \in H_{loc}^2$  since

$$-\Delta G_i + \lambda G_i = \bar{a} G_{i-1} + g$$

and because  $\bar{a}(x), g(x) \in L^\infty$ . Therefore

$$(1.17) \quad -\Delta G + \lambda G - \bar{a} G = g \quad (\text{in the sense of distributions}).$$

Since  $\bar{a}(x), g(x), G(x) \in L^\infty$  it follows from a result of

Friedrichs' that (1.17) implies

$$G \in H_{loc}^2.$$

All that now remains is to consider the case  $\bar{a}(x)$ ,

$g(x) \in C^\infty$  also. In this case, however, it follows by induction that

$$G \in H_{loc}^{2l} \quad \text{for all positive integers } l.$$

This completes the proof of the lemma.

We end this section with the proof of Theorem 1.1.

Proof of Theorem 1.1. It will be sufficient to prove that for every  $\phi \in C_0^\infty$

$$(1.18) \quad (\nabla \phi, \nabla \phi) + \lambda_0 (\phi, \phi) - (\phi, \phi)_\mu \geq (1 - \theta_0) (\nabla \phi, \nabla \phi).$$

We shall prove (1.18) by constructing special functions

$W_j(x) \in C^\infty$  and decomposing  $\phi$  as

$$(1.19) \quad \phi = \sum_j W_j \psi_j,$$

and then expanding  $(\nabla(W_j \psi_j), \nabla(W_j \psi_j))$  appropriately (using the properties of  $W_j$ ).

We define  $W_j(x)$  as follows. Let  $\{a_j\}_{j=1}^\infty$  be the sequence of functions whose existence is guaranteed by (i) of Lemma 1.3.

We let  $G_j$  be the solution of

$$(1.20) \quad -\Delta G_j + \lambda G_j - a_j G_j = a_j f_\lambda \quad \text{for any } \lambda > \lambda_0,$$

$$\text{where } f_\lambda(x) = \frac{1}{2n} \left( \sum_{k=1}^n \sqrt{\lambda} x_k + e^{-\sqrt{\lambda} x_k} \right),$$

which we can do by (ii) of Lemma 1.3 and where  $\bar{a} = a_j$  and

$g = a_j f_\lambda$ . Finally, we define  $W_j$  as

$$W_j \equiv G_j + f_\lambda.$$

Notice that by (1.20) and (ii) of Lemma 1.3

$$(1.21) \quad \begin{cases} W_j \in L^\infty \cap C^\infty, & W_j(x) \geq 1, \\ -\Delta W_j + \lambda W_j - a_j W_j = 0. \end{cases}$$

We now evaluate  $(\nabla\phi, \nabla\phi)$  defining  $\psi_j \in C_0^\infty$  by (1.19).

We have using integration by parts,

$$\begin{aligned} (\nabla\phi, \nabla\phi) &= (\nabla(W_j \psi_j), \nabla(W_j \psi_j)) \\ &= (W_j \psi_j, -\Delta(W_j \psi_j)) \\ &= (W_j \psi_j, -W_j (\Delta\psi_j) - 2(\nabla W_j, \nabla\psi_j)) \\ &\quad + (W_j \psi_j, (-\Delta W_j) \psi_j) \\ &= (\psi_j, -W_j^2 (\Delta\psi_j) - \nabla(W_j^2) \nabla\psi_j) + (W_j \psi_j, (-\Delta W_j) \psi_j). \end{aligned}$$

Using integration by parts again and (1.21) we get

$$\begin{aligned} (\nabla\phi, \nabla\phi) + \lambda(\phi, \phi) &= (W_j (\nabla\psi_j), W_j (\nabla\psi_j)) + (W_j \psi_j, a_j W_j \psi_j) \\ &\geq (\phi, a_j \phi), \\ &\geq (\phi^2, a_j). \end{aligned}$$

By (i) of Lemma 1.3  $(a_j)_{j=1}^\infty$  converges in  $D'(\mathbb{R}^n)$  to  $\mu$  and so

$$(\nabla\phi, \nabla\phi) + \lambda(\phi, \phi) \geq (\phi^2, 1)_\mu = (\phi, \phi)_\mu.$$

Next by noting that the above inequality holds for all

$\lambda > \lambda_0$  we see

$$(1.22) \quad (\nabla\phi, \nabla\phi) + \lambda_0(\phi, \phi) \geq (\phi, \phi)_\mu.$$

Now if (1.2) holds for  $\mu$  with  $\theta_0 < 1$  then (1.2) holds for  $\tilde{\mu} = \theta_0^{-1} \mu$  with  $\tilde{\theta}_0 = 1$ . Thus (1.22) holds for  $\mu$  replaced by  $\tilde{\mu}$  and this gives us

$$(1.23) \quad (\nabla\phi, \nabla\phi) + \lambda_0(\phi, \phi) \geq (\phi, \phi)_\mu.$$

Using the definition of  $\tilde{\mu}$  and rearranging (1.23) we get

$$(\nabla\phi, \nabla\phi) + \lambda_0(\phi, \phi) - \theta_0^{-1}(\phi, \phi)_\mu \geq 0.$$

From this we have

$$\theta_0 (\nabla\phi, \nabla\phi) + \lambda_0 \theta_0 (\phi, \phi) - (\phi, \phi)_\mu \geq 0$$

and by adding  $(1 - \theta_0) (\nabla\phi, \nabla\phi)$  to both sides of the above inequality we get

$$(\nabla\phi, \nabla\phi) + \lambda_0 \theta_0 (\phi, \phi) - (\phi, \phi)_\mu \geq (1 - \theta_0) (\nabla\phi, \nabla\phi).$$

From this (1.3) follows easily and this completes the proof of Theorem 1.1.

## 2. Uniqueness of solutions in $L^2$

From our previous work we shall prove two uniqueness results for functions in  $L^2$  which satisfy an appropriate partial differential equation or inequality. A version of our second result has been proved in a recent paper by Gaveau.

We let  $q(x) \in L_{loc}^1$ ,  $q(x) \geq 0$ . To such a function we associate a non-negative measure  $\mu_q$  by setting

$$\mu_q(G) \equiv \int_G q(x) dx.$$

When we say (1.2) holds for  $q$  we mean (1.2) holds for  $\mu_q$ .

For  $q(x) \in L^1_{loc}$ ,  $q(x) \geq 0$  we also define  $q_j(x)$  by

$$(2.1) \quad q_j(x) = \begin{cases} q(x) & \text{if } q(x) < j \\ j & \text{if } q(x) \geq j \end{cases}$$

Theorem 2.1. Let  $q(x) \in L^1_{loc}$ ,  $q(x) \geq 0$  and suppose (1.2)

holds for  $q$  with  $\theta_0 < 1$ . If  $u(x) \in L^2$ ,  $u(x) \geq 0$ ,

$qu(x) \in L^1_{loc}$ ,  $\lambda > \lambda_0$  and

$$(2.2) \quad -\Delta u + \lambda u - qu \leq 0 \quad (\text{in the sense of distributions}),$$

then  $u(x) \equiv 0$ .

Proof. The idea of the proof is to solve a sequence of approximate adjoint equations to (2.2) for functions in some appropriate class and then show  $u \equiv 0$  by using these with inequality (2.2). Let  $g \in C^\infty_{0+} = \{\psi \in C^\infty_0 \mid \psi \geq 0\}$ ; for each positive integer  $j$  let  $G^j$  be the solution of

$$(2.3) \quad -\Delta G^j + \lambda G^j - q_j G^j = g,$$

(where the  $q_j(x)$  are defined by (2.1)) which we know we can solve uniquely by Corollary 1.2. By Corollary 1.2 and Lemma 1.3

$$(2.4) \quad \begin{cases} \|G^j\|_{H^1} \leq C(\theta_0, \lambda) \|g\|_2, \\ 0 \leq G^j(x) \leq K, \text{ a constant independent of } j. \end{cases}$$

We note also that since  $q_j \in L^\infty$

$$(2.5) \quad G^j \in H^2,$$

however, the norm of  $G^j$  in  $H^2$  may explode as  $j \rightarrow \infty$ .

We give a modification due to H. Brezis of our original proof.

We introduce the functions  $\zeta_R(x) \in C^\infty_0$

$$(2.6) \quad \begin{cases} 0 \leq \zeta_R(x) \leq 1, \\ \zeta_R(x) = 0 & \text{if } |x| > R+1, \\ \zeta_R(x) = 1 & \text{if } |x| < R, \end{cases}$$

and  $\zeta_R(x)$  has first and second derivatives uniformly bounded in  $R$  for all  $R > 0$ .

Thus by (2.3)

$$(g, u) = (u, -\zeta_R \Delta G^j + \zeta_R \lambda G^j - \zeta_R q_j G^j),$$

for  $R$  sufficiently large. From this we get

$$(2.7) \quad (g, u) = (u, -\Delta(\zeta_R G^j) + \lambda(\zeta_R G^j) - q(\zeta_R G^j)) + (u, (\Delta \zeta_R) G^j + 2(\nabla \zeta_R) \nabla G^j) + (u, (q - q_j) \zeta_R G^j).$$

Now,  $\zeta_R G^j(x) \in H^2 \cap L^\infty$ ,  $\zeta_R G^j(x) \geq 0$  and  $\zeta_R G^j(x)$  has compact support. Choosing

$$\psi_l \rightarrow \zeta_R G^j \text{ in } H^2 \cap L^\infty,$$

and by integrating (2.2) multiplied by  $\psi_l$  we conclude upon letting  $l \rightarrow \infty$  that,

$$(u, -\Delta(\zeta_R G^j) + \lambda(\zeta_R G^j) - q(\zeta_R G^j)) \leq 0.$$

From this and (2.7) we get

$$(u, g) \leq (u, (\Delta \zeta_R) G^j + 2(\nabla \zeta_R) \nabla G^j) + (u, (q - q_j) \zeta_R G^j).$$

For  $R$  fixed we see that

$$(2.8) \quad (u, (q - q_j) \zeta_R G^j) \rightarrow 0 \text{ a.e. as } j \rightarrow \infty.$$

By (2.1) and (2.4) we have

$$|u(q - q_j) \zeta_R^j| \leq K |u \zeta_R|.$$

From this, (2.8), since  $u \zeta \in L^1_{loc}$  and because  $\zeta_R \in C^\infty_0$  we may apply the Lebesgue dominated convergence theorem to conclude

$$(u, (q - q_j) \zeta_R^j) \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Using this in (2.7) gives us

$$(2.9) \quad (u, g) \leq (u, (\Delta \zeta_R) G + 2(\nabla \zeta_R) \nabla G).$$

By hypothesis  $\Delta \zeta_R, \nabla \zeta_R$  are uniformly bounded independent of  $R$  and so there is a positive constant  $C'$  such that  $C'$  is independent of  $R$  and

$$(2.10) \quad \max\{\|\Delta \zeta_R\|_{L^\infty}, \|\nabla \zeta_R\|_{L^\infty}\} \leq C'.$$

We see from (2.6) that

$$\text{supp}(|\nabla \zeta_R| + |\Delta \zeta_R|) \subset A_R \equiv \{x \in \mathbb{R}^n \mid R < |x| < R + 1\}.$$

Using this fact, (2.4) and (2.10) we derive

$$(2.11) \quad (u, g) \leq C' C(\theta, \lambda) \|g\|_{L^2 A_R} \int_{L^2 A_R} |u|^2(x) dx,$$

from (2.9). By assumption  $u \in L^2$  and therefore

$$\int_{A_R} |u|^2(x) dx \rightarrow 0 \text{ as } R \rightarrow \infty.$$

Using this in (2.11) we get

$$(u, g) \leq 0.$$

Since the above inequality is true for all  $g \in C^\infty_{0+}$  we see that

$$u(x) \equiv 0.$$

This completes the proof of Theorem 2.1.

In order to prove our next theorem we require the use of a lemma due to Kato. With the aid of this lemma we can reduce the proof of Theorem 2.2 to the case just handled. For the rest of this paper functions and constants will be complex valued unless specifically stated otherwise.

[2]; Lemma A, Pg. 138. Let

$$(2.12) \quad L = \sum_{j,k=1}^n \left( \frac{\partial}{\partial x_j} - i b_j(x) \right) a_{jk}(x) \left( \frac{\partial}{\partial x_k} - i b_k(x) \right),$$

where  $a_{jk}$  and  $b_j$  are real valued functions in  $C^1(\Omega)$ ,  $\Omega$  being an open set in  $\mathbb{R}^n$ . Assume that  $(a_{jk}(x))$  is a positive definite symmetric matrix for each  $x \in \Omega$ . If  $u$  and  $Lu$  are in  $L^1_{loc}(\Omega)$  then

$$(2.13) \quad L_0 |u| \geq \text{Re}\{(\text{sign } \bar{u}) Lu\}$$

where  $L_0$  is the operator  $L$  with  $b_j = 0$ ,  $j = 1, \dots, n$ . (By definition  $\text{sign}(\bar{u}(x)) = \bar{u}(x)/|u(x)|$  if  $u(x) \neq 0$  and  $\text{sign}(\bar{u}(x)) = 0$  if  $u(x) = 0$ .) Note that we are now allowing  $u(x)$  to be a complex valued function.

The following result is now easily proved.

Theorem 2.2. Let  $q_1(x) \geq 0$  and  $q_2(x) \geq 0$  be real valued and measurable on  $\mathbb{R}^n$  and suppose (1.2) holds for  $q_2(x)$ . If  $u(x) \in L^2$ ,  $(q_1 + q_2)u \in L^1_{loc}$ ,  $\text{Re}(\zeta) > \lambda_0$ ,  $\theta_0 < 1$  and

$$(2.14) \quad \left( - \sum_{j=1}^n \left( \frac{\partial}{\partial x_j} - i b_j(x) \right)^2 + \zeta + q_1(x) - q_2(x) \right) u = 0$$

(in the sense of distributions),

(where  $b_j(x)$ ,  $j = 1, \dots, n$  are real valued continuously

differentiable) then

$$u(x) \equiv 0.$$

Proof. By Lemma A (from [2]) with  $\Omega = \mathbb{R}^n$  and

$a_{jk}(x) = \delta_{jk}$  and by (2.14) we conclude

$$\operatorname{Re}(-\Delta|u| + \zeta|u| + q_1|u| - q_2|u|) \leq 0$$

(in the sense of distributions).

This easily yields

$$(2.15) \quad -\Delta|u| + \operatorname{Re}(\zeta)|u| - q_2|u| \leq 0$$

(in the sense of distributions).

Now we merely apply Theorem 2.1 to  $|u|(x)$  with  $q(x) = q_2(x)$

and  $\lambda = \operatorname{Re}(\zeta)$ . We conclude

$$|u|(x) \equiv 0,$$

and this completes the proof of the theorem.

### 3. Application to Schrödinger operators

Consider the Schrödinger operator

$$(3.1) \quad T \equiv \sum_{j=1}^n \left( \frac{\partial}{\partial x_j} - ib_j(x) \right)^2 + q_1(x) - q_2(x)$$

where  $b_j$  and  $q_i$  are real valued functions on  $\mathbb{R}^n$  such that

$$q_i \in L_{loc}^2, \quad b_j \in C^1, \quad i = 1, 2; \quad j = 1, \dots, n.$$

Denote by  $T_0$  the operator  $T$  with  $b_j = 0$ ,  $j = 1, \dots, n$ . We suppose  $T$  is defined on  $L^2$  with values in  $D'$  (the space of distributions on  $\mathbb{R}^n$ ).

\*\*This argument was suggested by H. Brezis.

Let  $\tilde{T} = T|_{C_0^\infty}$  i.e. the restriction of  $T$  to  $D = C_0^\infty$ . It is

clear that  $\tilde{T}$  is a symmetric operator and in [2] Kato has shown that under certain conditions on  $q_1$  and  $q_2$ ,  $\tilde{T}$  is essentially selfadjoint. The conditions we are interested in are

$$(3.2) \quad q_1(x) \geq -q^*(|x|), \quad \text{where } q^*(x) \geq 0 \text{ is monotone nondecreasing in } x > 0 \text{ and}$$

$$q^*(x) = O(x^2) \text{ as } x \rightarrow \infty.$$

$$(3.3) \quad q_2 \in L_{loc}^2, \text{ with}$$

$$\int_{|x| \leq r} |q_2(x)|^2 dx \leq K r^{2s} \quad 1 \leq r < \infty$$

where  $K$  and  $s$  are some constants and

$$(3.4) \quad \int_{|y| \leq r} |q_2(x-y)| |y|^{2-n} dy \rightarrow 0 \text{ as } r \rightarrow 0$$

uniformly in  $x \in \mathbb{R}^n$ ,

where  $|y|^{2-n}$  should be replaced by  $|\log|y||$  if  $n = 2$ .

We shall show that (3.3) and (3.4) can be replaced by the simpler condition

$$(3.5) \quad q_2 \in L_{loc}^2 \text{ and there exist constants } r_0 > 0, 1 > \alpha_0 > 0$$

such that

$$\frac{\Gamma(\frac{n}{2})}{2(n-2)\pi^{\frac{n}{2}}} \int_{|y| \leq r_0} |q_2(x-y)| |y|^{2-n} dy \leq \alpha_0 \text{ for all } x \in \mathbb{R}^n,$$

where  $|y|^{2-n}$  should be replaced by  $|\log|y||$  if  $n = 2$  and the leading constant appropriately modified. This leads to the following theorem.

Theorem 3.1. Let  $\tilde{T}$  be the operator defined on  $D$ . If (3.2) holds for  $q_1(x)$  and if (3.5) holds for  $q_2(x)$  then  $\tilde{T}$  is essentially selfadjoint.

Remark 3. Kato notes that if  $n \geq 5$  then (3.3) and (3.4) may be replaced by

$$(3.6) \quad q_2 \in L^{n/2}.$$

In a private communication Brezis has shown that for  $n \leq 4$  there are conditions analogous to (3.6) which may be used to replace (3.3) and (3.4). For example, if  $n = 3$  then we may replace (3.3) and (3.4) by

$$(3.7) \quad q_2 \in L^{3/2} \cap L_{loc}^2.$$

His proof depends on a modification of the argument used to prove Theorem 2.1 and the results in [2].

Proof of Theorem 3.1 for  $q^*$  = constant. We shall give an outline of our method of attack. We show that if  $\operatorname{Re}(\zeta)$  is sufficiently large then  $T + \zeta : L^2 \rightarrow D'$  is a one to one map. As in [2] this implies that  $\tilde{T}$  is essentially selfadjoint. In order to show  $T + \zeta$  is one to one we shall prove that if (3.5) holds then there exists a  $\lambda_0 \geq 0$  such that for any  $\lambda > \lambda_0$

$$(3.8) \quad (-4 + \lambda i)^{-1} (|q_2|)(x) \leq \alpha_0 + \frac{1 - \alpha_0}{2}.$$

We may then apply Theorem 2.2 to  $T + \zeta$  to achieve the desired result. We now verify the steps in the outline via a series of lemmas. (For convenience we assume  $n \geq 3$  in what follows.)

Lemma 3.2. Assume (3.5) holds for  $q_2(x)$ . Then there exists a positive constant  $K$  such that

$$(3.9) \quad \int_{|y| < r} |q_2(x-y)| dy \leq Kr^n \quad \text{for all } x \in \mathbb{R}^n, \quad r \geq r_0.$$

Proof. Set  $r_1 = \frac{r_0}{\sqrt{n}}$ , then

$$\begin{aligned} \int_{|y_1| < r_1, \dots, |y_n| < r_1} |q(x-y)| dy &\leq \int_{|y| < \sqrt{n} r_1} |q(x-y)| dy \\ &\leq C_n^{-1} (\sqrt{n} r_1)^{n-2} \int_{|y| < \sqrt{n} r_1} C_n |q(x-y)| |y|^{-n+2} dy \end{aligned}$$

where  $C_n = \frac{\Gamma(\frac{n}{2})}{2(n-2)\pi^{n/2}}$ . By (3.5) and the definition of  $r_1$

we have for some constants  $K_1$  and  $K_2$

$$\int_{|y_1| < r_1, \dots, |y_n| < r_1} |q(x-y)| dy \leq K_1 r_1^{n-2} \alpha_0 \leq K_2 r_1^n.$$

Using the fact that  $K_2$  is independent of  $x$  we see

$$\int_{|y_1| < jr_1, \dots, |y_n| < jr_1} |q(x-y)| dy \leq K_2 j^n r_1^n = K_2 (jr_1)^n.$$

Finally, by increasing  $K_2$  if necessary we conclude that there is a constant  $K$  such that

$$\int_{|y| < r} |q(x-y)| dy \leq Kr^n \quad \text{for all } x \in \mathbb{R}^n, \quad r \geq r_0.$$

This completes the proof of the lemma.

We remind the reader of the definition in section 1 of  $J_\lambda(x)$ . It is verified by change of variables that

$$(3.10) \quad J_\lambda^2(x) = \lambda^{n-2} J_1(\lambda x).$$

In [3] it is shown that there exist positive constants  $\delta$  and  $R$  such that

$$(3.11) \quad \text{if } |x| > R \text{ then } J_1(x) \leq e^{-\delta|x|}.$$

Lemma 3.3. Assume (3.5) holds for  $q_2(x)$ . Then there exists a constant  $\lambda_0 > 0$  such that

$$(3.12) \quad (-\Delta + \lambda_0 I)^{-1}(|q_2|)(x) \leq \alpha_0 + \frac{1 - \alpha_0}{2}.$$

Proof. We must show that for some  $\lambda_0 > 0$

$$(3.13) \quad \int_{\mathbb{R}^n} J_{\lambda_0}(y) |q_2(x-y)| dy \leq \alpha_0 + \frac{1 - \alpha_0}{2}.$$

Now

$$\int_{\mathbb{R}^n} J_{\lambda_0}(y) |q_2(x-y)| dy = \int_{|y| < r_0} J_{\lambda_0}(y) |q_2(x-y)| dy + \int_{|y| \geq r_0} J_{\lambda_0}(y) |q_2(x-y)| dy.$$

Using (3.5), the equality above, and since  $0 \leq J_{\lambda_0}(y) \leq C_n |y|^{-n+2}$  we conclude that it is sufficient to prove

$$(3.14) \quad \int_{|y| \geq r_0} J_{\lambda_0}(y) |q_2(x-y)| dy \leq \frac{1 - \alpha_0}{2}, \text{ for some } \lambda_0.$$

We claim

$$(3.15) \quad \int_{|y| \geq r_0} J_{\lambda_0}(y) |q_2(x-y)| dy \rightarrow 0 \text{ as } \lambda \rightarrow \infty.$$

Denote the integral above by  $I_\lambda$ ; we have

$$I_\lambda = \sum_{i=0}^{\infty} \int_{2^i r_0 \leq |y| < (2^{i+1}) r_0} |q_2(x-y)| J_{\lambda_0}(y) dy.$$

We may assume without loss of generality that  $\lambda r_0 > R$  (where  $R$  is the constant appearing in (3.11)) and thus conclude

$$I_\lambda \leq \sum_{i=0}^{\infty} \lambda^{n-2} \int_{2^i r_0 \leq |y| < (2^{i+1}) r_0} e^{-\delta\lambda|y|} |q_2(x-y)| dy.$$

Since  $|y| \geq r_0$  there are constants  $\delta' > 0$  and  $K' > 0$  such that

$$I_\lambda \leq \sum_{i=0}^{\infty} K' \lambda^{-1} \int_{2^i r_0 \leq |y| < (2^{i+1}) r_0} e^{-\delta'|y|} |q_2(x-y)| dy.$$

Further if  $\lambda > \lambda'$  then  $\delta'$  and  $K'$  depend only on  $\lambda'$ , i.e. we may assume  $\delta'$  and  $K'$  are fixed as  $\lambda \rightarrow \infty$ . We now have

$$I_\lambda \leq \sum_{i=0}^{\infty} K' \lambda^{-1} \int_{2^i r_0 \leq |y| < (2^{i+1}) r_0} e^{-\delta' 2^i r_0} |q_2(x-y)| dy.$$

Using the estimate (3.9) of Lemma 3.2 for the ball of radius  $(2^{i+1}) r_0$  we conclude

$$I_\lambda \leq \sum_{i=0}^{\infty} K' \lambda^{-1} K(2^{i+1} r_0)^n e^{-\delta' 2^i r_0} \leq K'' \lambda^{-1} \int_0^{\infty} 2^{n(\xi+1)} r_0^n e^{-\delta' 2^{\xi-1} r_0} d\xi.$$

We make the change of variables,  $\tau = 2^\xi$ , and lump constants into one term to obtain

$$I_\lambda \leq K \lambda^{-1} \int_0^{\infty} \tau^{n-1} e^{-\frac{\delta' r_0}{2} \tau} d\tau.$$

It is now obvious that the integral above is bounded independent of  $\lambda$  and therefore there exists a constant  $C > 0$  such that

$$I_\lambda \leq C\lambda^{-1}.$$

This completes the proof of (3.15), which proves Lemma 3.3.

We now complete the proof of Theorem 3.1 for  $q^* = \text{constant}$ .

Let  $\lambda$  be a constant  $\lambda > \lambda_0$  ( $\lambda_0$  determined by Lemma 3.3).

It follows from Lemma 3.3 that

$$(3.16) \quad (-\Delta + \lambda I)^{-1}(q_2^+)(x) \leq \alpha_0 + \frac{1 - \alpha_0}{2} < 1,$$

where  $q_2^+(x) = \max\{q_2(x), 0\}$ . Now consider

$$T + q^* \equiv \sum_{j=1}^n \left( \frac{\partial}{\partial x_j} - ib_j \right)^2 + (q_1(x) + q^* + q_2^-(x)) - q_2^+(x).$$

By (3.16) and assumptions on  $T$  we may apply Theorem 2.2 to

$T + q^*$  to conclude that if  $u \in L^2$ ,

$$(T + q^* + \zeta)u = 0, \quad (\text{in the sense of distributions})$$

and  $\text{Re}(\zeta) > \lambda_0$  then  $u(x) \equiv 0$ . Thus it immediately follows that if  $u \in L^2$ ,

$$(T + \zeta)u = 0, \quad (\text{in the sense of distributions})$$

and  $\text{Re}(\zeta) > \lambda_0 + q^*$  then  $u(x) \equiv 0$ . This proves  $T$  is essentially selfadjoint in the case  $q^* = \text{constant}$ .

In order to complete the proof of Theorem 3.1 we shall have to prove a result analogous to Proposition 4, pg. 141 of [2]. We introduce the function space  $H_b^1$ .  $H_b^1$  is defined as the completion of  $C_0^\infty$  in the norm

$$\|\phi\|_{H_b^1}^2 \equiv \sum_{j=1}^n \left\| \left( \frac{\partial \phi}{\partial x_j} - ib_j \phi \right) \right\|_{L^2}^2 + \|\phi\|_{L^2}^2.$$

As noted in [2]  $H_b^1$  is a Hilbert space

$$H_b^1 \subset L^2 \subset H_b^{-1} \quad \text{and} \quad \|\phi\|_{H_b^1} \geq \|\phi\|_{H_b^{-1}}.$$

It is also clear that

$$(T\phi, \phi) = \|\phi\|_{H_b^1}^2 - \|\phi\|_{L^2}^2 + ((q_1 - q_2)\phi, \phi).$$

Lemma 3.4. Let  $u \in L^2$ ,  $Tu \in H_b^{-1}$  and  $q^* = \text{constant}$ . Then  $u \in H_b^1$  and there are constants  $C_1$  and  $C_2$  such that

$$(3.17) \quad \|u\|_{H_b^1} \leq C_1 \|Tu\|_{H_b^{-1}} + C_2 (1 + (q^*)^{\frac{1}{2}}) \|u\|_{L^2}.$$

Proof. For any  $\phi \in C_0^\infty$  we have

$$((T + \lambda)\phi, \phi) = \|\phi\|_{H_b^1}^2 + (\lambda - 1)\|\phi\|_{L^2}^2 + ((q_1 - q_2)\phi, \phi).$$

We take  $\lambda = \lambda_0 + q^* + 1$  where  $\lambda_0$  is the constant from Lemma 3.3. Thus

$$(3.18) \quad ((T + \lambda)\phi, \phi) \geq \|\phi\|_{H_b^1}^2 + \lambda_0 \|\phi\|_{L^2}^2 - (q_2\phi, \phi).$$

By Lemma 3.3 and Theorem 1.1 we have

$$\left(1 - \frac{\alpha_0}{2}\right) \|\phi\|_{H_b^1}^2 + \lambda_0 \|\phi\|_{L^2}^2 - (|q_2|\phi, \phi) \geq 0,$$

and using this in (3.18) gives us (since  $\|\cdot\|_{H_b^1} \geq \|\cdot\|_{H_b^{-1}}$ )

$$((T + \lambda)\phi, \phi) \geq \left( \frac{1 - \alpha}{2} \right) \|\phi\|_{H_b^1}^2,$$

or

$$(T\phi, \phi) + \lambda \|\phi\|_{L^2}^2 \geq \left( \frac{1 - \alpha_0}{2} \right) \|\phi\|_{H_b^1}^2,$$

If  $\phi_i + w \in H_b^1$  then  $T\phi_i + Tw \in H_b^{-1}$  and so

$$(Tw, w) + \lambda \|w\|_{L^2}^2 \geq \left( \frac{1 - \alpha_0}{2} \right) \|w\|_{H_b^1}^2.$$

Consequently for any  $\epsilon > 0$  there is  $K_\epsilon$  such that

$$K_\epsilon \|Tw\|_{H_b^{-1}}^2 + \epsilon \|w\|_{H_b^1}^2 + (\lambda_0 + q^*) \|w\|_{L^2}^2 \geq \left( \frac{1 - \alpha_0}{2} \right) \|w\|_{H_b^1}^2$$

and therefore there are constants  $C_1$  and  $C_2$  such that

$$(3.19) \quad C_1 \|Tw\|_{H_b^{-1}}^2 + C_2 \left( 1 + (q^*)^{\frac{1}{2}} \right) \|w\|_{L^2}^2 \geq \|w\|_{H_b^1}^2.$$

for any  $w \in H_b^1$ .

It only remains to show that  $u \in H_b^1$ . This follows for exactly the same reason as in [2]; this completes the proof of the lemma.

Proof of Theorem 3.1. Using the result for the case  $q^* = \text{constant}$  and Lemma 3.4 we may now proceed exactly as in [2]. This completes the proof of the theorem.

#### Acknowledgements

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024 and the National Science Foundation under Grant No. MCS75-17385 A01.

The author wishes to thank H. Brezis for suggesting this problem and subsequent discussions of the material presented here.

#### References

- [1] A. Friedman, Partial Differential Equations, Holt, Rinehart and Winston, New York, N.Y. 1969.
- [2] T. Kato, Schrödinger operators with singular potentials, Israel J. Math. 13 (1972), 135-148.
- [3] E. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton Univ. Press, Princeton, N.J. 1970.
- [4] K. Yosida, Functional Analysis, Springer-Verlag, Berlin, W. Germany 1971.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER <b>(14) MRC-TSR-1827</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER <b>(9) Technical</b>	4. TITLE (and Subtitle) UNIQUENESS OF SOLUTIONS TO $-\Delta u - qu = \sigma$ . <b>↑ Delta</b>
5. AUTHOR(s) <b>(10) Robert Jensen</b>	6. PERFORMING ORG. REPORT NUMBER	7. CONTRACT OR GRANT NUMBER(s) <b>(15) DAAG29-75-C-0024, NSF-MCS75-17385-A01</b>	8. TYPE OF REPORT & PERIOD COVERED <b>Summary Report, no specific reporting period</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit # 1 (Applied Analysis)	11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.	12. REPORT DATE <b>(11) February 1978 (12) 16 p.</b>
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	14. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>	15. NUMBER OF PAGES <b>24</b>	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES U. S. Army Research Office National Science Foundation P. O. Box 12211 Washington, D. C. 20550 Research Triangle Park North Carolina 27709			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bessel potential Distribution Measure Schrödinger operator			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We give <sup>are given</sup> conditions for the function $q(x)$ which guarantee uniqueness of solutions to $-\Delta u(x) + \lambda u(x) - q(x)u(x) = f$ , where $\lambda_0 \geq 0$ is a constant. <sup>lambda</sup> <sub>sub phi</sub> <sup>sub phi</sup> We consider <sup>are considered</sup> separately the cases for solutions in $H_0^{(1)}$ and $L_0^{(2)}$ . In fact, we <sup>separately.</sup> even prove existence of $H_0^{(1)}$ solutions under the additional assumptions $q(x) \geq 0$ and $f \in L^2$ . <sup>These</sup> We apply <sup>are applied</sup> our results to the problem of essential $\rightarrow$ next page <sup>are given</sup> <sup>are applied</sup>			

> or =

can element of  $L^2$

20. ABSTRACT - Cont'd.

selfadjointness for a class of Schrodinger operators and ~~we are able to~~  
~~prove~~ an interesting generalization of a result due to Kato <sup>is proven.</sup>

Gaveau <sup>has</sup> proved a result similar to <sup>this</sup> ~~ours~~ on  $L^2$  uniqueness. His  
proof, however, depends <sup>ed</sup> on a probabalistic argument, while in contrast ~~we~~ <sup>there is</sup>

~~no~~ ~~do not use any~~ probability theory <sup>used these</sup> in ~~our~~ proofs.