

AD-A054 549

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER  
COMPLETELY INTEGRABLE SYSTEMS AND SYMPLECTIC ACTIONS. (U)  
FEB 78 M ADLER

F/G 12/1

DAA629-75-C-0024

NL


UNCLASSIFIED

MRC-TSR-1830

1 of 1  
AD  
A054549



END  
DATE  
FILMED  
7 -78  
DDC



FOR FURTHER TRAN

13 SC

AD A 054549

MRC Technical Summary Report #1830

COMPLETELY INTEGRABLE SYSTEMS AND SYMPLECTIC ACTIONS

M. Adler

Mathematics Research Center  
University of Wisconsin-Madison  
610 Walnut Street  
Madison, Wisconsin 53706

February 1978 •

(Received December 5, 1977)

DDC FILE COPY

DDC  
RECEIVED  
JUN 2 1978  
D

Approved for public release  
Distribution unlimited

Sponsored by  
U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

National Science Foundation  
Washington, D. C. 20550

See 1473

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

COMPLETELY INTEGRABLE SYSTEMS AND SYMPLECTIC ACTIONS

M. Adler

Technical Summary Report #1830  
February 1978

ABSTRACT

We study results on a class of completely integrable systems, for instance with Hamiltonian

$$H(x, y) = \frac{1}{2} \sum_{i=1}^n y_i^2 + \sum_{i < j} (x_i - x_j)^{-2} + \alpha \sum_{i=1}^n x_i^2,$$

using quotient manifolds induced by symplectic group actions, which enables us to integrate the systems and understand their complete integrability. In addition, we give a natural interpretation for the scattering maps associated with these systems.

AMS (MOS) Subject Classification: 34C35

Key Words: Hamiltonian systems, Symplectic group actions, Quotient manifolds, Completely integrable systems, Canonical transformations, Scattering maps

Work Unit Number 1 (Applied Analysis)

---

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024 and the National Science Foundation under Grant No. MCS75-17385 A01.

SIGNIFICANCE AND EXPLANATION

We study mechanical problems with high degrees of symmetry, or equivalently many constants of motion. An example of such a system would be  $n$  decoupled oscillators, since such a system would have the associated  $n$  energies of the oscillators as constants of the motion.

In mechanics, the process of studying systems by ignoring symmetries, or constants of the motion, for instance studying a system in its center of mass coordinates, is well-known. Using a modern abstraction of this old and valuable idea, we study certain systems of interest in mathematical physics, which have many symmetries. We are able to completely solve these systems using the above idea.

ACCESSION for	
RTIS	White Section <input checked="" type="checkbox"/>
ODS	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

---

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

COMPLETELY INTEGRABLE SYSTEMS AND SYMPLECTIC ACTIONS

M. Adler

(1) Introduction - In this short note we explain the results of "Some Finite Dimensional Integrable Systems and Their Scattering Behavior", [1], by the author, and of [5,6,7], in terms of the abstract machinery set up in the paper of D. Kazhdan, B. Kostant, and S. Sternberg, entitled "Hamiltonian Group Actions and Dynamical Systems of Calogero Type", [2], which explains systems first discovered by F. Calogero, C. Marchioro, [3], and first discussed by J. Moser [4].

Briefly, the systems to be discussed have the property that their equations of motion can be expressed as matrix differential equations which can be easily integrated, and moreover, the integration process is seen to occur naturally in a space of much higher dimensionality than the systems in question. The systems to be studied are thus interpreted as quotient systems, of the much larger systems, where the quotienting out process is performed by a symplectic action of the unitary group.

The process of quotienting out in mechanics, such as using center of mass coordinates, i.e. ignoring the position of the center of mass, is indeed a common practice. We point out that usually quotienting out, or ignoring certain data, is a way of ignoring the symmetries, or integrals of the system, so as to arrive at some basic equations to study. Here the quotienting out does not really involve the integrals, but enables us to pass to the ultimate system to be studied. The integrals are in fact generated in a much more trivial way, through the use of natural Lagrangian submanifolds and simple canonical maps, which of course makes use of the quotient structure. In addition, the so-called scattering maps of these systems have a natural interpretation in this context.

In the first section we merely summarize the abstract machinery of [2] of use in the discussion, referring the reader to [2] and the paper of J. Marsden and A. Weinstein [8], for a fuller discussion. We then discuss the results of [1], which entails referring to [1] frequently.

(2) The Symplectic Structures - We summarize and briefly discuss the necessary abstract machinery needed to discuss [1]. Let  $(M, \omega, G)$  be a triple, with  $M$  an (exact) symplectic manifold with nondegenerate closed two-form  $\omega = d\tau$ , and  $G$  a Lie group, with elements  $g$ , which acts on  $M$  with a symplectic action. If  $\mathfrak{L}$  is the Lie algebra of  $G$ , with elements denoted by  $\dot{g}$ , then the action of  $G$  associates with each  $\dot{g}$  the Hamiltonian vector-field  $\hat{g}$ , and the Hamiltonian function  $f_{\dot{g}}(\cdot) = -\tau(\hat{g})(\cdot)$ , yields a Lie homomorphism, i.e.,

$$(2.1) \quad \{f_{\dot{g}_1}, f_{\dot{g}_2}\} = f_{[\dot{g}_1, \dot{g}_2]}, \quad [ , ] \text{ the bracket in } \mathfrak{L},$$

where  $\{ , \}$  is just the usual Poisson bracket, i.e. if

$$(2.2) \quad X_f \lrcorner \omega = df, \text{ then } X_{f_2}(f_1) = \omega(X_{f_1}, X_{f_2}) \equiv \{f_1, f_2\}.$$

We define the moment map of Souriac,

$$(2.3) \quad \Phi : M \rightarrow \mathfrak{L}^*, \text{ by } \Phi(m)(\dot{g}) = f_{\dot{g}}(m),$$

with  $\mathfrak{L}^*$  the dual of  $\mathfrak{L}$ . The group  $G$  acts on itself by conjugation, hence on  $\mathfrak{L}$  by the linearization of conjugation,  $\text{Adj}$ , and on  $\mathfrak{L}^*$  by  $(\text{Adj})^*$ , and it's easy to see that (2.1) is just the infinitesimal, and hence equivalent version of the relation of equivariance,

$$(2.4) \quad \Phi \circ g = (\text{Adj } g^{-1})^* \circ \Phi.$$

We then form the orbit of  $\alpha \in \mathfrak{L}^*$  under  $(\text{Adj})^*$ ,  $\theta_\alpha$ , and assume  $V = \Phi^{-1}(\theta_\alpha)$  is a manifold. Then in fact it is a coisotropic manifold, i.e.  $(TV_x)^{\perp} \subset (TV_x)$ , for all  $x \in V$ , with  $\perp$  denoting perpendicularity with respect to  $\omega$ , and we can thus form  $S = V / (\text{leaves of the foliation induced by } (TV)^{\perp})$ , taking  $S$  to be connected and assuming it to be a manifold. Then as a direct consequence of (2.4), it's not hard to see (although it's not shown in [2,8]), that  $S$  is a covering space of  $\theta'_\alpha$  as follows:

$$(2.5) \quad S \cong \theta'_\alpha \times \theta_\alpha, \quad \theta'_\alpha = \frac{\Phi^{-1}(\alpha)}{G_\alpha},$$

where  $/G_\alpha$  means we identify elements  $x, y \in M$  if they lie on the same  $G_\alpha$  orbit,

with  $G_\alpha$  the isotropy group of  $\alpha$ , i.e. the connected subgroup of  $G$  which fixes  $\alpha$  by its action on  $M$ . Incidentally, this shows  $S$  is a manifold precisely if  $\theta'_\alpha$  is one. By the coisotropy of  $V$ , and the transitivity of  $G$  on the fibers  $\theta'_\alpha$ ,  $\omega$  induces a symplectic structure on  $\theta'_\alpha$ ,  $\omega_\alpha$ , (i.e. we shall tacitly assume  $\omega_\alpha$  is nondegenerate), where  $\{ , \} \mapsto \{ , \}_\alpha$  is a homomorphism. The structure  $(\theta'_\alpha, \omega_\alpha), G$  shall be our arena of activity.

We note that by (2.1), (2.2), functions in  $M$  which are  $G$  invariant, induce Hamiltonian flows on  $M$  with pointwise fix the image of  $M$  under  $\phi$ , and hence they induce Hamiltonian flows on  $\theta'_\alpha$ . In addition, such functions, if they are in involution with respect to  $\{ , \}$  on  $M$ , are via the homomorphism  $\omega \rightarrow \omega_\alpha$ , automatically in involution in  $\theta'_\alpha$ , thought of, by their  $G$  invariance, as functions on  $\theta'_\alpha$ . This ends our discussion of quotient structures.

In preparation we discuss the  $M$ 's which shall come up in the examples.

Let  $F$  be the linear manifold of  $n \times n$  matrices with complex coefficients, and  $T^*F = \phi$  be the cotangent bundle of  $F$ , where we shall identify  $T^*F \sim F \times F$  via the bilinear form  $\langle X, Y \rangle = \text{tr}XY$ . Then the complex symplectic 2-form  $\omega$ , naturally associated with  $T^*F$  is

$$\omega = \sum dx_{ij} \wedge dy_{ji} = \langle dx, dy \rangle,$$

or alternately, we write Hamilton's equations, with Hamiltonian  $H = H(X, Y)$ , as

$$(2.6) \quad \dot{X} = H_Y, \quad \dot{Y} = -H_X,$$

where  $[H_X]_{ij} = \frac{\partial H}{\partial x_{ji}}$ , etc. If we restrict  $\omega$  to

$$(2.7) \quad M_1 = T^*\mathcal{L} = \{(X, Y) \mid X = X^*, Y = Y^*\},$$

where  $*$  denotes taking the Hermitian adjoint, i.e.  $\mathcal{L}$  is just the self-adjoint matrices, (which we shall identify with the Lie algebra of the unitary group  $G = U(n, \mathbb{C})$ ),  $\omega$  yields a real symplectic structure, with Hamilton's equations remaining as given in (2.6), where it is understood that  $H$  is real.

It is interesting to map  $\phi \mapsto \phi$  via

$$(2.8) \quad \tau : \begin{pmatrix} X \\ Y \end{pmatrix} \mapsto \begin{pmatrix} X \\ XY \end{pmatrix} = \begin{pmatrix} Z_1 \\ Z_2 \end{pmatrix},$$

then as one computes

$$\left\{ \begin{array}{l} H_X = H_{Z_1} + YH_{Z_2}, \quad H_Y = H_{Z_2} X \\ \dot{Z}_1 = \dot{X} = H_Y, \quad \dot{Z}_2 = (XY)' = H_Y Y - XH_X, \end{array} \right\}$$

and so we may write Hamilton's equations in  $(Z_1, Z_2)$  coordinates as

$$(2.9) \quad \dot{Z}_1 = H_{Z_2} Z_1, \quad \dot{Z}_2 = [H_{Z_2}, Z_2] - Z_1 H_{Z_1}.$$

Let us now restrict  $\tau$  to  $T^*U(n)$ , i.e. we identify

$$(2.10) \quad M_2 = T^*U(n, \mathbb{C}) \cong \{(Z_1, Z_2) \mid Z_1 Z_1^* = I, Z_2^* = Z_2\},$$

and restrict  $\tau \rightarrow \tau|_{\tau^{-1}(M_2)} = \bar{\tau}$ . Note  $\bar{\tau}$  is invertible, and of course we may just

as well identify  $\tau^{-1}(M_2)$  with  $T^*U(n)$ . In that case, since by the pairing  $\langle \cdot, \cdot \rangle$ ,  $X, Y$  are the usual dual coordination of  $T^*U(n)$ , (2.9) restricted to  $M_2$  precisely yields Hamilton's equations for the natural symplectic structure of  $T^*U(n)$ ; that is, up to the factor  $i$  which must be put in due to our identification of the Lie algebra of  $U(n)$  with self-adjoint matrices, or equivalently we may think of time as being purely imaginary in (2.9). We omit the necessary, but easy verification that (2.9) restricted to  $M_2$  automatically preserves  $M_2$ , which is sufficient to insure the restricted  $\omega$  is symplectic.

(3) Equations of Motion for the Simplest System - We now apply the discussion in Section 2. We shall let  $(M, \omega, G)$  of Section 1 be  $(M_1, (dx, dy), U(n))$  of Section 1. Since  $M_1 = T^*L = L \times L$ ,  $L$  the Lie algebra of  $U(n)$ ,  $U(n)$  acts naturally on  $L$  via  $\text{Adj}$ , i.e. by conjugation which naturally extends to a Hamiltonian action on  $T^*L$  by (if  $U \in U(n, \mathbb{C})$ )

$$U : (X, Y) \mapsto (UXU^{-1}, UYU^{-1}),$$

of which the linear version is

$$\dot{U} : (X, Y) \mapsto ([\dot{U}, X], [\dot{U}, Y]) \in \mathfrak{TM}_{1(X, Y)},$$

hence, by (2.3), and the above,

$$\phi(X, Y)(\dot{U}) = f_{\dot{U}}(X, Y) = \langle [\dot{U}, X], Y \rangle = \langle [X, Y], \dot{U} \rangle,$$

and so by the identification of  $L$  with  $L^*$  through  $\langle \cdot, \cdot \rangle$ ,

$$(3.1) \quad \Delta_{\alpha} \equiv \phi^{-1}(\alpha) = \{(X, Y) \mid [X, Y] = \alpha\},$$

and we shall once and for all pick  $\alpha$  such that

$$[\alpha]_{jk} = i(1 - \delta_{jk}), \quad \text{i.e. } \alpha = i(v^* \otimes v),$$

with  $v \in \mathbb{R}^n$ ,  $v = (1, 1, 1, \dots, 1)^T$ . Note that the isotropy subgroup  $G_{\alpha} = \{U \mid U\alpha U^{-1} = \alpha\}$ , and we shall define the reduced subgroup

$$(3.2) \quad G_0 = \{U \mid U(v) = v\}, \quad \text{with Lie algebra } \mathfrak{f}_0 = \{B \mid B(v) = 0\}.$$

In [2] it is shown by a simple linear algebra argument that if  $[X, Y] = \alpha$ , we can always find a unique  $U \in G_0$  such that

$$(3.3) \quad \left\{ \begin{array}{l} UXU^{-1} = \text{diag}(x_1, x_2, \dots, x_n) \equiv \bar{x}, \quad x_i < x_{i+1}, \quad \text{all } i \\ [UYU^{-1}]_{jk} = \delta_{jk} y_j + i(1 - \delta_{jk})(x_j - x_k)^{-1} \equiv \bar{y} \end{array} \right\},$$

and hence  $\theta'_{\alpha} = \phi^{-1}(\alpha)/G_{\alpha}$  is effectively coordinatized by  $((x_1, x_2, \dots, x_n), (y_1, \dots, y_n)) = (x, y)$ , and moreover it is shown in [2], by a local argument, that

$$(3.4) \quad \omega \mapsto \omega_{\alpha} = \sum_{i=1}^n dx_i \wedge dy_i, \quad \text{i.e. } (x, y)$$

form a set of canonical coordinates. Hence in this case,  $\theta'_{\alpha}$ , and thus  $S$  is a manifold (see Section 2), and  $\omega_{\alpha}$  is nondegenerate, and hence symplectic.

We now wish to find functions on  $\theta'_\alpha$ , and by the discussion in Section 2, functions of the form

$$(3.5) \quad H = H(X,Y) = \text{tr } P(X,Y) ,$$

with  $P(\cdot, \cdot)$  a noncommuting polynomial in its arguments will certainly do. If we take (3.5) as a Hamiltonian function on  $M_1$ , then (2.6) yields for Hamilton's equations,

$$(3.6) \quad \dot{X} = h_1(X,Y), \quad \dot{Y} = h_2(X,Y) ,$$

with  $h_i(\cdot, \cdot)$ ,  $i = 1, 2$ , polynomials in their arguments, uniquely determined by  $P(\cdot, \cdot)$ . As mentioned,  $H = \text{tr } P(X,Y)$  automatically can be thought of as a function on  $\theta'_\alpha$ , in fact via

$$(3.7) \quad h(X,Y) = \text{Tr } P(\bar{X}, \bar{Y}), \quad (\text{see (3.11)}) ,$$

and we wish to determine the analog of (3.6) for the system on  $\theta'_\alpha$  with Hamiltonian (3.7), or to put it another way, we shall determine how (3.6) transforms in  $\theta'_\alpha$ .

So assume we are given initial data  $(X(0), Y(0))$  for Hamilton's equations with Hamiltonian  $h(x,y)$  in  $\theta'_\alpha$ , which corresponds to  $(\bar{x}(0), \bar{y}(0))$  which we may identify, and thus set equal to,  $(X(0), Y(0))$  in  $M_1$ . Under the Hamiltonian  $h(x,y)$ ,  $(\bar{x}(0), \bar{y}(0)) \mapsto (\bar{x}(t), \bar{y}(t))$ , and correspondingly under the Hamiltonian  $H(X,Y)$ ,  $(X(0), Y(0)) \mapsto (X(t), Y(t))$ . By the previous remarks, we must have

$$(3.8) \quad X(t) = U\bar{x}(t)U^{-1}, \quad Y(t) = U\bar{y}(t)U^{-1}, \quad U = U(t) ,$$

with  $U(t) \in G_0$ , (see (3.2)), uniquely defined, as the  $H(X,Y)$  flow in the big manifold  $M_1$  descends to the  $h(x,y)$  flow in the little manifold  $\theta'_\alpha$  through quotienting out via  $G_0$ . Define  $B(t) \in \mathcal{L}_0$ , (see (3.2)), by  $\dot{U} = -UB$ , and so by (3.6), (3.8)

$$\dot{X} = U\delta\bar{x}U^{-1} = h_1(X,Y) = Uh_1(\bar{x}, \bar{y})U^{-1} ,$$

where  $\delta\bar{x} = \dot{\bar{x}} - [B, \bar{x}]$ , and so we have

$$(3.9) \quad \delta\bar{x} = h_1(\bar{x}, \bar{y}), \quad \delta\bar{y} = h_2(\bar{x}, \bar{y}) ,$$

as a consequence of Hamilton's equations on  $\theta'_\alpha$ ,  $\dot{\bar{x}} = \frac{\partial}{\partial x} h(x,y)$ ,  $\dot{\bar{y}} = -\frac{\partial}{\partial y} h(x,y)$ , and thus we see how (3.6) is transformed in  $\theta'_\alpha$ . Note that from the definition of  $\delta$ , (3.9), and  $B(v) = 0$ , (see (3.2)), we can immediately compute the unexpected functional dependence,  $\delta\bar{x} = B(\bar{x}(t), \bar{y}(t))$ , since  $\bar{x}$  is a diagonal matrix.

We specialize to the case  $H = H_f = \text{tr } f(Y)$ , for which we compute, (see (3.6)),  
 $h_1(X, Y) = f'(Y)$ ,  $h_2 = 0$ , and thus conclude from (3.6), (3.9)

$$(3.10) \quad \dot{X} = f'(Y), \quad Y' = 0,$$

$$(3.11) \quad \delta \bar{x} = f'(\bar{y}), \quad \delta \bar{y} = 0.$$

Since (3.10) is immediately solvable, we have in fact solved (3.11) by the use of (3.8).

We also note that since clearly the  $H_f$ 's are in involution on  $M_1$ , being functions only of  $Y$ , that by the homomorphism  $\omega \mapsto \omega_\alpha$ , the  $h_f$ 's,  $h_f = \text{tr } f(\bar{y})$ , are in involution on  $\mathfrak{g}'_\alpha$ , and in fact are generated by  $n$  independent functions,  $h^{(j)} = \text{tr}(\bar{y})^j$ ,  $j = 1, 2, \dots, n$ . Thus  $h = \text{tr}(\frac{1}{2} \bar{y}^2)$  gives rise to a completely integrable Hamiltonian system.

(4) Scattering Maps - Upon inspection, one observes that the map

$$(4.1) \quad N : (X, Y) \mapsto (Y, X), \quad \eta|_{\Delta_\alpha} : \Delta_\alpha \mapsto \Delta_{-\alpha}, \quad (\text{see (3.1)}) ,$$

is a canonical map with multiplier  $-1$  in  $M_1$ , and hence so is its projection  $\eta_p$  in  $\theta'_\alpha$ , since  $\omega \mapsto \omega_\alpha$  is a homomorphism. This map,  $\eta_p$ , is precisely the scattering map for system (3.11), with  $f(s) = \frac{1}{2} s^2$ , which is discussed in [1], Theorem 6, and was observed by J. Moser. More precisely, in the above case, (3.11) with  $f(s) = \frac{1}{2} s^2$ , one shows the time evolution of the system is given by

$$(x, y) = (qt + p + O(t^{-1}), p + O(t^{-2})), \quad t \rightarrow +\infty ,$$

with

$$\eta_p : (x(0), y(0)) \mapsto (q, p) .$$

Similarly one defines the  $\frac{\pi}{4}$  rotation map

$$(4.2) \quad \hat{\eta} : (X, Y) \mapsto 2^{-\frac{1}{2}} (X + Y, X - Y), \quad \hat{\eta}|_{\Delta_\alpha} : \Delta_\alpha \mapsto \Delta_\alpha ,$$

which is easily seen to be canonical with multiplier  $-1$  on  $M_1$ , (here one uses that  $X, Y$  are Hermitian), and the corresponding canonical projection,  $\hat{\eta}_p$  on  $\theta'_\alpha$ . In fact, in (3.7), if  $h(x, y) = \frac{1}{2} \text{tr}(y^{-2} - x^{-2})$ , then the time evolution of the system given by (3.9) is given by

$$(x(t), y(t)) = 2^{-\frac{1}{2}} \cdot (q^+ e^{+it} + p^+ e^{-it}, q^+ e^{+it} - p^+ e^{-it}) + O(e^{-2|t|}), \quad t \rightarrow \pm\infty$$

with

$$\eta_p(x(0), y(x)) = (q, p) .$$

This is shown in [1], Theorem 4. Moreover, as one easily checks

$$\eta = \hat{\eta} \circ \rho \circ \hat{\eta}^{-1}, \quad \rho : (X, Y) = (X, -Y) ,$$

hence

$$\eta_p = \hat{\eta}_p \circ \rho_p \circ \hat{\eta}_p^{-1}, \quad \text{and by (3.3), } \rho_p(x, y) = (x, -y) .$$

Note that we have shown, by the time reversibility of system (3.9), that

$$\eta_p : (q^-, p^-) \mapsto (q^+, p^+) . \quad \text{These latter statements are shown in [1], Theorems 5, 6.}$$

(5) Two Equivalent Systems - We now investigate the systems on  $\theta'_\alpha$  with Hamiltonians respectively

$$(5.1) \quad a) \quad h_1(x,y) = \text{tr } f(\bar{x}, \bar{y}), \quad b) \quad h_2(x,y) = \text{tr } f\left(\frac{1}{2} [\bar{x} + \bar{y}] \cdot [\bar{x} - \bar{y}]\right).$$

By the canonical map  $\hat{\eta}_p$  of Section 4, it is only necessary to investigate case a), and then via  $\hat{\eta}_p$ , transpose the results to case b). Also, via the transformation formalism (3.8), and (3.6)  $\mapsto$  (3.9), it is only necessary to study the equations on the full manifold  $M_1$ , rather than the quotient manifold  $\theta'_\alpha$ . The formalism says substitute  $\delta$  for  $\frac{d}{dt}$ ,  $(\bar{x}, \bar{y})$  for  $(X, Y)$ , which enables one to compute the generator  $B$ , and solve the equations on the big manifold,  $M_1$ , and via (3.8), pass to the solution on the quotient manifold  $\theta'_\alpha$ .

We thus need only study the system on  $M_1$  with Hamiltonian

$$(5.2) \quad H_1 = H_f(X, Y) = \text{tr } f(XY),$$

and we first show the  $H_f$ 's are in involution. Specifically assume

$$H^{(1)} = \text{tr } f_1(XY), \quad H_2 = \text{tr } f_2(XY),$$

then since, taking increments,

$$\delta H^{(1)} = \text{tr}(f'_1(XY) \cdot [\delta X \cdot Y + X \cdot \delta Y]) + O^{(2)},$$

where  $O^{(2)}$  means terms of at least second order, we have

$$(5.3) \quad H_X^{(1)} = Y \cdot f'_1(XY), \quad H_Y^{(1)} = f'_1(XY) \cdot X,$$

etc. for  $H^{(2)}$ . By (2.6), (2.2), the Poisson bracket  $\{ , \}$  of  $H^{(1)}, H^{(2)}$  is given by, (where  $(A, B) = \text{tr}(AB)$ ),  $\{H^{(1)}, H^{(2)}\} = (H_X^{(1)}, H_Y^{(2)}) - (H_Y^{(1)}, H_X^{(2)})$ , and thus substituting in (5.3), we find

$$\{H^{(1)}, H^{(2)}\} = (Y f'_1, f'_2 X) - (f'_1 X, Y f'_2) = \text{tr}(f'_1 f'_2 XY) - \text{tr}(f'_1 f'_2 XY) = 0,$$

where we have used  $[f'_2(XY), XY] = 0$ , and so  $\{H^{(1)}, H^{(2)}\} = 0$ .

We have thus through  $\hat{\eta}$ , shown the  $H_2$ 's,  $H_2 = H_f = \text{tr } f\left(\frac{1}{2} [X + Y] \cdot [X - Y]\right)$ , are in involution, and thus by the homomorphism  $\omega \mapsto \omega_\alpha$ , so are the  $\{\text{tr } f(\bar{x}\bar{y})\}$ 's,  $\{\text{tr } f\left(\frac{1}{2} [\bar{x} + \bar{y}] \cdot [\bar{x} - \bar{y}]\right)\}$ 's respectively, and so

$\text{tr}(\frac{1}{2} \bar{x}\bar{y})$ ,  $\text{tr}(\frac{1}{2} [\bar{x} + \bar{y}] \cdot [\bar{x} - \bar{y}])$  respectively give rise to completely integrable Hamiltonian systems on  $\theta'_\alpha$ , as observed and proven in [1]. This is thus the second, and more pleasant proof of that fact.

Now by (2.6), (5.3), Hamiltons equations for the system of (5.2) are

$$(5.4) \quad \dot{X} = f'(XY) \cdot X, \quad \dot{Y} = -Yf'(XY),$$

and since  $(XY)' = 0$  as a consequence of  $[f'(XY), XY] = 0$ , we immediately integrate

(5.4) to obtain

$$(5.5) \quad X = e^{\int f'(X_0 Y_0) dt} \cdot X_0, \quad Y = Y_0 \cdot e^{-\int f'(X_0 Y_0) dt},$$

where the subscript 0 denotes evaluation at  $t = 0$ . Thus by (3.9), the corresponding

equations on  $\theta'_\alpha$  for the systems of (5.1) are

$$(5.6) \quad \delta \bar{x} = f'(\bar{x}\bar{y}) \cdot \bar{x}, \quad \delta \bar{y} = -\bar{y}f'(\bar{x}\bar{y}),$$

while (3.8) implies the time evolution of (5.6) is given by

$$(5.7) \quad \text{diag}(x_1, x_2, \dots, x_n)(t) = U^{-1} e^{\int f'(X_0 Y_0) dt} \cdot X_0 U, \text{ etc. .}$$

The canonical map  $\hat{\eta}$  (5.4), implies the corresponding equations and time evolution for the system with Hamiltonian  $H_2 = \text{tr} f(\frac{1}{2} [X + Y] \cdot [X - Y])$  are

$$(5.8) \quad \left. \begin{aligned} (X + Y)' &= -f'(\frac{1}{2} [X + Y] \cdot [X - Y]) \cdot [X + Y], \\ (X - Y)' &= (X - Y) \cdot f'(\frac{1}{2} [X + Y] \cdot [X - Y]), \\ \text{with } \quad & \{\frac{1}{2} [X + Y] \cdot [X - Y]\}' = 0, \text{ and} \\ (X + Y) &= [\exp - \{t f'(\frac{1}{2} [X_0 + Y_0] \cdot [X_0 - Y_0])\}] \cdot [X_0 + Y_0], \\ (X - Y) &= [X_0 - Y_0] \cdot [\exp \{t f' [X_0 + Y_0] \cdot [X_0 - Y_0]\}], \end{aligned} \right\}$$

where the changing of minus signs,  $t \rightarrow -t$ , comes about because  $\hat{\eta}$  is canonical with multiplier  $-1$ . We now use the same procedure, (3.9), as above to transpose (5.8) to  $\theta'_\alpha$ , i.e. system (b) of (5.1), thus concluding

$$(5.9) \quad \partial \delta \bar{x} = [\bar{x}, f'] - [f', \bar{y}]_+, \quad 2 \delta \bar{y} = [\bar{y}, f'] - [f', \bar{x}]_+,$$

where  $[A, B]_+ = AB + BA$ ,  $f' = f'(\frac{1}{2} [\bar{x} + \bar{y}] \cdot [\bar{x} - \bar{y}])$ . Note the simplicity of (5.8), (5.9),

when  $f(s) = s$ . We could equally well study the systems gotten by 'stretching'  
 $X \mapsto \mu \cdot X$ , in (5.1), (5.2), which tend to have compact behavior and thus give rise to  
periodic solutions for  $\mu$  purely imaginary, see [1], in fact for  $f(s) = s$ ,  $\mu = \sqrt{-1}$ ,  
all solutions are periodic with one and the same nonprimitive period.

Note that our studying the case,  $H = \text{tr } f(XY)$ ,  $h = \text{tr } f(\overline{xy})$ , makes it unnecessary  
to study the (Sutherland) case where our manifold is

$$T^*U(n, \mathbb{C}) \cong \{(U, R) \mid UU^* = I, R = R^*\},$$

and our Hamiltonian is  $H(U, R) = H_f = \text{tr } f(R)$ , with Hamilton's equations given by (2.9),  
 $(U, R) = (z_1, z_2)$ , for after the change  $t \rightarrow it$ , we would get the same formal results as  
(5.4)-(5.7), including the involution statement, via the map  $\tau$ , (2.8), where we  
identify  $(X, XY) = (z_1, z_2)$  with  $(U, R)$ . We note that the condition  $(X, Y) \in \Delta_\alpha$ ,  
namely  $[X, Y] = \alpha$  is transformed into  $[U, U^{-1}R] = \alpha$ , i.e.  $R - U^{-1}RU = \alpha$ .

(6) Yet Another System on  $\theta_\alpha$  - We now discuss the system of Section 6 of [1].

One may either regard the Hamiltonian of this system on  $\theta'_\alpha$  as

$$(6.1) \quad h_1 = \text{tr} \left( \frac{1}{2} (\bar{xy})^2 + \bar{x} \right) ,$$

or

$$(6.2) \quad h_2 = \text{tr} \left( \frac{1}{2} \bar{xy}^2 \bar{x} + \bar{x} \right) ,$$

as  $h_1, h_2$  differ by a constant. Of course in the full manifold the corresponding Hamiltonians,

$$(6.3) \quad H_1 = h \left[ \frac{1}{2} (XY)^2 + X \right] ,$$

$$(6.4) \quad H_2 = \text{tr} \left[ \frac{1}{2} (XY^2X) + X \right] ,$$

are far from identical. Although it is shown in [1] that (6.1,2) is a completely integrable system, we shall not show (6.3,4) are completely integrable systems, in fact we have not been able to do this.

We shall study both (6.3,4), and then relate them in case the associated differential equations on  $\theta'_\alpha$  have the same initial data. Since the calculations are so similar to those of Section 5, we just give the results. For simplicity we set  $XY = Z$ . Then with the Hamiltonian of (6.3) we calculate, from (2.6),

$$\dot{X} = ZX, \quad \dot{Z} = -X ,$$

and since  $[X,Y] = \alpha$ ,  $(YX)^\cdot = -X$ , we have  $\frac{1}{2} (XY^2X) + X = e_1$ ,  $e_1$  a constant, and thus we arrive at, again using  $[X,Y] = \alpha$ ,

$$(6.5) \quad \frac{1}{2} Z^2 - \frac{1}{2} ZX - \dot{Z} = e_1 .$$

Letting  $Z = -2a_1^{-1} \dot{a}_1$ , we find  $\ddot{a}_1 = \frac{1}{2} a_1 e_1 - \frac{1}{2} \dot{a}_1 \alpha$ , hence we have

$$(6.7) \quad \left\{ \begin{array}{l} \begin{pmatrix} a_1 \\ \dot{a}_1 \end{pmatrix} = \begin{pmatrix} a_1(0) \\ \dot{a}_1(0) \end{pmatrix} \cdot \exp C_1 t , \\ C_1 = D_1(X,Y) = \begin{bmatrix} 0 & 1 \\ \frac{1}{4} (XY^2X) + \frac{1}{2} X & -\frac{\alpha}{2} \end{bmatrix} \text{ at } t = 0 . \end{array} \right.$$

For the Hamiltonian of (6.4), we find

$$\begin{aligned}\dot{Z} - \left[\frac{1}{2} YX, Z\right] &= -X \\ \dot{X} - \left[\frac{1}{2} YX, X\right] &= \frac{1}{2} (XZ + ZX),\end{aligned}$$

which motivates us to define the derivation  $\delta$ ,

$$\delta(\cdot) = \frac{d}{dt}(\cdot) - \left[\frac{1}{2} YX, \cdot\right],$$

and thus we have from the above,

$$(6.8) \quad \delta Z = -X, \quad \delta X = \frac{1}{2} (XZ + ZX).$$

Clearly if  $U(0) = I$ ,  $\dot{U} = -U\left(\frac{1}{2} YX\right)$ , we have the following rule of transformations for matrices  $A = A(t)$ : if  $\hat{A} \equiv UAU^{-1}$ , then  $\left(\frac{d}{dt}\right)^j \hat{A} = U\delta^j AU^{-1}$ . From (6.8) we conclude, using that  $\delta$  is a derivation,

$$(6.9) \quad \delta\left(\frac{1}{2} Z^2 + X\right) = 0, \quad \delta\left(\frac{1}{2} Z^2 - \delta Z\right) = 0,$$

hence by our rule of transformation,

$$\frac{1}{2} \hat{Z}^2 - \dot{\hat{Z}} = e_2, \quad e_2 \text{ a constant},$$

and thus letting  $\hat{Z} = -2a_2^{-1}\dot{a}_2$ , we find

$$\ddot{a}_2 = \frac{1}{2} a_2 e_2, \quad \text{i.e.}$$

$$(6.10) \quad \left\{ \begin{array}{l} \begin{pmatrix} a_2 \\ \dot{a}_2 \end{pmatrix} = \begin{pmatrix} a_2(0) \\ \dot{a}_2(0) \end{pmatrix} \cdot \exp C_2 t, \\ C_2 = D_2(X, Y) = \begin{bmatrix} 0 & I \\ \frac{1}{4} (XY)^2 + \frac{1}{2} X & 0 \end{bmatrix} \text{ at } t = 0. \end{array} \right\}$$

We now consider the case where the  $X(0)$  of both systems are the same in  $\theta'_\alpha$ , and moreover  $x_1(0) > 0$ . Then since both equations (6.3,4) are the same as seen in  $\theta'_\alpha$ , they both must have the same long term behavior of  $X$  as projected down into  $\theta'_\alpha$ . In [1], Section 6, it is shown that systems (6.3,4) have the following long term 'scattering' behavior:

$$(\log x, xy) = (\lambda^{\pm} t + \beta^{\pm} + O(t^{-1}), \pm \lambda + O(t^{-2})), \quad t \rightarrow \pm \infty,$$

where  $\log x = (\log x_1, \dots, \log x_n)$ ,  $xy = (x_1 y_1, \dots, x_n y_n)$ . By arguments in that same section, it's clear that the spectrum of  $C_1$ , or alternately  $C_2$ , precisely carry the data  $\lambda^{\pm}$ , and hence we must have

$$D_1(X, Y) \sim D_2(X, Y),$$

where  $\sim$  denotes spectral equivalence. By (6.7,10), (3.8), this implies

$$(6.11) \quad D_1(\bar{x}, \bar{y}) \sim D_2(\bar{x}, \bar{y}).$$

Moreover it follows from (6.5,9) that  $D_1(X, Y)$ ,  $D_2(X, Y)$  are isospectral matrices of the differential equations (6.3,4), and so in particular  $D_1(\bar{x}, \bar{y})$ ,  $D_2(\bar{x}, \bar{y})$  are for the (6.1,2) flow, and thus we arrive at

$$(6.12) \quad D_1(\bar{x}(0), \bar{y}(0)) \sim \lim_{t \rightarrow \pm \infty} D_1(\bar{x}(t), \bar{y}(t)) \sim \lim_{t \rightarrow \pm \infty} D_2(\bar{x}(t), \bar{y}(t)),$$

which yields the scattering behavior of system (6.1,2) as discussed in Corollaries 11.2, 11.3, of [1], which essentially maintains that the system scatters as if it is completely decoupled, and just constrained to maintain a fixed order on the line.

I wish to thank J. Moser, who suggested this research and encouraged it, and C. Conley, at those seminar at the University of Wisconsin at Madison, these results were first presented.

#### REFERENCES

- [1] M. Adler, Some Finite Dimensional Integrable Systems and Their Scattering Behavior  
Comm. Math. Phys. 55, 295-230 (1977).
- [2] D. Kazhdan, B. Kostant, S. Sternberg, Hamiltonian Group Actions and Dynamical  
Systems of Calogero type, to appear in C.P.A.M.
- [3] F. Calogero, C. Marchioro, Exact Solution of a One Dimensional Three-body Scattering  
Problem with Two Body and/or Three-body Inverse Square Potential, J. Math. Phys.  
15, 1425-1430 (1974).
- [4] J. Moser, Three Integrable Hamiltonian Systems Connected with Isospectral Deforma-  
tions, Advanc. Math. 16, 1-23 (1975).
- [5] M. A. Olshanetsky and A. M. Perelomov, Explicit Solutions of Some Completely  
Integrable Systems, Lettere Al Nuovo Cimento, Vol. 17, No. 3, 97-101 (1976).
- [6] M. A. Olshanetsky and A. M. Perelomov, Completely Integrable Classical Systems  
Connected with Semisimple Lie Algebras, II. Moscow Institute of Theoretical and  
Experimental Physics, ITEP-27 (1976).
- [7] K. Sawada and T. Kotera, Integrability and a Solution for One Dimensional N-Particle  
Systems with Inversely Quadratic Pair Potentials, preprint, Tokyo University of  
Ed. TUETP-75-10.
- [8] S. Marsden and A. Weinstein, Reduction of Symplectic Manifolds with Symmetry,  
Reports on Math. Physics, No. 1, Vol. 5, (1974).

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER <b>14</b> MRC-TSR-1830 <sup>v</sup>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER <b>9</b> Technical	
4. TITLE (and Subtitle) COMPLETELY INTEGRABLE SYSTEMS AND SYMPLECTIC ACTIONS.		5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period	
7. AUTHOR(s) M./Adler <b>10</b>		6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) <b>15</b> DAAG29-75-C-0024 <sup>v</sup> NSF-MCS75-17385	
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit # 1 - Applied Analysis	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE <b>11</b> February 1978	
		13. NUMBER OF PAGES <b>12</b> 19p. 15	
		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hamiltonian systems Symplectic group actions Quotient manifolds Completely integrable systems Canonical transformations Scattering maps			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We study results on a class of completely integrable systems, for instance with Hamiltonian $H(x, y) = \frac{1}{2} \sum_{i=1}^n y_i^2 + \sum_{i < j} (x_i - x_j)^{-2} + \alpha \sum_{i=1}^n x_i^2,$ using quotient manifolds induced by symplectic group actions, which enables us to integrate the systems and understand their complete integrability. In addition, we give a natural interpretation for the scattering maps associated with these systems.			