

AD-A056 344

MINNESOTA UNIV MINNEAPOLIS DEPT OF THEORETICAL STATISTICS F/G 12/1
A NOTE ON ANCILLARITY AND INDEPENDENCE VIA MODEL-PRESERVING TRA--ETC(U)
MAY 78 S D GUPTA DAAG29-76-G-0038

UNCLASSIFIED

UMN/DTS/TR-319


ARO-13149.6-M

NL

1 OF 1
ADA
056344



END
DATE
FILMED
8-78
DDC



NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

ARO 13149.6-M

(12)

LEVEL II

AD A 056344

AU NO.
DDC FILE COPY

UNIVERSITY OF MINNESOTA

SCHOOL OF
STATISTICS



DDC
RECEIVED
JUL 18 1978
JC B

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

78 07 00 088

AD A056344

12

LEVEL II

A NOTE ON
ANCILLARITY AND INDEPENDENCE
VIA
MODEL-PRESERVING TRANSFORMATIONS.

by

Somesh Das/Gupta*

Technical Report No. 319

May 1978

AD No. _____
DDC FILE COPY

UMN/DTS/IR-319

May 78

12 p.

AR5

13149.6-M

DDC
RECEIVED
JUL 18 1978
B

AMS Classification: 62B05, 62H10

Key Words: Sufficiency, Ancillarity, model-preserving transformations, compact group, invariant probability measure.

*Partially supported by a grant from the Mathematics Division, U.S. Army Research Office, Durham, N.C.; Grant No. DAAG29-76-G-0038

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

78 07 00 028

410 022

alt

Summary

Let G be a compact group of transformations which preserves a family $\mathcal{P}$ of probability measures. Let T be a statistic such that its space $\mathcal{J}$ is a G -space. If T is equivariant and G is transitive on $\mathcal{J}$ then T is ancillary for $\mathcal{P}$ and distributed independently of a maximal G -invariant statistic T^* which also happens to be sufficient for $\mathcal{P}$. This result is then extended to a group G satisfying the Hunt-Stein Condition.


1. Introduction

This note originated from the following question Professor M. Eaton asked me. "Is $\underline{X}/\|\underline{X}\|$ ancillary and distributed independently of $\|\underline{X}\|$ when the distribution of $\underline{X}$ on $\mathbb{R}^n$ is preserved under orthogonal linear transformations?" The answer to this is known to be in the affirmative when the distribution of $\underline{X}$ is also normal. The following fact seems to be related to this issue. If the distribution of $\underline{X}$ on $\mathbb{R}^n$ is continuous and preserved under all permutations of its coordinates then the set of order statistics is sufficient and distributed independently of the ranks. We shall recall the standard proof of this result in order to answer Eaton's question in the affirmative in a broader framework.

Consider a family $\mathcal{P}$ of probability measures on a measure space $(\mathcal{X}, \mathcal{G})$. Let G be a (one-to-one bimeasurable) group of transformations on $\mathcal{X}$ to $\mathcal{X}$. We say that $\mathcal{P}$ is preserved under G , if for all $A \in \mathcal{G}$, $g \in G$, $P \in \mathcal{P}$

$$(1) \quad P(A) = P(g^{-1}A).$$

Let $\mathcal{G}^*$ be the sub- σ -field of all G -invariant measurable sets. Suppose (for simplicity) that T^* is a maximal G -invariant statistic.



We shall find conditions for which (i) T^* is sufficient for $\mathcal{P}$,
(ii) a given statistic T is ancillary for $\mathcal{P}$, and (iii) T and T^*
are independently distributed.

2. The Main Result.

We assume that the group G is locally compact with a given topology
and $\mathcal{G}$ is the collection of all its Borel sets. We assume that the transfor-
mation $(g, x) \rightarrow g(x)$ is jointly measurable in the product space
 $(G \times \mathcal{X}, \mathcal{G} \times \mathcal{G})$.

Consider a statistic T with the associated space $(\mathcal{T}, \mathcal{B})$. We shall
use the following two conditions in proving our results.

Condition 1. For every $g \in G$

$$T(x_1) = T(x_2) \Rightarrow T(gx_1) = T(gx_2) .$$

Condition 2. Given any x_1 and x_2 in $\mathcal{X}$ there exists a g
(depending on x_1 and x_2) in G such that

$$T(x_1) = T(gx_2) .$$

Suppose that Condition 1 holds. Then every $g \in G$ induces a
transformation $\tilde{g}$ on $\mathcal{T}$ to $\mathcal{T}$ given by

$$\tilde{g} T(x) = T(gx) .$$

Let $\tilde{\mathcal{G}}$ be the collection of all such $\tilde{g}$'s. Then Condition II
simply means that

Condition 3. $\tilde{\mathcal{G}}$ is transitive on $\mathcal{T}$.

Theorem 1. Suppose that a family $\mathcal{P}$ of probability measures is preserved
under a compact transformation group G . Let T^* be a maximal G -invariant
statistic and T be a statistic satisfying Conditions 1 and 2. Then
(i) T^* is sufficient for $\mathcal{P}$, (ii) T is ancillary for $\mathcal{P}$, and (iii) T
and T^* are independently distributed.

Proof. There exists a left-invariant probability measure ν on $(G, \mathcal{Q})$. For $A \in \mathcal{Q}$, $A^* \in \mathcal{Q}^*$, $P \in \mathcal{P}$, $g \in G$

$$(2.1) \quad P(A \cap A^*) = P(gA \cap A^*)$$

Hence

$$(2.2) \quad P(A \cap A^*) = \int_G P(gA \cap A^*) d\nu(g)$$

Define

$$(2.3) \quad \tilde{A} = \{(x, g) \in X \times G: g^{-1}x \in A\},$$

and

$$(2.4) \quad \tilde{A}_x = \{g \in G: g^{-1}x \in A\}.$$

Applying Fubini's Theorem to the left-hand side of (2.2) we get

$$(2.5) \quad P(A \cap A^*) = \int_{A^*} \nu(\tilde{A}_x) dP(x).$$

Since for $h \in G$

$$(2.6) \quad \tilde{A}_{hx} = h \tilde{A}_x,$$

we find that $\nu(\tilde{A}_x)$ is G -invariant. Hence T^* is sufficient for $\mathcal{P}$.

Now choose and fix any arbitrary $x_0 \in X$. By Condition II there exists $g_x \in G$ such that

$$(2.7) \quad T(x) = T(g_x^{-1}x_0).$$

By Condition I

$$(2.8) \quad T(g^{-1}x) = T(g^{-1}g_x^{-1}x_0)$$

for every $g \in G$. For $B \in \mathcal{B}$ write $A = T^{-1}(B)$. Then

$$\begin{aligned} \tilde{A}_x &= \{g \in G: g^{-1}x \in T^{-1}B\} \\ &= \{g \in G: g^{-1}g_x^{-1}x_0 \in T^{-1}B\}. \end{aligned}$$

$$(2.9) \quad = g_x^{-1} \tilde{A}_{x_0} .$$

Since ν is left-invariant

$$(2.10) \quad \nu(\tilde{A}_x) = \nu(\tilde{A}_{x_0}) .$$

This shows that the conditional probability of $T \in B$ given G^* is a constant free from $T^*(x)$. The results (ii) and (iii) now follow.

Note that the conditions 1 and 2 are implied also by the following:

Condition 4. G^* is a transitive group of (one-to-one onto bimeasurable) transformations on X to X such that G^* can be factored directly as $G^* = G \times H$, and T is a maximal H -invariant statistic.

Theorem 1 (i) was proved by Farrell [4].

Remark 1. Usually bounded completeness and sufficiency of T^* is used to show that T is ancillary is equivalent to independence of T and T^* [6]. Theorem 1 would be useful in many cases where the above result does not hold, or it is difficult to verify the bounded completeness of T^* . To apply Theorem 1 one needs only to check the simple conditions 1 and 2. Moreover, Theorem 1 generalizes many well-known results without assuming the existence of density functions. We shall illustrate these remarks by some examples.

Remark 2. Suppose both X and Y are G -spaces. Then the mapping defined by (8) is called equivariant. In that case, let $C \subset X$ be a cross-section of $X \rightarrow X/G$. Let $\varphi: C \rightarrow Y$ be a map such that the isotropy group $G_c \subset G_{\varphi(c)}$ for all $c \in C$. Then there is a unique extension of φ to an equivariant map $T: X \rightarrow Y$ [3].

Remark 3. It was proved by Farrell [4] and later by Basu [1] that G^* is sufficient when G is countable. In the dominated case Theorem 1 (i) can be proved more easily [1,2].

3. Examples

(a) Let $\mathcal{X} = \{x \in \mathbb{R}^n: x \neq 0\}$, and $\mathcal{G}$ be the class of all Borel subsets of $\mathcal{X}$. Let G be the group of all orthogonal linear transformations. Condition 3 can be verified easily. Then Theorem 1 holds with

$$T^*(x) = \|x\|, \quad T(x) = x/\|x\|.$$

In fact, the condition (4) holds with $H = \{h_\tau: \tau > 0\}$, $h_\tau x = \tau x$.

(ii) Consider the special case of the above example when $n = 1$. In that case G is the group of all sign transformations, and $T^*(x) = |x|$, $T(x) = \text{sign}(x)$. It may be of interest to note the following example:

$$P(X = 1) = P(X = -1) = \theta/2, \quad P(X = 2) = P(X = -2) = (1-\theta)/2,$$

where $0 \leq \theta \leq 1$. Let

$$T_1(x) = \begin{cases} 1, & \text{if } x = 1, -2 \\ 0, & \text{if } x = -1, 2. \end{cases}$$

Then T_1 is not a function of $\text{sign}(x)$; but T_1 is ancillary and distributed independently of $|x|$.

(iii) Let S be a symmetric $n \times n$ random matrix. Consider a family $\mathcal{P}$ of distributions of S which is preserved under orthogonal linear transformations, i.e., S and gSg' have the same distribution for every $n \times n$ orthogonal matrix g . Note that S can be decomposed (uniquely with some conventions) as

$$S = L_S D_S L_S',$$

where L_S is orthogonal and D_S is diagonal with its diagonal elements as the eigenvalues of S . Let T^* be the vector consisting of the diagonal elements of S , and $T = L_S$. Since the condition 3 holds in this case, Theorem 1 also holds with the above T and T^* .

(iv) Let $\mathcal{X} = \{x \in \mathbb{R}^n: x \neq 0\}$ and $\mathcal{G}$ be the class of all Borel subsets of $\mathcal{X}$. Suppose $\mathcal{P}$ is preserved by the Haggstorm subgroup of $n \times n$ orthogonal matrices (i.e., $n \times n$ orthogonal matrices with unit row sums and unit column sums). An example of such a $\mathcal{P}$ is $\eta_n(\mu \underline{1}_n, \Sigma)$, where $\Sigma = [\sigma_{ij}]$ is positive definite with $\sigma_{ij} = \rho\sigma^2$ for $i \neq j$ and $\sigma_{ij} = \sigma^2$ for $i = j$, and $\underline{1}_n$ is the $n \times 1$ vector with all 1's. Then Theorem 1 holds with.

$$T^*(X) = (\bar{X} = \sum_{i=1}^n X_i/n, \sum_{i=1}^n X_i^2),$$

$$T(X) = (X_1 - \bar{X}, \dots, X_n - \bar{X}) / [\sum_{i=1}^n (X_i - \bar{X})^2]^{\frac{1}{2}}.$$

(v) Let $\mathbb{R}$ be the additive group of reals with the usual topology and $\mathbb{Z}$ be the subgroup of all integers. Then the quotient group $G = \mathbb{R}/\mathbb{Z}$ is compact with respect to the quotient topology [7]. It can be seen that G is isomorphic to the group $G_c = \{g_c: 0 \leq x < 1\}$ with $g_{c_1} + g_{c_2} = g_c$, where $c = c_1 + c_2 \pmod{1}$.

Let $\mathcal{P}$ be the family of uniform distributions on $[0, \theta)$, with θ being a positive integer. Consider

$$g_c x = [x] + \{x - [x] + c \pmod{1}\},$$

where $[x]$ is the integer part of x (see [1]). Then Theorem 1 holds with

$$T^*(x) = [x], T(x) = x - [x].$$

4. An Extension of Theorem 1.

Next we shall use Hunt-Stein's condition [6] given below to extend Theorem 1.

Condition 5. There exists a sequence $\{\nu_n\}$ of probability measures on $(G, \mathcal{G})$ such that for any $g \in G, B \in \mathcal{G}$

$$\lim_{n \rightarrow \infty} |\nu_n(gB) - \nu_n(B)| = 0 .$$

Theorem 2. Theorem 1 holds if the compactness of G is replaced by Condition 5, and X is Euclidean.

Proof. Proceeding as before we get

$$(4.1) \quad P(A \cap A^*) = \int_{A^*} \nu_n(\tilde{A}_x) dP(x)$$

for $A \in \mathcal{G}, A^* \in \mathcal{G}^*$ and $\tilde{A}_x$ given by (2.4). From (2.9) we get

$$(4.2) \quad \nu_n(\tilde{A}_x) = \nu_n(g_x^{-1} \tilde{A}_{x_0}) ,$$

where $A = T^{-1}B, B \in \mathcal{B}$.

Condition 5 implies (see [6], p. 337) that for any fixed x and $g \in G$

$$(4.3) \quad \nu_n(g \tilde{A}_x) - \nu_n(\tilde{A}_x) \rightarrow 0$$

as $n \rightarrow \infty$. Hence

$$(4.4) \quad \nu_n(\tilde{A}_x) - \nu_n(\tilde{A}_{x_0}) \rightarrow 0$$

as $n \rightarrow \infty$. By Lebesgue dominated convergence theorem

$$(4.5) \quad \int_{A^*} [\nu_n(\tilde{A}_x) - \nu_n(\tilde{A}_{x_0})] dP(x) \rightarrow 0 .$$

Thus from (4.1) we get

$$(4.6) \quad P(A \cap A^*) = P(A^*) \lim_{n \rightarrow \infty} \nu_n(\tilde{A}_{x_0}) .$$

Clearly now

$$(4.7) \quad P(A) = \lim_{n \rightarrow \infty} \nu_n(\tilde{A}_{x_0}) .$$

This shows $P(A)$ is a constant and $P(A \cap A^*) = P(A) \cdot P(A^*)$. The results (ii) and (iii) now follow. The fact that $\mathcal{G}^*$ is sufficient for $\mathcal{P}$ can be proved easily proceeding exactly as in Lehmann ([6], pp. 336-7).

References

- [1] Basu, D. (1965). On sufficiency and invariance.
- [2] Basu, D. (1965). Sufficiency and model-preserving transformations.
Inst. of Statist. Mimeo 420, Univ. of N.C., Chapel Hill.
- [3] Benson, C. T. (1971). Finite Reflection Groups. Bogden and Quigley,
New York.
- [4] Farrell, R. H. (1962). Representation of invariant measures.
Ill. J. Math. 6, 447-467.
- [5] Halmos, P. (1950). Measure Theory. Van Nostrand, New York.
- [6] Lehmann, E. (1959). Testing Statistical Hypotheses, Wiley, New York.
- [7] Pontryagin, L. S. (1966). Topological Groups, Gordon and Breach,
New York.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
A NOTE ON ANCILLARITY AND INDEPENDENCE VIA MODEL-PRESERVING TRANSFORMATIONS.		Technical Report
6. AUTHOR		6. PERFORMING ORG. REPORT NUMBER
Somesh Das Gupta		TR 319
7. PERFORMING ORGANIZATION NAME AND ADDRESS		8. CONTRACT OR GRANT NUMBER(s)
Department of Theoretical Statistics University of Minnesota Minneapolis, MN 55455		DAAG 29-76-G-0038
9. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Army Research Office Box 13311 Fort Monmouth, NJ 07709		
11. DISTRIBUTION STATEMENT (of the Report)		12. REPORT DATE
Approved for public release; distribution unlimited.		May, 1978
13. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		13. NUMBER OF PAGES
NA		9
14. SUPPLEMENTARY NOTES		15. SECURITY CLASS. (of this report)
The findings in this report are not to be construed as an official position of the Army position, unless so designated by other authorized authorities.		Unclassified
16. ABSTRACT (Continue on reverse side if necessary and identify by block number)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
Sufficiency, Ancillarity, model-preserving transformations, compact group, invariant probability measure.		NA
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
NA		
18. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Let G be a compact group of transformations which preserves a family P of probability measures. Let T be a statistic such that its space $\mathcal{T}$ is a G -space. If T is equivariant and G is transitive on $\mathcal{T}$ then T is ancillary for P and distributed independently of a maximal G -invariant statistic T^* which also happens to be sufficient for P . This result is then extended to a group G satisfying the Hunt- Stein Condition.		