

AD-A056 713

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY
THE STUDY OF THE INTRAMOLECULAR HEAVY ATOM EFFECT IN 9,10-DICHL--ETC(U)
JUN 78 T AKASHEH


F/G 20/10
N00014-75-C-0602

NL

UNCLASSIFIED


TR-30

1 OF 1
AD
A056 713



END
DATE
FILMED
9-78

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A056713

AD No.
 DC FILE COPY

LEVEL III

12

OFFICE OF NAVAL RESEARCH

Contract No. N00014-75-C-0602

Task No. NR-056-498

TECHNICAL REPORT NO. 30

THE STUDY OF THE INTRAMOLECULAR HEAVY ATOM EFFECT IN
9,10-DICHLOROPHENANTHRENE AND 1,2,3,4-
TETRACHLORONAPHTHALENE USING PMDR TECHNIQUES

by

Talal Akasheh
Department of Chemistry
University of California
Los Angeles, California 90024

DDC
JUL 26 1978
F

Prepared for Publication

in

Chemical Physics Letters

Reproduction in whole or in part is permitted for any
purpose of the United States Government

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

June, 1978

78 07 25 041

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 30✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ⑥ The Study of the Intramolecular Heavy Atom Effect in 9,10-Dichlorophenanthrene and 1,2,3,4-Tetrachloronaphthalene Using PMDR Techniques		5. TYPE OF REPORT & PERIOD COVERED ⑨ Interim Technical Report
6. PERFORMING ORG. REPORT NUMBER		7. CONTRACT OR GRANT NUMBER(s) ⑬ N00014-75-C-0602
8. AUTHOR(s) ⑩ Talal Akasheh		9. PERFORMING ORGANIZATION NAME AND ADDRESS Regents of the University of California University of California, 405 Hilgard Ave. Los Angeles, California 90024
10. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Branch Arlington, Virginia 22217 ⑪ 30 Jun 78		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-056-498
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91106 ⑫ 13 p		12. REPORT DATE June 30, 1978
13. DISTRIBUTION STATEMENT (of this Report) Distribution of this document is unlimited.		13. NUMBER OF PAGES 13
14. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) ⑭ TR-30		15. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. SUPPLEMENTARY NOTES To be published in Chemical Physics Letters		
17. KEY WORDS (Continue on reverse side if necessary and identify by block number) phosphorescence PMDR heavy atom effect Sigma, pi* and pi, pi*		
18. ABSTRACT (Continue on reverse side if necessary and identify by block number) The heavy atom effect in triplet 9,10-dichlorophenanthrene and 1,2,3,4-tetrachloronaphthalene is investigated by PMDR techniques. As in the parent hydrocarbons, spin-orbit and spin-orbit vibronic mechanisms are invoked to explain the results. The relative radiative rates of the triplet zero-field levels are determined by static distortions of the chlorine atoms, while the intersystem crossing rates from S_1 are mainly governed by the enhancement of spin-orbit vibronic coupling via C-Cl out-of-plane modes that mix the relevant σ, π^* and π, π^* states. In both processes the top two levels (τ_A and τ_B) are more active, with τ_B the most favoured of the two. Tau(B)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)
Tau(A) and Tau(B)

072 255

ACCESSION for	Write Section	☒
NTIS	Buff Section	☐
DDC		
UNANNOUNCED		
JUSTIFICATION		
BY	DISTRIBUTION/AVAILABILITY CODES	
Dist.		SPECIAL
A		

The Study of the Intramolecular Heavy Atom Effect in
9,10-Dichlorophenanthrene and 1,2,3,4-
Tetrachloronaphthalene Using PMDR Techniques

Talal Akasheh*
Department of Chemistry
University of California
Los Angeles, California 90024

ABSTRACT

The heavy atom effect in triplet 9,10-dichlorophenanthrene and 1,2,3,4-tetrachloronaphthalene is investigated by PMDR techniques. As in the parent hydrocarbons, spin-orbit and spin-orbit vibronic mechanisms are invoked to explain the results. The relative radiative rates of the triplet zero-field levels are determined by static distortions of the chlorine atoms, while the intersystem crossing rates from S_1 are mainly governed by the enhancement of spin-orbit vibronic coupling via C-Cl out-of-plane modes that mix the relevant σ, π^* and π, π^* states. In both processes the top two levels (τ_A and τ_B) are more active, with τ_B the most favoured of the two.

*Permanent address: Chemistry Department, Yarmouk University, Irbid, Jordan

78 07 25 041

1. Introduction:

It is well known that the internal heavy atom effect (HAE) on the dynamic properties of triplet aromatic molecules exhibits itself by enhancing multiplicity forbidden transitions via increased spin-orbit coupling routes with singlet states. $^1_{\sigma,\pi}^*$ states play the most prominent role in this scheme since coupling between $^1_{\pi,\pi}^*$ and $^3_{\pi,\pi}^*$ states is vanishingly small (1-3). Experimentally, an enhanced $S_1 \rightarrow T_1$ intersystem crossing rate, a shortening of the T_1 lifetime and an increased $T_1 \rightarrow S_0$ radiative decay are associated with the HAE (4). Considerations of the phosphorescence polarization and the anisotropy of the dynamic behaviour of individual zero-field levels of T_1 in haloaromatics require the involvement of spin-vibronic and spin-orbit vibronic interactions (5-7).

The use of PMDR techniques (3) has proven to be a powerful tool for investigating the dynamics of individual zero-field level behaviour. The radiative properties of the three levels for the 0-0 optical transitions have led to predictions concerning static molecular distortions (8,9) in haloaromatics as well as assignments of the orbital symmetries of T_1 in benzene (10), phenanthrene (11), naphthalene (12) and tetrachlorobenzene (13). Furthermore, the radiative properties of the zero-field levels observed by monitoring vibronic bands together with their relative intersystem crossing rates (12,14-16) established the importance of out-of-plane vibrational modes in the spin-orbit vibronic coupling scheme.

Recently, the dynamic properties of the triplet state of various haloaromatic compounds were studied (17). In this paper we report similar results for 9,10-dichlorophenanthrene (DCP) and 1,2,3,4-tetrachloronaphthalene (TCN) in *n*-hexane at 1.6°K. Experimental techniques have been thoroughly described in the literature (3,18-22), and will not be discussed here.

2. Results:

2.1.1. 9,10-Dichlorophenanthrene:

The uncorrected 1.5°K phosphorescence spectrum is shown in Fig. 1 while the calibrated (using Hg lines) band locations are given in Table 1. Two sites, 82 cm^{-1} apart, are observed with 0-0 bands at 20842 and 20760 cm^{-1} , respectively. The two sites are also observed in the 0-260 cm^{-1} vibrational bands. The stronger 0-0 band (20760 cm^{-1}) was monitored throughout the experiment. The zero-field transitions (D-E and 2E) that are shown in Table 2 are close to those known for phenanthrene. The 2E transition monitored at 20760 cm^{-1} is slightly different from that at 20842 cm^{-1} , thus confirming the existence of two sites. The D+E transition can only be observed in EEDOR (23). The relative radiative rates k_i^r (0-0), the lifetimes τ_i , the relative populating rates, K_i , and the relative populations, N_i , are also shown in Table 2. For comparison purposes, results for phenanthrene found in the literature (11,15) are included.

2.1.2. 1,2,3,4-Tetrachloronaphthalene:

The phosphorescence spectrum and calibrated bands for TCN are shown in Fig. 2 and Table 3, respectively. The zero-field transition energies (also presented in Fig. 2) are typical of naphthalene and azanaphthalene systems. Table 4 gives the dynamic properties. These are compared with naphthalene results (12, 15), and it is noticed that TCN follows the same general trends as in the parent hydrocarbon.

3. Discussion:

3.1. Relative radiative rates:

In both chlorinated molecules it is observed that τ_B (total symmetry = $\text{spin}(A_2) \times \text{orbital}(B_2) = B_1$) is the most radiative level in the 0-0. This is also the case for the parent hydrocarbons. The group theoretical predictions of spin-orbit coupling (SOC) in C_{2v} symmetry (see ref. 3) confirm that this level is the one that mixes with the $^1\sigma, \pi^*$ states (1B_1) and hence benefits most from spin-orbit coupling. The lowest level, τ_N , mixes with the $^1\pi, \pi^*$ states (1A_1) and hence is the least emissive. Similar arguments indicate that the orbital symmetry of the lowest triplet state of TCN and DCP is of B_2 species, since otherwise τ_A would be the most radiative level; if the orbital symmetry were A_1 , then τ_A would have total symmetry B_1 and it would mix with the $^1\sigma, \pi^*$ (B_1) state. Comparison of DCP with phenanthrene shows that the level τ_A is emissive in the substituted compound though the parent molecule doesn't show such activity. This can be explained by static in-plane distortion of the two chlorine atoms in a manner that destroys the C_2 and reflection symmetry perpendicular to the plane of the molecule. This mixes the τ_B character into the τ_A level and causes the observed radiative activity of the τ_A . In TCN static out-of-plane distortions of the crowded chlorine atoms seem to be the cause of the radiative character of the lowest level τ_N , which is dark in naphthalene. Such behavior has also been observed for 2,3-dibromonaphthalene (8).

3.2. Intersystem Crossing Rates:

The trend in the relative intersystem crossing rates shows the same behavior for the DCP and TCN as for the parent hydrocarbons. The top two levels, τ_A and τ_B , are most favoured with τ_B having the higher rate of the two. As in the parent hydrocarbons (3, 15) the results can be explained by spin-orbit

vibronic coupling (see refs. 3 and 15). Intersystem crossing from $S_1(^1\pi,\pi^*)$ to $T_1(^1\pi,\pi^*)$ is not favourable by SOC alone. Vibronic coupling via a_2 or b_1 vibrations could contaminate the $S_1(^1B_2)$ with $^1\sigma,\pi^*$ character. The contaminated singlet has better SOC ability with the T_1 state. Alternatively the vibronic mixing could occur in the triplet manifold with exactly the same predictions. The a_2 vibration enhances level τ_A while the b_1 vibration enhances the population of τ_B (e.g., $\tau_A: {}^3B_2 \xrightarrow{\text{SOC}} {}^1B_1 \xrightarrow{a_2} {}^1B_2$). However, compared to the parent hydrocarbons τ_B is even more favoured than τ_A . This indicates that the heavy-atom enhances the ability of the b_1 vibration to couple the electronic levels more than the a_2 mode. The π,π^* and σ,π^* states are orthogonal and possess different symmetries with respect to the molecular reflection plane. Thus the a_2 and b_1 modes are out-of-plane vibrational modes. Their enhancement by heavy atom substitution results in the enhancement of a strong nontotally symmetric progression in the phosphorescence spectrum. This has been observed at 77 °K for various halophenanthrenes (30). Furthermore, in tetrachlorobenzene (TCB) (14) the most effective out-of-plane mode was found to be the b_{2g} which corresponds to C-Cl out-of-plane vibration. Thus, the active b_1 vibration in DCP and TCN can be safely assumed to be the out-of-plane C-Cl vibrations corresponding to the b_{2g} out-of-plane C-H or C-C bending vibration in naphthalene and the b_{2g} out-of-plane C-Cl vibration in TCB.

ACKNOWLEDGMENT: The financial support of the Office of Naval Research is gratefully acknowledged.

REFERENCES

1. D. S. McClure, J. Chem. Phys., 20, 282(1952).
2. S. K. Lower and M. A. El-Sayed, Chem. Rev., 66, 199(1966).
3. M. A. El-Sayed, in Excited States, Vol. 1, p. 35, Academic Press, Inc., 1974.
4. C. T. Lin, Thesis, UCLA, 1974.
5. M. A. El-Sayed, Acta Phys. Polonica, XXXIV, 649(1968).
6. M. A. El-Sayed, Proceedings for the Int. Conf. on Lum., p. 373(1966).
7. N. K. Chaudhuri and M. A. El-Sayed, J. Chem. Phys., 47, 2566(1967).
8. M. A. El-Sayed, M. Leung and C. T. Lin, Chem. Phys. Lett., 14, 329(1972).
9. G. Kothandaraman and D. S. Tinti, Chem. Phys. Lett., 19, 225(1972).
10. A. A. Gwaiz, M. A. El-Sayed and D. S. Tinti, Chem. Phys. Lett., 9, 454(1971).
11. A. L. Kwiram, Chem. Phys. Lett., 1, 272(1967).
12. a. H. Sixl and M. Schwoerer, Chem. Phys. Lett., 6, 21(1970).
b. M. Schoerer and H. Sixl, Chem. Phys. Lett., 2, 14(1968)
13. C. R. Chen and M. A. El-Sayed, Chem. Phys. Lett., 10, 307(1971).
14. W. Pitts and M. A. El-Sayed, Chem. Phys., 19, 289(1977).
15. M. A. El-Sayed, W. R. Moomaw and J. B. Chodak, Chem. Phys. Lett., 20, 11(1973).
16. M. A. El-Sayed and C. R. Chen, Chem. Phys. Lett., 10, 313(1971).
17. C. T. Lin, J. of Luminescence, 12/13, 375(1976).
18. C. J. Wiscom and A. H. Maki, Chem. Phys. Lett., 12, 264(1971).
19. A. L. Shain and M. Sharnoff, J. Chem. Phys., 59, 2335(1973).
20. J. Schmidt, D. A. Antheunis and J. H. van der Waals, Mol. Phys., 22, 1(1971).
21. M. A. El-Sayed and J. Olmsted III, Chem. Phys. Lett., 11, 568(1971).
22. J. B. Chodak, Thesis, UCLA, 1974.
23. J. Turner and M. A. El-Sayed, unpublished results.
24. J. K. Roy and L. Goodman, J. Mol. Spect., 19, 389(1966).

Table 1. Calibrated phosphorescence bands of 9,10-dichlorophenanthrene

λ	cm^{-1}	Intensity*	$\Delta\nu(\text{cm}^{-1})$	
			Site I	Site II
4798	20842	M	0	
4817	20760	S		0
4861	20574	VW	268	
4875	20512	VW	330	
4881	20490	M		270
4896	20426	M	416	334
4899	20413	M	429	
4915	20346	M		414
4918	20334	M		426
4923	20313	VW	529	
4943	20231	W		529
4942	20236	W		
4996	20015	VW	827	
5000	20002	W	840	
5017	19932	W		828
5020	19922	M		838
5025	19900	VW	942	
5046	19819	VW		941
5047	19813	W	1029	
5069	19729	W		1031

* M = Medium; S = Strong; V = Very; W = Weak

Table 2. Zero-field transitions, the relative radiative rates, $k_i^r(0-0)$, the lifetimes, τ_i , and relative populations of the zero-field levels of 9,10-dichlorophenanthrene and phenanthrene (in parentheses)

	Phenanthrene*		9,10-Dichlorophenanthrene					2E	D-E
	$k_i^r(0-0)$	K_i	$\tau_i(\text{sec})$	$k_i^r(0-0)^*$	K_i	N_i			
τ_A	0	0.70	0.48	0.46	0.60	0.43		2.864 (2.802) [†]	
τ_B	1	1	0.14	1	1	0.20			1.732 (1.731) [†]
τ_N	0.25	0.32	2.0	0	0.12	0.37			

* See Refs. 11,15

[†] M. A. de Groot and J. H. van der Waals, *Physica* 19, 1128 (1963).

Table 3. Calibrated phosphorescence bands of 1,2,3,4-tetrachloronaphthalene

<u>Wavelength ($\text{\AA}$)</u>	<u>Frequency (cm^{-1})</u>	<u>Relative Intensity</u>	<u>$\Delta\nu(\text{cm}^{-1})$</u>
5058	19770	VS	0
5100	19609	VW	161
5137	19466	W	304
5142	19448	M	322
5144	19440	S	330
5150	19418	M	352
5161	19377	M	393
5175	19324	MW	446
5200	19230	M	540
5217	19168	W	602
5234	19105	M	665
5239	19088	M	682
5256	19026	W	744
5262	19003	MW	767
5281	18937	W	833
5291	18901	W	869
5320	18799	W	971
5342	18720	W	1050
5372	18614	W	1156
5434	18402	W	1368
5440	18381	M	1389

Table 4. The dynamic properties of 1,2,3,4-tetrachloronaphthalene and naphthalene.


<u>Naphthalene (N)</u>			<u>1,2,3,4-tetrachloronaphthalene</u>			
	<u>$k_1^r(0-0)^{(a)}$</u>	<u>$K_1^{(b)}$</u>	<u>$\tau_1(\text{sec})$</u>	<u>$k_1^r(0-0)$</u>	<u>K_1</u>	<u>N_1</u>
τ_B	1	1	0.055	1	1	0.29
τ_A	0.25-0.22	0.39	0.200	0.13	0.25	0.26
τ_N	~ 0	0.17	1.50	0.13	0.05	0.45


a) Ref. 12. b) Ref. 15

FIGURE CAPTIONS

Fig. 1 9,10-dichlorophenanthrene phosphorescence spectrum (uncorrected) at 1.5°K in n-hexane. The axes system and the zero-field energy level diagram are also shown.

Fig. 2 Phosphorescence spectrum (uncorrected) at 1.5°K in n-hexane, energy level diagram, axis system and zero-field transitions (in GHz) of 1,2,3,4-tetrachloronaphthalene.





TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
Office of Naval Research Arlington, Virginia 22217 Attn: Code 102IP	6	U.S. Army Research Office P.O. Box 12211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. Jerry Smith	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Weapons Center China Lake, California 93555 Attn: Head, Chemistry Division	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Civil Engineering Laboratory Port Hueneme, California 93041 Attn: Mr. W. S. Haynes	1
ONR Branch Office 760 Market Street, Rm. 447 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Professor O. Heinz Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
ONR Branch Office, Bldg. 114, Section D 496 Summer Street 666 Summer St. Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Office of Naval Research Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
The Asst. Secretary of the Navy (R&D) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1		
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1		

TECHNICAL REPORT DISTRIBUTION LIST

No. Copies

No. Copies

~~Dr. M. A. El-Sayed~~
~~University of California~~
~~Department of Chemistry~~
~~Los Angeles, California 90024~~ 1

Dr. M. W. Windsor
 Washington State University
 Department of Chemistry
 Pullman, Washington 99163 1

Dr. E. R. Bernstein
 Colorado State University
 Department of Chemistry
 Fort Collins, Colorado 80521 1

Dr. C. A. Heller
 Naval Weapons Center
 Code 6059
 China Lake, California 93555 1

Dr. M. H. Chisholm
 Princeton University
 Department of Chemistry
 Princeton, New Jersey 08540 1

Dr. J. R. MacDonald
 Naval Research Laboratory
 Chemistry Division
 Code 6110
 Washington, D.C. 20375 1

Dr. G. B. Schuster
 University of Illinois
 Chemistry Department
 Urbana, Illinois 61801 1

Dr. E. M. Eyring
 University of Utah
 Department of Chemistry
 Salt Lake City, Utah 84112 1

Dr. A. Adamson
 University of Southern California
 Department of Chemistry
 Los Angeles, California 90007 1

Dr. M. S. Wrighton
 Massachusetts Institute of Technology
 Department of Chemistry
 Cambridge, Massachusetts 02139 1

Dr. M. Rauhut
 American Cyanamid Company
 Chemical Research Division
 Bound Brook, New Jersey 08805 1

Dr. G. Jones, II
 Boston University
 Department of Chemistry
 Boston, Massachusetts 02215 1