


D-A056 815

TECHNICAL UNIV OF NORTHERN SWEDEN LULEA DIV OF SOILS--ETC F/G 8/7
CLAY MICROSTRUCTURE AND ICE NUCLEATION.(U)
APR 78 R PUSCH

DA-ERO-77-G-033
NL

UNCLASSIFIED

1 OF
AD
A056815



END
DATE
FILMED
9-78
DDC

AD A 056815

AD No. _____
DDC FILE COPY

12

LEVEL II

AD

6 CLAY MICROSTRUCTURE AND ICE NUCLEATION.

9 Final technical report, Aug 77 - Apr 78,

10 by Roland/Pusch

11 Apr 78

12 24 p.

16 IT 161142 BH 57 17 41

EUROPEAN RESEARCH OFFICE

United States Army

London England

15 GRANT NUMBER DA-ER0-77-G-033

Professor Roland Pusch
Div Soil Mechanics
University of Luleå
S-951 87 Luleå, Sweden

Approved for Public Release

Distribution unlimited

DDC
REPRODUCTION
JUL 28 1978
B

SIC 393238

78 07 17 013

mit

S/C 393 238

Technical Univ. of Northern Sweden, Luleå.
Div. of Soils Mechanics.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CLAY MICROSTRUCTURE AND ICE NUCLEATION		5. TYPE OF REPORT & PERIOD COVERED FINAL TECHNICAL REPORT Aug 77 - Apr 78
7. AUTHOR(s) ROLAND PUSCH		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Lulea Lulea, Sweden		8. CONTRACT OR GRANT NUMBER(s) DAERO-77-G-033 <i>new</i>
11. CONTROLLING OFFICE NAME AND ADDRESS US Army R&S Gp (Eur) Box 65, FPO New York 09510		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 6.11.02A-1T161102BH57-01-00-624
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE APRIL 1978
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ice Nucleation; clay microstructure		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The microstructural difference between weakly aggregated ("dispersed") and strongly aggregated clays means that a larger part of the water is structurally unaffected by clay mineral lattices in the last-mentioned type of clay which should therefore have a lower content of unfrozen water. This hypothesis was tested and found to be valid in an investigation where two such clays were actually compared with reference to the content of unfrozen water. The results from dilatometer and spin echo tests clearly show that this content is <u>CONT</u>		

DD FORM 1473 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

78 07 17 013

PTO

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ADN1 → dependent not only on the specific surface area but also on the microstructural constitution expressed in terms of the degree of aggregation.



ACCESSION FOR	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	ORAL COPY or SPECIAL
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

CLAY MICROSTRUCTURE AND ICE NUCLEATION ¹⁾

TABLE OF CONTENTS

	Page
SUMMARY	
INTRODUCTION, SCOPE OF INVESTIGATION	1
CLAY TYPES	2
Geotechnical properties	2
Grain size distribution	3
Mineralogy	5
Specific surface area	6
Salinity	6
Conclusions	6
TESTS	7
Equipment	7
Program	9
Accuracy	10
Test results	10
Spin echo technique test	15
MEANS OF DETERMINING AND EXPRESSING THE DEGREE OF AGGREGATION	16
CONCLUSIONS	18
REFERENCES	18

¹⁾ In this report ice nucleation stands for the formation of ice at very moderate negative temperatures in clays under closed conditions.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

SUMMARY

The microstructural difference between weakly aggregated ("dispersed") and strongly aggregated clays means that a larger part of the water is structurally unaffected by clay mineral lattices in the last-mentioned type of clay which should therefore have a lower content of unfrozen water. This hypothesis was tested and found to be valid in an investigation where two such clays were actually compared with reference to the content of unfrozen water. The results from dilatometer and spin echo tests clearly show that this content is dependent not only on the specific surface area but also on the microstructural constitution expressed in terms of the degree of aggregation.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

INTRODUCTION, SCOPE OF INVESTIGATION

The main characteristic feature of illite clay microstructure is aggregation, which is very strong in marine clays but less obvious in fresh water-deposited clays. Aggregation means that the particles form fairly stable groups (secondary particles) separated by voids. In principle, the microstructural difference between fresh water clays and marine clays is as shown in Fig. 1 and this has also been confirmed by a number of investigators (cf. PUSCH 1970, BARDEN 1973).




Fig. 1 Schematic picture of the microstructural pattern of natural illitic clay a) deposited in fresh water, b) deposited in salt water.

Fig. 1 means that the particles in fresh-water clay are collected in a large number of small but fairly porous aggregates separated by small voids, while they form big and dense aggregates separated by large voids in marine clay. A larger fraction of the soil water is located within the very small intra-aggregate voids in weakly aggregated ("dispersed"¹⁾) than in strongly aggregated clays. This can also

¹⁾ This description is commonly used for clays with a fairly uniform particle distribution. There is no single particle behaviour, however.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

be expressed by saying that a major part of the water is structurally unaffected by clay mineral lattices in the last-mentioned type of clay. The consequence is that a weakly aggregated clay should have a higher percentage of unfrozen water than a strongly aggregated clay with the same electrolyte, mineralogical and granulometrical compositions. Ice nucleation should occur more easily and rapidly in the "free water" in the large voids of strongly aggregated clay.

Previous investigations (PUSCH, 1977) support this hypothesis. Thus, electron microscopical (SEM) studies on freezing clay have shown that many particle aggregates are moved without much internal distortion, which indicates that much of the intra-aggregate water does not freeze in the temperature interval 0° to -5°C. The conclusion of this work was that the amount of unfrozen water should be large when the particle aggregates are small and fairly porous but closely spaced since a considerable mineral surface is then exposed to the water. In principle, this is in agreement with the statement by DILLON & ANDERSLAND (1966) and ANDERSON & TICE (1972) that the unfrozen water is a function of the specific surface area. However, since the degree of aggregation in fact determines to what extent this area is exposed to water it should be equally important. The scope of the present investigation was to investigate this experimentally. For this purpose the unfrozen water content of two soft, illitic clays formed in fresh and salt water was determined. The clays had quite different microstructural constitutions but otherwise similar compositions and properties.

CLAY TYPES

Geotechnical properties

A fresh-water clay from Skå-Edeby (Stockholm area) and a marine clay from Lilla Edet (Swedish west-coast) were used for the investigation. Their main geotechnical properties are given in Table 1. The sampling depth was 5 m at the Skå-Edeby and 3 m at the Lilla Edet site.

The fresh-water clay from Skå-Edeby is normally sensitive while the marine clay from Lilla Edet is quick. The high sensitivity of the latter clay is because it has been leached by percolating fresh water in nature.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

Table 1 Geotechnical data

Clay	ρ t/m ³	τ_{fu} kPa	S_t	w %	W _L %	w _p %	Organic content %
Skå-Edeby	1.42	8-10	12	82-106	93	29	~1
Lilla Edet	1.48	13-18	42	97-101	71-73	33	~1

Grain size distribution

The plastic limit is of the same order of magnitude for the two clays while the Skå-Edeby clay has a somewhat higher liquid limit which indicates a higher clay content (Table 2).

Table 2 Grain size distribution

Clay	Weight percentage of material finer than				
	0.04 mm	0.02 mm	0.01 mm	0.006 mm	0.002 mm
Skå-Edeby	82	80	79	79	77
Lilla Edet	99	99	90	88	67

Previous electron microscopical investigations have shown that the average particle size of the clay fraction is much smaller in the Lilla Edet clay than in the Skå-Edeby clay (PUSCH, 1970). Thus, using the definition of particle size given by Fig. 2, it was found that the median value a was only 0.018 μm for the first-mentioned clay while it was 0.037 μm for the Skå-Edeby clay. A microstructural study showed that a large number of particles smaller than 0.02 μm act as linking units in the particle network. Hence, they were not formed after the deposition of the sediment and they are not a product of the preparation of the clay suspensions for the granulometrical investigation. A detailed description of the granulometrical and morphological properties of the clay fractions is given in Table 3.

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 2 Example of histogram of particle size for the clay fraction.

Table 3 Median values, quartiles and percentiles of **a** and **a/c** for clay particles

Clay	a in μm						a/c		
	M ¹	Q ₁ ²	Q ₃ ³	P ₁₀ ⁴	S _K ⁵	log S ₀ ⁶	M	Q ₁	Q ₃
Skå-Edeby	0.037	0.026	0.083	0.021	1.58	0.252	4.0	2.9	5.9
Lilla Edet	0.018	0.014	0.025	0.012	1.08	0.126	2.7	1.8	3.3

¹ Median value

² Lower quartile

³ Upper quartile

⁴ 10th percentile (finest part of the cumulative curve)

⁵ Skewness (TRASK 1932), $(Q_1 \cdot Q_3) / M^2$

⁶ S_0 = sorting coefficient $\sqrt{Q_3/Q_1}$; log S_0 -values are directly comparable

The presence of many particles of colloidal size in the clay fraction of the Lilla Edet clay is of great importance as concerns the specific surface area. As will be shown later in the text this means that although the clay content (weight percentage of material finer than 2 μm) is higher for the Skå-Edeby clay, the specific surface area is much higher for the Lilla Edet clay.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

Mineralogy

X-ray diffraction analyses (cf. Fig. 3) of particles finer than 10 μm have shown that the main minerals in the Skå-Edeby clay are carbonate minerals, (mainly calcite), illite, quartz, kaolinite, feldspars and chlorite, the three first-mentioned being dominant. In the Lilla Edet clay, which is rich in rock flour, the main minerals are feldspars, quartz, illite, kaolinite, hornblende and rock-forming micas. Illite is the dominant clay mineral in both clays. The clay fraction of the Lilla Edet clay contains considerable amounts of quartz and feldspars.




Fig. 3 Representative diffractometer records showing d-values (horizontal axis gives 2θ). Upper diagram Skå-Edeby. Lower diagram Lilla Edet.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

Specific surface area

The specific surface area was determined in two ways. Firstly, it was calculated on the basis of Table 3 assuming the particles to have a cylindrical shape with a diameter equal to **a** and a height equal to **C**. The calculation gave 99 m²/g for the Skå-Edeby clay and 204 m²/g for the Lilla Edet clay. Secondly, it was determined experimentally by applying the BET method and MADSEN's (1977) glycerol sorption technique. This investigation gave 690 m²/g for the Skå-Edeby clay and 996 m²/g for the Lilla Edet clay. The latter values, which should be more reliable than the theoretically deduced ones, indicate rather extreme specific surface areas, almost of the order found for 100% Na-montmorillonite. The reason for the discrepancy is not known but it may partly be due to the action of organic substance and mineral weathering which are not accounted for in the theoretical deduction. It should be mentioned that the experimental values are not improbably high since the Skå-Edeby clay is known to be so fine-grained that it does not obey Darcy's law (HANSBO, 1960). The important thing is that the specific surface area of the Lilla Edet clay is higher than that of the Skå-Edeby clay according to both determinations.

The average surface activity expressed by the cation exchange capacity is similar for both clays (29-33 meq/100 g dry material) despite the higher specific surface area of the Lilla Edet clay. This is explained by the fact that the Lilla Edet clay fraction is rich in inactive rock flour, mainly very fine-grained quartz and feldspars.

Salinity

The salinity of the water during the formation of the investigated clays has been estimated by applying diatom analyses (PUSCH, 1970). The original salt content of the Skå-Edeby clay must have been less than the present value 5%, while the Lilla Edet clay was formed under marine conditions with a salt content of about 35%. Today, this latter value has been reduced to about 5% due to leaching by fresh-water percolation which converted the Lilla Edet material to a quick clay.

Conclusions

The two clays have quite different microstructural constitutions but otherwise similar compositions and properties except for the specific surface area, which is higher for the marine Lilla Edet clay. This should give the Lilla Edet clay the highest content of unfrozen water if there were no influence of the microstructural geometry. A comparison between the contents of unfrozen water of the clays can there-

CLAY MICROSTRUCTURE AND ICE NUCLEATION

fore tell us whether this geometry - or rather the degree of aggregation - has any effect.

TESTS

Equipment

Dilatometer technique was applied for the determination of the amount of unfrozen water. The samples (40 mm diameter and 20-25 mm height) were mounted in closely fitting steel cylinders which allowed uni-axial swelling only and no uptake of external water (closed system). The design of the dilatometer is shown in Figs. 4 and 5.

The cylinder, which was equipped with a sharp edge, was pushed into the clay to a depth of 20-25 mm. After truncation with a thin steel wire the lids were applied and the device closed. This operation was made in a batch with paraffine oil (purum), which was the liquid used for measuring the volume changes. Since the whole device was submerged in the oil throughout the mounting operation there should be no air left in the cylinder and glass pipe or any other part of the measuring system. Thus, by calculating the temperature-dependent volume change of the steel cylinder and by determining experimentally the same property of the paraffine oil, the device was calibrated so that an observed displacement of the oil meniscus in the glass pipe could be expressed in terms of a volume change of the clay sample. The relationship turned out to be 29.3 mm^3 per mm displacement. The latter could be measured with an accuracy of $\pm 0.01 \text{ mm}$. Then, the amount of ice could be estimated as a function of temperature by assuming that water expands 9.1% when it freezes.

Two cells were mounted on a frame with a precision device (metal scale and lens) for measuring the temperature-dependent displacement of the oil meniscus (Fig. 6).

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 4 Section through dilatometer.

1. Steel cylinder for clay sample and paraffine oil.
2. Glass pipe for measuring the volume of extruded paraffine oil.
3. Insulation of plastic with copper needle for ice nucleation.




Fig. 5 The dilatometer. Left picture shows assembled device. Note the cup and needle for liquid air cooling (ice nucleation) in the right picture which shows the various components.

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 6 Frame with two cells ready to be submerged in the methyl alcohol bath.

The frame was then submerged in a Hetofrig CB-6 cooling device with methyl alcohol in its cooling chamber. Insulation material was applied around the glass pipe and the metal pieces extending from the chamber so that a uniform temperature of all the parts of the measuring device was obtained.

Program

The temperature was lowered stepwise to ± 0 , -1 , -2 , -3 , -4 , -5 , -6 , -7 , -8 , -9 , and -10°C and was then raised again in 1°C steps until $\pm 0^{\circ}\text{C}$ was reached and the samples thawed. Equilibrium after each temperature change was obtained within 3-6 hours.

Tests were made with undisturbed as well as with remoulded samples. To avoid the entrance of air into the samples at

CLAY MICROSTRUCTURE AND ICE NUCLEATION

the remoulding they were wrapped in thin plastic bags. They were then pressed and gently kneaded.

In all the tests ice nucleation was initiated at 0°C by cooling a very fine copper needle which penetrated the lid and a few millimeters of the clay sample. The needle, which was isolated from the metal cell by a plastic plug, was equipped with a small cup in which liquid air was poured. It was found that the temperature at the point of the needle decreased from room temperature to about -70°C almost instantaneously at the application of liquid air (cf. Figs. 4 and 5).

Accuracy

There are several possible errors which affect the accuracy of the measured content of unfrozen water. The main ones are:

1. The temperature of the methyl alcohol bath cannot be kept at the desired temperature with an accuracy better than $\pm 0.1^\circ\text{C}$.
2. The behaviour of the paraffine oil at its contact with the clay at sub-zero temperatures is not known. There may be a zone where water and oil form a mixture the temperature/volume properties of which are not known.
3. The ice formed in the clay has a density which is not a constant. It contains unfrozen water to an unknown extent and its density can therefore vary within a considerable interval.

The last-mentioned point is most important and it was concluded that the amount of unfrozen water can be determined with an accuracy of no better than about $\pm 10\%$ when applying the technique described here. This was considered to be sufficiently accurate for the present purpose. It was decided, however, to check the results by applying a different technique. This check, which is described later in the text, confirmed the main conclusions from the dilatometer tests.

Test results

Fig. 7 shows the unfrozen water content (w_u =ratio of unfrozen water mass and mass of solid substance) as a function of temperature for undisturbed samples.

The two curves show the relationship for a successively lowered temperature (0°C and lower). We see that the unfrozen water content of the Skå-Edeby clay is considerably higher than that of the Lilla Edet clay. Thus, the hypothesis that the "dispersed" fresh-water clay should have a higher percentage of unfrozen water than the leached marine clay is valid.

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 7 Undisturbed Skå-Edeby (+) and Lilla Edet (●) samples.

This hypothesis implies that if a structural change is caused so that the two clays become identical with respect to the degree of aggregation, they will also have the same amount of unfrozen water. Such a structural change can be obtained by remoulding (Fig. 8). Fig. 9, which shows the results from tests with remoulded samples, confirms that practically identical amounts of unfrozen water are obtained.




Fig. 8 Schematical picture of the microstructural change induced by large shear strain or remoulding (ARNOLD, 1967).

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 9 Remoulded Skå-Edeby (+) and Lilla Edet (●) samples.

The explanation is of course that the remoulding produces particle dispersion which increases the mineral surface exposed to pore water and thereby also the amount of unfrozen water just as we can see from Fig. 10.




Fig. 10 Lilla Edet samples. U = undisturbed, R = remoulded.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

In fresh-water clays with a high degree of particle dispersion, remoulding may improve the dispersion a little but it will also create a certain number of larger continuous pores. Since the majority of the water in such pores is readily frozen the net effect will be a slightly reduced amount of unfrozen water in the vicinity of 0°C (Fig. 11).




Fig. 11 Skå-Edeby samples. U = undisturbed, R = remoulded

The consolidation effects are very well illustrated by the hysteresis loops in Fig. 12. This picture shows that when temperature is lowered from -1°C to -10°C and then increased again to -1°C , the unfrozen water was reduced to about half the original amount. This is because consolidation produced zones or regions which behaved as very big, dense and fairly "dry" particle aggregates between ice accumulations. The water transported to freezing zones in connection with the formation of consolidated big aggregates (Fig. 13) is not readily brought back into the dense aggregates when the temperature increases and a net decrease in the amount of adsorbed water is therefore obtained.

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 12 Lilla Edet samples frozen to -10°C and then heated to 0°C . U = undisturbed, R = remoulded.




Fig. 13 Local consolidation produced by freezing.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

Spin echo technique test

Pulsed NMR technique has been applied at the U S Army Cold Regions Research and Engineering Laboratory (CRREL), Hanover, New Hampshire, USA, for several years and a practical method of determining the phase composition of frozen soil (unfrozen water content vs temperature) has been described recently by TICE, BURROUS and ANDERSON (1978). Samples of the Skå-Edeby and Lilla Edet clays were sent to CRREL¹⁾ for determination of the amount of unfrozen water.

Representative results of the CRREL study are given in Tables 4 and 5 together with the author's results.

Table 4 Unfrozen water content of undisturbed Skå-Edeby clay

Temperature °C	w _u % CRREL	w _u % AUTHOR
-1.0	47	57
-2.0	31-35	38
-3.0	26-34	26
-4.0	22-24	22
-5.0	20-21	18
-6.0	18-19	16

Table 5 Unfrozen water content of undisturbed Lilla Edet clay

Temperature °C	w _u % CRREL	w _u % AUTHOR
-1.0	36	47
-2.0	25	24.5
-3.0	20-21	18
-4.0	17-18	15
-5.0	15	12
-6.0	14	11

¹⁾ part of a current research cooperation.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

Tables 4 and 5 show that the agreement is good for the w_u -values determined by the two methods, especially in the temperature interval -2 to -4°C . It is concluded also that the CRREL tests confirm the author's result that the Skå-Edeby clay has a higher amount of unfrozen water than the Lilla Edet clay at each specific temperature.

MEANS OF DETERMINING AND EXPRESSING THE DEGREE OF AGGREGATION

At least for scientific purposes it should be possible to apply a method used for classifying clay microstructure to determine and express the degree of aggregation (PUSCH, 1970). It requires that the clay is impregnated and that ultra-thin sections are cut by applying ultra-microtomy, and that electron micrographs (TEM) are taken.

Micrographs obtained from ultra-thin sections (300 Å) represent two-dimensional cross sections through the clay and this offers a possibility of defining microstructural parameters and applying simple statistical methods to describe the patterns. The sectioned pore area P in percent of the total area T of the micrograph is a suitable parameter, which can be termed "two-dimensional porosity". The measurement of these quantities is made by using drawn images of the micrographs. In these drawings the pore area is marked white while particles and aggregates are marked black regardless of their density (Fig. 14).

For homogeneous clay it has been shown that this parameter, determined by using 5-10 micrographs, is representative of volumes of several cubic centimeters although the analyzed sections are very small. The very obvious difference in the degree of aggregation seen in Fig. 14 is illustrated by the P/T -values. Thus, fig. 15 shows P/T for the two salinity regions which are representative of the Skå-Edeby (lower histogram) and Lilla Edet (upper histogram) clays.

There is no simple relationship between P/T , the specific surface area and the amount of unfrozen water. It is reasonable to believe, however, that P/T or some function of it can be used to correct the specific surface area which is a basic measure of the unfrozen water content. The matter requires additional work before a reliable correction factor can be suggested, however.

CLAY MICROSTRUCTURE AND ICE NUCLEATION




Fig. 14 Micrographs (upper line) and drawn images of Skå-Edeby (left) and Lilla Edet clay samples.




Fig. 15 P/T versus salinity of the water in which the clays were formed. The histograms were based on several hundred micrographs.

CLAY MICROSTRUCTURE AND ICE NUCLEATION

CONCLUSIONS

The investigation shows that the content of unfrozen water is larger when the particle aggregates are small and fairly porous but closely spaced than if the aggregates are large and dense and separated by large pores. This is because the microstructural constitution determines to what extent the integrated mineral surface is exposed to the pore water. Thus, the specific surface area is not a sufficient measure of the content of unfrozen water. It should be corrected with reference to the degree of aggregation.

A practical consequence of this investigation is that determination of the content of unfrozen water of structured soils must be made by using soil samples with a defined state: undisturbed or remoulded. The latter includes all cases where the soil has a freeze history (previously frozen and thawed etc.)

REFERENCES

- ANDERSON, D.M. & TICE, A.R., Predicting unfrozen water contents in frozen soils from surface area measurements. Highway Research Record, no 393, 1972.
- ARNOLD, M., A study of thixotropic action in bentonite clay. Thesis, Faculty of Engineering, University of Adelaide, Australia, 1967, pp. 13-15.
- BARDEN, L., Macro- and microstructure of soils. Int. Symp. on Soil Structure, Gothenburg 1973, Appendix, pp. 24-26.
- DILLON, H.B. & ANDERSLAND, O.B., Predicting unfrozen water contents in frozen soils. Can. Geot. J., Vol. 3, No. 2, 1966, pp.53-60.
- HANSBO, S., Consolidation of clay, with special reference to influence of vertical sand drains. Swed. Geot. Inst. Proc. No. 18, 1960, pp. 48-49.
- MADSEN, F.T., Surface area measurements of clay minerals by glycerol sorption on a thermobalance. Thermo-

CLAY MICROSTRUCTURE AND ICE NUCLEATION

- chimica Acta, No. 21, 1977,
pp. 89-93.
- PUSCH, R., Clay microstructure. Document
D8:1970 Nat. Swed. Build. Res.
1970.
- PUSCH, R., Ice formation in clays with
special reference to their mic-
rostructural constitution. Proc.
Int. Symp. on Frost Action in
Soils. Luleå, Sweden Feb. 16-18
1977, Vol. 1 pp. 137-142.
- TICE, A.R., BURROUS, C.M., & Determination of unfrozen water
ANDERSON, D. M., in frozen soil by pulsed nuclear
magnetic resonance. 3rd Int.
Conf. on Permafrost, Edmonton,
Alberta, Canada. 1978 (in press).