

AD-A057 940

OKLAHOMA UNIV HEALTH SCIENCES CENTER OKLAHOMA CITY
'ESCHERICHIA COLI' SHOCK IN THE BABOON AND THE RESPONSE TO ADRE--ETC(U)
JUN 78 L B HINSHAW, J J COALSON, B A BENJAMIN N00014-76-C-0229
TR-127

F/8 6/13


UNCLASSIFIED

NL

| OF |
AD
A057940



END
DATE
FILMED
10-78
DDC



ADA 057940

AD NO.
DDC FILE COPY

LEVEL #

9
5c

OFFICE OF NAVAL RESEARCH
CONTRACT N00014-76-C-0229
PROJECT NO. NR 207-040

TECHNICAL REPORT NO. 127

**ESCHERICHIA COLI SHOCK IN THE BABOON
AND THE RESPONSE TO ADRENOCORTICOSTEROID TREATMENT**

L. B. Hinshaw, J. J. Coalson, B. A. Benjamin, L. T. Archer,
B. K. Beller, D. R. Kling, E. M. Hasser, R. W. Phillips

Prepared for Publication

in

Surgery, Gynecology, & Obstetrics

University of Oklahoma Health Sciences Center
Departments of Physiology & Biophysics, Pathology,
Surgery, and Gynecology & Obstetrics
Oklahoma City, Oklahoma

12 June 1978

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

Reproduction in whole or in part is permitted for
any purpose of the United States Government

DDC
RECEIVED
AUG 17 1978
A

78 07 13 033

OFFICE OF NAVAL RESEARCH

CONTRACT NO. 15 00014-76-C-0229

PROJECT NO. NR 207-040

9 TECHNICAL REPORT, NO. 127

6 'ESCHERICHIA COLI' SHOCK IN THE BABOON
AND THE RESPONSE TO ADRENOCORTICOSTEROID TREATMENT.

10 L. B. Hinshaw, J. J. Coalson, B. A. Benjamin, L. T. Archer
B. K. Beller, O. R. Kling, E. M. Hasser, R. W. Phillips

12 34 p. Prepared for Publication 14 TR-127

in

Surgery, Gynecology, & Obstetrics

407 464
University of Oklahoma Health Sciences Center
Departments of Physiology & Biophysics, Pathology,
Surgery, and Gynecology & Obstetrics
Oklahoma City, Oklahoma

11 12 Jun 78

Reproduction in whole or in part is permitted for
any purpose of the United States Government

UNCLASSIFIED	White Section	<input checked="" type="checkbox"/>
UNCLASSIFIED	Diff Section	<input type="checkbox"/>
UNCLASSIFIED		<input type="checkbox"/>
BY: <i>[Signature]</i>		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		

78 07 13 033

407 464

elt

The incidence of septic shock has increased significantly in the United States with the occurrence of over 100,000 deaths per year according to McCabe (27). Much effort has been directed toward the development of experimental animal models more closely resembling the clinical situation. The dog administered live *Escherichia coli* organisms has proven to be a useful shock model, as reported by Archer (1) and Griffiths (11), Groves (12) and Hinshaw (15) and their colleagues. Recently, however, emphasis has been placed on the nonhuman primate as an experimental septic shock model. Coalson (6), Cryer (10), Herman (13), Hinshaw (17,19) and Horwitz (22) and their co-workers have been instrumental in developing experimental procedures with baboons receiving lethal intravenous administrations of live *Escherichia coli* organisms. Results have demonstrated marked hemodynamic, respiratory, metabolic, hematologic and morphologic abnormalities bearing distinct resemblances to the human entity of septic shock. Attempts to treat this latter model, however, have been uniformly unsuccessful in terms of preventing the lethal endpoint of shock.

Corticosteroid administration in endotoxic, live *Escherichia coli*, or septic shock has yielded conflicting results. The studies of Herman (13) and Pingleton (29) and associates have been negative, whereas benefits have been shown by Schumer (35) and Hinshaw (14,16), Holtzman (21), Latour (24), Pitcairn (30), Rao (32), Sambhi (33), Schuler (34), Vaughn (37) and White (38) and their co-workers. Improvements in cardiac output, regional blood flow and metabolism, prevention of generalized intravascular clotting, and increased survival rate in these studies have been demonstrated by treatment with large doses of adrenocorticosteroids.

The purpose of the present study is two-fold; first, the response of the awake baboon to a slow infusion of *Escherichia coli* organisms will be examined

in order to evaluate its effectiveness as a model in approximating the condition of the human patient. Second, the effectiveness of the corticosteroid, methylprednisolone sodium succinate, administered alone as a treatment for septic shock in the unanesthetized baboon, will be evaluated with respect to various hemodynamic, metabolic and hematologic parameters, tissue morphology and survival.

METHODS

Experiments were carried out on 12 healthy baboons, *Papio anubis*, weighing between 9.6 and 19.1 kg. Animals were fasted overnight and given water *ad libitum*. The following morning they were restrained with a squeeze cage device and administered ketamine hydrochloride (Ketaset; Bristol Laboratories, Syracuse, N.Y.), 14 ± 0.5 mg/kg intramuscularly. Femoral vessels were exposed and cannulated in one hind limb for pressure measurements, fluid, live organism and steroid administration, and blood sampling. Animals were positioned comfortably in a primate restraining chair, inclined at 45° , and a Tele-Thermometer probe was inserted in the rectum. A 3-hour equilibration period post-ketamine was allowed to insure that animals were awake, had reached a steady state, and exhibited a quiet behavior. During the period of equilibration, saline was infused intravenously at the rate of 3 ml/kg/hour to satisfy minimum body fluid requirements, but infusions were decreased if the hematocrit fell excessively.

Mean arterial pressure and heart rate were monitored on a Sanborn recorder. Arterial blood samples were taken for control determinations of plasma glucose, serum insulin, hematocrit (Hct), leukocyte concentration, differential leukocyte concentrations, pH, lactate, sodium (Na^+), potassium (K^+), serum cortisol, serum glutamic pyruvic transaminase (SGPT) and arginase. Glucose, pH, Hct,

and leukocyte concentrations were recorded hourly while other parameters were collected at predetermined intervals.

Animals were paired in each group by infusing comparable intravenous doses of *Escherichia coli* organisms prepared as previously reported (18,19) and administered during a 5-hour period with one member of each pair receiving methylprednisolone sodium succinate (MP) (Solu-Medrol; The Upjohn Company, Kalamazoo, Mich.). Three groups were studied and mean *Escherichia coli* doses were as follows: Group A (N=4), 2.9×10^{10} ; Group B (N=2), 1.9×10^{10} ; and Group C (N=5), 5.6×10^{10} organisms/kg. Animals in Group B were administered one intramuscular dose of gentamicin sulfate (Garamycin; Schering Pharmaceutical Corp., Kenilworth, N.J.), 4.4 mg/kg, at 1.5 hours after the total *Escherichia coli* dose was administered. One additional baboon was administered saline in place of organisms and continuously observed for a 24-hour period. A summary of the *Escherichia coli* doses infused and therapy instituted is shown in Table I. Animals were continuously observed for a 24-hour period or until death. Tissues were removed for evaluation following nembutal administration from survivors at the 24th hour and from nonsurvivors when mean blood pressures fell to 25 mmHg, in order to eliminate tissue damage due solely to hypoxia. Specimens of heart, liver, kidney, adrenal, lung, intestine, skin and pancreas were obtained for morphologic evaluations and are the subject of a separate report (7).

Plasma glucose was determined by a glucose analyzer with an accuracy of ± 3 mg%, serum insulin was obtained by radioimmunoassay and pH was determined by a blood gas analyzer, as previously reported (18,19). Total leukocyte concentrations were measured with an automatic particle counter (Coulter ZF; Hialeah, Florida) and differential leukocyte concentrations by microscopic examination of blood smears stained with Wright's stain. Arginase concen-

trations were determined using "Chem assay" kits (Pitman-Moore, Inc.; Washington Crossing, N.J.) and SGPT levels were measured using a Serosonic test kit (Mallinckrodt, Inc.; St. Louis, Mo.). Sodium and potassium concentrations were determined with an Instrumentation Laboratories Model 143 Flame Photometer. Cortisol concentrations were measured by radioimmunoassay (23) and lactate concentrations were determined with perchloric acid, glycine-hydrazine buffer, NAD^+ and lactic dehydrogenase.

Baboons given MP received bolus injections of 30 mg/kg at 15 minutes after initiation of *Escherichia coli* infusion and subsequent 2-hour infusions of 15 mg/kg at 2-hour intervals. The initial MP bolus injection, given at 15 minutes, was considered essential since by that time a large number of organisms had been infused (2.0×10^9 organisms/kg body weight). Maintenance infusions were considered essential because of the short half-life of MP, determined to be 3-4 hours by DiSanto (The Upjohn Company; Kalamazoo, Mich.).

RESULTS

Figures 1A, 1B, 1C and 1D illustrate the responses of four baboons given a mean dose of 2.9×10^{10} live *Escherichia coli* organisms per kg (designated as low dose--Group A). Animals 1 and 2, 5 and 6 are paired and all are given comparable doses of *Escherichia coli* organisms with baboons 2 and 6 being administered methylprednisolone (MP). Figure 1A shows that mean systemic arterial pressure (MSAP) responses are variable with baboon 1 dying acutely with a rapidly developing hypotension and animals 5 and 6 surviving but exhibiting progressively decreasing pressures. Animal 2 died at 18 hours with the last measured parameters recorded at 8 hours. Blood glucose concentrations decreased whether the response was rapid, as with baboons 1 and 2, or prolonged, as with animals 5 and 6. The last recorded glucose values in

all animals varied between 21 and 55 mg%. Insulin values usually decreased and terminal hypoinsulinemia was seen in all baboons. Figure 1B shows that hematocrits were relatively stable except those of baboon 5, which decreased. Animals demonstrated maximal increases in temperature (T_R) 2 to 6 hours following initiation of *Escherichia coli* infusion. Arterial pH in each animal was relatively constant during the total observation period. Arginase concentrations increased sharply in animal 1 while SGPT values rose markedly in baboons 2 and 6, at 8 and 24 hours respectively (Figure 1C). Arterial potassium concentrations rose preterminally in all animals to values not exceeding 7 mEq/L, and sodium concentrations increased within 8 hours in baboons 1 and 2. Since the administered steroid MP is included in the adrenocortical assay procedure, concentrations are seen to be an order of magnitude higher in steroid-treated baboons compared with non-treated animals (Figure 1D). MP treatment did not alter survival rate in baboons receiving the lower dose of organisms and did not change any parameter measured in the untreated animals.

Figures 2A, 2B, 2C and 2D (Group B) show individual responses of baboons administered a mean dose of 1.9×10^{10} organisms/kg and one intramuscular dose of gentamicin (baboons 3 and 4), and one control baboon (animal 7) given saline only. Baboon 3 in this group was treated with MP. All three animals survived the 24-hour observation period. MSAP and blood glucose values were relatively constant for 24 hours although animal 4 had the lowest pressures and glucose values. Arterial insulin concentrations were variable, and baboon 4 had a markedly high insulin value of 140 μ U/ml at 15 hours after *Escherichia coli* infusion (Figure 2A). Figure 2B shows that Hct, T_R and pH values were relatively constant in all animals, while Figure 3A shows that the SGPT level

was markedly elevated at 24 hours in baboon 4. Arginase and SGPT concentrations remained low in baboon 3 administered *Escherichia coli*, MP and gentamicin, as was also seen in the saline control. Sodium levels rose abruptly in both *Escherichia coli*-treated animals while potassium concentrations increased only in baboon 4 (3.5 to 6.0 mEq/L). Both Na^+ and K^+ levels were constant in the control baboon for the 24-hour period. Glucocorticoid levels were approximately ten times higher in the MP-treated animal (baboon 3) compared with baboons 4 and 7 not receiving the steroid (Figure 2D).

Figures 3A, 3B, 3C and 3D demonstrate individual responses of Group C baboons infused with a mean of 5.6×10^{10} organisms/kg, a dose approximately twice as great as that in Group A, animals 9, 10 and 11 receiving MP (45-60 mg/kg). All animals died within 11 hours after the initiation of organism infusion. Three baboons (8, 9 and 10) showed progressively developing hypotension while baboons 11 and 12 demonstrated a rapid onset of severe hypotension (Figure 3A). Blood glucose levels are seen to fall continuously, after an initial hyperglycemia, to lethal hypoglycemic values in four animals (7 to 46 mg%). In contrast, baboon 10 died in 7 hours with a marked hyperglycemia (151 mg%). Arterial insulin values decreased rapidly in this group in all but one baboon, and terminal hypoinsulinemia was observed in all animals (0-8 $\mu\text{U}/\text{ml}$). Baboon 9 exhibited a blood glucose value of 28 mg% and a 5.5 $\mu\text{U}/\text{ml}$ insulin concentration at 4 hours following onset of *Escherichia coli* administration. Arterial pH values were relatively constant in these baboons except for terminal decreases in three animals and alkalosis in one animal. Hematocrit and temperature responses were variable although hemoconcentration was not observed in any animal (Figure 3B). Figure 3C illustrates marked increases in SGPT and arginase in baboon 10 and elevations of SGPT in animal

12. Arterial potassium concentrations rose in all animals to values not exceeding 6 mEq/L, while sodium concentrations were variable. Baboon 12 exhibited a decrease in plasma sodium concentration from 163 to 136 mEq/L in 11 hours. Figure 3D demonstrates that arterial glucocorticoid values were approximately ten times higher in steroid-treated animals.

Table II shows the effects of live *Escherichia coli* organisms on arterial blood lactate concentrations. Although lactate levels rose in all *Escherichia coli*-treated animals, in general increases are more marked in the baboons who died. Although there is an unexplained rise in lactate in the control animal at 5 hours, the subsequent values in this animal were relatively normal.

The average accumulated dose of *Escherichia coli* organisms administered prior to the first injection of MP was 2.0×10^9 organisms/kg in Groups A and C. Paired baboons, whether given *Escherichia coli* alone or *Escherichia coli* with steroid (MP), demonstrated no actual differences in physiologic, metabolic or morphologic findings. The pathologic findings in liver, kidney and adrenals are striking in all of the experimental animals. Lesions consisting of fibrin thrombi, edema and/or hemorrhage, and necrosis are present (7). The saline control showed no pathologic aberrations. All baboons except the control became markedly leukopenic, and similar degrees of leukopenia were observed from 2 to 15 hours in animals both with and without methylprednisolone. Differential cell evaluations were carried out and demonstrate a severe degree of neutropenia, including mature and immature cells, in all animals but the control. Small decreases in lymphocyte, monocyte and eosinophil concentrations were also noted from 2 to 15 hours after the onset of *Escherichia coli* infusion. The control baboon showed stable differential cell concentrations during the 24-hour observation period.

DISCUSSION

The present study was undertaken to evaluate the suitability of the unanesthetized baboon for studies relevant to human septic shock. Previous reports from this laboratory (6,17,19) have assayed hemodynamic, metabolic and morphologic responses of the anesthetized baboon to administrations of live *Escherichia coli* organisms or *Escherichia coli* endotoxin. The purpose of the present study was (a) to determine the responses of the unanesthetized instrumented baboon to 5-hour infusions of live *Escherichia coli* organisms at two dose ranges continuously observed during a 24-hour period or until death; and (b) to determine if treatment with pharmacological doses of the corticosteroid, methylprednisolone sodium succinate (MP), following infusions of *Escherichia coli* organisms is effective in improving the hemodynamic, metabolic, hematologic, morphologic and survival status of the animal. A 5-minute bolus injection of MP was given following the administration of 2.0×10^9 organisms/kg body weight, followed by infusions of MP at regular intervals in order to maintain its concentration at effective levels (half-life of MP, 3-4 hours). Administrations of MP were given beyond the 5-hour period in which the average infused organism quantity was 4.0×10^{10} organisms/kg, until 24 hours or death. No differences in physiologic, metabolic, hematologic or morphologic parameters were observed between treated and untreated animals, nor was survival time influenced by steroid administration. Since animals were carefully matched between treated and untreated groups, small improvements following steroid administration should have been detected. Since such was not the case, the reasons for the ineffectiveness of the steroid are in question.

Recent studies conducted by White and associates in this laboratory (38) have demonstrated clear protection against endotoxin shock by utilizing MP in dogs with a treatment regimen very similar to that used in the present study. White's group (38) showed that MP treatment prevented hypoglycemia, hemoconcentration and liver pathology, achieving an 80% survival rate with MP administration in LD₁₀₀ endotoxin shock. The findings of Pitcairn (30) and Schumer (35) and associates in animals given *Escherichia coli* and in septic patients demonstrated significant protection against shock when MP was combined with antibiotic therapy.

Systemic hypotension would be expected to be lessened by the cardiovascular actions of MP, reported to include increased cardiac output by Sambhi and colleagues (33), increased inotropic support of the heart by Rao and Cavanagh (32), increased coronary blood flow by Hinshaw and associates (16), and increased regional blood flow by Vaughn (37) and Hinshaw (16) and co-workers. However, MP did not prevent systemic hypotension in the present study.

Hypoglycemia has been reported in canine live *Escherichia coli*-induced shock by Archer (1) and by Griffiths (11) and Groves (12) and their associates; in baboons receiving live *Escherichia coli* by Hinshaw and co-workers (17,19); and in human septic shock by Yeung (39) and Berk (2), Rackwitz (31) and their associates. Problems of glucose metabolism have been suggested in clinical shock cases by Cowley and others (9). The hypoglycemia of canine live *Escherichia coli* shock has been ascribed to impaired gluconeogenesis by Groves and others (12). Liver dysfunction and abnormal hepatic morphology were observed in the present study. MP administration would be expected to alleviate the hypoglycemia in baboons inasmuch as Berry (3) and Holtzman and colleagues (21) have found that corticosteroid administration stimulates

hepatic gluconeogenesis and Schuler and associates (34) have reported that MP supports carbohydrate metabolism in shocked nonhuman primates. MP utilized in the present study, however, did not intensify the degree of early hyperglycemia, nor did it alleviate the subsequently observed hypoglycemia.

Hypoinsulinemia, prominently demonstrated in the present study, was also uninfluenced by treatment with MP. Decreased insulin concentrations were observed in previous baboon shock studies by Hinshaw (17,19) and Cryer (10) and colleagues in lethal *Escherichia coli* septicemia in baboons. Clowes and co-authors (5) have observed hypoinsulinemia in patients with low cardiac outputs in septic shock, and hypoinsulinemia has been observed by Carey's group (4) in clinical hypovolemic shock. Hypoinsulinemia has been proposed to result from the combined effects of alpha adrenergic suppression of insulin release from the pancreas as reported by Cryer and associates (10) and diminished pancreatic blood flow in shock as shown by Lau and colleagues (24). Insulin resistance may be an added complication although its documentation in *Escherichia coli* shock is lacking.

Abnormal intravascular coagulation defects, exemplified in the present study by fibrin thrombi deposition in liver, kidney and adrenals, has also been reported in previous reports in the baboon administered live *Escherichia coli* by Coalson (6), Hinshaw (17,19), Holcroft (20) and Horwitz (22) and their associates, and in patients subjected to septic shock as reported by Corrigan and Jordan (8), Matsuda and Shimada (26) and McGovern (28). Latour and co-workers (24) have reported that glucocorticoids prevent generalized intravascular clotting, and it would be expected that MP would alleviate the degree of fibrin thrombi deposition in the present study. That such did not occur is contrasted with recent findings that although heparin administration

prevents the formation of fibrin thrombi in baboons subjected to *Escherichia coli* shock (19) and prevents disseminated intravascular coagulation in septic shock patients as reported by Corrigan and Jordan (8), increased survival rate is not realized nor is survival time extended.

Sibbald and associates (36) have recently observed variations in adrenocortical responsiveness during sepsis. Although cortisol levels in septic patients were markedly elevated in some instances (from control values of 8-18 $\mu\text{g/dl}$ to 65 $\mu\text{g/dl}$), death occurred in each case. They propose that unrecognized adrenocortical insufficiency may be a critical factor in septic shock. Adrenal morphologic changes were striking in the present baboon study although cortisol concentrations were elevated above control in most instances. The effectiveness of these concentrations of cortisol and MP, however, may still not have been adequate in view of the findings of Sibbald and co-authors (36).

Since hemodynamic, metabolic, hematologic and morphologic defects were not prevented by methylprednisolone administration, it is not surprising that survival rate was not improved. However, steroid treatment alone in live *Escherichia coli* organism-induced shock in contrast to *Escherichia coli* endotoxin-induced shock may require other supportive measures in addition to MP, such as antibiotic administration.

SUMMARY

Recent studies from this laboratory have demonstrated that methylprednisolone sodium succinate (MP) increases survival rate in animals given LD₁₀₀ *Escherichia coli* endotoxin. The purpose of this study was to determine the effects of MP on the baboon infused with live *Escherichia coli* organisms.

Awake animals were paired by infusing comparable intravenous doses of *Escherichia coli* during a 5-hour period. Baboons given MP received bolus injections of 30 mg/kg at 15 minutes after initiation of *Escherichia coli* infusion and 2-hour infusions of 15 mg/kg at 2-hour intervals until death or for a 24-hour period. Mortality rate was unaltered by MP. Six of 7 animals dying became progressively hypoglycemic while hypoinsulinemia occurred in all animals within 6 hours and was sustained until death. Systemic hypotension was observed, although pressures were variable. Potassium and lactate concentrations increased while pH remained relatively constant in most animals. SGPT and arginase concentrations rose in most baboons dying in 18 hours. Morphologic studies revealed the presence of fibrin thrombi in the liver, kidney and adrenal tissue in most animals. No significant differences in physiologic, metabolic, hematologic or morphologic parameters were observed between treated and untreated animals.

REFERENCES

1. Archer, L. T. Hypoglycemia in conscious dogs in live *Escherichia coli* septicemia: A chronic study. *Circ. Shock*, 1976, 3: 93.
2. Berk, J. L., Hagen, J. F., Beyer, W. H., and Gerber, M. J. Hypoglycemia of shock. *Ann. Surg.*, 1970, 171: 400.
3. Berry, L. J. Metabolic effects of bacterial endotoxins. In: *Microbial Toxins*. Edited by S. Kadis, G. Weinbaum, and S. J. Ajl. Vol. V, pp. 165-208. New York: Academic Press, 1971.
4. Carey, L. C., Lowery, B. D., and Cloutier, C. T. Blood sugar and insulin response of humans in shock. *Ann. Surg.*, 1970, 172: 342.
5. Clowes, G. H. A., Jr., O'Donnell, T. F., Jr., Ryan, N. T., and Blackburn, G. L. Energy metabolism in sepsis: Treatment based on different patterns in shock and high output stage. *Ann. Surg.*, 1974, 179: 684.
6. Coalson, J. J., Hinshaw, L. B., Guenter, C. A., Berrell, E. L., and Greenfield, L. J. Pathophysiologic responses of the subhuman primate in experimental septic shock. *Lab. Invest.*, 1975, 32: 561.
7. Coalson, J. J., Benjamin, B. A., Archer, L. T., Beller, B. K., Spaet, R. H., and Hinshaw, L. B. *Escherichia coli* shock following adrenocorticosteroid treatment: Pathologic study in baboons. Submitted for publication.
8. Corrigan, J. J., Jr., and Jordan, C. M. Heparin therapy in septicemia with disseminated intravascular coagulation. Effect on mortality and on correction of hemostasis defects. *New Engl. J. Med.*, 1977, 283: 778.
9. Cowley, R. A., Attar, S., LaBrosse, E., McLaughlin, J., Scanlon, E., Wheeler, S., Hanashiro, P., Grumberg, I., Vitek, V., Mansberger, A., and Firminger, H. Some significant biochemical parameters found in 300 shock patients. *J. Trauma*, 1969, 9: 926.

10. Cryer, P. E., Coran, A. G., Sode, J., Herman, C. M., and Horwitz, D. L. Lethal *Escherichia coli* septicemia in the baboon: alpha-adrenergic inhibition of insulin secretion and its relationship to the duration of survival. *J. Lab. Clin. Med.*, 1972, 79: 622.
11. Griffiths, J., Groves, A. C., and Leung, F. Y. Hypertriglyceridemia and hypoglycemia in gram-negative sepsis in the dog. *Surg. Gynec. Obstet.*, 1973, 136: 897.
12. Groves, A. C., Woolf, L. I., O'Regan, P. J., Beach, C., Hasinoff, C., and Sutherland, W. H. Impaired gluconeogenesis in dogs with *E. coli* bacteremia. *Surgery*, 1974, 76: 533.
13. Herman, C. M., Oshima, O., and Erdos, E. G. The effect of adrenocorticosteroid pretreatment on kinin system and coagulation response to septic shock in the baboon. *J. Lab. Clin. Med.*, 1974, 84: 731.
14. Hinshaw, L. B., Solomon, L. A., Freeny, P. C., and Reins, D. A. Hemodynamic and survival effects of methylprednisolone in endotoxin shock. *Arch. Surg.*, 1967, 94: 61.
15. Hinshaw, L. B., Mathis, M. C., Nanaeto, J. A., and Holmes, D. D. Recovery patterns and lethal manifestations of live *E. coli* organism shock. *J. Trauma*, 1970, 10: 787.
16. Hinshaw, L. B., Archer, L. T., Black, M. R., and Greenfield, L. J. Effects of methylprednisolone sodium succinate on myocardial performance, hemodynamics and metabolism in normal and failing hearts. In: *Steroids and Shock*. Edited by T. M. Glenn. Pp. 253-273. Baltimore: University Park Press, 1974.
17. Hinshaw, L. B., Benjamin, B., Coalson, J. J., Elkins, R. C., Taylor, F. B., Jr., Price, J. T., Smith, C. W., and Greenfield, L. J. Hypoglycemia in lethal septic shock in subhuman primates. *Circ. Shock*, 1975, 2: 197.

18. Hinshaw, L. B., Benjamin, B., Archer, L. T., Beller, B., Coalson, J. J., and Hirsch, J. G. Physiopathologic responses of the rhesus monkey to live *Escherichia coli*. *Surg. Gynec. Obstet.*, 1974, 142: 893.
19. Hinshaw, L. B., Benjamin, B., Holmes, D. D., Beller, B., Archer, L. T., Coalson, J. J., and Whitsett, T. Responses of the baboon to live *Escherichia coli* organisms and endotoxin. *Surg. Gynec. Obstet.*, 1977, 145: 1.
20. Holcroft, J. W., Blaisdell, F. W., Trunkey, D. D., and Lim, R. C. Intra-vascular coagulation and pulmonary edema in the septic baboon. *J. Surg. Res.*, 1977, 22: 209.
21. Holtzman, S., Schuler, J., Earnest, W., Carbohydrate metabolism during endotoxemia. *Circ. Shock*, 1974, 1: 99.
22. Horwitz, D. L., Moquin, R. B., and Herman, C. M. Coagulation changes of septic shock in the sub-human primate and their relationship to hemodynamic changes. *Ann. Surg.*, 1972, 175: 417.
23. Kling, O. R., and Westphal, P. K. Steroid changes during the menstrual cycle of the baboon (*Papio cynocephalus*) and human. *Biol. Reprod.*, in press.
24. Latour, J. G., McKay, D. G., and Nasu, K. Prevention of the generalized Schwartzman reaction by glucocorticoids. *Am. J. Obstet. Gynecol.*, 1971, 113: 863.
25. Lau, T. S., Taubenfligel, W., Levene, R., Farago, G., Chan, H., Koven, I., and Drucker, W. R. Pancreatic blood flow and insulin output in severe hemorrhage. *J. Trauma*, 1972, 12: 880.
26. Matsuda, T., and Shimada, K. Shock in patients with disseminated intra-vascular coagulation (DIC). *Japan. Circ. J.*, 1977, 41: 376.
27. McCabe, W. R. Gram-negative bacteremia. *Adv. Intern. Med.*, 1974, 19: 135.
28. McGovern, V. J. The pathophysiology of gram-negative septicemia. *Pathology*, 1972, 4: 265.

29. Pingleton, W. W., Coalson, J. J., Hinshaw, L. B., and Guenter, C. A. Effects of steroid pretreatment on development of shock lung: Hemodynamic, respiratory, and morphologic studies. *Lab. Invest.*, 1972, 27: 445.
30. Pitcairn, M., Schuler, J., Erve, P. R., Holtzman, S., and Schumer, W. Glucocorticoid and antibiotic effect on experimental gram-negative bacteremic shock. *Arch. Surg.*, 1975, 110: 1012.
31. Rackwitz, R., Jahrmarker, H., Prechtel, K., Thiessen, K., and Grohmann, H. Hypoglykämie während kreislaufshock. *Klin. Wochenschr.*, 1974, 52: 605.
32. Rao, P. S., and Cavanagh, D. Endotoxin shock in the subhuman primate. *Arch. Surg.*, 1971, 102: 486.
33. Sambhi, M. P., Weil, M. H., and Udhoji, U. H. Acute pharmacodynamic effects of glucocorticoids: cardiac output and related hemodynamic changes in normal subjects and patients in shock. *Circulation*, 1967, 31: 523.
34. Schuler, J. J., Erve, P. R., and Schumer, W. Glucocorticosteroid effect on hepatic carbohydrate metabolism in the endotoxin-shocked monkey. *Ann. Surg.*, 1976, 183: 345.
35. Schumer, W. Steroids in the treatment of clinical septic shock. *Ann. Surg.*, 1976, 184: 333.
36. Sibbald, W. J., Short, A., Cohen, M. P., and Wilson, R. F. Variations in adrenocortical responsiveness during severe bacterial infections. Unrecognized adrenocortical insufficiency in severe bacterial infections. *Ann. Surg.*, 1977, 186: 29.
37. Vaughn, D., Kirschbaum, T., Bersentes, T., and Assali, N. S. Effects of corticosteroid hormones on regional circulation in endotoxin shock. *Proc. Soc. Exptl. Biol. Med.*, 1967, 124: 760.
38. White, G. L., Archer, L. T., Beller, B. K., and Hinshaw, L. B. Increased survival with methylprednisolone treatment in canine endotoxin shock. *J. Surg. Res.*, in press.
39. Yeung, C. Y. Hypoglycemia in neonatal sepsis. *J. Ped.*, 1970, 77: 812.

TABLE I. SUMMARY OF DOSES OF LIVE ESCHERICHIA COLI ORGANISMS, METHYLPREDNISOLONE SODIUM SUCCINATE (MP), GENTAMICIN, OR SALINE AND SURVIVAL TIMES IN BABOONS

Group	Baboon No.	Sex	Weight (kg)	E. coli (org./kg)	Methylprednisolone (mg total) (mg/kg)	Gentamicin (mg/kg)	Saline (ml total) (ml/kg)	Time of death (hr)
A	1	F	19.1	3.0×10^{10}			638	9
A	2*	F	12.6	2.4×10^{10}	737		486	18
A	5	F	12.6	3.4×10^{10}			900	Survivor
A	6	F	15.3	2.6×10^{10}	1610		1040	Survivor
B	3	F	16.2	1.7×10^{10}	1884	4.4	1520	Survivor
B	4	F	14.3	2.0×10^{10}		4.4	1500	Survivor
Control	7	M	15.5				960	Survivor
C	8	F	9.6	5.2×10^{10}			340	7
C	9	F	14.2	5.1×10^{10}	639		248	4
C	10	F	13.7	6.8×10^{10}	823		380	7
C	11	F	14.1	5.6×10^{10}	847		390	8
C	12	F	17.6	5.2×10^{10}			420	11

17

TABLE II. EFFECTS OF INFUSED LIVE ESCHERICHIA COLI ORGANISMS ON ARTERIAL BLOOD LACTATE CONCENTRATIONS IN BABOONS

Group	Baboon No.	Time during shock, hrs.															
		Control	2	4	5	6	7	8	10	11	15	20	24				
A	1	29.8	36.3	87.4		97.6		113.0 ^{d, 9*}									
A	2*	14.9	62.7	78.2		54.4		67.6 ^{d, 18}									
A	5	30.9	20.6		48.7				57.7	49.5							55.5
A	6	38.1	34.2		38.6				56.6	52.2	68.5						55.4
B	3	16.3	83.4		37.4				54.8	33.8	35.2						38.6
B	4	15.1	30.3		36.4				36.4	36.5	43.5						38.7
Control	7	+	11.9		47.0				9.7	7.4	6.4						8.2
C	8	5.0	22.2		40.4		42.0 ^{d, 7}										
C	9	9.1	37.6	136.0 ^{d, 4}													
C	10	11.6	47.9		80.6		223.0 ^{d, 7}										
C	11	20.8	30.7		50.6			97.6 ^{d, 8}									
C	12	12.2	22.5		25.1				26.0	48.4 ^{d, 11}							

*time of death

[†]observed for 8 hours

[‡]insufficient sample

Figures 1A, 1B, 1C and 1D (Group A):

Effects of methylprednisolone sodium succinate (MP) administration on baboons subjected to five-hour infusions of live *Escherichia coli* organisms (mean dose, 2.9×10^9 organisms/kg body weight).

*MP, 30 mg/kg, intravenously administered, 15 minutes following onset of *Escherichia coli* infusion.




FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 1D

Figures 2A, 2B, 2C and 2D (Group B):

Effects of methylprednisolone sodium succinate (MP) administration on baboons subjected to five-hour infusions of live *Escherichia coli* organisms (mean dose, 1.9×10^{10} organisms/kg body weight) and given gentamicin at 6.5 hours. One control animal received saline only.

*MP, 30 mg/kg intravenously administered, 15 minutes following onset of *Escherichia coli* infusion.

**Gentamicin, 4.4 mg/kg, intravenously administered, 1.5 hours following completion of *Escherichia coli* infusion.

FIGURE 2B





FIGURE 2A






FIGURE 2C




FIGURE 2D

Figures 3A, 3B, 3C and 3D (Group C):

Effects of methylprednisolone sodium succinate (MP) administration on baboons subjected to five-hour infusions of live *Escherichia coli* organisms (mean dose, 5.6×10^{10} organisms/kg body weight).

*MP, 30 mg/kg, intravenously administered, 15 minutes following onset of *Escherichia coli* infusion.




FIGURE 3A




FIGURE 3B




FIGURE 3C




FIGURE 3D

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 127 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ESCHERICHIA COLI SHOCK IN THE BABOON AND THE RESPONSE TO ADRENOCORTICOSTEROID TREATMENT	5. TYPE OF REPORT & PERIOD COVERED Technical Report	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) L. B. Hinshaw, J. J. Coalson, B. A. Benjamin, L. T. Archer, B. K. Beller, O. R. Kling, E.M. Hasser, R. W. Phillips (Colo. State Univ.)	8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0229 ✓	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Oklahoma Health Sciences Center Oklahoma City, Oklahoma	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, Virginia	12. REPORT DATE 12 June 1978	
	13. NUMBER OF PAGES 27	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Reproduction in whole or in part is permitted for any purpose of the United States Government		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <div style="border: 1px solid black; padding: 5px; display: inline-block;">DISTRIBUTION STATEMENT A Approved for public release Distribution Unlimited</div>		
18. SUPPLEMENTARY NOTES Prepared for publication in Surgery, Gynecology & Obstetrics		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) subhuman primate Escherichia coli shock adrenocorticosteroid treatment		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Recent studies from this laboratory have demonstrated that methylprednisolone sodium succinate (MP) increases survival rate in animals given LD ₁₀₀ Escherichia coli endotoxin. The purpose of this study was to determine the effects of MP on the baboon infused with live Escherichia coli organisms. Awake animals were paired by infusing comparable intravenous doses of Escherichia coli during a 5-hour period. Baboons given MP received bolus injections of 30 mg/kg at 15 minutes after (cont.) →		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

initiation of Escherichia coli infusion and 2-hour infusions of 15 mg/kg at 2-hour intervals until death or for a 24-hour period. Mortality rate was unaltered by MP. Six of 7 animals dying became progressively hypoglycemic while hypoinsulinemia occurred in all animals within 6 hours and was sustained until death. Systemic hypotension was observed, although pressures were variable. Potassium and lactate concentrations increased while pH remained relatively constant in most animals. SGPT and arginase concentrations rose in most baboons dying in 18 hours. Morphologic studies revealed the presence of fibrin thrombi in the liver, kidney and adrenal tissue in most animals. No significant differences in physiologic, metabolic, hematologic or morphologic parameters were observed between treated and untreated animals.

UNCLASSIFIED

OFFICE OF NAVAL RESEARCH
BIOLOGICAL SCIENCES DIVISION
BIOPHYSICS PROGRAM, Code 444
DISTRIBUTION LIST FOR TECHNICAL, ANNUAL AND FINAL REPORTS

Number of Copies

- (12) Administrator, Defense Documentation Center
Cameron Station
Alexandria, Virginia 22314
- (6) Director, Naval Research Laboratory
Attention: Technical Information Division
Code 2627
Washington, D. C. 20375
- (6) Office of Naval Research
Attention: Code 102IP (ONRL DOC)
800 N. Quincy Street
Arlington, Virginia 22217
- (3) Office of Naval Research
Biophysics Program
Code 444
Arlington, Virginia 22217
- (1) Commanding Officer
Naval Medical Research and Development Command
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Chief, Bureau of Medicine and Surgery
Department of the Navy
Washington, D. C. 20375
- (2) Technical Reference Library
Naval Medical Research Institute
National Naval Medical Center
Bethesda, Maryland 20014
- (1) Office of Naval Research Branch Office
495 Summer Street
Boston, Massachusetts 02210
- (1) Office of Naval Research Branch Office
536 South Clark Street
Chicago, Illinois 60605

- (1) Office of Naval Research Branch Office
1030 East Green Street
Pasadena, California 91106
- (1) Commanding Officer
Naval Medical Research Unit No. 2
Box 14
APO San Francisco 96263
- (1) Commanding Officer
Naval Medical Research Unit No. 3
FPO New York 09527
- (1) Officer in Charge
Submarine Medical Research Laboratory
Naval Submarine Base, New London
Groton, Connecticut 06342
- (1) Scientific Library
Naval Medical Field Research Laboratory
Camp Lejeune, North Carolina 28542
- (1) Scientific Library
Naval Aerospace Medical Research Institute
Naval Aerospace Medical Center
Pensacola, Florida 32512
- (1) Commanding Officer
Naval Air Development Center
Attn: Aerospace Medical Research Department
Warminster, Pennsylvania 18974
- (1) DIRECTOR
Naval Biosciences Laboratory
Building 844
Naval Supply Center
Oakland, California 94625
- (1) Commander, Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709
- (1) DIRECTOR OF LIFE SCIENCES
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, D. C. 20332

- (1) Commanding General
Army Medical Research and Development Command
Forrestal Building
Washington, D. C. 20314
- (1) Department of the Army
U. S. Army Science and
Technology Center - Far East
APO San Francisco 96328
- (1) Assistant Chief for Technology
Office of Naval Research, Code 200
800 N. Quincy Street
Arlington, Virginia 22217