

AD-A058 571

AEROSPACE CORP EL SEGUNDO CALIF IVAN A GETTING LABS F/G 7/2
EFFECT OF VIBRATIONAL EXCITATION IN REACTIONS OF OH WITH H₂.(U)
AUG 78 G C LIGHT, J H MATSUMOTO F04701-77-C-0078

UNCLASSIFIED

TR-0078(3970-40)-3

SAMSO-TR-78-78


NL

| OF /

AD
A058 571



END
DATE
FILMED
11-78
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL II

12
na


AD A058571

Effect of Vibrational Excitation in Reactions of OH with H₂

G. C. LIGHT and J. H. MATSUMOTO
Chemistry and Physics Laboratory
The Ivan A. Getting Laboratories
The Aerospace Corporation
El Segundo, Calif. 90245

15 August 1978

Interim Report



APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED

Prepared for
SPACE AND MISSILE SYSTEMS ORGANIZATION
AIR FORCE SYSTEMS COMMAND
Los Angeles Air Force Station
P.O. Box 92960, Worldway Postal Center
Los Angeles, Calif. 90009


78 09 07 024


DDC FILE COPY

This interim report was submitted by The Aerospace Corporation, El Segundo, CA 90245, under Contract No. F04701-77-C-0078 with the Space and Missile Systems Organization, Deputy for Advanced Space Programs, P. O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009. It was reviewed and approved for The Aerospace Corporation by S. Siegel, Director, Chemistry and Physics Laboratory. Lieutenant Arturo G. Fernandez, SAMSO/YCPT, was the project officer for Advanced Space Programs.


This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication. Publication of this report does not constitute Air Force approval of the report's findings or conclusions. It is published only for the exchange and stimulation of ideas.


Arturo G. Fernandez, Lt, USAF
Project Officer


Robert W. Lindemuth, Lt Col, USAF
Chief, Technology Plans Division

FOR THE COMMANDER


FLOYD R. STUART, Col, USAF
Deputy for Advanced Space Programs

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SAMSO TR-78-78	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) EFFECT OF VIBRATIONAL EXCITATION IN REACTIONS OF OH WITH H ₂ [†]	5. TYPE OF REPORT & PERIOD COVERED Interim Repts	
7. AUTHOR(s) Glenn C. Light and James H. Matsumoto	6. PERFORMING ORG. REPORT NUMBER TR-0078(3970-40)-3	
	7. CONTRACT OR GRANT NUMBER(s) F04701-77-C-0078	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Aerospace Corporation El Segundo, Calif. 90245	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Space and Missile Systems Organization Air Force Systems Command Los Angeles, Calif. 90009	12. REPORT DATE 15 August 1978	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 11	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 12 14 p.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Hydroxyl Molecule Hydrogen Molecule Chemical Reaction Vibrational Excitation Laser-Induced Fluorescence Detection Rate Constants		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Upper limits have been deduced at T ≈ 298 K for the rates of chemical reactions of hydroxyl OH(v = 0) and OH [†] (v = 1) with vibrationally excited hydrogen H ₂ [†] (v = 1). These are 3.5 × 10 ¹² and 6.2 × 10 ¹² cc/mol-sec, respectively.		

DD FORM 1473
(FACSIMILE)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

409 944

alt

PREFACE

The authors are glad to acknowledge stimulating discussions with Dr. R. R. Herm and Dr. R. J. McNeal during the analysis of the results.

Section ☒	
B II Section ☐	
☐	
DISTRACTION/AVAILABILITY CODES	
OF SPECIAL	
A	

REPORT

The following data is submitted for the purpose of the report.

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48	49	50
51	52	53	54	55	56	57	58	59	60
61	62	63	64	65	66	67	68	69	70
71	72	73	74	75	76	77	78	79	80
81	82	83	84	85	86	87	88	89	90
91	92	93	94	95	96	97	98	99	100

blank
2

CONTENTS

PREFACE	1
I. INTRODUCTION	5
II. EXPERIMENT	5
III. RESULTS AND ANALYSIS	6
IV. DISCUSSION	10
REFERENCES	13

FIGURE


1. Dependence of Relative Hydroxyl Density $\text{OH}(v = 0)$ and $\text{OH}^\dagger(v = 1)$ on Absolute Density of Vibrationally Excited Hydrogen $\text{H}_2^\dagger(v = 1)$	9
--	---

blank
4


I. Introduction

Recently, a measurement of the rate of the reaction between $O(^3P)$ and $H_2^{\dagger}(v = 1)$ has appeared in the literature,¹ and an upper limit to the rate of $OH^{\dagger}(v = 1) + H_2(v = 0)$ has been given.²

These reports are beginning to characterize the effects of vibrational energy in the diatomic molecules of the H_2/O_2 system. This work extends this beginning by reporting upper limits to the reaction rates for two reactions:




and




II. Experiment


The experimental apparatus has been fully described elsewhere.^{1,3} Briefly, the measurements were obtained in a laboratory flow tube apparatus at a total pressure of 3 Torr consisting of equal parts N_2 and H_2 at a temperature of 298 K. Vibrationally excited hydroxyl was produced entirely by reaction (3)



Vibrational ground state hydroxyl was produced predominantly by physical quenching of $OH^{\dagger}(v = 1)$. At lower concentrations of $H_2^{\dagger}(v = 1)$ there were minor contributions from reaction (4),



Atomic oxygen was produced by reaction (5), for which the N atoms were provided by a microwave discharge in N_2





The flow rate of nitric oxide was adjusted so that operation at the visible titration end point was achieved. Vibrationally excited hydrogen* was generated by the flow of hydrogen over a heated tungsten filament. The density of $H_2^\dagger$ was determined from spectral absorption measurements in the Lyman band system of hydrogen. Hydroxyl densities were deduced from laser induced fluorescence (LIF) in the ($A^2\Sigma \rightarrow X^2\Pi$) system of OH. Measurements were conducted under conditions where energy transfer^{1,3} from active nitrogen to form $H_2^\dagger$ was a negligible process in comparison with the formation of $H_2^\dagger$ on the heated filament.

The primary independent variable in these measurements was the density of $H_2^\dagger$. This was varied by changing the temperature of the tungsten heating element.






III. Results and Analysis

Fig. 1 shows data for the dependence of OH and $OH^\dagger$ densities on $[H_2^\dagger]$. It is seen that the density of OH is first order in $[H_2^\dagger]$ at the larger values of $[H_2^\dagger]$ but tends toward zeroth order as $[H_2^\dagger]$ decreases. The OH at low values of $[H_2^\dagger]$ comes from reaction (4). The density of $OH^\dagger$ behaves oppositely, being essentially first order in $[H_2^\dagger]$ at low values but tending toward zeroth order at higher values. The kinetics of this system have been modelled using a computer code to integrate the rate equations. Individual vibrational levels of $OH^\dagger$ ($v \leq 3$) and $H_2^\dagger$ were treated as separate entities with generation and removal processes individually specified. Analysis of the computer results showed the densities of OH and $OH^\dagger$ to be steady state with the most important loss processes for which rate constants or limits are available as follows; for OH:

* Hereinafter, the symbols $H_2^\dagger(v=1)$ and $OH^\dagger(v=1)$ will be simplified to $H_2^\dagger$ and $OH^\dagger$, respectively unless otherwise specified. The symbols H_2 and OH shall denote the vibrational ground state species only.



and for $\text{OH}^\ddagger$:



In addition, of course, it is necessary to consider the possibility that the reactions of interest, i. e., (1) and (2), contribute significantly. The important generation processes for OH were all found to be physical deactivation of $\text{OH}^\ddagger$. These are (8a), (9a), and (10). For $\text{OH}^\ddagger$, the important source terms are reactions (3) and (-9a).

Using the reaction scheme outlined one can form the steady state expressions for the densities of OH and $\text{OH}^\ddagger$. Taking logarithms and differentiating with respect to $[\text{H}_2^\ddagger]$, expressions (a) and (b) for the slopes of plots of $\log [\text{OH}(v)]$ versus $\log [\text{H}_2^\ddagger]$ are derived.

$$\frac{d(\log[\text{OH}^\ddagger])}{d(\log[\text{H}_2^\ddagger])} \equiv m^\ddagger = 1 - \left(\frac{k_2 + k_{10} \left(\partial[\text{H}] / \partial[\text{H}_2^\ddagger] \right)}{k_8[\text{O}]/[\text{H}_2^\ddagger] + k_9[\text{H}_2]/[\text{H}_2^\ddagger] + k_{10}[\text{H}]/[\text{H}_2^\ddagger] + k_2} \right) \quad (a)$$

$$\frac{d(\log[\text{OH}])}{d(\log[\text{H}_2^\ddagger])} \equiv m = m^\ddagger +$$

$$\left(\frac{k_{10} \partial[\text{H}] / \partial[\text{H}_2^\ddagger]}{k_{8a}[\text{O}]/[\text{H}_2^\ddagger] + k_{10}[\text{H}]/[\text{H}_2^\ddagger] + k_{9a}[\text{H}_2]/[\text{H}_2^\ddagger]} \right) - \left(\frac{k_1}{k_1 + k_7[\text{H}_2]/[\text{H}_2^\ddagger] + k_6[\text{O}]/[\text{H}_2^\ddagger]} \right) \quad (b)$$

The partial differential ($\partial[H]/\partial[H_2^\ddagger]$) appears in both expressions and requires some explanation. $[H]$ will depend on $[H_2^\ddagger]$ in two ways; first, directly as a result of reaction (3) followed by (6) through (10), and secondly, in an indirect way, since H_2 is thermally partially dissociated in contact with the heated tungsten filament. Analysis has shown that for large $[H_2^\ddagger]$ the second source of H atoms is considerably more important than the first if one assumes that both H and $H_2^\ddagger$ are present in amounts determined by equilibrium at the filament temperatures. Because of this and because the density of H is not measured in these experiments it is only possible to deduce upper limits to the rate constants k_1 and k_2 .

The analysis proceeds by determining experimental values of m and $m^\ddagger$ from the data in Fig. 1 at the maximum value of $[H_2^\ddagger]$ where data were obtained. These values of m and $m^\ddagger$ are then used in expressions (a) and (b) to find mutually consistent sets of values for $[H]$, $\partial[H]/\partial[H_2^\ddagger]$, and the unknown rate constants k_1 , k_2 , k_{9a} and k_{9b} . These unknown values were allowed to vary throughout their respective ranges of uncertainty. It was found that there were certain values of k_1 and k_2 which could never be exceeded irrespective of the values assigned to the other uncertain quantities in expressions (a) and (b). These values of k_1 and k_2 are necessarily upper limits and are the values reported herein.

In this analysis values for $[H]$ and $\partial[H]/\partial[H_2^\ddagger]$ were allowed to vary from zero upwards to the values dictated by thermal equilibrium at the filament temperature. However, the ratio of the terms was constrained by expression (c), which is a consequence only of the assumption of thermochemical equilibrium irrespective of temperature.

$$\partial[H]/\partial[H_2^\ddagger] = \frac{(\Delta H^\circ)_r}{2G_0(1)} \left(\frac{[H]}{[H_2^\ddagger]} \right) \quad (c)$$

where $(\Delta H^\circ)_r$ is the heat of reaction for dissociation of H_2 and $G_0(1)$ is the energy of the first vibrational state of H_2 above that of the ground state.





Fig. 1. Dependence of Relative Hydroxyl Density $OH(v = 0)$ and $OH^\dagger(v = 1)$ on Absolute Density of Vibrationally Excited Hydrogen $H_2^\dagger(v = 1)$. (Δ) $OH(v = 0)$; ($\odot$) $OH^\dagger(v = 1)$.

The value of k_{9a} was allowed to vary between zero and the upper limit reported by Spencer et. al.,⁽²⁾ i.e., $k_9 = k_{9a} + k_{9b} \leq 6 \times 10^9$ cc/mole-sec. In the case of k_{9b} however, it was felt that the minimum value should be that of the fully ground state reaction, (7), since vibrational excitation is unlikely to decrease that value. At room temperature, $k_7 = 3.6 \times 10^9$ which is not too different from the upper limit to k_9 . For purposes of this analysis therefore, k_{9b} was assumed to be fixed at 6×10^9 . Values for k_1 and k_2 were constrained only to be positive numbers.

From Fig. 1 we obtain $m^\ddagger = 0.60$, and $m \geq 0.8$, for $[H_2^\ddagger] = 4 \times 10^{-10}$ and 3×10^{-10} moles/cc, respectively. Consistent with these values of m and $m^\ddagger$ the largest value of k_2 which permits a positive value of k_1 for all possible values of $[H]$, $\partial [H]/\partial [H_2^\ddagger]$, and k_{9a} is: $k_2 \leq 6.2 \times 10^{12}$ cc/mole-sec. Likewise, the largest value of k_1 which permits a positive value of k_2 for all possible values of $[H]$, $\partial [H]/\partial [H_2^\ddagger]$ and k_{9a} is: $k_1 \leq 3.5 \times 10^{12}$ cc/mole-sec.

IV. Discussion

In the above analysis reaction of OH and deactivation and reaction of $OH^\ddagger$ at the wall are negligible processes. The time constants for homogeneous gas phase removal of OH is, $\tau \leq 1.6 \times 10^{-3}$ sec and of $OH^\ddagger$ is $\tau^\ddagger \leq 2.4 \times 10^{-4}$ sec. These are much faster processes than removal at the wall which has been measured by Spencer and Glass⁴ with a fluorocarbon wax such as used in the present experiments. Also, the homogeneous reaction (11) has been ignored in the above analysis.



It is similar to reaction (-9a) except that the vibrational energy exchange is further from resonance for (11) than it is for (-9a). Using the upper limit² for k_{9a} and the principle of microscopic reversibility it is necessary that

$$k_{(-9a)} \leq 10^{11} \text{ cc/mole-sec.}$$

It is not likely that reaction (11) would be greatly faster than this and so it is reasonable to ignore it in the above analysis.

The effect of vibrational excitation in OH on the reaction with H_2 was found by Spencer et. al.² to be minor, such that $(k_{9a} + k_{9b})/k_7 < 1.7$. We have not been able to measure or find lower limits for the effect of vibrational excitation in H_2 on this reaction. It seems likely that this reaction proceeds by breaking of the H_2 bond rather than of the OH bond since the latter course requires the insertion of O between H atoms in order to form water. If this is true one would expect that vibrational energy in hydrogen would be much more effective in promoting the reaction than would vibrational energy in OH. The results reported here and by Spencer et. al. are at least consistent with this view.

Our results indicate that $k_1/k_7 \leq 1000$ and that $k_2/k_7 \leq 1800$ at $T = 298$ K. Thus, there is room still for appreciable acceleration of the reaction by one quantum of vibrational energy in H_2 or by one quantum in both OH and H_2 . Zellner⁵ has determined a preliminary result of $k_1 \approx (3.6 \pm 1.8) \times 10^{11}$ cc/mole-sec, corresponding to an increase of roughly a factor of 100 over the ground state reaction. This result is consistent with our upper limits and with the view sketched above.

From the upper limits for k_1 and k_2 reported here and from the upper limit for k_{9b} it is clear that one quantum of vibrational energy in either OH or in H_2 or in both simultaneously is not as effective in promoting the reaction as is the same specific energy equipartitioned between all modes of both reactants.

References

1. G. C. Light, J. Chem. Phys., 68, 2831, (1978).
2. J. E. Spencer, H. Endo, G. P. Glass, Proc. 16th Symposium (International) on Combustion, The Combustion Institute, Pittsburgh, Pa., 15 - 20 August, (1976).
3. G. C. Light, Proc. 17th Symposium (International) on Combustion, The Combustion Institute, Pittsburgh, Pa. 21 - 25 August, (1978), to be published.
4. J. E. Spencer, G. P. Glass, Chem. Phys. 15, 35 (1976).
5. R. Zellner, private communication.

THE IVAN A. GETTING LABORATORIES

The Laboratory Operations of The Aerospace Corporation is conducting experimental and theoretical investigations necessary for the evaluation and application of scientific advances to new military concepts and systems. Versatility and flexibility have been developed to a high degree by the laboratory personnel in dealing with the many problems encountered in the nation's rapidly developing space and missile systems. Expertise in the latest scientific developments is vital to the accomplishment of tasks related to these problems. The laboratories that contribute to this research are:

Aerophysics Laboratory: Launch and reentry aerodynamics, heat transfer, reentry physics, chemical kinetics, structural mechanics, flight dynamics, atmospheric pollution, and high-power gas lasers.

Chemistry and Physics Laboratory: Atmospheric reactions and atmospheric optics, chemical reactions in polluted atmospheres, chemical reactions of excited species in rocket plumes, chemical thermodynamics, plasma and laser-induced reactions, laser chemistry, propulsion chemistry, space vacuum and radiation effects on materials, lubrication and surface phenomena, photo-sensitive materials and sensors, high precision laser ranging, and the application of physics and chemistry to problems of law enforcement and biomedicine.

Electronics Research Laboratory: Electromagnetic theory, devices, and propagation phenomena, including plasma electromagnetics; quantum electronics, lasers, and electro-optics; communication sciences, applied electronics, semiconducting, superconducting, and crystal device physics, optical and acoustical imaging; atmospheric pollution; millimeter wave and far-infrared technology.

Materials Sciences Laboratory: Development of new materials; metal matrix composites and new forms of carbon; test and evaluation of graphite and ceramics in reentry; spacecraft materials and electronic components in nuclear weapons environment; application of fracture mechanics to stress corrosion and fatigue-induced fractures in structural metals.

Space Sciences Laboratory: Atmospheric and ionospheric physics, radiation from the atmosphere, density and composition of the atmosphere, aurorae and airglow; magnetospheric physics, cosmic rays, generation and propagation of plasma waves in the magnetosphere; solar physics, studies of solar magnetic fields; space astronomy, x-ray astronomy; the effects of nuclear explosions, magnetic storms, and solar activity on the earth's atmosphere, ionosphere, and magnetosphere; the effects of optical, electromagnetic, and particulate radiations in space on space systems.

THE AEROSPACE CORPORATION
El Segundo, California