

AD-A058 599

ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND ABERD--ETC F/G 7/4
A MODIFIED LEPS SURFACE FOR THE GROUND STATE OF THE WATER MOLEC--ETC(U)
JUL 78 A GAUSS

UNCLASSIFIED

ARBRL-TR-02089

SBIE-AD-E430 092

NL

| OF |

AD
A058599



END


DATE

FILMED

11-78

DDC





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A058599

AD No. _____
100 FILE COPY

(12) ^{NW} **LEVEL** II

AD-E430092

TECHNICAL REPORT ARBRL-TR-02089

A MODIFIED LEPS SURFACE FOR THE GROUND
STATE OF THE WATER MOLECULE

Arthur Gauss, Jr.

July 1978



US ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND
BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

Approved for public release; distribution unlimited.

78 08 16 039

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating
or sponsoring activity is prohibited.

Additional copies of this report may be obtained
from the National Technical Information Service,
U.S. Department of Commerce, Springfield, Virginia
22161.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

*The use of trade names or manufacturers' names in this report
does not constitute indorsement of any commercial product.*

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TECHNICAL REPORT, ARBRL-TR-02089	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Modified LEPS Surface for the Ground State of the Water Molecule		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Arthur/Gauss, Jr		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Ballistic Research Laboratory ATTN: DRDAR-BLB Aberdeen Proving Ground, MD 21005		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Armament Research & Development Command US Army Ballistic Research Laboratory ATTN: DRDAR-BL, APG, MD 21005		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS RDT&E/1L161102AH43
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) SBIE AD-E430 092		12. REPORT DATE JULY 1978
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 13
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
18. SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Potential energy surfaces, water molecule, LEPS surfaces, semi-empirical potential surfaces, adjustable Sato parameter, Sato parameter function of coordinates, potential energy well		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A Modified LEPS potential surface using a Sato parameter which is a function of the internuclear coordinates has been developed for the ground state of the water molecule. By adjusting the Sato parameter many of the known properties of the water potential are fitted with this surface.		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

393 471

08 16 039

DDC
RECEIVED
SEP 15 1978
B

TABLE OF CONTENTS

	Page
I. INTRODUCTION.	5
II. MODIFIED LEPS SURFACE FOR WATER	5
III. DISCUSSION.	12
DISTRIBUTION LIST	13

ACCESSION for		
NTIS	White Section	☒
DDC	Buff Section	☐
UNANNOUNCED		☐
JUSTIFICATION _____		
BY _____		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL. and/or	SPECIAL
A		

I. INTRODUCTION

In the Born-Oppenheimer approximation for molecular collision problems the electronic potential energy serves as the potential energy for motion of the molecular nuclei. To facilitate dynamics calculations for these problems an analytical potential surface is a very valuable tool. An analytical surface provides the necessary derivatives of the potential with respect to the internuclear coordinates over all space. An analytical surface must also be flexible enough to provide a reasonable fit to the known properties of the potential surface derived from experiment or ab-initio calculations. Many molecules or molecular complexes are characterized by having their minimum electronic potential energy in a bent configuration ($\theta \neq 0, 180^\circ$; see Figure 1). Such molecules, of which water is an example, cannot be modeled using a standard LEPS (London, Eyring, Polanyi, Sato) surface. In this paper a modified LEPS surface is presented which can model the bent configured water molecule. While not as accurate as some other analytical methods^{1,2} the modified LEPS surface is easier to apply to dynamics calculations. Modified LEPS surfaces have been developed for other systems, HO_2 and CO_2 , in addition to water. Dynamics calculations are already complete for the HO_2 system.

II. MODIFIED LEPS SURFACE FOR WATER

An LEPS surface which approximates the equilibrium ground state for water (X^1A_1) has been developed. A single Sato parameter which is a function of the internuclear coordinates is used to alter the LEPS so that the water molecule in and near the equilibrium configuration can be modeled. An earlier study³ suggested the use of large values of the Sato parameter to produce wells or barriers along the reaction path. The general form of the LEPS potential⁴ is

$$V = Q_1' + Q_2' + Q_3' - [\alpha_1'^2 + \alpha_2'^2 + \alpha_3'^2 - \alpha_1'\alpha_2' - \alpha_1'\alpha_3' - \alpha_2'\alpha_3']^{1/2} + D_2^e,$$

1. K.S. Sorbie and J.N. Murrell, *Molecular Physics* 29 (1975), 1387.
2. S. Farantos, E.C. Leisegang, J.N. Murrell and K. Sorbie, *Molecular Physics* 34 (1977) 947.
3. A. Gauss Jr., *J. Chem. Phys* 65 (1976) 4365.
4. J.T. Muckerman, *J. Chem. Phys.* 56 (1972) 2997.




Figure 1. The internuclear distances (r_1, r_2, r_3) and the angle θ are defined as shown.

where

$$Q_i' = Q_i / (1 + \Delta) = D_i^e / 4(1 + \Delta) \\ \times \left\{ (3 + \Delta) \exp - 2\beta_i(r_i - r_{i0}) \right. \\ \left. - (2 + 6\Delta) \exp - \beta_i(r_i - r_{i0}) \right\}$$

and

$$\alpha_i' = \alpha_i / (1 + \Delta) = D_i^e / 4(1 + \Delta) \\ \times \left\{ (1 + 3\Delta) \exp - 2\beta_i(r_i - r_{i0}) \right. \\ \left. - (6 + 2\Delta) \exp - \beta_i(r_i - r_{i0}) \right\}, \quad i = 1, 2, 3.$$

The Morse parameters (D_i^e , β_i , r_{i0}) are displayed in Table 1. The Sato parameter is given by

$$\Delta = A \{f(\theta)\} \{(\sin 2 \text{ Th})^m\} \\ \times \left\{ (0.94r_1r_2/r_{10}^2)^B \exp \left(-0.94(r_1r_2/r_{10}^2)^B \right) \right\},$$

where $\text{Th} = \text{Arctan}(r_1/r_2)$ and the parameters A, m , and B are adjustable.

Table 1.

O-H	H-H
$D_1^e = D_2^e = 106.53 \text{ kcal/mole}$	$D_3^e = 109.5 \text{ kcal/mole}$
$\beta_1 = \beta_2 = 2.29 \text{ \AA}^{-1}$	$\beta_3 = 1.942 \text{ \AA}^{-1}$
$r_{10} = r_{20} = 0.9706 \text{ \AA}$	$r_{30} = 0.7417 \text{ \AA}$

This form of the Sato parameter could be applied to other three-atom complexes where two of the atoms are identical. It has the proper symmetry; that is, it is invariant to interchange of r_1 and r_2 . A single Sato parameter has been employed here. Single Sato parameters have been used successfully by other workers on three-atom

systems where two of the atoms are identical^{4,5}. The Sato parameter ranges from near zero for $r_1 = r_{10}$, $r_2 \rightarrow \infty$ (or $r_2 = r_{20}$, $r_1 \rightarrow \infty$) to slightly greater than two near the minimum energy configuration ($r_1 = r_2 \approx 0.87 \text{ \AA}$, $\theta = 106^\circ$).

The function $f(\theta)$ contains additional adjustable parameters. In the case of the water molecule $f(\theta) = [\sin(\text{Th1})]^6 + \frac{3}{4} [\sin(\text{Th2})]^4$, where $\text{Th1} = \frac{6}{7} \theta$ and $\text{Th2} = \frac{1}{2} \theta$. The exponents, the constant multiplying the $[\sin(\text{Th2})]^4$ term, and the constants multiplying θ in Th1 and Th2 are the adjustable parameters. These parameters were varied so that the potential minimum was at $\theta = 106^\circ$, close to the actual angle of $104\text{--}105^\circ$. In addition $f(\theta)$ was adjusted so that the difference between the depth of the potential well in the linear configuration and the depth in the 106° configuration was $\approx 1.8 \text{ eV}$ (compare figs. 2 and 3), close to the 1.5 eV difference shown in the potential plot by Herzberg⁷.

The second bracketed term in Δ , $(\sin 2\text{Th})^m$, prevents the potential well from extending to large values of r_1 and r_2 along the entrance or exit valleys; it also helps eliminate the double minima near the center of the potential well (a problem when large (> 1) values of the Sato parameter are employed). A large value of the exponent, $m = 16$, was needed for the water potential surface.

The third bracketed term in Δ eliminates the large negative values of the potential that occur for small values of r_1 and r_2 when Δ is large. This term approaches zero as r_1 and r_2 approach zero causing the Sato parameter to approach zero and the potential (V) to form a high wall (see Figures 2,3 and 4). It goes to zero for large values r_1 and r_2 due to the exponential. By adjusting B one can ensure that the potential minimum will be near $r_1 = r_2 \approx r_{10}$. For the water molecule $B=4$ was chosen. For this H_2O surface the minimum is near $r_1=r_2 \approx 0.87 \text{ \AA}$ at $\theta = 106^\circ$; the actual surface has its minimum at $r_1=r_2 \approx 0.96 \text{ \AA}$ (at $\theta = 104\text{--}105^\circ$)⁶.

The parameter A multiplying the whole expression is chosen to insure that the depth of the potential well (at $\theta \approx 106^\circ$) is close to the experimental value. In the case of the water molecule $A = 10^2$ was chosen. This value makes the well depth $\approx 5 \text{ eV}$ below the zero of

5. M. Baer, *J. Chem. Phys.* 60 (1974) 1057.
6. B.J. Rosenberg, W.C. Ermler and I. Shavitt, *J. Chem. Phys.* 65 (1976) 4022.
7. G. Herzberg, *Electronic spectra of polyatomic molecules* (Van Nostrand, Princeton, 1966).




Figure 2. H_2O ground state surface for $\theta = 170^\circ$ (contours in eV).




Figure 3. H_2O ground state surface at $\theta = 105^\circ$ (contours in eV). The hole in the surface is nearly 5 eV deep at this angle.




Figure 4. H_2O ground state surface for $\theta = 60^\circ$ (contours in eV).

the potential, the separated $\text{OH } (^2\Pi) + \text{H } (^2\text{S})$ state. Thus the overall binding energy is ≈ 9.5 eV, which is close to the experimental estimate of 10 eV⁸.

In the $\text{O} + \text{H}_2$ channel the surface does not separate properly. It should go to $\text{O}(^1\text{D}) + \text{H}_2$; however, it actually separates to $\text{O}(^3\text{P}) + \text{H}_2$.

III. DISCUSSION

The modified LEPS surface for water makes use of only a single variable Sato parameter. Since there are two diatomics involved, OH and O_2 , two Sato parameters could have been used and would provide additional flexibility. Note that the force constants for H_2O were not fit using this surface. By using two Sato parameters matching the force constants of the LEPS surface to their true values should be possible. The two Sato parameter modification has already been applied to HO_2 with good results for the force constants of the complex.

The form of the Sato parameter used here is not the only possible functional form. Other forms could be adopted and will be in the future. In all cases, however, the functional form must have the symmetry of the molecule or molecular complex in question. In summary, the modified LEPS has already proved itself a useful analytical potential surface and with future modification will be further improved.

8. B.J. Rosenberg and I. Shavitt, *J. Chem. Phys.* 63 (1975) 2162.

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>	<u>No. of Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: DDC-TCA Cameron Station Alexandria, VA 22314	1	Commander US Army Missile Materiel Readiness Command ATTN: DRSMI-AOM Redstone Arsenal, AL 35809
1	Commander US Army Materiel Development and Readiness Command ATTN: DRCMD-ST, N. Klein 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander US Army Tank-Automotive Research and Development Command ATTN: DRDTA-UL Warren, MI 48090
1	Commander US Army Aviation Research and Development Command ATTN: DRS-AV-E P.O. Box 209 St. Louis MO 63166	2	Commander US Army Armament Research and Development Command ATTN: DRDAR-TSS Dover, NJ 07801
1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	Commander US Army Armament Materiel Readiness Command ATTN: DRSAR-LEP-L, Tech Lib Rock Island, IL 61299
1	Commander US Army Electronics Research and Development Command Technical Support Activity ATTN: DELSD-L Fort Monmouth, NJ 07703	1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SL, Tech Lib White Sands Missile Range NM 88002
1	Commander US Army Communications Research and Development Command ATTN: DRDCO-SGS Fort Monmouth, NJ 07703		<u>Aberdeen Proving Ground</u> Dir, USAMSAA Cdr, USATECOM ATTN: DRSTE-SG-H
1	Commander US Army Missile Research and Development Command ATTN: DRDMI-R Redstone Arsenal, AL 35809		

ED
78