

AD-A060 620

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
A DRAZIN INVERSE FOR RECTANGULAR MATRICES.(U)

F/G 12/1

JUN 78 R E CLINE, T N GREVILLE

DAAG29-75-C-0024

UNCLASSIFIED

MRC-TSR-1855

NL


| OF |
AD
A060620



END
DATE
FILMED

1-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A060620

MRC Technical Summary Report #1855


A DRAZIN INVERSE FOR RECTANGULAR MATRICES

R. E. Cline and T. N. E. Greville

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

June 1978

Received April 19, 1978



Approved for public release
Distribution unlimited

Sponsored by
U. S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

78 09 12 044

DDC FILE COPY

ERRATUM

MRC Technical Summary Report #1855

A DRAZIN INVERSE FOR RECTANGULAR MATRICES

R. E. Cline and T. N. E. Greville

In equation (3.7), page 8, the second = sign should be "minus".

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

⑥ A DRAZIN INVERSE FOR RECTANGULAR MATRICES.

⑩ R. E. Cline[†] and T. N. E. Greville

⑨ Technical Summary Report, #1855
June 1978

⑪ Jun 78

⑭ MRZ-TSR-1855

ABSTRACT

⑫ 14 p.

The definition of the Drazin inverse of a square matrix with complex elements is extended to rectangular matrices by showing that for any B and W , m by n and n by m , respectively, there exists a unique matrix, X , such that $(BW)^k = (BW)^{k+1}XW$, for some positive integer k , $XWBWX=X$ and $BWX=XWB$. Various expressions satisfied by B , W , X and related matrices are developed.

⑮ DAAG 29-75-C-0024

AMS(MOS) Subject Classification: 15A09

Key Words: Drazin inverse, rectangular matrix, weighted inverse

Work Unit No. 2 - Other Mathematical Methods

[†]Mathematics Department and the Computing Center
University of Tennessee, Knoxville, Tennessee 37916

SEARCHED	INDEXED
SERIALIZED	FILED
JUN 1978	
FBI - KNOX	
A	

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024.

221 200 12 044 24

SIGNIFICANCE AND EXPLANATION

Efficient methods for handling systems of linear simultaneous algebraic equations provide a fundamental tool in solving problems in almost every area of computing and applied mathematics. In matrix notation, the solution of $Ax = b$ is given by $x = A^{-1}b$, where A^{-1} denotes the inverse of A .

In classical matrix algebra, the inverse of A exists only if A is square and "nonsingular." The unique inverse then satisfies $AA^{-1} = A^{-1}A = I$. A singular or rectangular matrix has no inverse in this sense. However there may be associated with it a variety of "generalized inverses", each having some of the properties of the usual inverse. For example suppose that we have m equations in n unknowns, $Ax = b$, where A is $m \times n$, $m > n$, rank n . The solution in a least squares sense is

$$x = (A^T A)^{-1} A^T b$$

and the matrix multiplying b is a generalized inverse.

A certain type of generalized inverse, the Drazin inverse, has heretofore been defined only for square (usually singular) matrices and has found application to Markov processes and to the solution of systems of ordinary differential equations. In this paper the definition of the Drazin inverse is extended to rectangular matrices and its properties studied. (This extended Drazin inverse is defined in the Abstract.)

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

A DRAZIN INVERSE FOR RECTANGULAR MATRICES

R. E. Cline and T. N. E. Greville

1. INTRODUCTION.

Let A be any rectangular matrix with complex elements. Then the Moore-Penrose inverse of A is the unique matrix $X = A^+$ such that

$$(1.1) \quad AXA = A, \quad XAX = X, \quad (AX)^H = AX, \quad (XA)^H = XA$$

(where the superscript H denotes conjugate transpose). On the other hand, if A is square, the Drazin inverse of A is the unique matrix $X = A_d$ such that

$$(1.2) \quad A^k = A^{k+1}X, \text{ for some positive integer } k,$$

$$(1.3) \quad X = X^2A,$$

$$(1.4) \quad AX = XA,$$

It is the purpose of this paper to show that a Drazin inverse can be defined for rectangular matrices in such a way that both A^+ and A_d (when A is square) follow as special cases.

2. THE W-WEIGHTED DRAZIN INVERSE OF B .

Although the Drazin inverse was originally considered for elements in an associative ring [3] and Lemma 1 was established in that context [2], we use this result only for matrices and restate it accordingly.

LEMMA 1: For any matrices B and W , m by n and n by m , respectively,

$$(BW)_d = B(WB)_d^2 W.$$

The reader can also verify Lemma 1 by taking X equal to the right member of the equation and $A = BW$ and verifying that (1.2)-(1.4) are satisfied.

Using (1.4) to rewrite (1.3) as $A_d = A_d A A_d$, the expressions in Corollary 1.1 now follow at once by induction.

COROLLARY 1.1: For every positive integer p ,

$$W(BW)_d^p = (WB)_d^p W$$

and

$$B(WB)_d^p = (BW)_d^p B.$$

Our first result, Theorem 2, is established for an arbitrary positive integer p . As will be indicated following the proof, however, the general case can always be reduced to $p = 2$ by a simple transformation.

THEOREM 2: For any matrices B and W , and for every positive integer p , there is a unique X such that

$$(2.1) \quad (BW)_d^p XW = (BW)_d^p,$$

$$(2.2) \quad BWX = XWB$$

$$(2.3) \quad BW(BW)_d^p X = X.$$

Also, there is a unique X such that

$$(2.4) \quad XW = BW(BW)_d^p,$$

$$(2.5) \quad WX = WB(WB)_d^p$$

$$(2.6) \quad XW(BW)^{p-1}X = X.$$

The unique matrix X which satisfies both sets of equations is

$$(2.7) \quad X = B(WB)_d^p.$$

Proof: Using (1.3) and Corollary 1.1, it is easily seen that X in (2.7) satisfies (2.1) to (2.6).

To establish uniqueness, we show first that (2.1), (2.2) and (2.3) imply (2.4), (2.5) and (2.6), and then that (2.4), (2.5) and (2.6) imply (2.7).

Now $XW = BW(BW)_d^p XW = BW(BW)_d^p$, by (2.3) and (2.1). Thus (2.4) holds, and combines with (2.3) to give

$$XW(BW)^{p-1}X = BW(BW)_d^p (BW)^{p-1}X = BW(BW)_d^p X = X.$$

Thus (2.6) also holds. Finally, using (2.2), (2.3), (2.4) and Corollary 1.1 we have

$$\begin{aligned}
WX &= WBW(BW)_d^X = (WB)_d WBWX \\
&= (WB)_d WXWB = (WB)_d WBW(BW)_d^P B \\
&= (WB)_d^2 (WB)_d^{P+1} = (WB)_d^P (WB)_d^P,
\end{aligned}$$

that is, (2.5). Hence (2.1), (2.2) and (2.3) imply (2.4), (2.5) and (2.6).

If (2.4), (2.5) and (2.6) hold, then

$$\begin{aligned}
X &= XW(BW)^{P-1}X = BW(BW)_d^P (BW)^{P-1}X \\
&= (BW)_d BWX = (BW)_d BWB(WB)_d^P \\
&= BWB(WB)_d^{P+1} = B(WB)_d^P.
\end{aligned}$$

Hence (2.7) holds. ■

Observe next that if $p \geq 1$, $q \geq -1$ and $r \geq 0$ are integers such that $q+2r+2 = p$, and if we let $(WB)^q = (WB)_d$ when $q = -1$, then $B(WB)_d^P = B(WB)^q [((WB)^r W) (B(WB)^q)]_d^2$. Consequently, considerations of X in Theorem 2 can always be reduced to the case $p = 2$ if B is replaced by $B(WB)^q$ and W is replaced by $(WB)^r W$.

COROLLARY 2.1: The matrix $X = B(WB)_d^2$ is the unique solution to the equations

$$(2.8) \quad (BW)^k = (BW)^{k+1} XW, \text{ for some positive integer } k,$$

$$(2.9) \quad X = XWBWX,$$

$$(2.10) \quad BWX = XWB.$$

Proof: That $X = B(WB)_d^2$ is a solution is apparent by noting that the relations in (2.10) and (2.2) are identical, (2.9) is (2.6) when $p = 2$ and, with $XW = (BW)_d$ by (2.4), (2.8) is (1.2) for $A = BW$.

To show uniqueness, suppose both X_1 and X_2 are solutions to (2.8) for some positive integers k_1 and k_2 , respectively, (2.9) and (2.10). Let $\hat{k} = \max(k_1, k_2)$. Then using repeated applications of equations (2.8), (2.9), and (2.10) we have

$$\begin{aligned}
X_1 &= X_1 WBWX_1 = BWX_1 WX_1 = (BW)^2 (X_1 W)^2 X_1 \\
&= \dots = (BW)^{\hat{k}} (X_1 W)^{\hat{k}} X_1 = (BW)^{\hat{k}+1} X_2 W (X_1 W)^{\hat{k}} X_1 \\
&= X_2 (WB)^{\hat{k}+1} W (X_1 W)^{\hat{k}} X_1 = X_2 WBW (BW)^{\hat{k}} (X_1 W)^{\hat{k}} X_1 \\
&= X_2 WBWX_1.
\end{aligned}$$

Continuing in a similar manner, $X_2 WBWX_1$ can be reduced to X_2 . Thus $X = B(WB)_d^2$ is unique. ■

It should be noted in Corollary 2.1 that $B_d = B(WB)_d^2$ when B is square and $W = I$. Also, there is a direct correspondence between the relations in (2.8) and (1.2), (2.9) and (1.3) when written as $X = XAX$, and (2.10) and (1.4) in which the role of W in (2.8), (2.9) and (2.10) is to act as a "sandwich" matrix so that products such as BWX and XWB can be defined. In view of the correspondence between the defining equations for A_d and those in Corollary 2.1, we define the Drazin inverse of a rectangular matrix in the following manner:

DEFINITION 1: For any matrices B and W , m by n and n by m , respectively, the matrix $X = B(WB)_d^2$ is called the W -weighted Drazin inverse of B , and is written as $X = B_{d,W}$.

Given a square matrix, A , the smallest positive integer k such that (1.2) holds is called the index of A , and it can be shown that k is the minimal power for which A^k and A^{k+1} have the same rank. Moreover, $(A_d)_d = A_d^2$, so that $(A_d)_d = A$ if and only if A has index one, whereas $((A_d)_d)_d = A_d$, and A_d always has index one [3]. In Theorem 3 we characterize matrices, X , such that $X = B_{d,W}$ for some W in terms of matrices with index one.

THEOREM 3: Let B and X be any m by n matrices. Then $X = B_{d,W}$ for some W if and only if X has the form

$$(2.11) \quad X = BYBYB$$

for some matrix Y such that both BY and YB have index one.

Proof: Suppose $W = Y(BY)_d^2 = (YB)_d^2 Y$, where BY and YB have index one.

Then $WB = (YB)_d$ and, using (2.11),

$$WX = (YB)_d YBYB = YB = (WB)_d.$$

In a similar manner, $BW = (BY)_d$, and

$$XW = BY = (BW)_d.$$

Finally,

$$XWBWX = BYBYB = X,$$

so that (2.4), (2.5) and (2.6) hold for B, W, X and $p = 2$. Thus $X = B_{d,W}$.

Conversely, if $X = B_{d,W}$ for some W , then

$$\begin{aligned} X &= B(WB)_d^2 = (BW)_d B(WB)_d \\ &= BW(BW)_d^2 B(WB)_d^2 WB \\ &= B(WB)_d^2 WB(WB)_d^2 WB \end{aligned}$$

is of the form (2.11) where $Y = (WB)_d^2 W$. Moreover, $BY = (BW)_d$ and $YB = (WB)_d$ have index one. ■

For any matrix B in Theorem 3, both W and Y are n by m , and the following corollary shows that $W = Y$ and $X = B$ in (2.11) is necessary and sufficient to have B and $B_{d,W}$ equal.

COROLLARY 3.1: $B = B_{d,W}$ if and only if $B = BWBWB$.

Proof: If $B = BWBWB$, then $WB = (WB)^3$ implies $WB = (WB)_d$. Thus, $B = B_{d,W}$.

Conversely, if $B = B_{d,W} = B(WB)_d^2$, then $WB = (WB)_d$. ■

3. THE B-WEIGHTED DRAZIN INVERSE OF W .

For any B and W , m by n and n by m , respectively, interchanging the roles of B and W throughout §2 provides a completely dual set of results for $W_{d,B} = W(BW)_d^2$, the B-weighted Drazin inverse of W . However, whereas $B_{d,W}$ has the size of B , $W_{d,B}$ has the size of B^H and B^+ . Analogous to the observation in §2 that $B_d = B_{d,W}$ when B is square and $W = I$, we now consider choices of W for which $B^+ = W_{d,B}$.

Following the notation of [1], let $B^{(2)} = \{Z | ZBZ = Z\}$. Then $W_{d,B} \in B^{(2)}$ for any W such that $B^+ = W_{d,B}$.

LEMMA 4: $W_{d,B} \in B^{(2)}$ if and only if $W_{d,B} = W(BW)_d$.

Proof: If $W_{d,B} \in B^{(2)}$, then $W(BW)_d^3 = W(BW)_d^2$ so that $(BW)_d$ is idempotent.

Conversely, $W(BW)_d \in B^{(2)}$. ■

An immediate consequence of Lemma 4 is that only expressions of the form $B^+ = W(BW)_d$ must be considered when determining choices of W . Moreover, observe that since $A^+ = A_d$ for any Hermitian matrix and $(BB^H)^+ = B^{H+}B^+$ for any B , then

$$(3.1) \quad B^+ = B^+BB^+ = B^HB^H B^+ = B^H(BB^H)_d .$$

Therefore, one choice of W is B^H . To characterize the class of all matrices which can be used to form B^+ in this manner, it is convenient to replace W by C^H , where C has the size of B .

DEFINITION 2: For any m by n matrices B and C , C is said to be alias to B if $B^+ = C^H(BC^H)_d$.

Given any m by n matrix B with rank r , $B = EF$ is said to be a full rank factorization if E and F^H have r columns [1]. In this case $B^+ = F^+E^+$ with $E^+ = (E^HE)^{-1}E^H$ a left inverse of E and $F^+ = F^H(FF^H)^{-1}$ a right inverse of F .

Matrices alias to B are now characterized using a full rank factorization of B . In the proof of Theorem 5 we use the facts [4] that the general solution of a consistent system of equations $AXB = C$ can be written as

$$(3.2) \quad X = X_0 + Y - A^+ AYBB^+,$$

where X_0 is any particular solution and Y is arbitrary, and that necessary and sufficient conditions for the equations $AX = B$, $XC = D$ to have a common solution are that each is consistent and $AD = BC$ in which case

$$(3.3) \quad X_0 = A^+ B + DC^+ - A^+ ADC^+$$

is a solution.

THEOREM 5: Let $B = EF$ be any full rank factorization. Then C is alias to B if and only if C has the form

$$(3.4) \quad C = ESF + Z$$

where S is any nonsingular matrix and Z is any matrix such that $BZ^H = 0$ and $Z^H B = 0$.

Proof; If C has the form in (3.4) where S and Z satisfy the hypotheses, then

$$\begin{aligned} C^H (BC^H)_d &= (C^H B)_d C^H = (F^H S^H E^H EF)_d F^H S^H E^H \\ &= F^H S^H (E^H E F F^H S^H)_d E^H \\ &= F^H S^H (S^H)^{-1} (FF^H)^{-1} (E^H E)^{-1} E^H \\ &= F^+ E^+ = B^+. \end{aligned}$$

Thus C is alias to B .

To prove the converse, suppose first that B has either full row rank or full column rank: If B has full row rank and C is alias to B , then $B^+ = B^H (BB^H)^{-1} = C^H (BC^H)_d$ implies that BC^H is nonsingular. Hence $C = S_1 B$ where $S_1 = CB^H (BB^H)^{-1} = CB^+$ is nonsingular. By a similar type of argument, $C = BS_2$ with S_2 nonsingular if B has full column rank. Consequently, (3.4) holds with $Z = 0$ in both cases.

Suppose now that B has neither full row rank nor full column rank. With $B = EF$ a full rank factorization and C alias to B , then $F^+E^+ = C^H(BC^H)_d$ implies

$$E^+ = FC^H(EFC^H)_d$$

and

$$F^+ = C^H(EFC^H)_d E = C^H E (FC^H E)_d.$$

Therefore, CF^H is alias to E and $E^H C$ is alias to F , so that there exist nonsingular matrices Q and R , say, for which

$$(3.5) \quad FC^H = QE^H$$

and

$$(3.6) \quad C^H E = F^H R.$$

Using (3.3) it then follows that

$$(3.7) \quad X_0 = F^+QE^H + F^HRE^+ \stackrel{(3.5)}{=} F^+FF^HRE^+ = F^+QE^H$$

is a solution to (3.5), (3.6) and thus to

$$(3.8) \quad FC^H E = QE^H E.$$

But by (3.2), all solutions of (3.8) can be written as

$$(3.9) \quad C^H = X_0 + Z^H$$

where $Z^H = Y - F^+FYEE^+$ with Y arbitrary. Consequently those solutions in (3.9) such that (3.5) and (3.6) also hold must satisfy $FZ^H = 0$ and $Z^H E = 0$, which implies $BZ^H = 0$ and $Z^H B = 0$. Finally, the form in (3.4) is obtained from (3.9) by noting that the last expression for X_0 in (3.7) can be written as $X_0 = F^H S^H E^H$ with $S^H = (FF^H)^{-1} Q$ nonsingular. ■

In Corollary 5.1 we characterize those matrices in (3.4) with $Z = 0$.

COROLLARY 5.1: For any full rank factorization $B = EF$, the set of matrices alias to B of the form $C = ESF$ is an equivalence class.

Proof: The relation is reflexive with $S = I$, that is, (3.1). Since $C = E(SF)$ is a full rank factorization, symmetry follows from $B = ES^{-1}(SF)$. Moreover, if D alias to C has the form $D = ET(SF)$ with T nonsingular, then D is alias to B . ■

REFERENCES

1. A. Ben-Israel and T. N. E. Greville, Generalized Inverses: Theory and Applications, Wiley, New York, 1974.
2. R. E. Cline, An application of representations for the generalized inverse of a matrix, Tech. Summary Rep. 592, Mathematics Research Center, U.S. Army, University of Wisconsin, Madison, Wisconsin, 1965.
3. M. P. Drazin, Pseudo-inverses in associative rings and semigroups, Amer. Math. Monthly, 65 (1958), pp. 506-513.
4. R. Penrose, A generalized inverse for matrices, Proc. Camb. Phil. Soc., 51 (1955), pp. 406-413.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1855 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Drazin Inverse for Rectangular Matrices		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) R. E. Cline and T. N. E. Greville		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 2 - other mathematical methods
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE June 1978
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Drazin inverse, rectangular matrix, weighted inverse		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A Drazin inverse is defined for a rectangular matrix and its properties studied.		