

AD-A060 651


CARNEGIE-MELLON UNIV PITTSBURGH PA MANAGEMENT SCIENC--ETC F/G 12/1  
FUNCTIONS FOR MEASURING THE QUALITY OF APPROXIMATE SOLUTIONS TO--ETC(U)  
JUN 78 E ZEMEL N00014-75-C-0621  
MSRR-422-REV NL

UNCLASSIFIED

| DF |  
AD  
A060651



END  
DATE  
FILMED  
1-79  
DDC



**LEVEL II**

①  
NW

AD A 0 6 0 6 5 1




DDC FILE COPY

**Carnegie-Mellon University**

PITTSBURGH, PENNSYLVANIA 15213

**GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION**

WILLIAM LARIMER MELLON, FOUNDER



DDC  
RECEIVED  
NOV 1 1978  
REGULATED  
D

**DISTRIBUTION STATEMENT A**  
Approved for public release;  
Distribution Unlimited

78 10 23 025

ADDITION TO	
DTIC	White Section <input checked="" type="checkbox"/>
DDC	Dist Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY GROUP	
Dist.	AVAIL. and/or SPECIAL
A	

# LEVEL II



Management Sciences Research Report No. 422

AD A060651

DDC FILE COPY

FUNCTIONS FOR MEASURING THE  
QUALITY OF APPROXIMATE SOLUTIONS  
TO ZERO-ONE PROGRAMMING PROBLEMS.

by

10 Eitan/Zemel

14 MSRR-422-REV

April 1978

Revised June 1978

12 28p

13 N00014-75-C-0621  
NSF-MPS 73-08534

The research underlying this report was supported by Grant  
MPS 73-08534 A02 of the National Science Foundation. Reproduction  
in whole or in part is permitted for any purpose of the U.S.  
Government.

Management Science Research Group  
Graduate School of Industrial Administration  
Carnegie-Mellon University  
Pittsburgh, Pennsylvania 15213

DDC  
RECEIVED  
NOV 1 1978  
D

\*Graduate School of Management, Northwestern University, Evanston, Illinois

403 426

**DISTRIBUTION STATEMENT A**  
Approved for public release;  
Distribution Unlimited

78 10 23 025

JOB

ABSTRACT

↘ We point out the difficulties associated with measuring the quality of an approximate (heuristic) solution by "Percentage-Error" as is customary. We define a set of properties that are to be expected from a proper measure of solution quality and investigate the family of functions which satisfy those conditions. In particular, we show that for any class of 0-1 programming problems appropriate functions do exist and that they are uniquely defined up to monotone transformations. We conclude with several examples of linear functions which are suitable for certain classes of problems.

↖

## I. Introduction

Consider the 0-1 integer program

$$\begin{aligned}
 (p) \quad & \max \quad \sum_{j=1}^n c_j x_j \\
 & \text{s.t} \quad \sum_{j=1}^n a_{ij} x_j \begin{matrix} \leq \\ \text{or} \\ = \end{matrix} b_i \quad i \in M = \{1 \dots m\} \\
 & \quad \quad \quad x_j = 0 \text{ or } 1 \quad j \in N = \{1 \dots n\}
 \end{aligned}$$

We will denote by  $U$  the universe of problems of the type  $p$ . For

$p \in U$  let

$$F(p) = \{x \in \{0,1\}^n \mid \sum_{j=1}^n a_{ij} x_j \begin{matrix} \leq \\ \text{or} \\ = \end{matrix} b_i \quad i \in M\}$$

$$Z(p) = \max_{x \in F(p)} cx$$

$$O(p) = \{x \in F(p) \mid cx = Z(p)\}$$

be the feasible set, optimal value, and optimal set for  $p$  respectively.

Let now  $P \subseteq U$  be a particular class of problems (say knapsack, travelling salesman, set covering, etc.) and let  $A$  be an exact algorithm for  $P$ , i.e. such that given any problem  $p \in P$ ,  $A$  finds an optimal solution,  $x \in O(p)$ . The work of Cook [1], Karp [6], and others indicates that for many interesting classes of problems  $P$  (including the ones just mentioned), one is unlikely to find an exact algorithm  $A$  which will be able to solve every problem  $p \in P$  without, on occasion, investing a computational effort which is essentially equivalent to enumerating a significant fraction of the vertices of  $\{0,1\}^N$ . It is also well known that practitioners of Integer and Combinational Programming have often been confronted with individual and classes of problems which defy solution in "reasonable" amounts of time. Thus, it is not surprising that

a considerable amount of effort has been recently devoted to the analysis of approximate (heuristic) algorithms, e.g., [3], [5], [7]. In contrast to an exact one, an approximate algorithm finds a feasible solution,  $x \in F(p)$ , but not necessarily an optimal one. The effectiveness of such an algorithm with respect to the problem  $p$  must then be evaluated along two attributes: the computational effort involved in finding  $x$  and the "quality" of  $x$  as an approximation for the optimal solution. It is this second attribute which is our domain of interest in this paper.

It will be more convenient to measure the "quality" of a given solution on a negative scale, i.e., we will be concerned with measuring the "distance", so to speak, of  $x$  from an optimal solution. We will call this "distance" the Error associated with  $x$  and any function which attempts to quantify it, the error evaluation of  $x$  relative to  $p$ , denoted  $E_p(x,p)$ . The main purpose of this paper is then to investigate and characterize functions which can serve as error evaluators. (It should be noted that this discussion has nothing to contribute towards the resolution, and is independent of, the deeper question associated with aggregating the characteristics of a given algorithm with regard to the individual problems  $p \in P$ , into a measure of effectiveness relevant to the class  $P$  as a whole. It is only after a proper measure is established for the individual problems that one needs decide whether averaging or worst case analysis is the meaningful, tractable, practical, etc. method of aggregation.)

The organization of the paper is as follows. In section II we present a set of properties to be expected from an error evaluating function and examine them vis-a-vis the traditional "percentage error" measure. In section III we show that functions which satisfy the conditions introduced in section II do indeed exist. We then discuss the possible uniqueness of such functions. Finally, in section IV, we conclude with several examples of error evaluating functions for particular classes of problems.

## II. Basic Properties of Error Evaluating Functions

A given error evaluation function (E.E.F.) is not expected to perform properly unless it satisfies a set of reasonable properties. In this section, we attempt to propose such a set. To simplify notation, we will delete the parameter P from our notation for  $E_p(x,p)$ .

The first two properties, A and B, require hardly a comment. An E.E.F. should certainly be a monotone (at least weakly) function of the solution value  $cx$ . Also, the error associated with an optimal solution can be conveniently set to zero. Thus:

Condition A. Let  $p \in P, x^1, x^2 \in F(p)$   
 $cx^1 \geq cx^2 \Rightarrow E(x^1, p) \leq E(x^2, p).$

Condition B. Let  $p \in P$   
 $x \in O(p) \Rightarrow E(x, p) = 0$

An E.F.F. often used by researchers is given by

$$E_1(x, p) = (cx^P - cx) / |cx^P|$$

where  $x^P \in O(p)$  is an optimal solution for p.

If P is such that for every  $p \in P, cx^P \neq 0$ , then  $E_1(x, p)$  is easily seen to satisfy conditions A and B. So, of course, do the functions

$$E_2(x, p) = cx^P - cx$$

$$E_3(x, p) = (cx^P - cx) / |c_1|$$

where  $c_1$  is the first (index wise) non zero entry in the objective row.

$$E_4(x, p) = \begin{cases} 0 & \text{if } x \in O(p) \\ 1 & \text{otherwise} \end{cases}$$

etc.

Let us examine  $E_2(x, p)$  more carefully. One finds it hardly surprising

that this function is rarely proposed as an E.E.F. Intuitively, one feels that the difference in the objective values is meaningless by itself and can be gauged only relative to some other normalization factor. Put differently,  $E_2(x,p)$  is unacceptable since its value can be arbitrarily manipulated by changes of units in the objective function. Likewise,  $E_3(x,p)$  can be rejected on the grounds that its value depends on the particular order in which the variables are indexed. Several authors have observed that similar objections can be often raised with respect to  $E_1$ . For instance, Cornuejols, Fisher and Nemhauser [ 2 ] examined a set of problems of the plant location variety. Such problems are known to be invariant under addition of certain row constants to the cost matrix. Yet, as noted in [ 2 ], the value of the error for a given non-optimal solution, as measured by  $E_1(x,p)$ , does change under such transformations. To overcome this difficulty, an E.E.F. is proposed in [ 2 ] which is invariant under the particular cost transformations in question. An identical problem arises, of course, when one applies  $E_1(x,p)$  as an E.E.F. for transportation problems, general (as opposed to Euclidean) travelling salesman problems and the like

A related objection to  $E_1(x,p)$  is brought forward by Korte in [ 7 ]. It is well known that a given optimization problem can often be presented as a maximization or minimization problem, the two versions being virtually equivalent. Yet the equivalent does not always carry over to the corresponding approximation problems. For instance, the value assigned by  $E_1$  to a given non-optimal solution for the knapsack problem depends on the version chosen. Even worse, the computational complexity of the two versions may be different. This is the case for the travelling salesman problem where the maximization version can be solved by a guaranteed performance polynomial algorithm [ 7 ], yet the problem of solving the minimization version to any accuracy is NP hard [ 8 ]. As Korte suggests, this behavior casts some doubts as to the appropriateness of measuring errors by  $E_1(x,p)$  for this class of problems.

Condition C below is designed to take account of the ideas brought forward in the foregoing discussion. Basically, it enforces equality of the values

assigned by an E.E.F. to corresponding solutions in problems which are obviously 'equivalent', but may appear different when viewed superficially. By equivalent we mean here problems which are related to each other by permutations of the rows or columns, complementing of variables, and trivial algebraic manipulations of the objective function and constraints. But first we must digress for some notation and definitions.

For a problem  $p \in P$ , let us denote

$$c = (c_j)$$

$$A = (a_{ij}) \text{ with } a^j, (a_i) \text{ the } j\text{th column (ith row) of } A.$$

$$b = (b_i)$$

$$\rho = (\rho_i) \text{ is } "\leq" \text{ or } "=" \text{ as the case may be.}$$

We then have the more compact representation:

$$\begin{aligned} (p) \quad & \max \quad cx \\ & \text{s.t. } Ax \rho b \\ & x \in \{0,1\}^N \end{aligned}$$

On occasion, we will refer to  $p$  as  $(c, A, \rho, b)$ .

For definitions 1 - 4 below, let  $p = (c, A, \rho, b) \in U, x \in F(p)$ .

Definition 1 Let  $\varphi$  be a column permutation operator. By  $\varphi p$ , we mean the problem  $p' = (\varphi c, \varphi A, \rho, b)$ .  $x' = \varphi x$  is defined accordingly.

Definition 2 Let  $\pi$  be a row permutation operator. By  $\pi p$  we mean the problem  $p' = (c, \pi A, \pi \rho, \pi b)$ . By  $\pi x$  we simply mean  $x$ .

Definition 3 Let  $S \subseteq N$ . We will denote by  $D^S$  the operator which complements the variables  $x_j, j \in S$ , i.e.:

$$D^S(p) = p' = (c', A', \rho, b')$$

with

$$c'_j = \begin{cases} -c_j & j \in S \\ c_j & j \notin S \end{cases}$$

$$a'_j = \begin{cases} -a_j & j \in S \\ a_j & j \notin S \end{cases}$$

$$b'_i = b_i - \sum_{j \in S} a_{ij}$$

Also

$$D^S x = x' \quad \text{is given by}$$

$$x'_j = \begin{cases} 1-x_j & j \in S \\ x_j & j \notin S \end{cases}$$

Unless specifically required, we will suppress the parameter S and refer to the operator as D.

Definition 4 Let  $M_p = \{i \in M \mid \rho_i \text{ is "="}\}$  be the index set for the equality rows of p. Let also

$\alpha^0, \dots, \alpha^M \in R_+$  be any set of M+1 positive constants

$\beta^0, \dots, \beta^M \in R^{M_p}$  be any set of M+1 real,  $M_p$  dimensional vectors such

that the matrix  $\beta_j^i + \alpha^i \delta_{ij}$   $i \in M_p, j \in M_p$ , with  $\delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases}$  is non singular. Then

$$G_{\alpha, \beta} p = p' = (c', A', \rho, b')$$

Where

$$c'_j = \alpha^0 c_j + \sum_{i \in M_p} \beta_i^0 a_{ij}$$

$$a'_{kj} = \alpha^k a_{kj} + \sum_{i \in M_p} \beta_i^k a_{ij}$$

$$b'_k = \alpha^k b_k + \sum_{i \in M_p} \beta_i^k b_i$$

Obviously,

$$G_{\alpha, \beta} x = x$$

Again we will usually suppress the parameter  $\alpha, \beta$  and refer to the operator as  $G$ .

Definition 5 Let  $p^1 \in U$ , and let  $p^2 \in U$  be a problem obtainable from  $p^1$  by a sequence of  $\pi$ ,  $\varphi$ ,  $G$  and  $D$  operators applied in any order. We will then say that  $p^1$  is reducible to  $p^2$ , denoted  $p^1 \rightarrow p^2$ . Also let  $x^1 \in F(p^1)$ , and let  $x^2$  be obtained from  $x^1$  by an identical sequence of operators. Then we say that  $x^2$  corresponds to  $x^1$ ,  $x^1 \rightarrow x^2$ , relative to  $p^1 \rightarrow p^2$ . One can easily verify that in this case,  $x^2 \in F(p^2)$ .

We are now ready to define the third property of E.E.F's:

Condition C. Let  $p^1, p^2 \in P$  such that  $p^1 \rightarrow p^2$ . Further, let  $x^1 \in F(p^1)$  and let  $x^2$  correspond to  $x^1$  relative to  $p^1 \rightarrow p^2$ .

Then:

$$E(x^1, p^1) = E(x^2, p^2).$$

### III. Further Properties of Error Evaluating Functions

Condition C cuts significantly the number of available E.E.F.'s. For instance,  $E_1(x,p)$  is excluded for several frequently occurring classes of problems P. Naturally, for a given P, the questions arise as to the existence and uniqueness of an E.E.F. which satisfies conditions A-C. In this section, we address these questions. The first is settled in the affirmative (Theorem 1). The second in a qualified affirmative (Theorems 2 and 3). We open the discussion with some technical remarks concerning the relation  $\rightarrow$ .

Remark 1 It can be easily established that any of the row type operators,  $\pi$  and G, commutes with the column type operators,  $\varphi$  and D. On the other hand,  $\pi$  and G do not commute, nor do  $\varphi$  and D. However, the following weaker form of commutativity does hold:

(i) For every column permutation operator  $\varphi$  and every subset  $S \subseteq N$ , there exists  $S' \subseteq N$  such that

$$(1) \quad \varphi D^S = D^{S'} \varphi$$

The property goes the other way too, i.e., for every  $\varphi$  and every  $S' \subseteq N$ , there exists  $S \subseteq N$  such that (1) holds.

(ii) For every row permutation operator  $\pi$  and every set of constants  $\alpha, \beta$  as defined in condition C, there exists a set of  $\alpha', \beta'$  such that

$$(2) \quad \pi G_{\alpha, \beta} = G_{\alpha', \beta'} \pi$$

Again, the statement is true in the reverse direction too, i.e., for every set  $\alpha', \beta'$  there exists a set  $\alpha, \beta$  such that (2) holds.

It can also be easily verified that any sequence of  $\pi$  type operators is in itself a  $\pi$  type operator.  $\varphi, G$  and D operators behave likewise. Also, each of the four types of operators contains an identity operator. Thus, definition 5 can be stated equivalently:

Definition 5': Let  $p^1, p^2 \in U$ . Then  $p^1 \rightarrow p^2$  iff there exists  $\pi, G, \varphi, D$  such that

$$p^2 = \pi \varphi G D p^1$$

In such a case, we say that  $x^1 \rightarrow x^2$  relative to  $p^1 \rightarrow p^2$  if:

$$x^2 = \varphi D x^1$$

Proposition 1

The relation  $\rightarrow$  among problems is

- (i) Symmetric, i.e.,  $p^1 \rightarrow p^2 \Rightarrow p^2 \rightarrow p^1$
- (ii) Transitive, i.e.,  $p^1 \rightarrow p^2, p^2 \rightarrow p^3 \Rightarrow p^1 \rightarrow p^3$
- (iii) Reflexive, i.e.,  $p \rightarrow p$

A corresponding set of statements can be made with respect to the relation  $\rightarrow$  among the corresponding solutions  $x^1, x^2, x^3$ .

Proof:

- (i) follows from the fact that each operator has an inverse,
- (iii) follows from the fact that each operator type has an identity, and
- (ii) follows by the same reasoning used in the paragraph preceding definition 5' Q.E.D.

Proposition 2

Let  $P \subseteq U$  be arbitrary. Then  $P$  has a unique partition:

$$P = \bigcup_{l \in K} P_l \quad \text{such that}$$

- (i)  $P_l \cap P_{l'} = \emptyset$  for  $l \neq l'$
- (ii) Let  $p^1 \in P_l, p^2 \in P_{l'}$ . Then  $p^1 \rightarrow p^2 \Leftrightarrow l = l'$

Proof immediate from proposition 1. Q.E.D.

We will refer to the  $P_l$ 's as the equivalence classes within  $P$ . We will call two individual problems equivalent if they belong to the same equivalence class.

Remark 2

Let  $p^i \in P$ ,  $i = 1, 2$  such that  $p^1 \rightarrow p^2$ . The transformations involved in this relation are not necessarily unique. It follows then that there may be more than one distinct  $x^2 \in F(p^2)$  such that  $x^1 \rightarrow x^2$  relative to  $p^1 \rightarrow p^2$ . For example, let

$$\begin{aligned} (p^1) \quad & \max x_1 + x_2 \\ & \text{s.t.} \\ & x_1 + 2x_2 \leq 2 \\ & 2x_1 + x_2 \leq 2 \\ & x_1, x_2 = 0 \text{ or } 1 \end{aligned}$$

and let  $x^1 \in F(p^1)$  be  $(1, 0)$ . Consider

$$\begin{aligned} (p^2) \quad & \max 2x_1 + 2x_2 \\ & \text{s.t.} \\ & x_1 + 2x_2 \leq 2 \\ & 2x_1 + x_2 \leq 2 \\ & x_1, x_2 = 0 \text{ or } 1 \end{aligned}$$

One can observe that  $p^2 = \pi \varphi G_{\alpha, \beta} D^S p^1$  where

$S = \emptyset$ ;  $\alpha^0 = 2$ ;  $\alpha^i = 1$ ,  $i = 1, 2$ ;  $\pi$  and  $\varphi$  are the identity permutations. (Since in this case  $M_p = \emptyset$ ,  $\beta^i = \emptyset$  too.) The solution which corresponds to  $x^1$  is then  $x^2 = \varphi D x^1 = (1, 0)$ . But another set of possible operators which transforms  $p^1$  to  $p^2$  is given by

$$p^2 = \pi' \varphi' G_{\alpha, \beta} D^S p^1$$

where  $G$  and  $D$  are as before, but where  $\varphi'$  permutes the columns of  $p^1$  and  $\pi'$  permutes the rows. Under these operators, the solution which corresponds to  $x^1$  relative to  $p^1 \rightarrow p^2$  is  $y^2 = \varphi' x^1 = (0, 1) \neq x^2$ . However, there is a considerable similarity between the feasible sets of equivalent problems as the following three propositions indicate.

Proposition 3

For  $i = 1, 2$  let

$$p^i = (c^i, A^i, \rho^i, b^i) \in P$$

be such that  $p^1 \rightarrow p^2$ . In particular, let

$$p^2 = \pi \varphi G D p^1$$

For an arbitrary  $x \in F(p^1)$  let  $y \in F(p^2)$  be given by

$$y = \varphi D x$$

Then,

$$c^2 y = k_1 c^1 x + k_2 \quad \text{where } k_1 > 0, k_2 \text{ depend on the transformations}$$

$\pi, \varphi, G$  and  $D$ , but not on the particular solution  $x$ .

Proof Let us examine the effect of each of the transformations in question on the value of the objective function of the corresponding solutions.

(i) Let  $p = (c, A, \rho, b)$ ,  $p' = D^S p$ ,  $x \in F(p)$ ,  $x' = D^S x$  as in definition 3.

$$\begin{aligned}
 c'x' &= \sum_{j \in N} c'_j x'_j = \sum_{j \in N \setminus S} c_j x_j + \sum_{j \in S} (-c_j) (1-x_j) \\
 &= \sum_{j \in N} c_j x_j - \sum_{j \in S} c_j = k_1 c x + k_2
 \end{aligned}$$

where  $k_1 = 1$  and  $k_2 = -\sum_{j \in S} c_j$  do not depend on  $x$ .

(ii) Let  $p = (c, A, \rho, b)$ ,  $p' = G_{\alpha, \beta} p$ ,  $x \in F(p)$ ,  $x' = x$  as in definition 4.

$$\begin{aligned}
 c'x' &= \sum_{j \in N} c'_j x'_j = \sum_{j \in N} (\alpha^0 c_j + \sum_{i \in M_p} \beta_i^0 a_{ij}) x_j \\
 &= \alpha^0 \sum_{j \in N} c_j x_j + \sum_{i \in M_p} \beta_i^0 \sum_{j \in N} a_{ij} x_j = \\
 &= \alpha^0 c x + \sum_{i \in M_p} \beta_i^0 b_i = k_1 c x + k_2
 \end{aligned}$$

where  $k_1 = \alpha^0 > 0$ , and  $k_2 = \sum_{i \in M_p} \beta_i^0 b_i$  do not depend on  $x$ .

(iii) It can be easily verified that  $\pi$  and  $\varphi$  do not have any effect on the objective value. Q.E.D.

Proposition 4

Let  $p^i \in P, i = 1, 2$  such that  $p^1 \rightarrow p^2$ . Let  $Z_1^i > Z_2^i \dots > Z_{r(i)}^i$  be the distinct values of  $\{c^i x, | x \in F(p^i)\}$ , with the corresponding solution sets  $N_j^i, 1 \leq j \leq r(i)$ , and multiplicities  $n_j^i = |N_j^i|$ . Then

(i)  $r(1) = r(2) = r$

(ii)  $n_j^1 = n_j^2$  for  $j = 1, \dots, r$

Proof Let  $p^2 = \pi \varphi G D p^1$  be a particular transformation of  $p^1$  to  $p^2$ . Let  $x^1, y^1 \in F(p^1)$ ,  $x^2 = \varphi D x^1$ ,  $y^2 = \varphi D y^1$ . It follows directly from proposition 3 that  $c^1 x^1 > c^1 y^1 \Leftrightarrow c^2 x^2 > c^2 y^2$ . Thus,  $x \in N_1^1 \Leftrightarrow \varphi D x = y \in N_1^2$ . Since  $\varphi D$  is a one to one transformation of  $F(p^1)$  onto  $F(p^2)$ , we have that  $n_1^1 = n_1^2$ . To complete the proof, we have to repeat the argument with respect to  $N_j^1$ ,  $j = 2, \dots, r(1)$ . Q.E.D.

Proposition 5

Let  $p^1, p^2 \in U$  be such that  $p^1 \rightarrow p^2$  and assume that there is more than one set of transformations which induce this relation. In particular,

$$p^2 = \pi \varphi G D p^1$$

but also

$$p^2 = \pi' \varphi' G' D' p^1$$

Let

$$x^1 \in F(p^1); x^2 = \varphi D x^1; y^2 = \varphi' D' x^1.$$

Then

$$c^2 y^2 = c^2 x^2$$

Proof Using the notation of proposition 4, let  $x^1 \in N_j^1$  for some  $1 \leq j \leq r$ .

Then by the same logic applied in the proof of that proposition, we have that  $x^2 \in N_j^2$  for the same index  $j$ . But then also  $y^2 \in N_j^2$  by the same argument. Thus,

$$c^2 x^2 = c^2 y^2 = z_j^2 \quad \text{Q.E.D.}$$

We are now ready to address the existence and uniqueness issues for those E.E.F.'s which satisfy conditions A-C for a given class P. We will call such E.E.F.'s proper for P. It should be noted that this concept is defined relative to a given class P since condition C is in effect among problems within P only. Thus, it is entirely possible to have an E.E.F. which is proper for one class of problems P, but improper for another. Indeed, on occasion, one may wish to override the equivalence between problems implied by the relation  $\rightarrow$ . For instance, one may feel that a standard transportation problem, whose constraints have been manipulated by G and D type transformations, may not be easily recognized as a transportation problem and therefore need not necessarily be evaluated

identically. The same distinction can be made between general and Euclidean cost traveling salesman problems, etc. Such cases can be conveniently handled by a proper choice of the class  $P$ . (One would choose  $P$  to be the class of standard format transportation problems in the first case, Euclidean traveling salesman problems in the second). Naturally, the larger is  $P$ , the fewer are the functions which are proper for it. One may then be concerned that if  $P$  is large enough, a proper E.E.F. may not exist. This possibility is ruled out by

Theorem 1 For every class  $P \subseteq U$ , there exists a proper E.E.F.

Proof Let  $p \in P$  and let  $z_1 > z_2 \dots > z_r$  be the distinct values of the objective function with the corresponding solution sets  $N_1, N_2, \dots, N_r$ . Consider the E.E.F.

$$E^*(x,p) = j-1 \quad \text{if} \quad x \in N_j, \text{ i.e. if } cx = z_j$$

It can be easily seen that  $E^*(x,p)$  satisfies conditions A and B. The fact that it satisfies condition C as well follows from propositions 4 and 5. Q.E.D.

Can we expect uniqueness of  $E^*$  as a proper E.E.F. for  $P$ ? The following Theorem shows that every proper E.E.F. yields in a natural way a vast number of new proper E.E.F.'s.

Theorem 2 Let  $E(x,p)$  be a proper E.E.F. for  $P$ . Let  $P_1, \dots, P_\ell$  be the partition of  $P$  to equivalence classes established in proposition 2. For each  $1 \leq k \leq \ell$ , let  $b_k(\cdot)$  be any non-decreasing function of one argument such that  $b_k(0) = 0$ .

Then

$$E'(x,p) = b_k(E(x,p)) \quad \text{for } p \in P_k$$

is also a proper E.E.F.

Proof We have to show that  $E'(x,p)$  satisfies conditions A, B and C. That it satisfies A follows from the facts that  $E(x,p)$  satisfies A and that  $b_k(\cdot)$  is

monotone.  $E'(x,p)$  satisfies B since  $E(x,p)$  does and since  $b_k(0) = 0$ . It remains to be seen that  $E'(x,p)$  satisfies condition C. Let then  $p^1, p^2 \in P$ , such that  $p^1 \rightarrow p^2$ . It follows that  $p^1, p^2 \in P_k$  for some  $1 \leq k \leq l$ . Hence, the same function,  $b_k(\cdot)$ , is used to define  $E'(x,p)$  for both  $p^1$  and  $p^2$ . Let  $x^1 \rightarrow x^2$  relative to  $p^1 \rightarrow p^2$ . Since  $E(x,p)$  is proper

$$E(x^1, p^1) = E(x^2, p^2)$$

but then

$$E'(x^1, p^1) = b_k(E(x^1, p^1)) = b_k(E(x^2, p^2)) = E'(x^2, p^2) \quad \text{Q.E.D.}$$

In light of Theorem 2, the uniqueness of proper E.E.F. is ruled out. It is interesting to note, however, that every proper E.E.F. is accounted for by applying a monotone transformations of the type referred to in Theorem 2 to  $E^*(x,p)$ , the E.E.F. referred to in the proof of Theorem 1. Thus, as the following theorem indicates, we have uniqueness of  $E^*(x,p)$  up to monotone transformations:

Theorem 3 Let  $P \subseteq U$  and let  $E(x,p)$  be a proper E.E.F. for  $P$ . Then  $E(x,p)$  can be represented as

$$E(x,p) = b_k(E^*(x,p))$$

where  $E^*(x,p)$  is the E.E.F. referred to in the proof of Theorem 1 and  $b_k(\cdot)$  are the monotone functions referred to in Theorem 2.

Proof The proof is by constructing the functions  $b_k(\cdot)$ ,  $1 \leq k \leq l$ . For every  $1 \leq k \leq l$ , let us choose a representative problem  $p^k \in P_k$  arbitrarily. Let  $N_1^k, N_2^k \dots N_{r(k)}^k$  be the partition of the feasible set of  $p^k$  according to the objective value as in the proof of Theorem 1. For each  $1 \leq j \leq r(k)$ , choose  $x^j$  arbitrarily such that  $x^j \in N_j^k$ . By definition

$$E^*(x^j, p^k) = j - 1$$

Let  $E(x^j, p^k) = \alpha_{kj}$

Since  $E(x,p)$  is proper,  $\alpha_{kj}$  is well defined, in the sense that its value depends on the class  $P_k$  and on the index  $j$ , but not on the particular representative problem  $p^k \in P_k$  and solution  $x^j \in N_j^k$  chosen. Since  $E^*(x,p^k)$  ranges over the integers  $0 - r(k) - 1$  only, it suffices to define  $b_k$  on those values. Define

$$b_k(j-1) = \alpha_{jk} \text{ for } j = 1, r(k)$$

It then follows by construction that

$$E(x,p^k) = b_k(E^*(x,p^k)).$$

Since  $p^k \in P_k$  was chosen arbitrarily, we have that

$$E(x,p) = b_k(E^*(x,p)) \text{ for } p \in P_k.$$

It remains to be seen that  $b_k(\cdot)$  does have the properties asserted. Both follow the fact that  $E(x,p)$  is proper. Thus,  $b_k(\cdot)$ ,  $k = 1 \dots l$  is monotone since condition A implies that the  $\alpha_{kj}$  are monotone with  $j$ . Also, condition B implies that for  $x \in 0(p^k)$ ,  $E(x,p^k) = 0$ . Thus,  $\alpha_{1,k} = 0$  and hence  $b_k(0) = 0$ ,  $k = 1 \dots l$  Q.E.D.

IV. Linear E.E.F.'s: Examples

The construction of Theorem 3 accounts for all the proper E.E.F.'s for a given class P. However, this construction is intractable computationally, since it calls for determination of the complete solution structure (i.e., the sequence of the  $z_i$ 's), and for the identification of the appropriate monotone function,  $b_k(\cdot)$ , applicable to the problem in question. In this section, we examine some linear proper E.E.F.'s which can be calculated on the basis of considerably less information. One type of such functions uses the concept of a reference point, brought forward by Cornuejols, Fisher and Nemhauser [ 2 ]. Because of the vast richness of the family of proper linear E.E.F.'s, the section is intended to be a sketchy, rather than systematic, discussion of some examples.

By a linear E.E.F., we mean a function of the form

$$E(x,p) = \beta_p - \alpha_p c^p x$$

where  $c^p$  is the cost vector for p,  $\beta_p$  and  $\alpha_p \geq 0$  are constants which depend on p, but not on x. We will restrict our attention to the non-trivial case  $\alpha_p > 0$ . From condition B, we get an alternative expression for E

denoting 
$$E(x,p) = \alpha_p (z(p) - c^p x)$$

we have 
$$\gamma_p = 1/\alpha_p$$

$$E(x,p) = (z(p) - c^p x)/\gamma_p .$$

We refer to  $\gamma_p$  as the normalization factor. In order that E(x,p) be proper, the  $\gamma_p$ 's should be chosen in such a way that condition C is enforced. This can usually be done in several ways as the following proposition implies.

Proposition 7. Let  $\alpha_i > 0$ ,  $w_i$ ,  $i = 1 \dots k$  be arbitrary such that  $\sum_{i=1}^k w_i = 1$ . Let also  $E^i(x,p) = (z(p) - c^p x)/\gamma_p^i$ ,  $i = 1 \dots k$  be proper linear E.E.F.'s for P.

Then, so are

$$\tilde{E}(x,p) = (z(p) - c^p \cdot x) / \sum_{i=1}^k \alpha_i \gamma_p^i$$

$$\bar{E}(x,p) = (z(p) - c^p \cdot x) / \prod_{i=1}^k \alpha_i (\gamma_p^i)^{w_i}$$

and

$$\hat{E}(x,p) = (z(p) - c^p \cdot x) / \max_{i=1,k} \alpha_i \gamma_p^i$$

Proof. It is obvious that multiplying a proper E.E.F. by  $1/\alpha$  for  $\alpha > 0$ , leaves the E.E.F. proper (Theorem 2). Therefore, it is enough to consider the case  $\alpha_i = 1, i=1, \dots, k$ . Consider  $\tilde{E}$  first. It is enough to examine the case  $k=2$  since the general case follows by induction. Let then

$$\tilde{E}(x,p) = (z(p) - c^p \cdot x) / (\gamma_p^1 + \gamma_p^2)$$

Let  $p, q \in P$ , s.t.  $p \rightarrow q$  and let  $x \in F(p)$ ,  $x \rightarrow y$  relative to  $p \rightarrow q$ . We have to show that

$$\tilde{E}(x,p) = \tilde{E}(y,q)$$

The statement is trivial if  $x \in 0(p)$ . Assume, then, that  $z(p) - c^p \cdot x > 0$ .

Since  $E^i$  are proper for  $i = 1, 2$  we know that

$$(z(p) - c^p \cdot x) / \gamma_p^i = (z(q) - c^q \cdot x) / \gamma_q^i \quad i=1,2$$

since the numerator of the two expressions is non-zero

$$\gamma_p^i / (z(p) - c^p \cdot x) = \gamma_q^i / (z(q) - c^q \cdot x)$$

Adding the two equations for  $i=1,2$ , and flipping the numerator and denominator back, we get the required result for  $\tilde{E}$ .

Let us now examine  $\bar{E}$  with  $p, q, x$  and  $y$  as before. Since  $E^i$  are proper:

$$(z(p) - c^p \cdot x) / \gamma_p^i = (z(q) - c^q \cdot x) / \gamma_q^i \quad i=1, \dots, k$$

$$(z(p) - c^p \cdot x) = (z(q) - c^q \cdot x) \cdot (\gamma_p^i / \gamma_q^i)$$

Raising each side to the power  $w^i$ , and multiplying the terms together, we get:

$$\prod_{i=1}^k (z(p) - c^P \cdot x)^{w_i} = \prod_{i=1}^k (z(q) - c^Q \cdot x)^{w_i} \cdot \prod_{i=1}^k (\gamma_p^i / \gamma_q^i)^{w_i}$$

or since

$$\sum_{i=1}^k w_i = 1$$

$$(z(p) - c^P \cdot x) = (z(q) - c^Q \cdot x) / \prod_{i=1}^k (\gamma_p^i / \gamma_q^i)^{w_i}$$

Rearranging terms, we get the desired result.

Finally, consider  $\hat{E}$ . We have to show that

$$(z(p) - c^P \cdot x) / \max_{i=1, \dots, k} \gamma_p^i = (z(q) - c^Q \cdot y) / \max_{i=1, \dots, k} \gamma_q^i$$

Let  $\gamma_p^j = \max_{i=1, \dots, k} \gamma_p^i$ . It is enough to show that  $\gamma_q^j = \max_{i=1, \dots, k} \gamma_q^i$ . Assume,

on the contrary, that  $\gamma_q^\ell > \gamma_q^j$  for some  $1 \leq \ell \leq k$ . It then follows that

$$(z(p) - c^P \cdot x) / \gamma_p^j \leq (z(p) - c^P \cdot x) / \gamma_p^\ell = (z(q) - c^Q \cdot y) / \gamma_q^\ell < (z(q) - c^Q \cdot y) / \gamma_q^j$$

But then, since  $E^j$  is proper,

$$(z(p) - c^P \cdot x) / \gamma_p^j = (z(q) - c^Q \cdot y) / \gamma_q^j, \text{ a contradiction.}$$

Q.E.D.

Proposition 7 suggests a strong relationship between normalization factors for Proper E.E.F.'s and homogeneous functions of degree 1. This relationship is further established in

Proposition 8. Let  $P \subseteq U$ . For  $p \in P$ , let  $c^P$  the cost vector of  $p$ . Let

$L^P$  be the row space spanned by the equality rows,  $i \in M_p$ , of  $A$

$c_L^P$  be the orthogonal projection of  $c^P$  on  $L^P$

$$d^P = c^P - c_L^P$$

Then

$$E(x, p) = (z(p) - c^P \cdot x) / f(d^P)$$

is a proper E.E.F. for  $p$  whenever  $f(d^P)$  satisfies:

(i) for every  $i \in \{1, \dots, N\}$

$$f(d_1^P \dots d_i^P \dots d_n^P) = f(d_1^P, \dots, d_i^P, \dots, d_n^P)$$

(ii) for every  $i, j \in \{1, \dots, N\}$

$$f(d_1^P \dots d_i^P \dots d_j^P \dots d_n^P) = f(d_1^P \dots d_j^P \dots d_i^P \dots d_n^P)$$

(iii)  $f(d)$  is homogeneous of degree one, i.e.,

$$f(\alpha d_1^P, \alpha d_2^P, \dots, \alpha d_n^P) = \alpha f(d_1^P, \dots, d_n^P)$$

Sketch of Proof and Discussion. Consider first the case  $M_p = \emptyset$ , i.e., there are no equality constraints in  $p$ . In this case,  $L^P$  and  $c_L^P$  are both equal to  $(0, \dots, 0)$  and  $d^P = c^P$ . If  $M_p \neq \emptyset$ , then  $d^P$  is an invariant under  $G$  type transformations with  $\alpha^0 = 1$ , i.e., under additions of multiples of the equality constraints to the objective row. The case of  $\alpha^0 \neq 1$ , (i.e., rescaling of the objective row) is taken care of by requirement (iii) which assures that such scaling will affect the numerator and denominator proportionally. Condition (i) is designed to accommodate complementing variables (i.e.,  $D$  type operators). It can be (fully or partially) relaxed if  $P$  itself excludes such operators. In other words, if  $P$  does not contain any two problems which are obtainable from each other by complementing a given column, then (i) can be relaxed with respect to this column. For instance, one can argue that for, say, travelling salesman problems, the operation of complementing variables is rather unnatural and therefore problems with complemented variables should be excluded from  $P$ . In such cases, there is no need to insist on the symmetry imposed by (i). (Allowing for complementing variables, however, comes in very handy in the context of enumeration algorithms).

Condition (ii) ensures the invariance of  $E$  under permutation of columns ( $\phi$  type operators). Again, it can be often relaxed in response to the structure of  $P$ . For example, if  $P$  is the set of, say, plant location problems, there is

a natural distinction between the variables which represent the on-off status of a plant and those which correspond to the distribution pattern. Clearly, there is no reason to impose condition (ii) between these two sets of variables. (Strictly speaking, in this case,  $f$  is not a function of  $d$  alone since specific entries in  $d$  are identified by their respective columns in  $A$ . Nevertheless, we will refer to the normalization factor as  $f(d)$ ).

Examples.

(1) Let  $d$  be rearranged such that

$$|d_{\rho_1}| \geq |d_{\rho_2}| \dots \geq |d_{\rho_n}| .$$

Let

$$f^k(d) = \sum_{i=1}^k |d_{\rho_i}|$$

be the sum of  $k$  largest elements (in absolute value) in the (modified) objective row  $d$ .  $f^k(d)$  satisfies (i), (ii) and (iii) and is then appropriate for a normalization factor. For instance, if  $P$  is the class of (binary) knapsack problems, then  $f^1(d)$ , which in this case reduces to  $|c_{\max}| = \max |c_i|$ , is a proper normalization factor. Note that  $|c_{\max}|$  represents, in a way, a measure for what the algorithm is all about since by rounding the optimal solution for the continuous relaxation of the problem we get a solution which is at most  $|c_{\max}|$  units away from the optimum. Thus,  $E(x,p) = (z(p) - c^p x) / |c_{\max}|$  is, in my opinion, a meaningful measure for the quality of a solution to the knapsack problem. I am not aware of any approximate algorithm for this problem which can guarantee a performance better than the obvious level of 1 with respect to this measure.

(2) If the equalities defining the individual problems  $p \in P$  are of the transportation type, then  $d$  can be obtained from  $c$  by simply adding row and column constants in such a way that the row and column sums of  $d$  are zeroes.

(This can be accomplished by a one pass scan of the rows and columns of  $c$ ). If we exclude from  $P$  problems in which some of the variables are complemented, we can relax requirement (i) of Proposition 8. Let  $d$  be arranged such that

$$d_{\pi_1} \geq d_{\pi_2} \dots \geq d_{\pi_n}$$

and consider the range of (modified) costs

$$f^R(d) = d_{\pi_1} - d_{\pi_n}$$

$f^R(d)$  measure the possible variability of the per unit cost over the different routes. Multiplied by the total amount shipped, it yields an upper bound on the difference in the objective function between any two feasible solutions to  $p$ , and may be therefore meaningful as a normalization factor. This measure can be refined in several ways by considering the ranges of the individual rows and columns and aggregating them as in Proposition 7.

(3) We finally consider a class of normalization factors obtained by using a reference point, as suggested in [2]. For every  $p \in P$ , let  $x^p \in [0,1]^N$ , the Unit Cube, be a reference point (not necessarily feasible, not necessarily integer). Let the operators  $\varphi$  and  $D$  be extended in a natural way to the Unit Cube so it will make sense to refer to the correspondence  $x \rightarrow y$  relative to  $p \rightarrow q$ , for  $x \in [0,1]^N$  even though  $x \notin F(p)$ . (In this case, of course,  $y \notin F(q)$ .)

Proposition 9. Let  $P \subseteq U$ . For each  $p \in P$ , let  $x^p \in [0,1]^N$  be such that:

- (i)  $z(p) \neq c^p x^p$
- (ii)  $a_i^p x^p = b_i, \quad i \in M_p$  i.e.,  $x^p$  satisfies the equality constraints of  $p$
- (iii)  $p \rightarrow q \Rightarrow x^p \rightarrow x^q$  relative to  $p \rightarrow q$ .

Then

$$E(x,p) = (z(p) - c^p x) / (z(p) - c^p x^p)$$

is a proper E.E.F. for P.

Proof. The proof resembles closely that of Proposition 8 (which was sketched only). The only comment in order is the fact that it is condition (ii), rather than feasibility or integrality of x, which enforces the invariance of E under G type transformations. Q.E.D.

There are a number of ways to choose  $x^p$  to satisfy (i), (ii) and (iii). Clearly, one can use the solution of a given reference heuristic algorithm for P which is invariant under the transformations in question. For instance, for the knapsack problem,  $x^p$  can be chosen as the solution of the greedy (in the sense of the cost to weight ratios) algorithm, provided ties are broken properly. (E.g., breaking ties according to the size of the cost coefficients is o.k., however, breaking them according to the index of the variables in question is not.)

Alternatively, a generalization of the origin can be used. For instance, if  $M_p = \emptyset$ , one can set

$$(1) \quad x_j^p = \begin{cases} 0 & c_j \geq 0 \\ 1 & c_j < 0 \end{cases}$$

Similarly, if  $M_p$  is composed of a set of disjoint multiple choice constraints:

$$\sum_{j \in s_i} x_j = k_i,$$

then  $x^p$  can be obtained by picking the worst (cost-wise)  $k_i$  variables from each set  $s_i$ .

It is interesting to note that the traditional percentage error measure can be viewed as using the origin for a reference point. Since the origin does not necessarily satisfy the conditions of Proposition 9, this measure may turn out to be improper. However, it can often be modified in such a way so as to make it proper while still retaining some of its basic properties. This is what is actually happening when one uses  $x^p$  as defined in (1) for the case  $M_p = \emptyset$ . Clearly, if for some  $p \in P$ ,  $c_j \geq 0$  for all  $j \in N$ , then this measure coincides with the percentage error measure for this problem. However, the modified measure is different from the original one if some of the  $c_j$  are negative.

REFERENCES

- [1] Cook, S. A., "The Complexity of Theorem Proving Procedures", Proc. 3rd Annual ACM Symp. on the Theory of Computing (1971), pp. 151-158.
- [2] Cornuejols, G., M. L. Fisher and G. L. Nemhauser, "Location of Bank Accounts to Optimize Float: An Analytic Study of Exact and Approximate Algorithms", Mang. Sci. 23 (1977), pp. 789-810.
- [3] Garey, M. R. and D. S. Johnson, "Approximation Algorithm for Combinational Problems: An Annotated Bibliography" in Traub, S. F. (ed), "Algorithm and Complexity: New Directions and Recent Results", Academic Press, New York, 1976.
- [4] Ibarra, O. H. and Ch. E. Kim, "Fast Approximation Algorithms for the Knapsack and Sum of Subsets Problems", JACM 22 (1975), pp. 463-468.
- [5] Johnson, D. S., "Approximation Algorithms for Combinatorial Problems", J. of Compt. and Sys. Sci. 9 (1974), pp. 256-278.
- [6] Karp, R. M., "On the Computational Complexity of Combinatorial Problems" Networks 5 (1975), pp. 45-68.
- [7] Korte, B., "Approximation Algorithms for Discrete Optimization Problems", Presented at Discrete Optimization 77, Vancouver, British Columbia, August, 1977.
- [8] Sahni, S. and T. Gonzales, "P Complete Approximation Problems", JACM 23 (1976), pp. 555-565.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
Technical Report No. 422		
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
Functions for Measuring the Quality of Approximate Solutions to Zero-One Programming Problems		Technical Report June 1978
		6. PERFORMING ORG. REPORT NUMBER
		MSRR 422
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Eitan Zemel		N00014-75-C-0621 MPS73-08534 A02 NSF
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Graduate School of Industrial Administration Carnegie-Mellon University Pittsburgh, Pennsylvania 15213		NR 047-048
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Personnel and Training Research Programs Office of Naval Research (Code 434) Arlington, Virginia 22217		June 1978
		13. NUMBER OF PAGES
		23
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Algorithms, performance measures, heuristics		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
<p>We point out the difficulties associated with measuring the quality of an approximate (heuristic) solution by "Percentage-Error" as is customary. We define a set of properties that are to be expected from a proper measure of solution quality and investigate the family of functions which satisfy those conditions. In particular, we show that for any class of 0-1 programming problems appropriate functions do exist and that they are uniquely defined up to monotone transformations. We conclude with several examples of linear functions which are suitable for certain classes of problems.</p>		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)