


AD-A067 596

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/6 7/2
PHOTOCHEMISTRY AND ELECTRONIC STRUCTURE OF BIS(DICARBONYL(PI5-C--ETC(U)
APR 79 H B ABRAHAMSON, M C PALAZZOTTO N00014-75-C-0880
TR-15 NI

UNCLASSIFIED

| OF |

AD
A067596



END
DATE
FILMED
6-79
DDC

UNCLASSIFIED

LEVEL II

12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER OAR-TR-15	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Photochemistry and Electronic Structure of Bis(dif- carbonyl(η^5 -cyclopentadienyl)ruthenium) and Its Iron Analogue		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) H. B. Abrahamson, M. C. Palazzotto, C. L. Reichel, and M. S. Wrighton		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0880
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-579
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, Virginia 22217		12. REPORT DATE April 3, 1979
		13. NUMBER OF PAGES 24
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; reproduction is permitted for any purpose of the United States government; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Distribution of this document is unlimited.		
18. SUPPLEMENTARY NOTES Prepared for publication in the Journal of the American Chemical Society.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Metal-Metal Bonds, Bridged Metal Bonds, Photochemistry, Metal-Centered Radicals		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The photochemistry and electronic spectra of $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ ($M = Fe, Ru$) are reported. Each complex undergoes efficient $M-M$ bond cleavage subsequent to electronic excitation in CCl_4 solution to yield $[M(\eta^5-C_5H_5)(CO)_2Cl]$ as the only M containing product. The disappearance quantum yield at 366 nm for $M = Fe$ and Ru is 0.23 and 0.44, respectively, in CCl_4 where the Fe species is fully in the bridged form and the Ru species is a mixture of the bridged and non-bridged form. The quantum yield for disappearance of the Ru species is the same in CH_3CN and hydrocarbon solution of 0.1M CCl_4 , where the structure in solution is essentially		

ADA067596

DDC FILE COPY

DDC

APR 23 1979

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ADDITIONAL TO	
DTIC	Write Section <input checked="" type="checkbox"/>
DDC	Diff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

OFFICE OF NAVAL RESEARCH

Contract No. ⁽¹⁵⁾ ~~NO0014-75-C-0880~~

⁽⁹⁾ Interim ⁽¹⁴⁾ TR-
TECHNICAL REPORT, NO. 15

⁽⁶⁾ Photochemistry and Electronic Structure of
Bis(dicarbonyl(π -cyclopentadienyl)ruthenium) and Its
Iron Analogue.

^(pi 5)

by

⁽¹⁰⁾ Harmon B. / Abrahamson, Michael C. / Palazzotto, Carol L. / Reichel and Mark S. / Wrighton

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

⁽¹¹⁾ 3 Apr ~~11-3~~, 1979

⁽¹²⁾ 28p.

Reproduction in whole or part is permitted for any purpose of the United States Government.

Approved for public release; distribution unlimited.



220 007
79 04 17 007 LB


Photochemistry and Electronic Structure of Bis(dicarbonyl)(η^5 -cyclopentadienyl)ruthenium) and Its Iron Analogue

Harmon B. Abrahamson, Michael C. Palazzotto, Carol L. Reichel, and
Mark S. Wrighton*


Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Abstract: The photochemistry and electronic spectra of $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ ($M = Fe, Ru$) are reported. Each complex undergoes efficient M-M bond cleavage subsequent to electronic excitation in CCl_4 solution to yield $[M(\eta^5-C_5H_5)(CO)_2Cl]$ as the only M containing product. The disappearance quantum yield at 366 nm for $M = Fe$ and Ru is 0.23 and 0.44, respectively, in CCl_4 where the Fe species is fully in the bridged form and the Ru species is a mixture of the bridged and non-bridged form. The quantum yield for disappearance of the Ru species is the same in CH_3CN and hydrocarbon solution of 0.1M CCl_4 where the structure in solution is essentially fully bridged (CH_3CN) or ~50/50 bridged/non-bridged (hydrocarbon). The results support the conclusion that carbonyl bridged metal-metal bonds can be efficiently cleaved by optical excitation. The quantum yields for both $M=Fe$ and Ru are somewhat wavelength dependent, with higher energy excitation giving modestly increased quantum yields. The electronic spectrum of the Ru species is very solvent and temperature sensitive in accord with known effects on the equilibrium between the bridged and non-bridged form. The $\sigma_b \rightarrow \sigma^*$ absorption in the bridged form is at ~265 nm and in the non-bridged form it is at ~330 nm representing a difference of ~7000 cm^{-1} in energy. This result is similar to that reported previously for the bridged and non-bridged forms of $[Co_2(CO)_8]$. $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ is fully bridged under all conditions under consideration here and exhibits a $\sigma_b \rightarrow \sigma^*$ absorption at ~350 nm.

The photochemistry of a number of dinuclear metal-metal bonded organometallic complexes is dominated by cleavage of the metal-metal bond.⁷⁻⁸ This chemistry is in accord with lowest excited states which involve the population of an orbital which is strongly antibonding (σ^*) with respect to the metal-metal bond.^{7,9} All of the complexes for which detailed studies have been reported involve systems for which the metal-metal bond is not bridged. A number of qualitative observations have been described which strongly implicate symmetrical cleavage reactions of bridged systems subsequent to electronic excitation. One good example is represented by equation (1);⁸



$[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ is believed to be greater than 99% in the bridged form, I, at room temperature, equation (2).¹⁰ However, there are no quantum yield data



for the reaction given in equation (1) and such data are necessary in order to assess the reactivity of the lowest electronic excited states of form I. Further, based on the recent report¹¹ concerning the electronic spectral changes accompanying a temperature induced shift in the equilibrium distribution of the bridged and non-bridged forms of $[\text{Co}_2(\text{CO})_8]$,¹² a significant shift in the equilibrium indicated in equation (2) should result in a large electronic spectral change. In this connection $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ represents an interesting substance, since it is roughly a one-to-one mixture of forms I and

II at room temperature in solution¹⁰ but is exclusively bridged in the solid state¹³ or at low temperature in solution.¹⁰ Thermodynamic data for equilibrium (2) have been reported previously.^{10b}

In this report we describe results concerning the photochemistry and electronic spectroscopy of $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ ($M = Fe, Ru$). Quantum yield data support the conclusion that the complexes are efficiently cleaved to yield mononuclear products when irradiation is carried out in the presence of CCl_4 . The spectral studies show a large change in the absorption spectrum of the Ru complex upon lowering the temperature or solvent polarity, in accord with shifts in the equilibrium represented in (2). Changes in the spectrum of $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ with variation in temperature or solvent polarity are modest by comparison to the Ru analogue, consistent with essentially a fully bridged structure under all conditions.


Results

Electronic and Infrared Absorption Spectra. $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ exhibits a remarkably temperature and solvent sensitive absorption spectrum, Figures 1 and 2. As seen in Figure 1, an intense band grows at ~265 nm at the expense of an absorption feature at ~330 nm as the sample is cooled from 298 to 77°K. The large spectral changes are consistent with the equilibrium indicated in (2) where I is the exclusive low temperature form.¹⁰ Smooth variation in the sample temperature from 298 to 77°K yields substantial spectral changes short of the glassy state of the solvent, and there is fair preservation of an isosbestic point at ~300 nm when solvent contraction is taken into account.

Figures 2 and 3 show comparisons of the uv-vis and ir spectra in isooctane and CH_3CN . EtOH solvent yields a spectrum similar to that obtained in CH_3CN , while CCl_4 solvent yields a spectrum resembling that found in the alkane solvent. Note that the 298°K CH_3CN spectrum is very similar to that in EPA at 77°K; also, the 298°K spectrum in EtOH does not undergo the dramatic change upon cooling the sample to 77°K that is found for EPA solutions. The ir spectrum, Figure 3, is quite different in CH_3CN and isooctane. The differences appear to reflect a change in the ratio of I/II such that more of the bridged species is present in CH_3CN , owing to the larger integrated area under the 1781 cm^{-1} absorption which has been associated with the bridging carbonyls. Table I lists uv-vis, and ir absorptions and associated absorptivities for the Ru species.

By way of contrast to the Ru species, the electronic spectrum of $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ is only modestly affected by solvent polarity, Figure 4, or by cooling an EPA solution from 298 to 77°K, Figure 5. In particular, the salient near-uv maximum near 350 nm behaves in a manner quite different than for the band near 330 nm in the analogous Ru complex. Uv-vis and ir absorption maxima and molar absorptivities for the Fe complex are included in Table I.


Photochemistry. Irradiation of $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ in degassed CCl_4 solution at 298°K results in reaction according to equation (3). The reaction can be



followed quantitatively by ir using the absorptivities and band positions set out in Table I. For M=Fe there is measurable thermal reaction at 298°K to give the same product, but there is little, if any, detectable reaction for the Ru species on the timescale of the photochemical experiments reported here. Quantum yield data for both complexes are given in Table II.

Irradiation of $[Ru_2(\eta^5-C_5H_5)_2(CO)_4]$ has also been carried out in dilute solutions of CCl_4 in different solvents. In particular, experiments have been carried out in CH_3CN and in C_6H_6 /isooctane (1/24) solutions containing 0.1M CCl_4 . These solvents represent the extremes with respect to the electronic spectral changes observed for the Ru complex, Figure 2. The quantum yield data are included in Table II; the only metal carbonyl product observed is $[Ru(\eta^5-C_5H_5)(CO)_2Cl]$. Irradiation of $[Ru_2(\eta^5-C_5H_5)_2(CO)_4]$ in the presence of 1- IC_5H_{11} yields only one infrared detectable product which is assigned as $[Ru(\eta^5-C_5H_5)(CO)_2I]$; the quantum yield is nearly the same as for reaction with CCl_4 .

Irradiation of $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ in the presence of 0.1 M PPh_3 in degassed benzene solution results in the formation of the monosubstituted dinuclear species, equation (4).¹⁴ Quantum yield data are included in Table II.



Further irradiation of $[Fe_2(\eta^5-C_5H_5)_2(CO)_3(PPh_3)]$ in the presence of 0.1 M PPh_3 in methylcyclohexane solution leads to no rapid spectral changes. Irradiation of $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ in methylcyclohexane solutions of 0.1 M $P(OCH_3)_3$ results in at least two primary products. One of the products is definitely

the monosubstitution product, $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{P}(\text{OCH}_3)_3)]$, based on an infrared spectral comparison with an authentic sample.¹⁴ A second carbonyl product also is a primary product and it is very likely $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_2(\text{P}(\text{OCH}_3)_3)_2]$ with bands at 2014 and 1716 cm^{-1} ; the same product bands are obtained by irradiating $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{P}(\text{OCH}_3)_3)]$ in the presence of $\text{P}(\text{OCH}_3)_3$. Multiple substitution products were noted previously¹⁴ but were not characterized. The important finding here is that the bands at 1752 cm^{-1} and 1716 cm^{-1} grow in together and the formation of the multiple substitution does not require irradiation of the monosubstituted species.

Irradiation of $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in degassed benzene solution containing 0.1 M PPh_3 and 0.1M CCl_4 yields no ir detectable $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{PPh}_3)]$. The dominant product under these conditions is $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Cl}]$. Figure 6 shows the results found for suppression of the formation of $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{PPh}_3)]$ by using 0.1 M 1- IC_5H_{11} instead of 0.1 M CCl_4 . In this case the major product is still $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{PPh}_3)]$, but $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{I}]$ is formed to the small extent that formation of $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{PPh}_3)]$ is suppressed.


Discussion

The photochemistry found here appears to parallel that found for all other dinuclear metal-metal bonded carbonyl complexes.¹⁻⁷ However, what is remarkable here is the fact that the $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ species have bridging CO's to a greater or lesser extent depending on M.¹⁰ Further, the large spectral changes for M=Ru with variation in solvent or temperature are in accord with the notion that the equilibrium ratio of the non-bridged to the bridged form decreases at lower temperature¹⁰ and apparently with more polar solvents as well. We assign the intense near-uv absorption feature at ~330 nm in the Ru complex at 298°K to the $\sigma_b \rightarrow \sigma^*$ excitation of the non-bridged form. Such a spectral feature has come to be regarded⁹ as a characteristic of M-M bonded complexes; the mononuclear $[Ru(\eta^5-C_5H_5)CO_2X]$ species have no such low energy intense absorption. Upon cooling the solution to 77°K the 330 nm feature significantly diminishes and growth of the absorption at ~265 nm is substantial. We assign the 265 nm feature to a $\sigma_b \rightarrow \sigma^*$ excitation of the CO bridged Ru species. Likewise, the relative absorption (uv-vis and ir) spectra in isoctane (~50/50 bridged/non-bridged)¹⁰ and CH_3CN (apparently fully bridged) are in accord with this assignment. Consistently, the Fe complex does not exhibit substantial temperature or solvent effects, since it is ~99% bridged in solution at 298°K in alkane solvent.¹⁰ We adopt the earlier assignment that the absorption feature at ~350 nm is due to the $\sigma_b \rightarrow \sigma^*$ of the bridged Fe species.^{9c} We previously argued^{1a} that the $\sigma_b \rightarrow \sigma^*$ position of the non-bridged Fe species would be in the vicinity of ~420 nm and the differences in the spectrum in isoctane and CH_3CN in that region may in fact be due to the small amount of non-bridged material present in alkane solvent which is not present in the more polar CH_3CN .

The $\sigma_b \rightarrow \sigma^*$ positions for the bridged and non-bridged forms of the Ru species differ by ~ 7000 cm^{-1} . This difference is very similar to that found in the $[Co_2(CO)_8]$ system¹¹ where an equilibrium as in (2) also obtains.¹²

The higher energy $\sigma_b \rightarrow \sigma^*$ absorption in the bridged species is possibly a consequence of a shorter metal-metal bond distance. Also, it should be indicated that the assignment of the absorption band as $\sigma_b \rightarrow \sigma^*$ for the bridged form is made with the understanding that the bridging CO's seriously alter the orbital scheme. But the appearance of the spectral feature (solvent and temperature dependence) is consistent with the $\sigma_b \rightarrow \sigma^*$ type assignment. The weaker shoulders and lower energy maxima for the metal-metal bonded species are attributable to $\pi-d \rightarrow \sigma^*$ transitions but cannot be assigned in detail.

The photochemistry of the $[M_2(\eta^5-C_5H_5)_2(CO)_4]$ is consistent with electronic transitions which terminate in a σ^* orbital with respect to the M-M bond. We have previously reported the synthesis of heterodinuclear metal-metal bonded complexes by simultaneous irradiation of $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ and a second homodinuclear metal-metal bonded complex.^{1a} Equation (5) is



representative of the chemistry; such is consistent with the photogeneration of $[Fe(\eta^5-C_5H_5)(CO)_2]$ radicals from $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$. In the presence of CCl_4 the metal-centered radicals abstract Cl to form $[Fe(\eta^5-C_5H_5)_2(CO)_2Cl]$. Photogeneration of P-donor substitution products likely occurs via substitution of CO by the P-donor at the radical stage rather than by dissociative loss of CO from the excited $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$. This interpretation is supported by the observation that the presence of halocarbons can quench the formation of the simple substitution products. Neither CCl_4 nor $1-IC_5H_{11}$ should be competitive with an equal concentration of PPh_3 for coordinatively unsaturated $[Fe_2(\eta^5-C_5H_5)_2(CO)_3]$. The lower retardation of the substitution by $1-IC_5H_{11}$

vs. CCl_4 is in accord with the lower reactivity of 1- IC_5H_{11} .^{1a} Further, the observation of a multiple substitution product as a primary photoproduct from the photolysis of the Fe species in the presence of $\text{P}(\text{OCH}_3)_3$ is inconsistent with the dissociative loss of CO as the primary photoprocess, whereas such a product could result from the coupling of two substituted radicals.¹⁵ The substitution lability of $17e^-$ metal-centered radicals is well established.⁷

The reaction quantum yields are only modestly affected by variation in the excitation wavelength which suggests that the upper excited states are more reactive than the lowest excited states. This trend is consistent with the fact that the lowest excited states are $d\pi \rightarrow \sigma^*$ type and the M-M bond order is not as diminished as in the $\sigma_b \rightarrow \sigma^*$ excited states populated at the higher energies. Such wavelength effects on M-M bond cleavage have been noted previously.^{1c}

The quantum efficiency for the Fe-Fe bond homolysis is quite high despite the fact that the species is principally CO bridged under the conditions of the irradiation. Irradiation of the Ru complex provides further evidence for the efficient scission of a bridged metal-metal bond.

The reaction quantum yields in CH_3CN vs. the hydrocarbon solvent are essentially the same, despite the fact that the equilibrium indicated in (2) is substantially altered. In fact, the differences in quantum yield are not outside what could be interpreted as just a "solvent effect". The bridged complexes have reaction quantum yields which are similar to those for non-bridged complexes.¹ Unfortunately, quantum yield data do not provide absolute rate constants for reaction, and it is consequently impossible to make a detailed comparison of excited state reactivities of bridged and non-bridged complexes. It would be of interest, too, to learn the position of equilibrium (2) in the excited state. Since the $\sigma_b \rightarrow \sigma^*$ for the non-bridged is at lower energy, excitation of the bridged species may yield an excited, non-bridged complex which either nonradiatively decays to the ground state or fragments to two radicals.

Experimental

Materials. All solvents for photochemical and spectroscopic studies were used as received in commercially available spectroquality. $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was used as obtained from commercial sources after recrystallization from hexane/ CH_2Cl_2 . Its spectral properties are consistent with those previously reported. $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was prepared according to the published literature procedure.¹⁶ Commercially available $[\text{RuCl}_3 \cdot n\text{H}_2\text{O}]$ was converted to $[\text{RuI}_3]$ by dissolving 7.3 g of $[\text{RuCl}_3 \cdot n\text{H}_2\text{O}]$ in a minimum amount of H_2O and mixing with ~40 ml of saturated aqueous KI. The solution was heated gently while stirring and black crystals formed. The crystals were collected by filtration and washed with dilute KI solution and then with absolute EtOH. The resulting RuI_3 was placed in a vacuum dessicator overnight to remove any residual H_2O . Conversion of $[\text{RuI}_3]$ to $[\text{Ru}(\text{CO})_2\text{I}_2]_n$ was effected by heating solid $[\text{RuI}_3]$ in a boat in a tube furnace at 220°C while CO was passed through the tube. The reaction continued until the color of the solid changed from black ($[\text{RuI}_3]$) to orange-red ($[\text{Ru}(\text{CO})_2\text{I}_2]_n$). The $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was finally obtained from reaction of $\text{Na}[\text{C}_5\text{H}_5] \cdot \text{DME}$ with $[\text{Ru}(\text{CO})_2\text{I}_2]_n$. 10.9 g of $[\text{Ru}(\text{CO})_2\text{I}_2]_n$ and 10 g of $\text{Na}[\text{C}_5\text{H}_5] \cdot \text{DME}$ were dissolved in 100 ml of DME and refluxed for ~15 hrs after which time the solution had turned very dark and ir indicated the presence of absorptions attributable to $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$. The solution was filtered and the solvent was removed under vacuum. The solid was taken up in benzene leaving a large amount of an insoluble dark solid. The benzene solution contained the desired product and $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)_2]$. The mixture was chromatographed on alumina with $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)_2]$ eluting first with benzene followed by $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$. The benzene was removed under vacuum. Final purification was by chromatography on alumina under Ar eluting with benzene. The final material was handled under inert atmosphere and the golden orange solid was stored in a Schlenk tube under Ar. An elemental analysis of the

$[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was satisfactory (Alfred Bernhardt, West Germany); for $\text{C}_{14}\text{H}_{10}\text{O}_4\text{Ru}_2$ calcd (found): %C, 37.84 (38.04); %H, 2.27 (2.26); %O, 14.40 (14.62). Spectral properties are in accord with those previously reported.^{10,16}

An authentic sample of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Cl}]$ was prepared by irradiation of $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in CCl_4 . 0.2 g of $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was dissolved in 125 ml of CCl_4 and placed in an Erlenmeyer flask. The solution was purged continuously with prepurified N_2 and irradiated with a G. E. Blacklite equipped with two 15 W bulbs with principal output in the near-uv centered at 355 nm. The photoreaction was followed by ir until ~90% consumption of starting material obtained (~2 hrs). The solution showed ir absorptions at 2055 and 2004 cm^{-1} corresponding to $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Cl}]$. The solution was rotary evaporated to dryness and the residue was chromatographed on alumina eluting with CH_2Cl_2 . By ir, the first compound eluted was $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ and the desired compound eluted second. Solvent (CH_2Cl_2) was removed by rotary evaporation. An elemental analysis for $[\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Cl}]$ was satisfactory (Alfred Bernhardt, West Germany); for $\text{C}_7\text{H}_5\text{O}_2\text{ClRu}$ calcd. (found): %C, 32.63 (32.51); %H, 1.96 (2.09); %Cl, 13.76 (13.81); %O, 12.42 (12.58). The $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Cl}]$ sample was that used in earlier studies.^{1a} Samples of the P-donor substituted Fe dimers were prepared according to the literature procedure and ir spectral data were consistent with the previous report.¹⁴

Spectra. All ir spectra were recorded using a Perkin-Elmer 180 spectrometer using matched 0.1 or 1.0 mm path NaCl or KBr cells. Electronic absorption spectra were recorded using a Cary 17 uv-vis-nir spectrophotometer. Low temperature spectra were recorded using an all quartz liquid N_2 dewar with optical quality flats for windows.

Photochemistry. Photochemical experiments were carried out using either a G. E. Blacklite (355 nm) or an appropriately filtered (313, 366, or 436 nm) 450 or 550 W Hanovia medium pressure Hg lamp with a merry-go-round.¹ The light intensity was determined by ferrioxalate actinometry¹⁸ and was in the range of 1.85×10^{-6} to 5×10^{-9} ein/min. Samples for quantum yield determinations were freeze-pump-thaw degassed, hermetically sealed samples in 13 mm o.d. Pyrex ampules. Concentration of $[\text{M}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ was in the range 10^{-4} - 2×10^{-3} M to insure ~100% light absorption and quantum yields are for conversions of <20%.

Acknowledgements. We thank the National Science Foundation and the Office of Naval Research for support of this research. HBA acknowledges support as an NSF Predoctoral Fellow, 1975-1978 and MSW as a Dreyfus Teacher-Scholar Grant recipient, 1975-1980. We thank Professor Harry B. Gray of the California Institute of Technology for a preprint of the work described in ref. 15.

References

1. (a) H. B. Abrahamson and M. S. Wrighton, Inorg. Chem., 17, 1003 (1978) and J. Amer. Chem. Soc., 99, 5510 (1977); (b) M. Wrighton and D. Bredesen, J. Organometal. Chem., 50, C35 (1973); (c) M. S. Wrighton and D. S. Ginley, J. Amer. Chem. Soc., 97, 2065, 4246, 4908 (1975); (d) D. L. Morse and M. S. Wrighton, ibid., 98, 3931 (1976).
2. (a) J. L. Hughey, IV, C. R. Bock, and T. J. Meyer, J. Amer. Chem. Soc., 97, 4440 (1975); (b) J. L. Hughey, IV, C. P. Anderson, and T. J. Meyer, J. Organometal. Chem., 125, C49 (1977).
3. A. R. Burckett, T. J. Meyer, and D. G. Whitten, J. Organometal. Chem., 67, 67 (1974).
4. R. M. Laine and P. C. Ford, Inorg. Chem., 16, 388 (1977).
5. (a) A. Hudson, M. F. Lappert, P. W. Lednor, and B. K. Nicholson, J. C. S. Chem. Comm., 966 (1974); (b) A. Hudson, M. F. Lappert, and B. K. Nicholson, J. C. S. Dalton, 551 (1977).
6. D. M. Allen, A. Cox, T. J. Kemp, Q. Sultane, and R. B. Pitts, J. C. S. Dalton, 1189 (1976).
7. (a) B. H. Byers and T. L. Brown, J. Amer. Chem. Soc., 97, 947, 3260 (1975); 99, 2527 (1977); (b) M. Absi-Halabi and T. L. Brown, ibid., 99, 2982 (1977).
8. C. Giannotti and G. Merle, J. Organometal. Chem., 105, 97 (1976).
9. (a) R. A. Levenson, H. B. Gray, and G. P. Caesar, J. Amer. Chem. Soc., 92, 3653 (1970); (b) R. A. Levenson and H. B. Gray, ibid., 97, 6042 (1975); (c) D. C. Harris and H. B. Gray, ibid., 97, 3073 (1975) and Inorg. Chem., 14, 1215 (1975).
10. (a) R. D. Fischer, A. Vogler, and K. Noack, J. Organometal. Chem., 7, 135 (1967); (b) K. Noack, ibid., 7, 151 (1967); (c) P. McArdle and A. R. Manning, J. Chem. Soc. (A), 2128 (1970).
11. H. B. Abrahamson, C. C. Frazier, D. S. Ginley, H. B. Gray, J. Lilienthal, D. R. Tyler, and M. S. Wrighton, Inorg. Chem., 16, 1554 (1977).
12. (a) A. Noack, Helv. Chim. Acta, 47, 1064 (1964); (b) G. Bor and K. Noack, J. Organometal. Chem., 64, 367 (1974); (c) G. Bor, U. K. Dietler, and K. Noack, J. Chem. Soc. Chem. Commun., 914 (1976); (d) S. Onaka and D. F. Shriver, Inorg. Chem., 15, 915 (1976); (e) R. L. Sweany and T. L. Brown, ibid., 16, 415 (1977).
13. O. S. Mills and J. P. Nice, J. Organometal. Chem., 9, 339 (1967).
14. R. J. Haines and A. L. duPreez, Inorg. Chem., 9, 1459 (1969).

15. In contrast to results for $P(OCH_3)_3$, the irradiation of $[Fe_2(\eta^5-C_5H_5)_2(CO)_4]$ in the presence of $P(O-i-Pr)_3$ yields only monosubstitution and evidence has been obtained for an intermediate dinuclear iron complex which is CO-bridged but not Fe-Fe bonded; D. R. Tyler, M. A. Schmidt, and H. B. Gray, submitted for publication and private communication.
16. E. O. Fischer and A. Vogler, Z. Naturforsch., 17b, 421 (1962).
17. F. G. Moses, R. S. H. Liu, and B. M. Monroe, Mol. Photochem., 1, 245 (1969).
18. C. G. Hatchard and C. A. Parker, Proc. Roy. Soc. London, Ser. A, 235, 518 (1956).

Table I. Spectral Properties of Relevant Complexes.^a

Complex	Ir, cm ⁻¹ (ε)	Uv-Vis, nm (ε)
[Ru ₂ (η ⁵ -C ₅ H ₅) ₂ (CO) ₄]	2026(180), 2016(530), 2009(1150), 1971(4320), 1963(1880), 1939(4150), 1781(1262)	435(1280), 330(13,900) 265(10,950)
[Ru(η ⁵ -C ₅ H ₅)(CO) ₂ Cl]	2057(3030), 2009(3210)	285(2480)
[Ru(η ⁵ -C ₅ H ₅)(CO) ₂ I]	2050(-----), 2003(-----)	----
[Fe ₂ (η ⁵ -C ₅ H ₅) ₂ (CO) ₄]	2004(2630), 1958(3450), 1782(3160); 1996(2770) ^b , 1952(2440) ^b , 1781(4130) ^b	514(710) ^b , 410(1870) ^b , 346(9190) ^b
[Fe ₂ (η ⁵ -C ₅ H ₅) ₂ (CO) ₃ (P(OCH ₃))]	1965 ^c , 1945 ^c , 1752 ^c	565 ^c , 415 ^c , 353 ^c
[Fe ₂ (η ⁵ -C ₅ H ₅) ₂ (CO) ₂ (P(OCH ₃)) ₂]	2014 ^c , 1716 ^c	----
[Fe(η ⁵ -C ₅ H ₅) ₂ (CO) ₂ Cl]	2054(3290), 2011(3080); 2049(2270) ^b , 2003(2340) ^b	403(590), 339(880)
[Fe ₂ (η ⁵ -C ₅ H ₅) ₂ (CO) ₃ (PPh ₃)]	1961(1870) ^c , 1937(1507) ^c , 1740(3700) ^c ; 1950(1440) ^b , 1930(940) ^b , 1732(4870) ^b	615 ^c , 430 ^c , 360 ^c
[Fe(η ⁵ -C ₅ H ₅)(CO) ₂ I]	2037(2370) ^b , 1993(2490) ^b	

^aAll spectral data are for CCl₄ solutions at 298°K unless noted otherwise.

^bBenzene solution at 298°K.

^cMethylcyclohexane solution at 298°K.

Table II. Reaction Quantum Yields for $[M_2(\eta^5-C_5H_5)_2(CO)_4]$.

M	Conditions ^a	Product	$\phi_{dis.}$		$\phi_{appear.}$	
			313	366	313	366
Ru	neat CCl_4	$[Ru(\eta^5-C_5H_5)(CO)_2Cl]$	—	0.44	—	0.92
	0.1M CCl_4 in C_6H_6 /isooctane (1/24)	$[Ru(\eta^5-C_5H_5)(CO)_2Cl]$	0.46	0.35	0.36	—
	0.1M CCl_4 in CH_3CN	$[Ru(\eta^5-C_5H_5)(CO)_2Cl]$	0.55	0.37	0.28	—
	0.1M 1- IC_5H_{11} in C_6H_6 /isooctane (1/24)	$[Ru(\eta^5-C_5H_5)(CO)_2I]$	—	0.45	—	—
Fe	neat CCl_4	$[Fe(\eta^5-C_5H_5)(CO)_2Cl]$	0.38	0.23	0.21	0.42
	0.1M PPh_3 in benzene	$[Fe_2(\eta^5-C_5H_5)_2(CO)_3PPh_3]$	—	0.05	—	0.06

^aDegassed solutions of starting complex ($\sim 10^{-3}M$) irradiated at 366 nm in hermetically sealed ampules.

^bQuantum yields for disappearance ($\phi_{dis.}$) of starting complex and appearance ($\phi_{appear.}$) of product;

error is $\pm 10\%$.

Figure Captions

Figure 1. Electronic absorption spectra upon changing the temperature from 298 to 77°K. The spectral changes are not corrected for solvent contraction.


Figure-2. Comparison of electronic absorption spectra of $[\text{Ru}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in CH_3CN and isooctane at $9.25 \times 10^{-4}\text{M}$ at 298°K in 1.00 cm path cells.


Figure 3. Comparison of infrared spectra in CH_3CN and isooctane at $9.25 \times 10^{-4}\text{M}$ at 298°K in 1.00 mm path cells.


Figure 4. Comparison of electronic spectrum of $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in isooctane and CH_3CN . Concentration of the complex is $8.3 \times 10^{-5}\text{M}$ in each case and a 1.00 cm path cell was used.


Figure 5. Spectral changes upon lowering the temperature from 298 to 77°K. Changes are not corrected for solvent contraction.

Figure 6. Plot of ir absorption due to $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ (1781 cm^{-1} , $\epsilon = 4130$) and $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_3(\text{PPh}_3)_2]$ (1732 cm^{-1} , $\epsilon = 4870$) upon irradiation of 0.03 M $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in deoxygenated benzene solution in the presence of 0.1M PPh_3 (●) or 0.1M PPh_3 plus $0.1\text{M 1-IC}_5\text{H}_{11}$ (○). Absorptions were measured in a 0.1 mm path cell.












- isooctane
- - - CH_3CN







For NR 051-579

472:GAN:716:tam
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division	1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko	1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division	1
		Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division	1

Encl

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. A. El-Sayed University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1	Dr. M. Rauhut American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805	1
Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163	1	Dr. J. I. Zink University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1
Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. B. Schechtman IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. J. R. MacDonald Naval Research Laboratory Chemistry Division Code 6110 Washington, D.C. 20375	1		
Dr. G. B. Schuster University of Illinois Chemistry Department Urbana, Illinois 61801	1		
Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1		
Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007	1		
Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1		